7-Connected Graphs are 4-Ordered

Rose McCarty^{*}, Yan Wang[†], Xingxing Yu[‡]

School of Mathematics Georgia Institute of Technology Atlanta, GA 30332

Abstract

A graph G is k-ordered if for any distinct vertices $v_1, v_2, \ldots, v_k \in V(G)$, it has a cycle through v_1, v_2, \ldots, v_k in order. Let f(k) denote the minimum integer so that every f(k)-connected graph is k-ordered. The first non-trivial case of determining f(k) is when k = 4, where the previously best known bounds are $7 \leq f(4) \leq 40$. We prove that in fact f(4) = 7.

Keywords: cycles; connectivity; graph linkages; k-ordered graphs

1 Introduction

The problem of studying connectivity and cycles through specified vertices originates with the classic result [4] that k-connected graphs have a cycle through any set of k vertices for $k \ge 2$. Specifying the order the vertices must appear on the cycle is a generalization introduced in [18]. The case k = 4 is particularly interesting because it is the first case where the ordered and unordered versions differ. Surveys on the existence of cycles through specified vertices can be found in [6] and [9].

Following the terminology of Faudree [6], for a positive integer k, we say a graph is k-ordered if, for distinct vertices v_1, v_2, \ldots, v_k , the graph has a cycle through v_1, v_2, \ldots, v_k in order. We define f(k) as the smallest positive integer so that every f(k)-connected graph is k-ordered. Clearly, f(1) = f(2) = 2 and f(3) = 3. Faudree asks for the determination of f(4) in [6]. Goddard [8] and Mukae *et al.* [17] have short proofs showing that 4-connected triangulations of surfaces are 4-ordered.

The best known upper bounds for f(4), and for f(k) in general, follow from work on linkages. We say that a graph G is k-linked if for any collection of k pairs of vertices $\{\{s_i, t_i\} : i \in \{1, 2, ..., k\}\}$, there exists a collection $\{P_i : i \in \{1, 2, ..., k\}\}$ with the path P_i from s_i to t_i for all $i \in \{1, ..., k\}$ such that, for any distinct

Mathematics Subject Classification (2010): 05C38, 05C40.

^{*}Now at the Department of Combinatorics and Optimization, University of Waterloo. Partially supported by NSF grant DMS-1265564 through X. Yu

[†]Partially supported by NSF grant DMS-1600738 through X. Yu

[‡]Partially supported by NSF grants DMS-1265564 and DMS-1600738

 $i, j \in \{1, 2, ..., k\}$, no vertex of P_i is an internal vertex of P_j . Let g(k) denote the smallest positive integer so that every g(k)-connected graph is k-linked. From the definitions, if a graph is k-linked then it is also k-ordered (by considering the k-pairs $\{v_1, v_2\}, \{v_2, v_3\}, \ldots, \{v_k, v_1\}$). Thus it follows that $f(k) \leq g(k)$ for every k.

Bollobás and Thomason [1] were the first to show that g(k) is linear in k, with $g(k) \leq 22k$. Kawarabayashi, Kostochka, and G. Yu [11] improved this to $g(k) \leq 12k$, and Thomas and Wollan [22] showed further that $g(k) \leq 10k$. Better bounds on g(k) are known for k = 3 and large graphs of bounded tree-width, see [21] and [7].

For the general case where $k \ge 4$, the upper bound of 10k is currently the best known on both g(k) and f(k) as far as the authors are aware. So in particular the previously best known upper bounds for k = 4 are $f(4) \le g(4) \le 40$. Ellingham, Plummer, and G. Yu [5] proved the following result, implying $f(4) \ge 7$.

Theorem 1.1. [5] There exists a 6-connected graph G and distinct vertices v_1, v_2, v_3 , $v_4 \in V(G)$ so that G does not contain a path through the vertices v_1, v_2, v_3, v_4 in order.

The main result of this paper is the following which, combined with Theorem 1.1, shows that f(4) = 7.

Theorem 1.2. Every 7-connected graph is 4-ordered.

Using a precise structure theorem of X. Yu [25, 26, 27], Ellingham, Plummer, and G. Yu [5] also proved the following.

Theorem 1.3. [5] For every 7-connected graph G with distinct vertices $v_1, v_2, v_3, v_4 \in V(G)$, the graph G has a path through v_1, v_2, v_3, v_4 in order.

Theorem 1.2 shows that no additional connectivity is required to guarantee the existence of a cycle through four vertices in order.

Proof Overview

Suppose that G is a 7-connected graph with four distinct vertices $c_0, c_1, c_2, c_3 \in V(G)$ that has no cycle through c_0, c_1, c_2, c_3 in order. Our approach is to find vertex-disjoint cycles C_0 and C_2 and a set $\{z_0, z_1, z_2, z_3\} \subseteq V(G) \setminus (V(C_0) \cup V(C_2))$ so that $c_0, c_1 \in$ $V(C_0), c_2, c_3 \in V(C_2)$, and for i = 0, 1, 2, 3, the vertex z_i is a neighbor of c_i . Define H to be the graph $G - (V(C_0) \cup V(C_2))$. Then H does not have vertex-disjoint paths, with one from z_0 to z_3 and the other from z_1 to z_2 . So, applying Seymour's characterization on the existence of 2-linkages [19], the graph H is "almost" planar and thus has small cuts.

We will show how to extend these small cuts to a small cut of G in Section 2. We introduce the notion of "separating pairs" which are certain subpaths of the cycles C_0, C_2 . The ends of these paths will be used to extend cuts. We prove some lemmas that help us find paths through certain vertices in case cuts do not extend.

In Section 3 we introduce the notion of "3-planar graphs" to explain precisely what we mean by saying that "H is almost planar". We will state Seymour's characterization of 2-linked graphs, and prove some lemmas on 3-planar graphs.

In Section 4, we show that if G is 7-connected then it has a special structure called a " (c_0, c_1, c_2, c_3) -skeleton". This structure allows us to assume that H has no 1-cut separating two of the vertices in $\{z_0, z_1, z_2, z_3\}$ from the rest. Using this notion of skeletons, in Section 5 we show that we can assume H is 2-connected. In Section 6 we use different techniques to show that H is 3-connected.

In Section 7 we first prove a discharging lemma on plane graphs. Then we use the fact that H is 3-connected and 3-planar, and the lemmas from Section 3, to show that in fact H is planar, and all neighbors of C_0 and C_2 are on the boundary of a single face. We then use the discharging lemma to force a special configuration in H, which we can use to find the desired cycle in G. We conclude and give some additional remarks in Section 8.

Notation

We conclude this section with notation and terminology we need in the rest of the paper. Suppose G is a graph and P is a path in G. Then define end(P) to be the set of vertices of smallest degree of P. Define $int(P) := V(P) \setminus end(P)$. Notice that if P has two or fewer vertices, then $int(P) = \emptyset$. Given a cycle C and an orientation of C, for any distinct $u, v \in V(C)$, we use C[u, v] to denote the subpath of C from u to v in clockwise order. Let C(u, v] := C[u, v] - u, C[u, v] := C[u, v] - v, and $C(u, v) := C[u, v] - \{u, v\}$. We use similar notation for subpaths of P.

2 Separating Pairs

Definition 2.1. Let G be a graph, let C be a cycle in G, let $v_0, v_1 \in V(C)$ be distinct, and let $A \subseteq V(G) \setminus V(C)$. Then a (v_0, v_1, C, A) -separating pair is a set of paths $\{R_0, R_1\}$ such that there exists an orientation of C so that

- (i) for $i = 0, 1, R_i$ is a subpath of $C[v_i, v_{1-i}]$,
- (ii) for $i = 0, 1, N(A) \cap V(C[v_i, v_{1-i}]) \subseteq V(R_i)$, and
- (iii) the graph G has no edge uv such that $u \in int(R_0) \cup int(R_1)$ and $v \in V(C) \setminus (V(R_0) \cup V(R_1))$.

A minimum (v_0, v_1, C, A) -separating pair is a (v_0, v_1, C, A) -separating pair $\{R_0, R_1\}$ so that $|V(R_0)| + |V(R_1)|$ is minimum. For all $i, j \in \{0, 1\}$, if $R_i \neq \emptyset$ then define r_i^j to be the end of R_i closest to v_j on $C[v_i, v_{1-i}]$. (Thus, $r_i^0 = r_i^1$ if R_i consists of a single vertex.) Clearly, a (v_0, v_1, C, A) -separating pair exists as the two paths in C between v_0 and v_1 form such a pair. Later in the paper we often construct small cutsets containing the set $end(R_0) \cup end(R_1)$. The separating pairs we use will always be chosen to be minimum. The following lemma shows that if G[V(C)] contains no cycle through v_0 and v_1 that is shorter than C, then minimum separating pairs satisfy some additional properties.

Lemma 2.2. Let G be a graph, let C be a cycle in G, let $v_0, v_1 \in V(C)$ be distinct, and let $A \subseteq V(G) \setminus V(C)$. Let $\{R_0, R_1\}$ be a minimum (v_0, v_1, C, A) -separating pair. Suppose that G[V(C)] contains no cycle through v_0 and v_1 with fewer vertices than C. Then for each choice $i, j \in \{0, 1\}$, if $R_i \neq \emptyset$ then either

- (i) $N(r_i^j) \cap A \neq \emptyset$, or
- (*ii*) $R_{1-i} \neq \emptyset$, $N(r_i^j) \cap int(R_{1-i}) \neq \emptyset$, and $N(r_{1-i}^j) \cap A \neq \emptyset$.

Proof. For convenience, let P_0, P_1 denote the two paths in C between v_0 and v_1 . Without loss of generality, assume that $R_i \subseteq P_i$ for i = 0, 1. For $i, j \in \{0, 1\}$, if $R_i \neq \emptyset$ let r_i^j be the end of R_i closest to v_j on P_i (with possibly $r_i^0 = r_i^1$). Notice that by *(ii)* of the definition of a separating pair, for j = 0, 1, if $v_j \in N(A)$ then $v_j \in V(R_0) \cap V(R_1)$.

We claim that for $k, l \in \{0, 1\}$ for which r_k^l is defined and $N(r_k^l) \cap A = \emptyset$, we have $R_{1-k} \neq \emptyset$ and $N(r_k^l) \cap \operatorname{int}(R_{1-k}) \neq \emptyset$. By symmetry, we assume r_0^0 is defined and $N(r_0^0) \cap A = \emptyset$. By the choice of $\{R_0, R_1\}$, $\{R_0 - r_0^0, R_1\}$ is not a (v_0, v_1, C, A) separating pair. Hence, there exists $uv \in E(G)$ with $u \in \operatorname{int}(R_0 - r_0^0) \cup \operatorname{int}(R_1)$ and $v \in V(C) \setminus (V(R_0 - r_0^0) \cup V(R_1))$. In particular, $u \in \operatorname{int}(R_0) \cup \operatorname{int}(R_1)$. Hence, $v = r_0^0$, since $\{R_0, R_1\}$ is a (v_0, v_1, C, A) -separating pair. If $u \in \operatorname{int}(R_0 - r_0^0)$ then G[V(C)]contains a cycle through v_0 and v_1 shorter than C, a contradiction. So $u \in \operatorname{int}(R_1)$.

By symmetry, it suffices to prove the assertion for the case i = 0 and j = 0. Thus, assume that r_0^0 is defined and $N(r_0^0) \cap A = \emptyset$. Then by the above claim, $R_1 \neq \emptyset$ and $N(r_0^0) \cap \operatorname{int}(R_1) \neq \emptyset$. If $N(r_1^0) \cap A \neq \emptyset$ then we are done. So assume $N(r_1^0) \cap A = \emptyset$. Then by the above claim, $N(r_1^0) \cap \operatorname{int}(R_0) \neq \emptyset$.

Let $u_0 \in N(r_1^0) \cap \operatorname{int}(R_0)$ with $P_0[u_0, v_1]$ minimal, $u_1 \in N(r_0^0) \cap \operatorname{int}(R_1)$ with $P_1[u_1, v_1]$ minimal, and let C' be the cycle through v_0 and v_1 defined as follows

$$P_0[v_0, r_0^0] \cup r_0 u_1 \cup P_1[u_1, v_1] \cup P_0[v_1, u_0] \cup u_0 r_1^0 \cup P_1[r_1^0, v_0].$$

Then $V(C') \subseteq V(C)$, and $V(C) \setminus V(C') = \operatorname{int}(P_0[u_0, r_0^0]) \cup \operatorname{int}(P_1[u_1, r_1^0])$. Hence, $P_0[r_0^0, u_0] = r_0^0 u_0$ and $P_1[r_1^0, u_1] = r_1^0 u_1$, as G[V(C)] contains no cycle through v_0 and v_1 and shorter than C.

Let $R'_0 = R_0 - r_0^0$ and $R'_1 = R_1 - r_1^0$. Clearly, $R'_i \subseteq P_i$ for $i \in \{0, 1\}$, and $N(A) \cap V(C) \subseteq V(R'_0) \cup V(R'_1)$. By the choice of u_0 and u_1 , we see that G has no edge from $\operatorname{int}(R'_0) \cup \operatorname{int}(R'_1)$ to $V(C) \setminus V(R'_0 \cup R'_1)$. So $\{R'_0, R'_1\}$ is a (v_0, v_1, C, A) -separating pair, contradicting the choice of $\{R_0, R_1\}$.

Now we prove a technical lemma on the existence of several types of paths and cycles in $G[V(C) \cup A]$. The proof is tedious case analysis, but we will use this lemma frequently.

Lemma 2.3. Let G be a graph, let C be a cycle in G, let $v_0, v_1 \in V(C)$ be distinct, and let $A \subseteq V(G) \setminus V(C)$. Let $\{R_0, R_1\}$ be a minimum (v_0, v_1, C, A) -separating pair and let $u_0 \in int(R_0)$. Suppose that G[V(C)] contains no cycle through v_0 and v_1 with fewer vertices than C. Then

- (i) there exists $a \in V(C)$ with $N(a) \cap A \neq \emptyset$ such that the graph G[V(C)] contains a path through u_0, v_0, v_1, a in order,
- (ii) if G[A] is connected then $G[V(C) \cup A] int(R_0)$ contains a cycle though v_0 and v_1 , and
- (iii) if G[A] is connected then for every vertex $u_1 \in int(R_1)$, $G[V(C) \cup A]$ contains a path through u_0, v_0, v_1, u_1 in order.

Proof. For convenience, let P_0 , P_1 denote the two paths in C between v_0 and v_1 containing R_0, R_1 , respectively. For all $i, j \in \{0, 1\}$, if R_i is non-empty then let r_i^j be the end of R_i closest to v_j on P_i . Since we assume $u_0 \in int(R_0)$, r_0^0 and r_0^1 are defined and distinct. For any distinct $x, y \in V(C) \cap N(A)$, if G[A] is connected then we use A[x, y] to denote a path in $G[A \cup \{x, y\}]$ from x to y.

We prove (i) first. If $N(r_0^1) \cap A \neq \emptyset$ then $C - P_0(u_0, r_0^1)$ gives the desired path for (i). So assume $N(r_0^1) \cap A = \emptyset$. Then by Lemma 2.2, $r_1^1 \in N(A)$ and there exists $w_1 \in int(R_1) \cap N(r_0^1)$. So $(C - P_0(u_0, r_0^1) - P_1(w_1, r_1^1)) \cup w_1 r_0^1$ gives the desired path for (i).

To prove (ii), assume G[A] is connected. If $N(r_0^0) \cap A \neq \emptyset$ and $N(r_0^1) \cap A \neq \emptyset$ then $(C - P_0(u_0, r_0^1)) \cup A[r_0^0, r_0^1]$ gives the desired cycle for (ii). So by symmetry, we may assume $N(r_0^1) \cap A = \emptyset$. Then by Lemma 2.2, $N(r_1^1) \cap A \neq \emptyset$ and there exists $w_1 \in \operatorname{int}(R_1) \cap N(r_0^1)$. If $N(r_0^0) \cap A \neq \emptyset$ then $(C - \operatorname{int}(R_0) - P_1(w_1, r_1^1)) \cup A[r_0^0, r_1^1] \cup w_1 r_0^1$ gives the desired cycle for (ii). So assume $N(r_0^0) \cap A = \emptyset$. Then by Lemma 2.2, $N(r_1^0) \cap A \neq \emptyset$ and there exists $x_1 \in \operatorname{int}(R_1) \cap N(r_0^0)$. Now $(C - \operatorname{int}(R_0) - \operatorname{int}(R_1)) \cup A[r_1^0, r_1^1] \cup r_0^0 x_1 \cup P_1[x_1, w_1] \cup w_1 r_0^1$ gives the desired cycle for (ii).

To prove (iii), assume G[A] is connected, and let $u_1 \in \operatorname{int}(R_1)$. If $N(r_1^0) \cap A \neq \emptyset$ and $N(r_0^1) \cap A \neq \emptyset$ then $P_0[u_0, v_0] \cup P_1[v_0, r_1^0] \cup A[r_1^0, r_0^1] \cup P_0[r_0^1, v_1] \cup P_1[v_1, u_1]$ gives the desired path for (iii). So we may assume by symmetry that $N(r_0^1) \cap A = \emptyset$. Then by Lemma 2.2, $N(r_1^1) \cap V(A) \neq \emptyset$ and there exists a vertex $w_1 \in \operatorname{int}(R_1) \cap N(r_0^1)$. If $N(r_1^0) \cap A \neq \emptyset$ then $P_0[u_0, v_0] \cup P_1[v_0, r_1^0] \cup A[r_1^0, r_1^1] \cup P_1[r_1^1, v_1] \cup P_0[v_1, r_0^1] \cup r_0^1 w_1 \cup$ $P_1[w_1, u_1]$ gives the desired path for (iii). So we may assume $N(r_1^0) \cap A = \emptyset$. Then by Lemma 2.2, $N(r_0^0) \cap V(A) \neq \emptyset$ and there exists a vertex $w_0 \in \operatorname{int}(R_0) \cap N(r_1^0)$. Now $P_0[u_0, w_0] \cup w_0 r_1^0 \cup P_1[r_1^0, v_0] \cup P_0[v_0, r_0^0] \cup A[r_0^0, r_1^1] \cup P_1[r_1^1, v_1] \cup P_0[v_1, r_0^1] \cup r_1[w_1, u_1]$ gives the desired path for (iii).

3 3-Planar Graphs

In this section, we introduce the notion of 3-planar graphs, and state a characterization of 2-linked graphs.

Let G be a graph and let $s_1, t_1, s_2, t_2 \in V(G)$ be distinct vertices. Then an $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage is a set of two disjoint paths P_1 and P_2 such that for i = 1, 2, end $(P_i) = \{s_i, t_i\}$. To state a result on graphs without an $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage, we use the following notion due to Seymour [19], which can also be found in [25].

Definition 3.1. A 3-planar graph (G, \mathcal{A}) consists of a graph G and a family $\mathcal{A} = \{A_1, \ldots, A_k\}$ of pairwise disjoint subsets of V(G) (allowing $\mathcal{A} = \emptyset$) such that

- (i) for $1 \le i \ne j \le k$, $N(A_i) \cap A_j = \emptyset$.
- (ii) for $1 \leq i \leq k$, $|N(A_i)| \leq 3$, and
- (iii) if $p(G, \mathcal{A})$ denotes the graph obtained from G by (for each i) deleting A_i and adding new edges joining every pair of distinct non-adjacent vertices in $N(A_i)$, then $p(G, \mathcal{A})$ can be drawn in a closed disc D with no pair of edges crossing such that, for each A_i with $|N(A_i)| = 3$, $N(A_i)$ induces a facial triangle in $p(G, \mathcal{A})$.

If, in addition, b_1, \ldots, b_n are some vertices in G such that $b_i \notin A_j$ for any $A_j \in \mathcal{A}$ and b_1, \ldots, b_n occur on the boundary of D in that cyclic order, then we say that $(G, \mathcal{A}, b_1, \ldots, b_n)$ is 3-planar. We will say that such a drawing is a *plane drawing of* $(G, \mathcal{A}, b_1, \ldots, b_n)$. We will say that (G, b_1, \ldots, b_n) is 3-planar if there exists a collection \mathcal{A} so that $(G, \mathcal{A}, b_1, \ldots, b_n)$ is 3-planar. If $(G, \emptyset, b_1, \ldots, b_n)$ is 3-planar we will say that (G, b_1, \ldots, b_n) is planar.

The main tool we will use is the following theorem due to Seymour [19], while different versions are proved in [2, 20, 23].

Theorem 3.2. Let G be a graph and let s_1, t_1, s_2, t_2 be distinct vertices of G. Then G contains no $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage if and only if (G, s_1, s_2, t_1, t_2) is 3-planar.

It is convenient for us to develop some lemmas on 3-planar graphs before we begin the proof of our main Theorem 1.2. The main lemmas in this section are Lemmas 3.4 and 3.7, which we will use in later sections to show that a certain 3-planar graph is in fact planar. First we have a definition from [25].

Definition 3.3. Let (G, \mathcal{A}) be 3-planar, let $A \in \mathcal{A}$ with $N(A) = \{a_1, \ldots, a_m\}$ (where $m \leq 3$), and let $H = G[A \cup N(A)]$. We say that A is *minimal* if there is no collection \mathcal{H} of pairwise disjoint subsets of A such that $\mathcal{H} \neq \{A\}$ and $(H, \mathcal{H}, a_1, \ldots, a_m)$ is 3-planar. We say that \mathcal{A} is minimal if every member of \mathcal{A} is minimal.

From [25], we have the following.

Lemma 3.4. If $(G, b_0, ..., b_n)$ is 3-planar, then there is a collection \mathcal{A} of pairwise disjoint subsets of $V(G) \setminus \{b_0, ..., b_n\}$ such that $(G, \mathcal{A}, b_0, ..., b_n)$ is 3-planar and \mathcal{A} is minimal.

Now we give two propositions to help prove Lemma 3.7 when extending a linkage in $p(G, \mathcal{A})$ to a linkage in G.

Proposition 3.5. Let (G, \mathcal{A}) be 3-planar so that \mathcal{A} is minimal. Let $s_1, t_1, s_2 \in V(p(G, \mathcal{A}))$ be distinct, and let $t_2 \in V(G)$. Let $t_2^* = t_2$ when $t_2 \in V(p(G, \mathcal{A}))$, and let t_2^* be an arbitrary vertex in $N(\mathcal{A})$ when $t_2 \in \mathcal{A}$ for some $\mathcal{A} \in \mathcal{A}$. Suppose that $t_2^* \notin \{s_1, t_1, s_2\}$ and $p(G, \mathcal{A})$ contains an $(\{s_1, t_1\}, \{s_2, t_2^*\})$ -linkage. Then G contains an $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage.

Proof. Apply Proposition 3.2 of [25], with $b \coloneqq s_2$, $b' \coloneqq s_1$, $v \coloneqq t_1$, and $u \coloneqq t_2$. Note that condition (ii) for Proposition 3.2 is guaranteed by Proposition 3.1 of [25]. (The proof basically starts with disjoint paths S_1, S_2 in $p(G, \mathcal{A})$ from s_1, s_2 to t_1, t_2^* , respectively, replaces each edge $uv \in E(S_1 \cup S_2) - E(G)$ with a path in $G[\mathcal{A} \cup N(\mathcal{A})]$ (for some $\mathcal{A} \in \mathcal{A}$ with $u, v \in N(\mathcal{A})$), and extends the path from t_2^* to t_2 .)

Proposition 3.6. Fix a plane drawing of a 3-connected planar graph G with outer cycle Z. Let $s_1, t_1, s_2 \in V(Z)$ be distinct vertices and let $t_2^* \in V(G) \setminus V(Z)$. Then G has an $(\{s_1, t_1\}, \{s_2, t_2^*\})$ -linkage.

Proof. Let P denote the path in $Z - s_2$ between s_1 and t_1 . If G - V(P) contains a path Q from t_2^* to s_2 then P, Q form the desired $(\{s_1, t_1\}, \{s_2, t_2^*\})$ -linkage. Hence, we may assume that such Q does not exist. Let C denote the component of G - V(P) containing t_2^* . Then $s_2 \notin V(C)$. Since G is 3-connected, $|N(C) \cap V(P)| \ge 3$. So let $v_0, v_1 \in N(C) \cap V(P)$ with $P[v_0, v_1]$ maximal. Then by planarity of $G, \{v_0, v_1\}$ is a cut, a contradiction.

Now we are ready to prove the final lemma for this section.

Lemma 3.7. Let (G, \mathcal{A}) be 3-planar so that G is 3-connected and \mathcal{A} is minimal. Fix some plane drawing of $p(G, \mathcal{A})$ and let F be the set of vertices on the outer face. Suppose that $|F| \ge 4$. Let s_1, t_1, s_2 be three distinct vertices in F, and let $t_2 \in V(G) \setminus F$. Then G has an $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage.

Proof. Since G is 3-connected and $p(G, \mathcal{A})$ has at least four vertices, $p(G, \mathcal{A})$ is 3connected. Let Z be the outer cycle of the plane drawing of $p(G, \mathcal{A})$. Then Z is chordless in $p(G, \mathcal{A})$, and $|V(Z)| = |F| \ge 4$. If $t_2 \in V(p(G, \mathcal{A}))$, define $t_2^* = t_2$. Otherwise, there is an $A \in \mathcal{A}$ so that $t_2 \in A$. Then define t_2^* to be a vertex in $N(A) \setminus V(Z)$, which exists since |N(A)| = 3, Z is chordless in $p(G, \mathcal{A})$, and $|V(Z)| \ge 4$.

In either case, by Proposition 3.6, $p(G, \mathcal{A})$ has an $(\{s_1, t_1\}, \{s_2, t_2^*\})$ -linkage. Then by Proposition 3.5, G has an $(\{s_1, t_1\}, \{s_2, t_2\})$ -linkage.



Figure 1: A depiction of a (c_0, c_1, c_2, c_3) -skeleton.

4 Skeletons

The objective of this section is to find an intermediate structure in a graph G, which we call a "skeleton", that will be helpful for finding an ordered cycle in G.

Definition 4.1. Let G be a graph and let $\{c_0, c_1, c_2, c_3\} \subseteq V(G)$. Then a (c_0, c_1, c_2, c_3) -skeleton is an ordered list $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ such that

- (i) C_0, C_2 , and Z are vertex disjoint cycles, $c_0, c_1 \in V(C_0), c_2, c_3 \in V(C_2)$,
- (ii) for each $i \in \{0, 1, 2, 3\}$, P_i is a path in G from c_i to some vertex $z_i \in V(Z)$ that is internally disjoint from $V(C_0) \cup V(C_2) \cup V(Z)$,
- (iii) P_0, P_1, P_2, P_3 are pairwise vertex disjoint, and z_0, z_1, z_3, z_2 occur on Z in this cyclic order.

An illustration of a skeleton is given in Figure 1. We will view $S = C_0 \cup C_2 \cup Z \cup (\bigcup_{i=0}^3 P_i)$. There are several main lemmas we need on skeletons.

Lemma 4.2. Let G be a graph with distinct vertices $c_0, c_1, c_2, c_3 \in V(G)$ and let $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ be a (c_0, c_1, c_2, c_3) -skeleton in G. Suppose G has no cycle through c_0, c_1, c_2, c_3 in order. Then, for any path P in G from $V(C_0)$ to $V(C_2)$ that is internally disjoint from S, either $end(P) = \{c_0, c_2\}$ or $end(P) = \{c_1, c_3\}$.

Proof. Let P be a path in G from $V(C_0)$ to $V(C_2)$ that is internally disjoint from S. By symmetry, we may assume that either c_0 is an end of P, or $end(P) \cap \{c_0, c_1, c_2, c_3\} = \emptyset$. Then, we may assume $c_2 \notin end(P)$, as otherwise $end(P) = \{c_0, c_2\}$ and we are done.

For i = 0, 2, let v_i be the end of P in $V(C_i)$, and fix an orientation of C_i so that c_{i+1} is not in the path $C_i[v_i, c_i]$. Fix any orientation of the cycle Z. For j = 1, 2, let z_j be the end of P_j on Z. Now

$$C_0[v_0, c_0] \cup C_0[c_0, c_1] \cup P_1 \cup Z[z_1, z_2] \cup P_2 \cup C_2[c_2, c_3] \cup C_2[c_3, v_2] \cup P$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Lemma 4.3. Suppose that G is a 7-connected graph with distinct vertices $c_0, c_1, c_2, c_3 \in V(G)$ so that G has no cycle through c_0, c_1, c_2, c_3 in order. Then, up to cyclically permuting the labels of the vertices c_0, c_1, c_2, c_3 , the graph G contains a (c_0, c_1, c_2, c_3) -skeleton.

Proof. First, we may assume that

(1) $c_i c_{i+1} \notin E(G)$ for i = 0, 1, 2, 3 (with $c_4 = c_0$).

For, suppose (1) fails. Without loss of generality, assume $c_3c_0 \in E(G)$. By Theorem 1.3, G has a path P between c_0 and c_3 such that c_0, c_1, c_2, c_3 occur on P in that order. Now $P \cup c_3c_0$ is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

We may also assume that, by cyclically permuting the labels of the vertices c_0, c_1, c_2, c_3 if necessary,

(2) there exist a family B_0 of three internally disjoint paths from c_0 to c_1 and a family B_2 of three internally disjoint paths from c_2 to c_3 , such that no path in B_0 intersects a path in B_2 .

To see this, we do the following. Let G' be the graph obtained from G by adding two copies of each of c_0, c_1, c_2 , and c_3 . That is, for each $i \in \{0, 1, 2, 3\}$, each copy of c_i has the same neighborhood as c_i . Let $S_{0,2} \subseteq V(G')$ be the set consisting of c_0 and c_2 and all of their copies, and let $S_{1,3} \subseteq V(G')$ be the set consisting of c_1 and c_3 and all of their copies. So $|S_{0,2}| = |S_{1,3}| = 6$.

Since G is 7-connected, G' is 7-connected. Hence, G' contains a set of six pairwise vertex-disjoint paths from $S_{0,2}$ to $S_{1,3}$. These six paths in G' correspond to six internally disjoint paths in G with ends in $\{c_0, c_1, c_2, c_3\}$, and for i = 0, 1, 2, 3, let A_i be the set of all three corresponding paths in G with c_i as an end.

Suppose $A_0 \cap A_1 \neq \emptyset$ and $A_0 \cap A_3 \neq \emptyset$. Then $|A_0 \cap A_1| < 3$ and so $A_1 \cap A_2 \neq \emptyset$. Likewise $A_3 \cap A_2 \neq \emptyset$. But then the union of one path each from $A_0 \cap A_1$, $A_1 \cap A_2$, $A_2 \cap A_3$, and $A_3 \cap A_0$ is a cycle in G through c_0, c_1, c_2, c_3 in order, a contradiction.

So $A_0 = A_1$ or $A_0 = A_3$. By symmetry, we may assume $A_0 = A_1$ (by relabeling c_0, c_1, c_2, c_3 as c_0, c_3, c_2, c_1 in the reverse cyclic order). Then $A_2 = A_3$. Now A_0, A_2 give rise to the desired B_0, B_2 , completing the proof of (2).

We also view B_0 (respectively, B_2) as a subgraph of G which is the union of the three paths in B_0 (respectively, B_2).

(3) There exist vertex-disjoint paths R_0 and R_1 from $V(B_0) \setminus \{c_0, c_1\}$ to $V(B_2) \setminus \{c_2, c_3\}$ that are internally disjoint from $B_0 \cup B_2$.

By (1), $|V(B_0) \setminus \{c_0, c_1\}|, |V(B_2) \setminus \{c_2, c_3\}| \ge 3$. Then (3) follows since $G - \{c_0, c_1, c_2, c_3\}$ is 3-connected.

For every $i \in \{0,1\}$ and $j \in \{0,2\}$, let r_i^j be the end of the paths R_i in $V(B_j)$. We claim that (4) r_0^0 and r_1^0 are on the same path in B_0 and the r_0^2 and r_1^2 are on the same path in B_2 .

Suppose (4) fails and by symmetry between B_0 and B_2 , we may assume that the vertices r_0^0 and r_1^0 are on different paths in B_0 . Then B_0 contains both a path P_0 through r_0^0, c_0, c_1, r_1^0 in order, as well as a path P'_0 through r_0^0, c_1, c_0, r_1^0 in order. Note that B_2 contains either a path P_2 through r_0^2, c_2, c_3, r_1^2 in order or a path P'_2 through r_0^2, c_3, c_2, r_1^2 in order. Then either $P_2 \cup R_0 \cup P'_0 \cup R_1$ or $P'_2 \cup R_0 \cup P_0 \cup R_1$ is a cycle in G through c_0, c_1, c_2, c_3 in order, a contradiction.

For j = 0, 2, let P_0^j, P_1^j, P_2^j be the paths in B_j and, by (4), assume that the ends of R_0 and R_1 in $V(B_j)$ are on the path P_0^j . Relabel the paths R_0 and R_1 if necessary so that c_0, r_0^0, r_1^0, c_1 occur on P_0^0 in that order.

Note c_2, r_0^2, r_1^2, c_3 occur on P_0^2 in that order. For, otherwise, the cycle $P_0^0[r_0^0, c_0] \cup P_1^0 \cup P_0^0[c_1, r_1^0] \cup R_1 \cup P_0^2[r_1^2, c_2] \cup P_1^2 \cup P_0^2[c_3, r_0^2] \cup R_0$ is a cycle in *G* through c_0, c_1, c_2, c_3 in order, a contradiction.

Let $Z \coloneqq P_0^0[r_0^0, r_1^0] \cup R_1 \cup P_0^2[r_1^2, r_0^2] \cup R_0$. Then

$$S := (P_1^0 \cup P_2^0, P_1^2 \cup P_2^2, Z, P_0^0[c_0, r_0^0], P_0^0[c_1, r_1^0], P_0^2[c_2, r_0^2], P_0^2[c_3, r_1^2])$$

is a (c_0, c_1, c_2, c_3) -skeleton.

5 Skeletons with Connectivity Properties

This section is dedicated to proving the following.

Proposition 5.1. Suppose that G is a 7-connected graph, $c_0, c_1, c_2, c_3 \in V(G)$ are distinct, and G has no cycle through c_0, c_1, c_2, c_3 in order. Then, up to cyclically permuting the labels of the vertices c_0, c_1, c_2, c_3 , the graph G has a (c_0, c_1, c_2, c_3) -skeleton $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ such that

- (i) for every $i \in \{0, 2\}$, the graph $G[V(C_i)]$ has no cycle through c_i and c_{i+1} with fewer vertices than C_i ,
- (ii) for every $j \in \{0, 1, 2, 3\}, |V(P_i)| = 2$, and
- (iii) the graph $G (V(C_0) \cup V(C_2))$ is 2-connected.

Proof. By Lemma 4.3, some (c_0, c_1, c_2, c_3) -skeleton exists. Let $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ be a (c_0, c_1, c_2, c_3) -skeleton, $H = G - (V(C_0) \cup V(C_2))$, and B be the block of H containing Z. Let $B_1, B_2, \ldots B_m$ be the components of H - V(B) with non-empty intersection with V(S), such that $|V(B_1)| \ge |V(B_2)| \ge \ldots \ge |V(B_m)|$. Let $A_1, A_2, \ldots A_n$ be the components of H - V(B) with empty intersection with V(S), such that $|V(A_1)| \ge |V(A_2)| \ge \ldots \ge |V(A_n)|$. We choose S so that

(1) the sum $\sum_{i=0}^{3} |V(P_i)|$ is minimum,

- (2) subject to (1), |V(B)| is maximum,
- (3) subject to (2), $(|V(B_1)|, |V(B_2)|, \ldots, |V(B_m)|)$ is maximal with respect to lexicographic ordering, and
- (4) subject to (3), $(|V(A_1)|, |V(A_2)|, \dots, |V(A_n)|)$ is maximal with respect to lexicographic ordering.

For i = 0, 1, 2, 3, let z_i be the end of P_i on Z. We will show that the skeleton S satisfies conclusions (i), (ii), and (iii). First we show that it satisfies (i).

Claim 5.1.1. For each $i \in \{0, 2\}$, the graph $G[V(C_i)]$ has no cycle through c_i and c_{i+1} with fewer vertices than C_i .

Proof. By symmetry between C_0 and C_2 , it suffices to consider the case i = 0. Suppose C'_0 is a cycle in $G[V(C_0)]$ through c_0 and c_1 such that $|V(C'_0)| < |V(C_0)|$. Then $V(C'_0) \subsetneq V(C_0)$. Write $S' \coloneqq (C'_0, C_2, Z, P_0, P_1, P_2, P_3)$ and $H' \coloneqq G - (V(C'_0) \cup V(C_2))$, and let B' be the block of H' containing V(Z). Then S' is a (c_0, c_1, c_2, c_3) -skeleton and $V(H) \subsetneq V(H')$.

Note that (1) holds for S'. Since $V(B) \subseteq V(B')$, by (2) above, we have V(B) = V(B'). Let U be a component of $G[V(C_0)] - V(C'_0)$. If $N(U) \cap V(B_i) \neq \emptyset$ for some $i \in \{1, \ldots, m\}$ then S' contradicts the choice of S via (3). Otherwise, S' contradicts the choice of S via (4).

The following is a convenient step to take on the way to proving that conclusions (ii) and (iii) are satisfied.

Claim 5.1.2. There is no component of H - V(B) with empty intersection with V(S).

Proof. Suppose otherwise. Then the component A_n exists. Since B is a block of H and A_n is a component of H - V(B), it follows that A_n has no more than one neighbor in H. Then since G is 7-connected, A_n has at least three neighbors on C_0 or C_2 . By symmetry between C_0 and C_2 , we may assume that $|N(A_n) \cap V(C_0)| \ge 3$. Let $\{R_0, R_1\}$ be a minimum $(c_0, c_1, C_0, V(A_n))$ -separating pair.

Now define $X := \operatorname{int}(R_0) \cup \operatorname{int}(R_1) \cup A_n$, $T := \operatorname{end}(R_0) \cup \operatorname{end}(R_1) \cup (N(A_n) \cap V(H))$, and $Y := V(G) \setminus (X \cup T)$. Then $|Y| \ge 2$ and $|T| \le 5$. Thus since G is 7-connected, there exist edges $uv, u'v' \in E(G)$ with $u, u' \in X, v, v' \in Y, u \ne u'$, and $v \ne v'$. By Lemma 4.2 and since $|N(A_n) \cap V(C_0)| \ge 3$, we have that $N(A_n) \cap V(C_2) = \emptyset$. Then by part (ii) of the definition of a separating pair, $N(A_n) \cap V(G - H) \subseteq V(R_0 \cup R_1)$. Thus we have that $u, u' \in \operatorname{int}(R_0) \cup \operatorname{int}(R_1)$. Then by Lemma 4.2 and the definition of a separating pair, we have that $v, v' \in V(H) \setminus (X \cup T)$. Now, up to symmetry between R_0 and R_1 , we distinguish the following two cases.

First, suppose that either both $u, u' \in int(R_0)$, or $u \in int(R_0)$ and $v \notin V(B)$. By Claim 5.1.1 and part (ii) of Lemma 2.3, $G[V(C_0) \cup V(A_n)] - int(R_0)$ contains a cycle C'_0 through c_0 and c_1 . Write $S' := (C'_0, C_2, Z, P_0, P_1, P_2, P_3)$ and H' := $G - (V(C'_0) \cup V(C_2))$, and let B' be the block of H' containing V(Z). Then S' is a (c_0, c_1, c_2, c_3) -skeleton (so (1) holds for S') and $(V(H) \setminus V(A_n)) \cup \operatorname{int}(R_0) \subseteq V(H')$. If both $v, v' \in V(B)$ then both $u, u' \in \operatorname{int}(R_0)$ and so |V(B')| > |V(B)|. Otherwise, either $u, u' \in \operatorname{int}(R_0)$ and $\{v, v'\} \not\subseteq V(B)$, or $u \in \operatorname{int}(R_0)$ and $v \notin V(B)$. In either case, one of the components $B_1, B_2, \ldots, B_m, A_1, A_2, \ldots, A_{n-1}$ grows, a contradiction.

Otherwise, without loss of generality, we may assume that $u \in int(R_0)$, $u' \in int(R_1)$, and both $v, v' \in V(B)$. For i = 2, 3, let z'_i be the vertex in $V(P_i) \cap V(B)$ that is closest to c_i on P_i . Then since v and v' are distinct and B is 2-connected, B contains disjoint paths R_2 , R_3 from $\{v, v'\}$ to z'_2, z'_3 , respectively. By Claim 5.1.1 and part (iii) of Lemma 2.3, $G[V(C_0) \cup V(A_n)]$ contains paths P and P' between u and u', such that P goes through u', c_0, c_1, u in order and P' goes through u, c_0, c_1, u' in order. Then, fixing an arbitrary cyclic order of C_2 , if $v \in V(R_2)$ and $v' \in V(R_3)$ then

$$P \cup uv \cup R_2 \cup P_2[z'_2, c_2] \cup C_2[c_2, c_3] \cup P_3[z'_3, c_3] \cup R_3 \cup v'u'$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction. If $v' \in V(R_2)$ and $v \in V(R_3)$ then

$$P' \cup u'v' \cup R_2 \cup P_2[z'_2, c_2] \cup C_2[c_2, c_3] \cup P_3[z'_3, c_3] \cup R_3 \cup vu$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

We now prove that conclusion (ii) of the proposition holds.

Claim 5.1.3. For every $j \in \{0, 1, 2, 3\}, |V(P_j)| = 2.$

Proof. By symmetry, we prove the case for j = 0. Suppose for contradiction that $|V(P_0)| > 2$. Let X_0 be the component of $H - z_0$ containing $int(P_0)$, and let X_1 be the component of $H - z_0$ containing z_1 .

Suppose that $H - z_0$ has a component A other than X_0 and X_1 . Then since the graph $S - (V(C_0) \cup V(C_2) \cup \{z_0\})$ has two components, one containing $\operatorname{int}(P_0)$ and the other containing z_1 , it follows that $A \cap V(S) = \emptyset$. Since B is 2-connected, $z_0 \in V(B)$, and $(B - z_0) \cap V(S) \neq \emptyset$, $A \cap B = \emptyset$. Then A is a component of H - V(B) with $V(A) \cap V(S) = \emptyset$, a contradiction to Claim 5.1.2. So $H - z_0$ has no components other than X_0 and X_1 . Fix a cyclic ordering of Z so that z_0, z_1, z_3, z_2 occur in that cyclic order.

Case 1. $X_0 = X_1$.

Then let P be a path in $H - \{z_0\}$ of minimum length so that P has one end in $int(P_0)$ and one end in $(V(S) \cap V(H)) \setminus V(P_0)$. Let u be the end of P in $int(P_0)$ and let v be the other end of P.

Suppose $v \in V(P_3) \cup V(Z(z_1, z_2))$. Then there is a path R with ends c_0 and c_3 in $(P_0 \cup P_3 \cup Z(z_1, z_2) \cup P) - z_0$. Then, fixing arbitrary cyclic orderings of C_0 and C_2 , the graph $C_0[c_0, c_1] \cup P_1 \cup Z[z_2, z_1] \cup P_2 \cup C_2[c_2, c_3] \cup R$ is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

So $v \in (Z[z_2, z_1] \setminus \{z_0\}) \cup \operatorname{int}(P_1) \cup \operatorname{int}(P_2)$. In this case we will find a (c_0, c_1, c_2, c_3) -skeleton $S' = (C_0, C_2, Z', P'_0, P'_1, P'_2, P'_3)$ with $\sum_{j=0}^3 |V(P'_j)| < \sum_{j=0}^3 |V(P_j)|$, which is

a contradiction to the choice of S. If $v \in \operatorname{int}(P_2)$, then define $P'_0 \coloneqq P_0[c_0, u]$, $P'_1 \coloneqq P_1$, $P'_2 \coloneqq P_2[c_2, v]$, $P'_3 \coloneqq P_3$, and $Z' \coloneqq P_0[u, z_0] \cup Z[z_0, z_2] \cup P_2[z_2, v] \cup P$. If $v \in Z[z_2, z_0)$, then define $P'_0 \coloneqq P_0[c_0, u]$, $P'_j \coloneqq P_j$ for j = 1, 2, 3, and $Z' \coloneqq P_0[u, z_0] \cup Z[z_0, v] \cup P$. The remaining cases are similar.

Case 2. $X_0 \neq X_1$.

We have shown that X_0 and X_1 are the only components of $H - \{z_0\}$, and that they are distinct. Now let $\{R_0, R_1\}$ be a minimum $(c_0, c_1, C_0, V(X_0))$ -separating pair. Define $T := \operatorname{end}(R_0) \cup \operatorname{end}(R_1) \cup \{z_0, c_2\}, X := X_0 \cup \operatorname{int}(R_0) \cup \operatorname{int}(R_1)$, and $Y := V(G) \setminus (X \cup T)$. Then $|T| \leq 6$ and both X and Y are non-empty. So there is an edge $uv \in E(G)$ with $u \in X$ and $v \in Y$.

Suppose that $v \in V(C_2)$. Fix a cyclic ordering of C_0 so that $u \notin V(C_0[c_0, c_1])$ and a cyclic ordering of C_2 so that $v \notin V(C_2[c_2, c_3])$. Define $R := C_0[c_0, c_1] \cup P_1 \cup Z[z_1, z_2] \cup P_2 \cup C_2[c_2, c_3]$. Then R is a path with ends c_0 and c_3 that contains the vertices c_0, c_1, c_2, c_3 in order. Since $G - \operatorname{int}(R)$ has a path from c_0 to c_3 (using X and uv), the graph G has a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

By the definition of a separating pair, $v \notin V(C_0)$. Thus $v \in X_1$ and $u \in int(R_0) \cup int(R_1)$. For i = 2, 3, let u_i be the neighbor of c_i on P_i .

We claim that the graph $H - V(X_0)$ contains disjoint paths R_2, R_3 from $\{v, z_0\}$ to u_2, u_3 , respectively. To see this, let P be a minimum-length path in X_1 from v to a vertex $p \in V(S) \cap V(X_1)$. If $p \in V(Z(z_0, z_2)) \cup V(P_1) \cup V(P_3)$, then $H - V(X_0)$ contains a $(\{u_2, z_0\}, \{u_3, v\})$ -linkage. Otherwise, $p \in V(P_2) \cup V(Z[z_2, z_0))$ and $H - V(X_0)$ contains a $(\{u_2, v\}, \{u_3, z_0\})$ -linkage.

First suppose $v \in V(R_2)$ and $z_0 \in V(R_3)$. Fix a cyclic ordering of C_0 so that $u \notin V(C[c_0, c_1])$ and fix an arbitrary cyclic ordering of C_2 . Then

$$C_0[c_0, u] \cup uv \cup R_2 \cup u_2c_2 \cup C_2[c_2, c_3] \cup c_3u_3 \cup R_3 \cup P_0$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

So assume $z_0 \in V(R_2)$ and $v \in V(R_3)$. Fix any cyclic ordering of C_2 . By Claim 5.1.1 and part (i) of Lemma 2.3, there is an edge $ax \in E(G)$ with $x \in X_0$ so that the graph $G[V(C_0)]$ contains a path P with ends u and a through u, c_0, c_1, a in order. Let P' be a path in $G[V(X_0) \cup \{z_0\}]$ with ends x and z_0 . Then

$$P \cup ax \cup P' \cup R_2 \cup u_2c_2 \cup C_2[c_2, c_3] \cup c_3u_3 \cup R_3 \cup vu$$

is a path through c_0, c_1, c_2, c_3 in order. This is a contradiction, and completes the proof of the claim.

Now we are ready to finish the proof of Proposition 5.1. Parts (i) and (ii) hold by Claims 5.1.1 and 5.1.3, respectively. Also by Claim 5.1.3, $V(S) \cap V(H) \subseteq V(B)$. Then every component of H - V(B) has empty intersection with V(S). So by Claim 5.1.2, V(H) = V(B) and thus H is 2-connected; so (iii) holds.

6 Raising the Connectivity

In this section we strengthen conclusion (iii) of Proposition 5.1 so that we will be able to apply Lemma 3.7. That is, we prove the following:

Proposition 6.1. Suppose that G is a 7-connected graph with distinct vertices c_0 , $c_1, c_2, c_3 \in V(G)$ so that G has no cycle through c_0, c_1, c_2, c_3 in order. Then, up to cyclically permuting the labels of the vertices c_0, c_1, c_2, c_3 , the graph G has a (c_0, c_1, c_2, c_3) -skeleton $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ so that:

- (i) for every $i \in \{0, 2\}$, the graph $G[V(C_i)]$ has no cycle through c_0 and c_1 with fewer vertices than C_i ,
- (ii) for every $j \in \{0, 1, 2, 3\}, |V(P_j)| = 2$, and
- (iii) the graph $G (V(C_0) \cup V(C_2))$ is 3-connected.

Proof. Let $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ be a (c_0, c_1, c_2, c_3) -skeleton as in Proposition 5.1. For every $i \in \{0, 1, 2, 3\}$, let z_i be the end of P_i on Z. Define $H := G - (V(C_0) \cup V(C_2))$. The only thing we need to prove is that H is 3-connected. Suppose H is not 3-connected. Let $T \subseteq V(H)$ and A be a component of H - T so that $|T| \leq 2, H - T$ is not connected, and $|V(A) \cap \{z_0, z_1, z_2, z_3\}|$ is minimum. Since G is 7-connected, there exists $i \in \{0, 2\}$ so that $N(A) \cap (V(C_i) \setminus \{c_i, c_{i+1}\})$ is non-empty. So by symmetry, we may assume that either $A \cap \{z_0, z_1, z_2, z_3\} = \emptyset$ and $N(A) \cap (V(C_0) \setminus \{c_0, c_1\})$ is non-empty, or $z_0 \in V(A)$. First we prove two claims.

Claim 6.1.1. $z_2, z_3 \notin V(A)$.

Proof. Suppose otherwise. Then $V(A) \cap \{z_0, z_1, z_2, z_3\} \neq \emptyset$, and so $z_0 \in V(A)$. Then by the choice of A and since H-T is disconnected, $|V(A) \cap \{z_0, z_1, z_2, z_3\}| = 2$. Indeed, $V(A) \cap \{z_0, z_1, z_2, z_3\} = \{z_0, z_2\}$. For, otherwise, $V(A) \cap \{z_0, z_1, z_2, z_3\} = \{z_0, z_3\}$. Then, since A is connected and H - V(A) is connected, H has a $(\{z_0, z_3\}, \{z_1, z_2\})$ linkage; so G has a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

For i = 0, 2, let $\{R_0^i, R_1^i\}$ be a minimum $(c_i, c_{i+1}, C_i, V(A))$ -separating pair. Define

$$T' \coloneqq (T \cup \operatorname{end}(R_0^0) \cup \operatorname{end}(R_1^0) \cup \operatorname{end}(R_0^2) \cup \operatorname{end}(R_1^2)) \setminus \{c_0, c_2\}.$$

By (ii) of the definition of separating pairs, for i = 0, 2 the vertex c_i is an end of both R_0^i and R_1^i . So $|T'| \leq 6$. Define

$$X \coloneqq \operatorname{int}(R_0^0) \cup \operatorname{int}(R_1^0) \cup \operatorname{int}(R_0^2) \cup \operatorname{int}(R_1^2) \cup V(A) \cup \{c_0, c_2\}$$

and let $Y \coloneqq V(G) \setminus (X \cup T')$.

Since G is 7-connected, T' is not a cut separating X and Y. So there exists an edge $uv \in E(G)$ with $u \in X$ and $v \in Y$. From (ii) of the definition of a separating pair, we have that $u \notin V(A)$. So by symmetry between C_0 and C_2 , we may assume that $u \in int(R_0^0) \cup int(R_1^0) \cup \{c_0\}$. Then by (iii) of the definition of a separating pair

and Lemma 4.2, we have that $v \in V(H) \setminus (V(A) \cup T)$. Recall that $G[V(C_0)]$ contains no cycle through c_0 and c_1 with fewer vertices than C_0 . Then by (i) of Lemma 2.3, there exists an edge $aa' \in E(G)$ with $a \in V(C_0)$ and $a' \in V(A)$ so that $G[V(C_0)]$ contains a path P with ends a and u which goes through the vertices u, c_0, c_1, a in order.

Since A and H - V(A) are both connected, the graph H has a $(\{z_2, a'\}, \{z_3, v\})$ linkage S_2, S_3 so that z_2 is an end of S_2 and z_3 is an end of S_3 . Then, fixing an arbitrary cyclic ordering of C_2 ,

$$P \cup aa' \cup S_2 \cup z_2c_2 \cup C_2[c_2, c_3] \cup c_3z_3 \cup S_3 \cup vu$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Now, let $\{R_0, R_1\}$ be a minimum $(c_0, c_1, C_0, V(A))$ -separating pair in $G - (T \cup C_2)$. Next, we show another claim.

Claim 6.1.2. There is an edge $uv \in V(G)$ with $u \in int(R_0) \cup int(R_1)$ and $v \in V(H) \setminus (T \cup V(A))$.

Proof. Define $W := \{c_{i+2} : z_i \in V(A)\}$. So $W \subseteq \{c_2, c_3\}$ by Claim 6.1.1. Define $T' := T \cup \operatorname{end}(R_0) \cup \operatorname{end}(R_1) \cup W$, $X := \operatorname{int}(R_0) \cup \operatorname{int}(R_1) \cup V(A)$, and $Y := V(G) \setminus (T' \cup X)$. Then $|T'| \leq 6$ since, for i = 0, 1, if $z_i \in V(A)$ then c_i is an end of both R_0 and R_1 . Furthermore, both X and Y are non-empty and V(G) is the disjoint union of X, Y, and T'. Since G is not 7-connected, T' is not a cut separating X and Y. So there is an edge $uv \in E(G)$ with $u \in X$ and $v \in Y$.

If $v \in V(H) \setminus (T \cup V(A))$, then $u \in int(R_0) \cup int(R_1)$ and we are done. By the definition of a separating pair, $v \notin V(C_0) \setminus (V(R_0) \cup V(R_1))$. So $v \in V(C_2) \setminus W$, and by Lemma 4.2, $u \in V(A)$.

First suppose $V(A) \cap \{z_0, z_1, z_2, z_3\} = \emptyset$. Then since H - V(A) is connected, it has (not necessarily disjoint) paths Q_0 with ends z_0 and z_3 and Q_1 with ends z_1 and z_2 . Recall that we assumed $(N(A) \cap V(C_0)) \setminus \{c_0, c_1\} \neq \emptyset$. So the graph $G[V(A) \cup V(C_0)]$ contains (not necessarily disjoint) paths S_0 with ends u and c_0 containing c_1 , and S_1 with ends u and c_1 containing c_0 . Then $S_0 \cup c_0 z_0 \cup Q_0$ is a path in $G - C_2$ that goes through u, c_1, c_0, z_3 in order. Similarly $S_1 \cup c_1 z_1 \cup Q_1$ is a path contained in $V(G) \setminus V(C_2)$ that goes through u, c_0, c_1, z_2 in order. Hence, $C_2 \cup S_0 \cup c_0 z_0 \cup Q_0 \cup uv$ or $C_2 \cup S_1 \cup c_1 z_1 \cup Q_1 \cup uv$ contains a cycle through c_0, c_1, c_2, c_3 in order in G, a contradiction. (When $v \in \{c_2, c_3\}$, only one of these works.)

Now suppose that $V(A) \cap \{z_0, z_1, z_2, z_3\} = \{z_0\}$. Then $c_2 \in W$ and $v \neq c_2$. Fix a cyclic ordering of C_2 so that $c_3 \in V(C[c_2, v])$. Let S_0 be a path contained in A with ends u and z_0 , and let S_1 be a path contained in H - A with ends z_1 and z_2 . Fix an arbitrary cyclic ordering of C_0 . Then

$$C_0[c_0, c_1] \cup c_1 z_1 \cup S_1 \cup z_2 c_2 \cup C[c_2, v] \cup v u \cup S_0 \cup z_0 c_0$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Finally, let $V(A) \cap \{z_0, z_1, z_2, z_3\} = \{z_0, z_1\}$. Then $W = \{c_2, c_3\}$; so $v \in V(C_2) \setminus \{c_2, c_3\}$. In this case it suffices to show that H has either a $(\{z_0, u\}, \{z_1, z_2\})$ -linkage or a $(\{z_0, z_3\}, \{z_1, u\})$ -linkage, as such a linkage, S, and uv give a cycle through c_0, c_1, c_2, c_3 in order, a contradiction. Fix a cyclic ordering of Z so that the vertices z_0, z_1, z_3, z_2 occur on Z in that order. Since $V(A) \cap \{z_0, z_1, z_2, z_3\} = \{z_0, z_1\}, T$ must contain exactly one vertex in $Z(z_1, z_3]$ and one vertex in $Z[z_2, z_0)$. Let P be a shortest path in A from u to a vertex $x \in V(Z) \cap V(A)$, and let x be the end of P on Z. Then $x \in V(Z(z_2, z_3))$. If $x \in V(Z(z_2, z_0))$, then $P \cup Z[x, z_0], Z[z_1, z_2]$ form a $(\{z_0, u\}, \{z_1, z_2\})$ -linkage. If $x \in V(Z(z_0, z_3))$, then $Z[z_3, z_0], P \cup Z[x, z_1]$ form a $(\{z_0, z_3\}, \{z_1, u\})$ -linkage. This completes the proof of the claim.

Now we show that there exists $t \in T$ so that H - V(A) contains disjoint paths Q_2, Q_3 from $\{t, v\}$ to z_2, z_3 , respectively. Otherwise, by Menger's Theorem, there is a separation (X, Y) of H - V(A) of order one or less so that $\{z_2, z_3\} \subseteq X$ and $\{v\} \cup T \subseteq Y$. Then $|X \setminus Y| \ge 1$ and $|(Y \cup V(A)) \setminus X| \ge |V(A)| \ge 1$. So $(X, Y \cup V(A))$ is a non-trivial separation of H of order one or less, a contradiction since H is 2-connected.

By (i) of Lemma 2.3, for i = 0, 1 there is an edge $a_i a'_i \in E(G)$ so that $a_i \in V(C_0)$, $a'_i \in V(A)$, and $G[V(C_0)]$ contains a path S_i with ends u and a_i going through u, c_{1-i}, c_i, a_i in order. For i = 0, 1, let Q_i be a path in A with ends a'_i and t.

Fix any cyclic ordering of C_2 . If $v \in V(Q_2)$ and $t \in V(Q_3)$, then

$$Q_2 \cup z_2 c_2 \cup C_2[c_2, c_3] \cup c_3 z_3 \cup Q_3 \cup Q_0 \cup a'_0 a_0 \cup S_0 \cup uv$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction. If $t \in V(Q_2)$ and $v \in V(Q_3)$, then

$$Q_3 \cup z_3 c_3 \cup C_2[c_2, c_3] \cup c_2 z_2 \cup Q_2 \cup Q_1 \cup a_1' a_1 \cup S_1 \cup uv$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

7 Discharging and Proof of Theorem 1.2

First we prove a discharging lemma on planar graphs, and then we complete the proof of the main theorem.

Lemma 7.1. Let H be a 3-connected planar graph with some fixed planar drawing. Let Z be the outer cycle of H and let x and y be distinct vertices in V(Z). Then either

- (i) there exists $v \in V(H) \setminus V(Z)$ with $d(v) \leq 6$, or
- (ii) there exists $uv \in E(Z)$ so that $\{u, v\} \cap \{x, y\} = \emptyset$ and $d(u) + d(v) \le 7$.

Proof. Let \mathcal{F} denote the set of all facial cycles of G. Then $2|E(H)| = \sum_{f \in \mathcal{F}} |V(f)| \ge |V(Z)| + 3(|\mathcal{F}| - 1)$. By Euler's Formula, we have $12 - 6|V(H)| + 6|E(H)| = 6|\mathcal{F}| \le 4|E(H)| + 6 - 2|V(Z)|$. Thus,

$$2|E(H)| \le 4|V(H)| + 2|V(H) \setminus V(Z)| - 6.$$

Now for every vertex $v \in V(Z)$, define $ch_0(v) \coloneqq d(v) - 4$ and for every $v \in V(H) \setminus V(Z)$, define $ch_0(v) \coloneqq d(v) - 7$. Then by the last inequality and since x and y have degrees at least three,

$$\sum_{v \in V(H) \setminus \{x,y\}} ch_0(v) \le 2 + 2|E(H)| - 4|V(H)| - 3|V(H) \setminus V(Z)|$$
$$\le -|V(H) \setminus V(Z)| - 4.$$

Now, for all distinct vertices $v, u \in V(Z) \setminus \{x, y\}$ so that $uv \in E(Z)$ and $d(u) \ge 6$, give one unit of charge from u to v. For all distinct vertices $v, u \in V(Z) \setminus \{x, y\}$ so that $uv \in E(Z)$ and d(u) = 5, give 1/2 unit of charge from u to v. Denote the charge function obtained from ch_0 this way by ch. Observe that $V(H) \setminus V(Z) \neq \emptyset$ as H is 3-connected, and hence

$$\sum_{v \in V(H) \setminus \{x,y\}} ch(v) = \sum_{v \in V(H) \setminus \{x,y\}} ch_0(v) < -4.$$

Suppose that conclusion (i) does not hold. Then for every vertex $v \in V(H) \setminus V(Z)$, $ch(v) = ch_0(v) = d(v) - 7 \ge 0$. For every vertex $v \in V(Z) \setminus \{x, y\}$ with $d(v) \ge 6$, $ch(v) \ge ch_0(v) - 2 = d(v) - 6 \ge 0$. Likewise every vertex $v \in V(Z) \setminus \{x, y\}$ of degree four or five has $ch(v) \ge 0$. If $v \in V(Z) \setminus \{x, y\}$ with degree less than four, then d(v) = 3 and $ch(v) \ge -1$.

Therefore, since $\sum_{v \in V(H) \setminus \{x,y\}} ch(v) < -4$, there exists a vertex $v \in V(Z) \setminus \{x,y\}$ with negative charge so that neither of its neighbors on Z are x or y. Let u and u' be the neighbors of v on Z. Then since ch(v) < 0, v has degree three and u or u' has degree no more than four. Thus case (ii) of the lemma holds.

Now we are ready to complete the proof of Theorem 1.2.

ι

Proof. Let G be a 7-connected graph with distinct vertices $c_0, c_1, c_2, c_3 \subseteq V(G)$ and, going for a contradiction, suppose that G has no cycle through c_0, c_1, c_2, c_3 in order. Let $S = (C_0, C_2, Z, P_0, P_1, P_2, P_3)$ be a (c_0, c_1, c_2, c_3) -skeleton as in Proposition 6.1. For every $i \in \{0, 1, 2, 3\}$, let z_i be the end of P_i on Z. Define $H \coloneqq G - (V(C_0) \cup V(C_2))$. First we prove a claim.

Claim 7.1.1. There is a plane drawing of the graph (H, z_0, z_1, z_3, z_2) with outer cycle Z' such that $N(C_0 \cup C_2) \subseteq V(Z')$.

Proof. First of all, since G has no cycle through c_0, c_1, c_2, c_3 in order, the graph H has no $(\{z_0, z_3\}, \{z_1, z_2\})$ -linkage. So by Theorem 3.2, we have that (H, z_0, z_1, z_3, z_2) is 3-planar. Then by Lemma 3.4, there is a collection \mathcal{A} of pairwise disjoint subsets of $V(H) \setminus \{z_0, z_1, z_2, z_3\}$ so that $(H, \mathcal{A}, z_0, z_1, z_3, z_2)$ is 3-planar and \mathcal{A} is minimal. Fix a plane drawing of $p(H, \mathcal{A})$, and let F be the set of vertices on its outer face (of $p(H, \mathcal{A})$).

We claim that $N(C_0 \cup C_2) \subseteq F$. If this is true, then since G is 7-connected and for every $A \in \mathcal{A}$, $|N_H(A)| = 3$, we will have $\mathcal{A} = \emptyset$ and be done. So suppose that there exists $u \in N(C_0 \cup C_2) \setminus F$. By symmetry between C_0 and C_2 , we may assume that $u \in N(C_0)$. By symmetry between c_0 and c_1 on C_0 , we may assume that there is a vertex $v \in V(C_0) \setminus \{c_1\}$ so that $vu \in E(G)$.

By Lemma 3.7 applied to $(H, \mathcal{A}, z_0, z_1, z_3, z_2)$, the graph H has a $(\{z_1, z_2\}, \{z_3, u\})$ linkage R_1, R_3 , where R_1 has ends z_1 and z_2 and R_3 has ends z_3 and u. Fix any cyclic order of C_2 , and a cyclic order of C_0 so that $v \notin \operatorname{int}(C_0[c_0, c_1])$. Then

$$C_0[c_0, c_1] \cup c_1 z_1 \cup R_1 \cup z_2 c_2 \cup C_2[c_2, c_3] \cup c_3 z_3 \cup R_3 \cup uv \cup C_0[v, c_0]$$

is a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Fix a plane drawing of (H, z_0, z_1, z_3, z_2) as in the claim, and fix a cyclic ordering of Z' so that z_0, z_1, z_3, z_2 occur on Z' in that order. We have one more claim:

Claim 7.1.2. There exist vertices $x_0, x_1 \in V(Z')$ so that $N(C_0) \cap V(H) \subseteq Z'[x_0, x_1]$ and $N(C_2) \cap V(H) \subseteq Z'[x_1, x_0]$. (So $x_0, z_0, z_1, x_1, z_3, z_2$ occur on Z' in order.)

Proof. We may assume that $N(c_0) \cap Z'(z_1, z_2) = \emptyset$ and $N(c_1) \cap Z'(z_3, z_0) = \emptyset$; for, otherwise, $G[V(C_0 \cup C_2 \cup Z')]$ contains a cycle through c_0, c_1, c_2, c_3 in order. Let x_0 be the neighbor of c_0 on Z' so that $Z'[z_2, x_0]$ is shortest possible. Let x_1 be the neighbor of c_1 on Z' so that $Z'[x_1, z_3]$ is shortest possible. We now show that x_0, x_1 satisfy Claim 7.1.2.

First suppose that there is a vertex $u \in Z'(x_1, x_0)$ with a neighbor $v \in V(C_0)$. By symmetry between c_0 and c_1 , we may assume that $v \neq c_1$. If $u \in Z'(x_1, z_2)$, the graph H has a $(\{z_1, z_2\}, \{u, z_3\})$ -linkage and thus a cycle through c_0, c_1, c_2, c_3 in order, a contradiction. So we may assume that $u \in Z'[z_2, x_0)$. By the choice of x_0 , $v \neq c_0$. Then H has a $(\{z_0, z_3\}, \{u, z_2\})$ -linkage, and thus a cycle through c_0, c_1, c_2, c_3 in order.

Now suppose that there is a vertex $u \in Z'(x_0, x_1)$ with a neighbor $v \in V(C_2)$. By symmetry between c_2 and c_3 we may assume that $v \neq c_3$. Then H contains a $(\{x_0, z_3\}, \{u, z_1\})$ -linkage, and thus G contains a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Since G is 7-connected and by Claim 7.1.2, every vertex in $V(H) \setminus V(Z')$ has degree at least seven in H. Thus by Lemma 7.1, there exists $uv \in E(Z')$ so that $\{u, v\} \cap \{x_0, x_1\} = \emptyset$ and $d_H(u) + d_H(v) \leq 7$. By symmetry between the cycles C_0

and C_2 , we may assume that both u and v are vertices in $Z'(x_0, x_1)$, and thus only have neighbors in $V(H) \cup V(C_0)$.

Let $\{R_0, R_1\}$ be a minimum $(c_0, c_1, C_0, \{u, v\})$ -separating pair. Define $A \coloneqq \operatorname{int}(R_0) \cup \operatorname{int}(R_1)$ and $T \coloneqq \operatorname{end}(R_0) \cup \operatorname{end}(R_1) \cup \{u, v\}$.

We may assume that $A \neq \emptyset$. Suppose otherwise, since $d_{C_0}(u) + d_{C_0}(v) \geq 7$ and $|V(C_0) \cap T| \leq 4$, there exists $i \in \{0, 1\}$ so that both ends of R_i are adjacent to both u and v. Since H either contains a $(\{u, z_3\}, \{v, z_2\})$ -linkage or a $(\{v, z_3\}, \{u, z_2\})$ -linkage, the graph G contains a cycle through c_0, c_1, c_2, c_3 in order, a contradiction.

Now since G is 7-connected and $|T| \leq 6$, there exists an edge $xy \in E(G)$ so that $x \in A$ and $y \in V(G) \setminus (A \cup T)$. Since $\{R_0, R_1\}$ is a separating pair, $y \notin V(C_0)$. By Lemma 4.2 and since $x \in V(C_0) \setminus \{c_0, c_1\}$, we have $y \notin V(C_2)$. So $y \in V(H) \setminus \{u, v\}$.

Since *H* is 3-connected, H - v is 2-connected and, hence, has two disjoint paths from $\{u, y\}$ to $\{z_2, z_3\}$. By (i) of Proposition 6.1 and (i) of Lemma 2.3, and since $uv \in E(G)$, for every $i \in \{0, 1\}$ the graph $G[V(C_0) \cup \{u, v\}]$ has a path through u, c_i, c_{1-i}, x in order. Thus *G* has a cycle through c_0, c_1, c_2, c_3 in order.

8 Concluding Remarks

Recall that f(k) is the minimum connectivity for a graph to be k-ordered, and g(k) is the minimum connectivity for a graph to be k-linked. Kostochka and G. Yu [14] asked the following.

Problem 8.1. Is it true that f(k) < g(k) for all $k \ge 2$?

It is not hard to show that f(2) = 2 and f(3) = 3 as there is only one cyclic ordering of three or fewer vertices. Jung showed that g(2) = 6 [10]. It follows that f(2) < g(2) and $f(3) < g(2) \le g(3)$. As observed in the literature, the graph obtained from the complete graph on 3k - 1 vertices by removing a matching of size k is not k-linked [21]. Thus as a corollary of our main Theorem 1.2, we have the next case that $f(4) = 7 < 10 \le g(4)$.

We also ask for a structural characterization when a graph G with four fixed vertices v_1, v_2, v_3, v_4 has no cycle through v_1, v_2, v_3, v_4 in order. By this we mean something similar to the Two Paths Theorem [19] and [23], which we rely on in this paper and introduced in Section 3. The theorem of X. Yu [25, 26, 27] characterizing when a graph has a path through four given vertices in a specific order also motivates our approach. We hope that some of the work in this paper can be used towards finding such a characterization. Much of the structure we expect to see appears in our proof (see Claim 7.1.1).

The techniques we use to prove Proposition 5.1 build upon work on the following:

Lovász Path Removal Conjecture. [15] For every positive integer k, there is an integer h(k) so that for every h(k)-connected graph G and all vertices s and t in G, there is an induced path P with ends s and t so that the graph G-V(P) is k-connected.

Kawarabayashi and Ozeki [12] made the following related conjecture.

Conjecture 8.2. [12] There exists a function f(k, l) such that the following holds. For every f(k, l)-connected graph G and two distinct vertices s and t in G, there are k internally disjoint paths P_1, \ldots, P_k with endpoints s and t such that $G - \bigcup_{i=1}^k V(P_i)$ is l-connected.

The above conjecture is implied by the Lovász Path Removal Conjecture. This can be seen by making copies of s and t (where all copies of s have the same neighborhood as s, and likewise for t) and repeatedly finding an induced path P between a copy of s and a copy of t that is internally disjoint from all copies of s and t. Kawarabayashi and Ozeki [12] proved that $f(k,1) \leq 2k + 1$ and $f(k,2) \leq 3k + 1$. Furthermore, J. Ma proved that with more connectivity a stronger conclusion holds [16]. Our proof of Proposition 5.1 uses similar techniques to [12] and [16] for the case of f(2,2).

Acknowledgments

The authors would like to thank Runrun Liu and the anonymous referees for helpful comments throughout, especially in corrections to Section 2.

References

- [1] B. Bollobas and C. Thomason, Highly linked graphs, *Combinatorica* **16** (1996) 313–320.
- [2] K. Chakravarti and N. Robertson, Covering three edges with a bond in a nonseparable graph, in: Deza and Rosenberg (Eds.), Ann. of Discrete Math., (1979) p. 247.
- [3] G. Chen, R. Gould, X. Yu, Graph connectivity after path removal, *Combinatorica* 23 (2003) 185–203.
- [4] G. A. Dirac, In abstrakten Graphen vorhandene vollstndige 4-Graphen und ihre Unterteilungen, Math. Nachr. 22 (1960) 61–85.
- [5] M.N. Ellingham, M.D. Plummer, and G. Yu, Linkage for the diamond and the path with four vertices, J. of Graph Theory 70 (2011) 241–261.
- [6] R.J. Faudree, Survey on results on k-ordered graphs, Discrete Math. 229 (2001) 73–87.
- [7] J. Fröhlich, K. Kawarabayashi, T. Müller, J. Pott, and P. Wollan, Linkages in large graphs of bounded tree-width, arXiv:1402.5549 (manuscript).
- [8] W. Goddard, 4-connected maximal planar graphs are 4-ordered, Discrete Math. 257(2-3) (2002) 405-410.
- [9] R. Gould, A look at cycles containing specified elements of a graph, Discrete Math. 309(29) (2009) 6299-6311.
- [10] H. Jung, Eine Verallgemeinerung des n-fachen zusammenhangs f
 ür Graphen, Math. Ann. 187 (1970) 95–103.
- [11] K. Kawarabayashi, A. Kostochka, G. Yu, On Sufficient Degree Conditions for a Graph to be k-linked, Combin. Probab. Comput. 15 (2006) 685–694.
- [12] K. Kawarabayashi and K. Ozeki, Non-separating subgraphs after deleting many disjoint paths, J. Combin. Theory Ser. B 101 (2011) 54–59.
- [13] M. Kriesell, Induced paths in 5-connected graphs, J. Graph Theory 36 (2001) 52-58.

- [14] A. Kostochka and G. Yu, An extremal problem for H-linked graphs, J. Graph Theory 50 (2005) 321–339.
- [15] L. Lovász, Problems in graph theory, in: M. Fielder (Ed.), Recent Advances in Graph Theory, Acadamia Prague (1975).
- [16] J. Ma, A note on Lovász removable path conjecture, J. Comb. 2(1) (2011) 103–109.
- [17] R. Mukae and K. Ozeki, 4-connected triangulations and 4-orderedness, Discrete Math. 310 (17-18) (2010) 2271-2272.
- [18] L. Ng and M. Schultz, k-ordered Hamiltonian graphs, J. Graph Theory 24 (1997) 45–57.
- [19] P. D. Seymour, Disjoint Paths in graphs, *Discrete Math* **29** (1980) 371–378.
- [20] Y. Shiloach, A polynomial solution to the undirected two paths problem, J. Assoc. Comput. Mach. 27 (3) (1980) 445–456.
- [21] R. Thomas and P. Wollan, The extremal function for 3-linked graphs, J. Combin. Theory Ser. B 98 (2008) 939–971.
- [22] R. Thomas and P. Wollan, An improved linear edge bound for graph linkage, Europ. J. Combinatorics 26 (2005) 309–324.
- [23] C. Thomassen, 2-linked graphs, Europ. J. Combin. 1 (1980) 371–378.
- [24] W. Tutte, How to draw a graph, Proc. London Math. Soc. 13 (1963) 743-767.
- [25] X.Yu, Disjoint paths in graphs I, 3-planar graphs and basic obstructions, Ann. Comb. 7(1) (2003) 89–103.
- [26] X. Yu, Disjoint paths in graphs II, a special case, Ann. Comb. 7(2) (2003) 105–126.
- [27] X. Yu, Disjoint paths in graphs III, Characterization, Ann. Comb. 7(2) (2003) 229–246.