# PRIMITIVE ROOT BIAS FOR TWIN PRIMES II: SCHINZEL-TYPE THEOREMS FOR TOTIENT QUOTIENTS AND THE SUM-OF-DIVISORS FUNCTION

STEPHAN RAMON GARCIA, FLORIAN LUCA, KYE SHI, AND GABE UDELL

ABSTRACT. Garcia, Kahoro, and Luca showed that the Bateman–Horn conjecture implies  $\varphi(p-1) \ge \varphi(p+1)$  for a majority of twin-primes pairs p, p+2 and that the reverse inequality holds for a small positive proportion of the twin primes. That is, p tends to have more primitive roots than does p + 2. We prove that Dickson's conjecture, which is much weaker than Bateman–Horn, implies that the quotients  $\frac{\varphi(p+1)}{\varphi(p-1)}$ , as p, p+2 range over the twin primes, are dense in the positive reals. We also establish several Schinzel-type theorems, some of them unconditional, about the behavior of  $\frac{\varphi(p+1)}{\varphi(p)}$  and  $\frac{\sigma(p+1)}{\sigma(p)}$ , in which  $\sigma$  denotes the sum-of-divisors function.

#### 1. INTRODUCTION

The number of primitive roots modulo a prime p is  $\varphi(p-1)$ , in which

$$\varphi(n) = \left| \left\{ i \in \{1, 2, \dots, n\} : (i, n) = 1 \right\} \right| = n \prod_{q \mid n} \left( 1 - \frac{1}{q} \right)$$
(1.1)

is the Euler totient function. In other words,  $\varphi(p-1)$  is the number of generators of the multiplicative group  $(\mathbb{Z}/p\mathbb{Z})^{\times}$ . We reserve p, q for prime numbers and use (a, b) to denote the greatest common divisor of a and b.

For twin primes p, p + 2, it is natural to ask about the relationship between  $\varphi(p-1)$  and  $\varphi(p+1)$ . Assuming the Bateman–Horn conjecture, Garcia, Kahoro, and Luca proved that

$$\varphi(p-1) \geqslant \varphi(p+1) \tag{1.2}$$

for a majority of the twin primes [10]. Such proportions are computed relative to the conjectured twin-prime counting function

$$\pi_2(x) \sim 2C_2 \int_2^x \frac{dt}{(\log t)^2}$$

in which

$$C_2 = \prod_{p \ge 3} \frac{p(p-2)}{(p-1)^2} \approx 0.660161815$$

is the twin primes constant [1, 13]. Here ~ stands for asymptotic equivalence:  $f \sim g$  means  $\lim_{x\to\infty} \frac{f(x)}{g(x)} = 1$ . The proportion of twin primes that satisfy (1.2) is at least 65% (assuming the Bateman–Horn conjecture), although computations

<sup>2010</sup> Mathematics Subject Classification. 11A07, 11A41, 11N05, 11N37, 11N56.

Key words and phrases. prime, twin prime, primitive root, Bateman–Horn conjecture, twin prime conjecture, prime bias, Dickson's conjecture, totient function, Chen's theorem.

SRG supported by NSF grant DMS-1800123.

p	$rac{\varphi(p+1)}{\varphi(p-1)}$	p	$\frac{\varphi(p+1)}{\varphi(p-1)}$	p	$\frac{\varphi(p+1)}{\varphi(p-1)}$	p	$\frac{\varphi(p+1)}{\varphi(p-1)}$
2381	1.03125	119771	1.0234	230861	1.03125	348461	1.02981
3851	1.06	126491	1.03986	232961	1.02648	354971	1.07174
14561	1.05208	129221	1.06786	237161	1.01061	356441	1.04177
17291	1.00309	134681	1.08247	241781	1.05652	357281	1.05826
20021	1.11806	136991	1.03558	246611	1.0571	361901	1.09268
20231	1.02941	142871	1.05983	251231	1.00926	362951	1.03542
26951	1.06857	145601	1.05313	259211	1.01637	371141	1.02375
34511	1.06845	150221	1.03489	270131	1.03752	399491	1.04795
41231	1.05926	156941	1.04382	274121	1.06364	402221	1.09235
47741	1.08	165551	1.0946	275591	1.01252	404321	1.01206
50051	1.12	166601	1.03296	278741	1.07537	406631	1.00558
52361	1.13594	167861	1.06481	282101	1.08833	410411	1.13514
55931	1.02446	173741	1.00101	282311	1.00772	413141	1.02876
57191	1.05026	175631	1.05845	298691	1.037	416501	1.03179
65171	1.02608	188861	1.04087	300581	1.03534	418601	1.1011
67211	1.01413	197891	1.02266	301841	1.04082	424271	1.16905
67271	1.0043	202931	1.05743	312551	1.04783	427421	1.00958
70841	1.11799	203771	1.01071	315701	1.09613	438131	1.03357
82811	1.02747	205031	1.0169	316031	1.05385	440441	1.15852
87011	1.07857	205661	1.05097	322631	1.07177	448631	1.10491
98561	1.0694	206081	1.00692	325781	1.05864	454721	1.00694
101501	1.00679	219311	1.05694	328511	1.05523	464171	1.00607
101531	1.00714	222041	1.02361	330821	1.04042	464381	1.01407
108461	1.00871	225611	1.0726	341321	1.02666	465011	1.06779
117041	1.12882	225941	1.00577	345731	1.04732	470471	1.1837
Table 1	. The rat	io $\frac{\varphi(p+1)}{\varphi(p-1)}$	for the f	irst 100	twin-prime	pairs wi	th $p \ge 5$

for which (1.2) fails. The initial numerical evidence suggests that  $\frac{\varphi(p+1)}{\varphi(p-1)}$  is bounded above as p, p+2 runs over the twin primes. Dickson's conjecture (and more extensive computation) suggest otherwise (Theorem 1).

suggest something around 98%. Moreover, at least 0.46% of the twin primes satisfy the reverse inequality  $\varphi(p-1) \leq \varphi(p+1)$  [10]. Analogous results for prime pairs p, p+k were obtained by Garcia, Luca, and Schaaff [12]. Garcia and Luca showed unconditionally that the split is 50/50 if only p is assumed to be prime [11].

A glance at the numerical evidence suggests that  $\frac{\varphi(p+1)}{\varphi(p-1)}$  is bounded as p, p+2 range over the twin primes; see Table 1. Our first theorem, whose proof is in Section 2, demonstrates that this is far from the truth.

**Theorem 1.** Dickson's conjecture implies that

$$\left\{\frac{\varphi(p+1)}{\varphi(p-1)}\,:\,p,\,p+2 \text{ are prime}\right\} \text{ is dense in } [0,\infty).$$

Before proceeding, we require a few words about Dickson's conjecture. The assertion that there are infinitely many twin primes is the twin prime conjecture, which remains unresolved despite significant recent work [16,18,19,27]. Thus, some unproved conjecture must be assumed to say anything nontrivial about the large-scale behavior of the twin primes. Dickson's conjecture is among the weakest general assertions that implies the twin prime conjecture [1,6,20].

 $\mathbf{2}$ 

p	$\frac{\varphi(p+1)}{\varphi(p-1)}$	p	$\frac{\varphi(p+1)}{\varphi(p-1)}$
2381	1.03125	17497481	1.27427
3851	1.06	69989921	1.27484
20021	1.11806	78278201	1.28693
50051	1.12	183953771	1.30984
52361	1.13594	242662421	1.32797
424271	1.16905	468818351	1.34577
470471	1.1837	2156564411	1.37262
602141	1.18793	24912037151	1.37901
2302301	1.2058	43874931101	1.37949
6806801	1.23097	73769375681	1.39837
16926911	1.23678	131104243271	1.42545

**Table 2.** Running leaders among twin primes p, p + 2 with  $p \ge 5$  for which (1.2) fails. Theorem 1 suggests that  $\frac{\varphi(p+1)}{\varphi(p-1)}$  can be arbitrarily large.

**Dickson's Conjecture.** If  $f_1, f_2, \ldots, f_k \in \mathbb{Z}[t]$  are linear polynomials with positive leading coefficients and  $f = f_1 f_2 \cdots f_k$  does not vanish identically modulo any prime, then  $f_1(t), f_2(t), \ldots, f_k(t)$  are simultaneously prime infinitely often.

The twin prime conjecture is the special case  $f_1(t) = t$  and  $f_2(t) = t + 2$ . Dickson's conjecture is weaker than the Bateman–Horn conjecture, which concerns polynomials of arbitrary degree and makes asymptotic predictions [1–3]. More extensive computations suggest the truth of Theorem 1; see Table 2.

Totient quotients have a long and storied history [21, Ch. 1]. Schinzel established a curious result in 1954 [22], when he showed that

$$\left\{\frac{\varphi(n+1)}{\varphi(n)}: n = 1, 2, \ldots\right\} \text{ is dense in } [0, \infty).$$
(1.3)

This inspired later research by Schinzel, Sierpiński, Erdős, and others [7,8,23–26].

The prime analogue of (1.3) is false since  $\limsup_{p\to\infty} \frac{\varphi(p+1)}{\varphi(p)} \leq \frac{1}{2}$  because p+1 is even when p is odd. Taking this into account, we prove in Section 3 that the following modified analogue of Schinzel's theorem holds unconditionally. The main ingredient is a generalization of Chen's theorem [9, Thm. 25.11].

Theorem 2. (Unconditional)

$$\left\{\frac{\varphi(p+1)}{\varphi(p)} : p \text{ prime}\right\} \text{ is dense in } \left[0, \frac{1}{2}\right].$$

The corresponding twin-prime analogue of Schinzel's theorem (1.3) is the following result, whose proof is in Section 4.

**Theorem 3.** Dickson's conjecture implies that

$$\left\{\frac{\varphi(p+1)}{\varphi(p)}\,:\,p,\,p+2 \text{ prime}\right\} \text{ is dense in } \left[0,\frac{1}{3}\right].$$

Our proofs are transparent enough to permit the construction of striking numerical examples that cannot be obtained easily through brute force alone. For example, the twin primes

 $p = 7642856398602124688629749934198565871312540429046303895770916192 \\951906734870659150561934966844646027084062815031558255187497845592422 \\263591099165956612523533130293687015551343007872097253311773591110917 \\$ 

 $528188316414599672498791126998631571669927514554696197257278634275128\\724279128189209746271781127728971190835766821004705695431931600462680\\599536653440211216644627374103340174280330773185320397138921513211749\\572729188424740776331734373496916520151521002589283157543831117944688\\070785572501766739882905425624193096685596320850078698276901411040453\\944941282023066485167044354346570749231118823789564541354721325254528\\236757389662894019979784752005836753790957246918204110303873795632580\\172005942425384506696512023749191214954151888293569682377697953981506\\694554048884224168645065101150446101000833424954728109752271343988672\\243937380333654916021273646288933510923983231864735583985112984704333\\900355465514181712760200823033200490940934697445099785257641185993998\\522771636033743449797832811518401162205004091941408725193787590452993\\251183485650458246694103753512762453446893673884455566405601755508021$ 

and p+2 yield a ratio  $\frac{\varphi(p+1)}{\varphi(p-1)} = 3.11615...$ , which is far larger than those displayed in Table 2. As another example, consider  $\frac{\pi}{10} = 0.31415... \in [0, \frac{1}{3}]$ . The method of proof of Theorem 3 (with slight modifications) and a computer search yields the the twin prime pair p = 7726274821004474852086566160138278575763701613133157and p+2, which satisfies (the underlined digits agree with those of  $\frac{\pi}{10}$ )

$$\frac{\varphi(p+1)}{\varphi(p)} = \underline{0.314159265358979}21341\dots$$

Theorem 1, Theorem 2, and Theorem 3 each have analogues for the sum-ofdivisors function  $\sigma(n) = \sum_{d|n} d$ . We collect these results in the following theorem, whose proof is in Section 5.

### Theorem 4.

4

(a) Dickson's conjecture implies that

$$\left\{\frac{\sigma(p+1)}{\sigma(p-1)}:\,p,\,p+2 \text{ prime}\right\} \text{ is dense in } [0,\infty).$$

(b) (Unconditional)

$$\left\{\frac{\sigma(p+1)}{\sigma(p)}: p \text{ prime}\right\} \text{ is dense in } \left[\frac{3}{2}, \infty\right).$$

(c) Dickson's conjecture implies that

$$\left\{\frac{\sigma(p+1)}{\sigma(p)}: p, p+2 \text{ prime}\right\} \text{ is dense in } [2,\infty).$$

#### 2. Proof of Theorem 1

A folk lemma. Mertens' third theorem asserts that

$$\prod_{q \leqslant x} \left( 1 - \frac{1}{q} \right) \sim \frac{e^{-\gamma}}{\log x},\tag{2.1}$$

in which  $\gamma$  is the Euler–Mascheroni constant [14, 17]. A more elementary proof of the following lemma can be based on [5, Prop. 8.8] instead.

**Lemma 5.** Let  $\mathcal{P}$  denote a finite set of primes. Then

$$\left\{\frac{\varphi(n)}{n}: n \text{ squarefree, } p \nmid n \text{ for all } p \in \mathcal{P}\right\} \text{ is dense in } [0,1].$$

*Proof.* Let  $\xi \in (0,1]$  and  $n_t = \prod_{e^{\xi t} \leq q < e^t} q$ , in which  $t > \frac{1}{\xi} \log \max \mathcal{P}$ . Then  $n_t$  is squarefree,  $p \nmid n_t$  for all  $p \in \mathcal{P}$ , and

$$\frac{\varphi(n_t)}{n_t} = \prod_{e^{\xi t} \leqslant q < e^t} \left( 1 - \frac{1}{q} \right) \sim \frac{e^{-\gamma}/\log(e^t)}{e^{-\gamma}/\log(e^{\xi t})} \sim \frac{\log(e^{\xi t})}{\log(e^t)} = \xi$$

as  $t \to \infty$ .

**Initial setup.** It suffices to show that Dickson's conjecture implies that for each fixed  $\xi \in (0,\infty)$  and  $0 < \delta < 1$ , there is a twin-prime pair p, p + 2 such that  $\frac{\varphi(p+1)}{\varphi(p-1)} \in (\xi(1-\delta), \xi(1+\delta)).$ 

Let  $0 < x \leq \min \left\{\frac{2}{3}, \xi\right\}$ . Lemma 5 provides a squarefree b such that

$$(b,6) = 1$$
 and  $\frac{\varphi(b)}{b} \in \left(\frac{x}{\xi}\left(\frac{1}{1+\delta}\right), \frac{x}{\xi}\right).$ 

A second appeal to Lemma 5 yields a squarefree a' such that

$$(a', 6b) = 1$$
 and  $\frac{\varphi(a')}{a'} \in \left(\frac{3x}{2}(1-\delta), \frac{3x}{2}\right)$ 

Our choice of x ensures that the intervals specified are contained in (0, 1). Let a = 3a' and observe that

$$\frac{\varphi(a)}{a} = \frac{2}{3} \frac{\varphi(a')}{a'} \in (x(1-\delta), x).$$

Consequently,

$$\frac{\varphi(a)}{a}\frac{b}{\varphi(b)} \in \left(\xi(1-\delta), \xi(1+\delta)\right).$$
(2.2)

**The polynomials.** Our strategy is to produce linear polynomials  $f_1, f_2, \ldots, f_4$  to which Dickson's conjecture can be applied, using  $f_1, f_2$  to produce twin primes p, p+2, and using  $f_3, f_4$  to ensure that  $\frac{\varphi(p+1)}{\varphi(p-1)}$  falls in the desired interval.

Since  $8, a^2, b^2$  are pairwise relatively prime, the Chinese remainder theorem provides c such that

$$c \equiv 5 \pmod{8},\tag{2.3}$$

$$c \equiv a - 1 \pmod{a^2}, \quad \text{and} \tag{2.4}$$

$$c \equiv b + 1 \pmod{b^2}.\tag{2.5}$$

Since  $3 \mid a$ , it follows from (2.4) that

$$c \equiv 2 \pmod{3}.\tag{2.6}$$

Define

$$h(t) = 24a^2b^2t + c$$

and let

$$\begin{split} f_1(t) &= h(t), \\ f_2(t) &= h(t) + 2, \\ f_3(t) &= \frac{h(t) - 1}{4b} = \frac{24a^2b^2t + (c - 1)}{4b} = 6a^2bt + \frac{c - 1}{4b}, \quad \text{and} \\ f_4(t) &= \frac{h(t) + 1}{2a} = \frac{24a^2b^2t + (c + 1)}{2a} = 12ab^2t + \frac{c + 1}{2a}, \end{split}$$

Clearly  $f_1, f_2 \in \mathbb{Z}[t]$ . Observe that (2.3) and (2.5) ensure that  $f_3$  has integral coefficients. Similarly, (2.3) and (2.4) ensure that  $f_4$  has integral coefficients. Thus, all four polynomials are in  $\mathbb{Z}[t]$  and have positive leading coefficients.

Nonvanishing product. We claim that  $f = f_1 f_2 f_3 f_4$  does not vanish identically modulo any prime. Since

$$f_1(t) \equiv f_2(t) \equiv c \equiv 1 \pmod{2}, \qquad \text{by (2.3)},$$
  

$$f_3(t) \equiv \frac{c-1}{4b} \equiv 1 \pmod{2}, \qquad \text{by (b, 2)} = 1 \text{ and (2.3)},$$
  

$$f_4(t) \equiv \frac{c+1}{2a} \equiv 1 \pmod{2}, \qquad \text{by (a, 2)} = 1 \text{ and (2.3)},$$

it follows that f does not vanish modulo 2. Similarly,

$$\begin{aligned} f_1(t) &\equiv c \equiv 2 \pmod{3}, & \text{by (2.6)}, \\ f_2(t) &\equiv c + 2 \equiv 1 \pmod{3}, & \text{by (2.6)}, \\ f_3(t) &\equiv \frac{c-1}{4b} \equiv b \not\equiv 0 \pmod{3}, & \text{by (2.6)}, \\ f_4(t) &\equiv \frac{c+1}{2a} \equiv 2 \pmod{3}, & \text{by (b,3)} = 1 \text{ and (2.6)}, \\ \end{aligned}$$

so f does not vanish modulo 3. The final statement perhaps deserves a bit of explanation. From (2.4) we have  $c + 1 \equiv a \pmod{a^2}$  and hence  $\frac{c+1}{a} \equiv 1 \pmod{a}$ . Since  $3 \mid a$ , it follows that  $\frac{c+1}{a} \equiv 1 \pmod{3}$  from which the desired statement follows.

For any prime  $q \ge 5$  such that  $q \nmid ab$ , the polynomial f has degree four and hence cannot vanish identically modulo q. Now suppose that  $q \ge 5$  is prime and  $q \mid ab$ . Then  $h(t) \equiv c \pmod{q}$ . Since (2.4) and (2.5) ensure that

$$c \equiv \begin{cases} -1 \pmod{q} & \text{if } q \mid a, \\ 1 \pmod{q} & \text{if } q \mid b, \end{cases}$$
(2.7)

it follows that  $f_1$  and  $f_2$  do not vanish modulo q. Similarly,

$$f_{3}(t) \equiv \frac{c-1}{4b} \equiv \begin{cases} -2^{-1}b^{-1} \pmod{q} & \text{if } q \mid a \ (\text{by } (2.7)), \\ 4^{-1} \pmod{q} & \text{if } q \mid b \ (\text{by } (2.5)), \end{cases}$$
$$f_{4}(t) \equiv \frac{c+1}{2a} \equiv \begin{cases} 2^{-1} \pmod{q} & \text{if } q \mid a \ (\text{by } (2.4)), \\ a^{-1} \pmod{q} & \text{if } q \mid b \ (\text{by } (2.7)). \end{cases}$$

Thus, f does not vanish identically modulo any prime.

**Conclusion.** Dickson's conjecture provides infinitely many T such that  $f_1(T)$ ,  $f_2(T)$ ,  $f_3(T)$ , and  $f_4(T)$  are prime. For such T, the primes

$$p = f_1(T)$$
 and  $p + 2 = f_2(T)$ 

satisfy

$$p+1 = 2a f_4(T)$$
 and  $p-1 = 4b f_3(T)$ .

Consequently, (2.2) ensures that

$$\frac{\varphi(p+1)}{\varphi(p-1)} = \frac{\varphi(p+1)}{p+1} \frac{p-1}{\varphi(p-1)} \frac{p+1}{p-1}$$

PRIMITIVE ROOT BIAS FOR TWIN PRIMES II

$$= \frac{\varphi(p+1)}{p+1} \frac{p-1}{\varphi(p-1)} (1+o(1))$$

$$= \frac{\varphi(2a f_4(T))}{2a f_4(T)} \frac{4b f_3(T)}{\varphi(4b f_3(T))} (1+o(1))$$

$$= \frac{\varphi(2) \varphi(a) \varphi(f_4(T))}{2a f_4(T)} \frac{4b f_3(T)}{\varphi(4) \varphi(b) \varphi(f_3(T))} (1+o(1))$$

$$= \frac{1\varphi(a)}{2a} \frac{f_4(T)-1}{f_4(T)} \frac{4b}{2\varphi(b)} \frac{f_3(T)}{f_3(T)-1} (1+o(1))$$

$$= \frac{\varphi(a)}{a} \frac{b}{\varphi(b)} (1+o(1))$$

belongs to  $(\xi(1-\delta), \xi(1+\delta))$  for large *T*. Here we have used the facts  $(2a, f_4(T)) = 1$  and  $(4b, f_3(T)) = 1$ , which follow from (2.3) and (2.4), and from (2.3) and (2.5), respectively.

### 3. Proof of Theorem 2

Chen's theorem asserts that every sufficiently large even number is the sum of two primes, or a sum of a prime and a semiprime (a number with precisely two prime factors) [4,15]. We require a generalization of Chen's theorem to linear forms. The version below is due to Friedlander and Iwaniec [9, Thm. 25.11].

**Theorem 6** (Chen, Friedlander–Iwaniec). Let  $a, c \ge 1$  and  $b \ne 0$  be pairwise coprime integers with  $2 \mid abc$ . For t sufficiently large (in terms of abc),

$$\left| \{ p \leqslant t : ap + b = cs \} \right| \ge \frac{W(abc)}{31c} \frac{Bt}{(\log t)^2},$$

in which s has at most two prime factors, each one larger than  $t^{3/11}$ ,

$$W(d) = \prod_{\substack{p|d \ p>2}} \left(1 - \frac{1}{p-1}\right)^{-1} \quad and \quad B = 2\prod_{p\geq 3} \left(1 - \frac{1}{(p-1)^2}\right).$$

Let  $\xi \in [0, \frac{1}{2}]$  and  $\delta > 0$ . For  $x \ge \frac{\log 2}{2\xi}$ , the integer Q = Q(x) defined by

$$Q = 2 \prod_{e^{2\xi x} < q \le e^z} q$$

is divisible by 2, but not 4. As  $x \to \infty$ , (1.1) and (2.1) imply

$$\frac{\varphi(Q)}{Q} = \frac{1}{2} \prod_{e^{2\xi x} < q \leqslant e^x} \left(1 - \frac{1}{q}\right) \sim \frac{1}{2} \frac{e^{-\gamma}}{\log(e^x)} \frac{\log(e^{2\xi x})}{e^{-\gamma}} = \xi.$$

For each x, apply Theorem 6 with a = b = 1 and c = Q to obtain an S = S(x) with at most two prime factors, both of which are greater than  $\max\{Q, x\}$ , such that p = p(x) = QS - 1 is prime. Then

$$\lim_{x \to \infty} \frac{\varphi(S)}{S} = \lim_{x \to \infty} \prod_{q|S} \left( 1 - \frac{1}{q} \right) = 1$$

and hence

$$\frac{\varphi(p+1)}{\varphi(p)} = \frac{\varphi(QS)}{\varphi(QS-1)} = \frac{\varphi(Q)\,\varphi(S)}{QS-2} = \frac{\varphi(Q)}{Q}\frac{\varphi(S)}{S}\frac{S}{S-2/Q} \to \xi$$

as  $x \to \infty$ . This concludes the proof.

### 4. Proof of Theorem 3

Fix  $\xi \in (0, \frac{1}{3})$  and let  $0 < \delta < \frac{1-3\xi}{3\xi}$ . Lemma 5 yields a squarefree Q' such that

$$(Q', 6) = 1$$
 and  $\frac{\varphi(Q')}{Q'} \in (3\xi(1-\delta), 3\xi(1+\delta)).$ 

Let Q = 6Q', and observe that

$$\frac{\varphi(Q)}{Q} = \frac{1}{3} \frac{\varphi(Q')}{Q'} \in \left(\xi(1-\delta), \xi(1+\delta)\right).$$

Define the polynomials

$$f_1(t) = t$$
,  $f_2(t) = Qt - 1$ , and  $f_3(t) = Qt + 1$ .

If  $q \ge 5$  and  $q \nmid Q$ , then  $f = f_1 f_2 f_3$  has degree three and cannot vanish identically modulo q. If  $q \mid Q$ , then  $f(1) = Q^2 - 1 \equiv -1 \pmod{q}$  and hence f does not vanish identically modulo q. In particular, f does not vanish identically modulo 2 or 3. Thus, f does not vanish identically modulo any prime.

Dickson's conjecture provides infinitely many T such that  $f_1(T)$ ,  $f_2(T)$ , and  $f_3(T)$  are prime. In particular, we may assume that the prime  $f_1(T) = T$  is greater than Q so that (Q,T) = 1. Then p = QT - 1 and p + 2 = QT + 1 are twin primes and p + 1 = QT. Then

$$\frac{\varphi(p+1)}{\varphi(p)} = \frac{\varphi(QT)}{\varphi(QT-1)}$$
$$= \frac{\varphi(Q)\,\varphi(T)}{\varphi(QT-1)}$$
$$= \frac{\varphi(Q)(T-1)}{QT-2}$$
$$= \frac{\varphi(Q)}{Q}\frac{T-1}{T-2/Q}$$
$$= \frac{\varphi(Q)}{Q}(1+o(1))$$

is in  $(\xi(1-\delta), \xi(1+\delta))$  for sufficiently large T.

### 5. Proof of Theorem 4

**Proof of Theorem 4a.** The proof of Theorem 4a is similar to the proof of Theorem 1. We first require the following version of Lemma 5 for the sum-of-divisors function.

**Lemma 7.** Let  $\mathcal{P}$  denote a finite set of primes. Then

$$\left\{\frac{\sigma(n)}{n}: n \text{ squarefree, } p \nmid n \text{ for all } p \in \mathcal{P}\right\} \text{ is dense in } [1,\infty).$$

*Proof.* Let  $Q = Q(t) = \prod_{q \leq t} q$ . Then Mertens' third theorem (2.1), the Euler product formula, and the evaluation  $\zeta(2) = \frac{\pi^2}{6}$  yield

$$\frac{\sigma(Q)}{Q} = \prod_{q \leqslant t} \frac{1+q}{q} = \prod_{q \leqslant t} \left( 1 + \frac{1}{q} \right) = \frac{\prod_{q \leqslant t} (1-1/q^2)}{\prod_{q \leqslant t} (1-1/q)} \sim \frac{6/\pi^2}{e^{-\gamma}/\log t} = \frac{6e^{\gamma}}{\pi^2} \log t$$

as  $t \to \infty$ . Let  $\xi \ge 1$  and define  $n_t = \prod_{e^t \le q \le e^{\xi t}} q$ , in which  $\log t > \max \mathcal{P}$ . Then  $n_t$  is squarefree,  $p \nmid n_t$  for all  $p \in \mathcal{P}$ , and

$$\frac{\sigma(n_t)}{n_t} \sim \frac{\log(e^{\xi t})}{\log(e^t)} = \xi$$

as  $t \to \infty$ .

Fix  $\xi \in (0, \infty)$  and  $0 < \delta < 1$ . Let  $x \ge \max\left\{\frac{4}{3}, \frac{7\xi}{6}\right\}$ . Then Lemma 7 provides a squarefree b such that

$$(b,6) = 1$$
 and  $\frac{\sigma(b)}{b} \in \left(\frac{6x}{7\xi}, \frac{6x}{7\xi}\left(\frac{1}{1-\delta}\right)\right).$ 

A second appeal to Lemma 7 yields a squarefree a' such that

$$(a', 6b) = 1$$
 and  $\frac{\sigma(a')}{a'} \in \left(\frac{3x}{4}, \frac{3x}{4}(1+\delta)\right).$ 

Our choice of x ensures that the intervals specified are contained in  $(1, \infty)$ . Let a = 3a' and observe that

$$\frac{\sigma(a)}{a} = \frac{4}{3} \frac{\sigma(a')}{a'} \in (x, x(1+\delta)).$$

Consequently,

$$\frac{\sigma(a)}{a}\frac{b}{\sigma(b)} \in \left(\frac{7}{6}\xi(1-\delta), \frac{7}{6}\xi(1+\delta)\right).$$
(5.1)

Define the polynomials  $f_1, f_2, f_3, f_4$  as in the proof of Theorem 1, in which we showed that the application of Dickson's conjecture to this family is permissible. Dickson's conjecture provides infinitely many T such that  $f_1(T), f_2(T), f_3(T)$ , and  $f_4(T)$  are prime. For such T, the primes

$$p = f_1(T)$$
 and  $p + 2 = f_2(T)$ 

satisfy  $p + 1 = 2a f_4(T)$  and  $p - 1 = 4b f_3(T)$ . Consequently, (5.1) ensures that

$$\begin{aligned} \frac{\sigma(p+1)}{\sigma(p-1)} &= \frac{\sigma(p+1)}{p+1} \frac{p-1}{\sigma(p-1)} \frac{p+1}{p-1} \\ &= \frac{\sigma(p+1)}{p+1} \frac{p-1}{\sigma(p-1)} (1+o(1)) \\ &= \frac{\sigma(2a \, f_4(T))}{2a \, f_4(T)} \frac{4b \, f_3(T)}{\sigma(4b \, f_3(T))} (1+o(1)) \\ &= \frac{\sigma(2) \, \sigma(a) \, \sigma(f_4(T))}{2a f_4(T)} \frac{4b \, f_3(T)}{\sigma(4) \, \sigma(b) \, \sigma(f_3(T))} (1+o(1)) \\ &= \frac{3\sigma(a)}{2a} \frac{f_4(T)+1}{f_4(T)} \frac{4b}{7\sigma(b)} \frac{f_3(T)}{f_3(T)+1} (1+o(1)) \\ &= \frac{6}{7} \frac{\sigma(a)}{a} \frac{b}{\sigma(b)} (1+o(1)) \end{aligned}$$

belongs to  $(\xi(1-\delta), \xi(1+\delta))$  for large T.

**Proof of Theorem 4b.** Since the proof of Theorem 4b is similar to the proof of Theorem 2, we only sketch the details. First, a simple modification of Lemma 7 shows that for any finite set  $\mathcal{P}$  of primes that does not contain 2, the set

$$\left\{\frac{\sigma(n)}{n}: n \text{ squarefree and even, } p \nmid n \text{ for all } p \in \mathcal{P}\right\}$$

is dense in  $\left[\frac{3}{2},\infty\right)$ . Let  $\xi \in \left[\frac{3}{2},\infty\right)$  and mimic the proof of Theorem 2 to find an even squarefree Q = Q(x) such that  $\frac{\sigma(Q)}{Q} \to \xi$  as  $x \to \infty$ . Apply Theorem 6 and obtain an S = S(x) with at most two prime factors, both of which are greater than  $\max\{Q, x\}$ , such that p = p(x) = QS - 1 is prime. Then  $\frac{\sigma(S)}{S} \to 1$  as  $x \to \infty$  and hence

$$\frac{\sigma(p+1)}{\sigma(p)} = \frac{\sigma(QS)}{\sigma(QS-1)} = \frac{\sigma(Q)\,\sigma(S)}{QS} = \frac{\sigma(Q)}{Q}\frac{\sigma(S)}{S} \to \xi. \quad \Box$$

**Proof of Theorem 4c.** Since the proof of Theorem 4c is similar to the proof of Theorem 3, we only sketch the details. Let  $\xi \in [2, \infty)$  and mimic the proof of Theorem 3 to find a squarefree Q = Q(x) which is divisible by 6 such that  $\frac{\sigma(Q)}{Q} \to \xi$  as  $x \to \infty$ .

Define the polynomials  $f_1, f_2, f_3$  as in the proof of Section 4 in which we showed that the application of Dickson's conjecture to this family is permissible. Thus, we can find arbitrarily large T such that  $f_1(T) = T$ ,  $p = f_2(T) = QT - 1$ , and  $p + 2 = f_3(T) = QT + 1$  are simultaneously prime and hence

$$\frac{\sigma(p+1)}{\sigma(p)} = \frac{\sigma(QT)}{\sigma(QT-1)} = \frac{\sigma(Q)\,\sigma(T)}{QT} = \frac{\sigma(Q)}{Q}\frac{T+1}{T} \to \xi. \quad \Box$$

### 6. Numerical examples

Our methods of proof are transparent enough that they permit us to construct numerical examples whose totient and divisor-sum quotients approximate various mathematical constants surprisingly well (much better than can be obtained by brute force alone). Tables 3, 4, 5, 6, 7, and 8 showcase various examples for each of the theorems proven above.

**Computational differences.** For the sake of optimization, our computation of numerical examples involves slightly different methods than those provided in the proofs. In particular, our provided proofs of Lemmas 5 and 7 construct a product of consecutive primes between  $e^{\xi t}$  and  $e^t$ . Our computation takes a more naïve but more efficient process: begin with 1, and repeatedly multiply by the next smallest  $q \notin \mathcal{P}$  so that  $\frac{\varphi(n)}{n} \ge \xi$  (resp., for Lemma 7,  $\frac{\sigma(n)}{n} \le \xi$ ); convergence of this process is guaranteed by the fact that  $\prod_q (1-1/q)$  diverges to 0 (resp.,  $\prod_q (1+1/q)$  diverges to  $\infty$ ), so the sequence we construct is monotonically decreasing (resp., increasing) and is bounded tightly below (resp., above) by  $\xi$ .

For Theorem 1, Theorem 3, Theorem 4a, and Theorem 4c, the method of construction is otherwise the same, relying on the same polynomial-based approach together with Dickson's conjecture. For Theorem 2 and Theorem 4b, instead of the unconditional method of proof based on Theorem 6 provided in the paper, we instead took a polynomial/Dickson approach similar to that of Theorem 3 and Theorem 4c based on Lemma 5 and Lemma 7, since we found no straightforward numerical implementation of Theorem 6.

## PRIMITIVE ROOT BIAS FOR TWIN PRIMES II

ξ	p	$rac{arphi(p+1)}{arphi(p-1)}$
$\gamma = 0.577215664901532960 \dots$	42372380883463421517 73231363103472579179 81529437	0.577215664901532895
$\frac{\pi}{10} = 0.31415926535897932\dots$	$\begin{array}{c} 82195174147573535757\\ 39578784498983486285\\ 59983838783756381656\\ 5056272351869\\ \end{array}$	<u>0.314159265358979</u> 29
$\frac{\pi}{2} = 1.570796326794\dots$	$\begin{array}{c} 16669991702466676639\\ 32927172834786166829\\ 99425902809932261 \end{array}$	<u>1.5707963267</u> 82
$\frac{e}{10} = 0.271828182845904523\dots$	34075187932170983182 24729850026082957686 06668541034063696514 57275739410260870451 9864613389	<u>0.2718281828459045</u> 01
$\frac{1}{e} = 0.36787944117144232\dots$	$\begin{array}{c} 11368365708750096478\\ 18990149600716232565\\ 77731144273048059639\\ 5907730326829\end{array}$	<u>0.367879441171442</u> 29
$\sqrt{2} = 1.414213562373095048\dots$	99936342969417150404 92969419440286922113 2671592712932741	$\underline{1.4142135623730950}34\dots$
$\sqrt{3} = 1.732050807568877\dots$	$\begin{array}{c} 68440491490666004609\\ 18767470146205475936\\ 30589653842274957065\\ 13562835250489629676\\ 61\end{array}$	1.732050807568862
$\frac{\sqrt{5}+1}{2} = 1.6180339887498948\dots$	13069649048652430795 14349754045887222621 81891430565948118397 38637429981	$1.6180339887498932\dots$
$\frac{\sqrt{5}-1}{2} = 0.61803398874989484\dots$	11381988572520783659 41813794511453008263 70298475149690990486 037	<u>0.618033988749894</u> 93
$\log 2 = 0.693147180 \dots$	76327808082936140771 2397	<u>0.6931471</u> 72
$\log 3 = 1.09861228866810969 \dots$	86435370621522915217 53627492119712964679 3034461	<u>1.098612288668109</u> 05

 ${\bf Table \ 3.} \ {\rm Numerical \ examples \ for \ Theorem \ 1}.$ 

ξ	p	$rac{\varphi(p+1)}{\varphi(p)}$	
$\frac{\pi}{10} = 0.314159265\dots$	1902037158772097	<u>0.3141592</u> 33	
$\frac{\pi}{20} = 0.1570796326794\dots$	65230510948153143551387418989	0.1570796169722	
$\frac{e}{10} = 0.2718281828\dots$	9240530296299581	0.2718281556	
$\frac{1}{e} = 0.367879441\dots$	5309646891817189	0.367879404	
$\frac{\sqrt{2}}{10} = 0.141421356\dots$	4002770936541226705231153047269	0.141421342	
$\sqrt{2} - 1 = 0.414213562\dots$	233570456771714761	0.414213520	
$\frac{\sqrt{3}}{10} = 0.173205080\dots$	721221963089661856995482309	0.173205063	
$\frac{\sqrt{5}}{10} = 0.223606797\dots$	1061017350953476949129	0.223606775	
$\frac{\sqrt{7}}{10} = 0.264575131\dots$	184295506315169	0.264575104	
$\frac{\sqrt{5}+1}{20} = 0.161803398\dots$	94721096130489558305686109	<u>0.1618033</u> 82	
The black New wind a second by The second			

 Table 4.
 Numerical examples for Theorem 2

ξ	p	$\frac{\varphi(p+1)}{\varphi(p)}$
$\frac{\pi}{10} = 0.314159265\dots$	1902037158772097	<u>0.3141592</u> 33
$\pi - 3 = 0.141592653$	37850921999916257860282849163969	0.141592639
$\frac{\pi}{20} = 0.15707963\dots$	2762774807373943331969	0.15707947
$\frac{e}{10} = 0.2718281828\dots$	9240621837106421	0.2718281556
$\frac{\sqrt{2}}{10} = 0.14142135623\dots$	4003599638847875898651948718589	0.14142134209
$\frac{\sqrt{3}}{10} = 0.173205080\dots$	721231627248456517343440289	0.173205063
$\frac{\sqrt{5}}{10} = 0.223606797\dots$	1061064248215841845709	0.223606775
$\frac{\sqrt{7}}{10} = 0.264575131\dots$	184327276293689	0.264575104
$\frac{\sqrt{5}+1}{20} = 0.161803398\dots$	94721096130489558305686109	0.161803382

**Table 5.** Numerical examples for Theorem  $\frac{3}{3}$ 

## PRIMITIVE ROOT BIAS FOR TWIN PRIMES II

ξ	p	$\frac{\sigma(p+1)}{\sigma(p-1)}$
$\gamma = 0.577215664901532$	2856597151653495728962024 3848158380563455580813926 3750257162232532026578053 941	<u>0.5772156649015</u> 27
$\pi = 3.1415\dots$	$\begin{array}{c} 2646921222256841983860921\\ 2640874686000298158238701\\ 0967174823412688455798495\\ 6967613160956654655736936\\ 7349279138922992270763544\\ 429 \end{array}$	<u>3.14</u> 07
$\frac{\pi}{2} = 1.57079632679489661\dots$	2714452215896122969174543 9828024978610834124808317 0868563698778989	$1.57079632679489675\dots$
e = 2.718281	2014265861114963900599828 5905237872952857257936121 6852839150744818500432120 229	<u>2.7182</u> 77
$\sqrt{2} = 1.414213562373095\dots$	$\begin{array}{c} 1300421337603424629894146\\ 7255162665877976721094905\\ 92798687340114429 \end{array}$	$1.414213562373089\dots$
$\sqrt{3} = 1.73205080756887729\dots$	5001304679832232346811636 9132922196818061901730529 1391240861141698770219064 8767132709	$\underline{1.732050807568877}44\dots$
$\sqrt{5} = 2.2360679774997\dots$	2013817027153422288458176 8517669808849964402993328 707829576024653820857989	<u>2.23606797749</u> 80
$\frac{\sqrt{5}+1}{2} = 1.618033988749894\dots$	$\begin{array}{c} 2768409745128994233528433\\ 3387767866165212510206810\\ 96425364422069\end{array}$	<u>1.6180339887498</u> 88
$\frac{\sqrt{5}-1}{2} = 0.6180339887498948\dots$	$\begin{array}{c} 1085428540928425422657480\\ 8906161529175312651234309\\ 7606604285214472315419401\\ 54141\end{array}$	<u>0.61803398874989</u> 54
$\log 2 = 0.6931471805599\dots$	$\begin{array}{c} 1583151329945483227597515\\ 3677620193994400861623554\\ 4504225821\end{array}$	<u>0.6931471805</u> 601
$\log 3 = 1.09861228866810969\dots$	$\begin{array}{c} 2122045296053250978265208\\ 8669266588247982830213119\\ 5312459828945357\end{array}$	<u>1.098612288668109</u> 89

 ${\bf Table \ 6.} \ {\rm Numerical \ examples \ for \ Theorem \ 4a}.$ 

ξ	p	$\frac{\sigma(p+1)}{\sigma(p)}$
$10\gamma = 5.77215\ldots$	8287834590808007955117965 6289238279189797435695514 9748729208974570049161227 89562389	<u>5.772</u> 21
$\pi = 3.1415926$	203351964077675489	3.1415957
$2\pi = 6.283185\ldots$	$\begin{array}{c} 6551931221328084483387020\\ 6608802113312867347600309\\ 2904430789475344820749138\\ 9591205917094426159824045\\ 6945706061680699198888379\\ 0296018540849009 \end{array}$	<u>6.2831</u> 91
$\frac{\pi}{2} = 1.57079632\dots$	22955076440560177	1.57079648
$e = 2.7182818\ldots$	1716574977543884369	2.7182821
$\sqrt{3} = 1.73205080\dots$	156513047792653	$\underline{1.732050}98$
$\sqrt{5} = 2.236067\dots$	4852141797161	2.236070
$\frac{\sqrt{5}+1}{2} = 1.6180339\dots$	18106083326748793	1.6180341

Table 7. Numerical examples for Theorem 4b.

ξ	p	$\frac{\sigma(p+1)}{\sigma(p)}$
$10\gamma = 5.7721566\dots$	$\begin{array}{c} 2595684868506848043313701\\ 5582183699928655989702069\\ 4446153232144556788787804\\ 2009056424590269\end{array}$	<u>5.77215</u> 72
$\pi = 3.14159265$	2007224256303311429	3.14159296
$2\pi = 6.28318530\dots$	$\begin{array}{c} 6561189247647575857894512\\ 4170120719766272845928619\\ 4871167020968826558631970\\ 9280421920002799868898085\\ 9337896045501356911046437\\ 15461617425356689 \end{array}$	<u>6.283185</u> 93
e = 2.7182818	1717018302510268229	<u>2.71828</u> 21
$\sqrt{5} = 2.2360679\dots$	285009842420045757101	<u>2.23606</u> 82

Table 8. Numerical examples for Theorem 4c.

### References

- [1] S.L. Aletheia-Zomlefer, L. Fukshansky, and S.R. Garcia. The Bateman-Horn conjecture: Heuristics, history, and applications. *Expo. Math.* (in press) https://arxiv.org/abs/1807.08899.
- [2] P.T. Bateman and R.A. Horn. A heuristic asymptotic formula concerning the distribution of prime numbers. *Math. Comp.*, 16:363–367, 1962.

- [3] P.T. Bateman and R.A. Horn. Primes represented by irreducible polynomials in one variable. In Proc. Sympos. Pure Math., Vol. VIII, pages 119–132. Amer. Math. Soc., Providence, R.I., 1965.
- [4] J.-R. Chen. On the representation of a larger even integer as the sum of a prime and the product of at most two primes. Sci. Sinica, 16:157–176, 1973.
- [5] J.-M. De Koninck and F. Luca. Analytic number theory, volume 134 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012. Exploring the anatomy of integers.
- [6] L.E. Dickson. A new extension of Dirichlet's theorem on prime numbers. Messenger of mathematics, 33:155–161, 1904.
- [7] P. Erdős. Some remarks on Euler's  $\varphi$  function. Acta Arith., 4:10–19, 1958.
- [8] P. Erdős, K. Győry, and Z. Papp. On some new properties of functions  $\sigma(n)$ ,  $\varphi(n)$ , d(n) and  $\nu(n)$ . Mat. Lapok, 28(1-3):125–131, 1980.
- J. Friedlander and H. Iwaniec. Opera de cribro, volume 57 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2010.
- [10] S.R. Garcia, E. Kahoro, and F. Luca. Primitive Root Bias for Twin Primes. Exp. Math., 28(2):151–160, 2019.
- [11] S.R. Garcia and F. Luca. On the difference in values of the Euler totient function near prime arguments. In *Irregularities in the distribution of prime numbers*, pages 69–96. Springer, Cham, 2018.
- [12] S.R. Garcia, F. Luca, and T. Schaaff. Primitive root biases for prime pairs I: Existence and non-totality of biases. J. Number Theory, 185:93–120, 2018.
- [13] G.H. Hardy and J.E. Littlewood. Some problems of 'Partitio numerorum'; III: On the expression of a number as a sum of primes. Acta Math., 114(3):215–273, 1923.
- [14] G.H. Hardy and E.M. Wright. An introduction to the theory of numbers. Oxford University Press, Oxford, sixth edition, 2008. Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles.
- [15] Chen J.-R. On the representation of a large even integer as the sum of a prime and the product of at most two primes. *Kexue Tongbao (Foreign Lang. Ed.)*, 17:385–386, 1966.
- [16] J. Maynard. Small gaps between primes. Ann. of Math. (2), 181(1):383-413, 2015.
- [17] F. Mertens. Ein Beitrag zur analytischen Zahlentheorie. J. Reine Angew. Math., 78:46–62, 1874.
- [18] D.H.J. Polymath. New equidistribution estimates of Zhang type. Algebra Number Theory, 8(9):2067–2199, 2014.
- [19] D.H.J. Polymath. Variants of the Selberg sieve, and bounded intervals containing many primes. Res. Math. Sci., 1:Art. 12, 83, 2014.
- [20] P. Ribenboim. The new book of prime number records. Springer-Verlag, New York, 1996.
- [21] J. Sándor, D.S. Mitrinović, and B. Crstici. Handbook of number theory. I. Springer, Dordrecht, 2006. Second printing of the 1996 original.
- [22] A. Schinzel. Generalisation of a theorem of B.S.K.R. Somayajulu on the Euler's function  $\varphi(n)$ . Ganita, 5:123–128 (1955), 1954.
- [23] A. Schinzel. Quelques théorèmes sur les fonctions  $\varphi(n)$  et  $\sigma(n)$ . Bull. Acad. Polon. Sci. Cl. III., 2:467–469 (1955), 1954.
- [24] A. Schinzel. On functions  $\varphi(n)$  and  $\sigma(n)$ . Bull. Acad. Polon. Sci. Cl. III., 3:415–419, 1955.
- [25] A. Schinzel and W. Sierpiński. Sur quelques propriétés des fonctions  $\varphi(n)$  et  $\sigma(n)$ . Bull. Acad. Polon. Sci. Cl. III., 2:463–466 (1955), 1954.
- [26] A. Schinzel and Y. Wang. A note on some properties of the functions  $\varphi(n)$ ,  $\sigma(n)$  and  $\theta(n)$ . Ann. Polon. Math., 4:201–213, 1958.
- [27] Y. Zhang. Bounded gaps between primes. Ann. of Math. (2), 179(3):1121-1174, 2014.

#### 16 STEPHAN RAMON GARCIA, FLORIAN LUCA, KYE SHI, AND GABE UDELL

Department of Mathematics, Pomona College, 610 N. College Ave., Claremont, CA 91711

*E-mail address*: stephan.garcia@pomona.edu *URL*: http://pages.pomona.edu/~sg064747

School of Mathematics, University of the Witwatersrand, Private Bag 3, Wits 2050, Johannesburg, South Africa, Research Group in Algebraic Structures and Applications, King Abdulaziz University, Jeddah, Saudi Arabia, Department of Mathematics, Faculty of Sciences, University of Ostrava, 30 dubna 22, 701 03 Ostrava 1, Czech Republic

*E-mail address*: Florian.Luca@wits.ac.za

Department of Mathematics, Harvey Mudd College, 340 East Foothill Boulevard, Claremont, CA 91711-5901

 $E\text{-}mail \ address: \ \texttt{kwshi@hmc.edu}$ 

Department of Mathematics, Pomona College, 610 N. College Ave., Claremont, CA 91711

*E-mail address*: grua2017@mymail.pomona.edu