BANACH SPACES OF ALMOST UNIVERSAL COMPLEMENTED DISPOSITION

JESÚS M .F. CASTILLO AND YOLANDA MORENO

ABSTRACT. We introduce and study the notion of space of almost universal complemented disposition (a.u.c.d.) as a generalization of Kadec space. We show that every Banach space with separable dual is isometrically contained as a 1-complemented subspace of a separable a.u.c.d. space and that all a.u.c.d. spaces with 1-FDD are isometric and contain isometric 1-complemented copies of every separable Banach space with 1-FDD. We then study spaces of universal complemented disposition (u.c.d.) and provide different constructions for such spaces. We also consider spaces of u.c.d. with respect to separable spaces.

1. INTRODUCTION

This paper can be considered a study of the properties that make the Kadec space [17] as it is: a separable Banach space containing complemented copies of every separable Banach space with the Bounded Approximation Property. This universal property, which apparently is a global property, is shown here to actually be a local property, called almost universal complemented disposition (a.u.c.d.) and shown to be very similar to Gurariy's almost universal disposition property [15]. To emphasize the knot between the Kadec and Gurariy spaces, let us briefly survey the history of (complementably) universal Banach spaces and spaces of universal (complemented) disposition.

The topic of Banach spaces of universal and almost universal disposition with respect to a class \mathcal{M} has its inception in the paper [15] of Gurariy, who constructed a separable space \mathcal{G} with the property that every into isometry from a finite dimensional space Finto \mathcal{G} can be extended to an into almost isometry $G \to \mathcal{G}$ on every finite dimensional superspace G of F. Several papers established the isometric uniqueness of \mathcal{G} [27], its maximality [34] and gave different descriptions for \mathcal{G} [28, 24, 37]. Gurariy also conjectured the existence of Banach spaces of universal disposition (without "almost") and of spaces of universal disposition with respect to the class \mathcal{S} of separable spaces. This conjecture was proved to be true in [1], where a general method to construct spaces of universal disposition with respect to different classes \mathcal{M} was presented. In particular, it was shown that the space that Gurariy conjectured is isometric to the Fraïssé limit, in the category of separable Banach spaces and into isometries, constructed by Kubis [22]. More recently, the papers [7] (resp. [5]) extend the method of [1] (resp. [22]) to the study of quasi-Banach (resp. Fréchet) spaces of (almost) universal disposition.

²⁰¹⁰ Mathematics Subject Classification. 46B03, 46M40.

This research has been supported in part by project MTM2016-76958-C2-1-P and Project IB16056 de la Junta de Extremadura.

Acknowledgement. The authors need to thank the truly remarkable job of the referee.

JESÚS M .F. CASTILLO AND YOLANDA MORENO

The topic of universal Banach spaces for a given class \mathcal{M} , i.e., Banach spaces in \mathcal{M} containing an isometric/isomorphic copy of every space in \mathcal{M} is another classical one (see the monograph of Dodos [12] to find updated information). The two germinal results are: the well known fact that C[0, 1] is isometrically universal for the class of separable spaces and Pełczyńsky's construction [33] of a space \mathcal{P} isomorphically universal for the class of Banach spaces with basis. Both results are relevant for the study in this paper. For instance, as the authors of [7] remark, the fact that a separable space of almost universal disposition is also separably universal (cf. Corollary 3.2 in this paper) depends, in principle, on the isometric uniqueness of the Gurariy space.

The topic of complementably universal space for a class \mathcal{M} , i.e., spaces in \mathcal{M} containing complemented copies of every space in \mathcal{M} , also contains many interesting results, sometimes requiring descriptive set theory techniques; see, for instance, [23, Theorem 1.2]. The topic emerges in 1969 when Pełczyński [33] shows that the space \mathcal{P} above mentioned is complementably universal for the class of Banach spaces with basis. In 1971 Kadec [17] obtains a complementably universal member \mathcal{K} for the class of separable Banach spaces with the Bounded Approximation Property (BAP); still in 1971 Pełczyński and Wojtaszczyk [34] prove that also the class of separably spaces with Finite Dimensional Decompositions has a complementably universal member \mathcal{PW} . The classical results of Pełczyński [32] (resp. Pełczyński-Wojtaszczyk) asserting that a separable Banach space has the BAP if and only if it is complemented in a space with basis (resp. FDD) implies that the spaces \mathcal{P}, \mathcal{K} and \mathcal{PW} contain complemented copies of all separable spaces with BAP. Kalton [19] performs a study of universal and complementably universal F-spaces, and remarks "there are a number of other existence and non-existence results known for other classes of separable spaces". It cannot go however without saying that Johnson and Szankowski [16] showed that no separable complementably universal space exists for the class of separable Banach spaces. A related topic is that of when a Banach space with a property P can be embedded into some Banach space with a finite dimensional decomposition and property P. See, e.g., [23, 30].

All spaces \mathcal{K} , \mathcal{P} and \mathcal{PW} are isomorphic (see Lemma 9.1). The isometric uniqueness of complementably universal spaces is a different thing. Garbulinska recovered in [13] the Fraïssé limit approach to construct first a complementably universal space \mathcal{G} for the class of separable spaces with FDD (thus isomorphic to the spaces of Kadec and Pełczyński) with a certain local isometric property. A closer inspection of the property that makes the space \mathcal{G} isometrically unique reveals that it is a local property, the one we have called almost universal complemented disposition and which is the object of study in this paper.

Sometimes the notation and results we present are rather technical. Thus, to encourage the reader and ease his way to the precise statements and proofs, let us briefly present intuitive versions of the main results in this paper. First of all, the key definition: A Banach space E will be called of *almost universal complemented disposition* (a.u.c.d. in short) if whenever one has an isometric embedding $F \to G$ between finite dimensional spaces with complemented range, every isometric embedding $F \to E$ with

complemented range admits an extension to an almost isometric embedding $G \to E$ with complemented range. Regarding separable a.u.c.d. spaces our main results are:

- There exist separable spaces of almost universal complemented disposition. In fact, every Banach space with separable dual can be isometrically embedded as a 1-complemented subspace of a separable a.u.c.d. space (Theorem 4.1).
- The separable spaces of a.u.c.d. are not unique. However, all separable spaces of a.u.c.d. with a 1-FDD are isometric (Proposition 5.3 and Theorem 5.4).
- Every separable space of a.u.c.d. with a 1-FDD contains isometric copies of every separable space and isometric 1-complemented copies of every separable space with a 1-FDD (Theorem 2.4).

The pièce of resistance of our analysis is the so-called Approximation Lemma 2.2 that roughly says that if E is Banach space of almost universal complemented disposition with a 1-FDD then every almost isometry $F \rightarrow E$ from a finite dimensional space F having almost complemented range can be approximated by an into isometry with complemented range.

Thus, since the Gurariy space \mathcal{G} is the only separable space of almost universal disposition while the Kadec space \mathcal{K} is the only separable space of almost universal complemented disposition with 1-FDD, these two spaces represent, in a sense, the same object in different categories. Indeed, if one moves from the category of Banach spaces and isometric embeddings to its "complemented" analogue, i.e., the category Banach spaces become the separable spaces with 1-FDD. It is clear than the "Gurariy objects" (i.e., the spaces of (almost) universal disposition) become the "Kadec objects" (the spaces of (almost) universal complemented disposition). Here it is the list of analogies:

- The Gurariy space:
 - (1) Is a space of almost universal disposition in the category of separable Banach spaces and single arrows (into isometries).
 - (2) It can be obtained as the Fraïssé limit of separable rational Banach spaces and single arrows.
 - (3) It can be constructed via an ω -times iterated push-out out from a countable dense set of single arrows between finite-dimensional Banach spaces.
 - (4) In the category, it is unique, up to isometries.
 - (5) It contains isometric copies of all separable Banach spaces.
 - (6) It is an \mathcal{L}_{∞} -space.
- The Kadec space:
 - (1) Is a space of almost universal disposition in the category of separable Banach spaces and double arrows (into isometries admitting norm one projections).
 - (2) It can be obtained as the Fraïssé limit of separable rational Banach spaces with 1-FDD and double arrows.
 - (3) It can be constructed via an ω -times iterated push-out out from a countable dense set of double arrows between finite-dimensional Banach spaces.
 - (4) In the category, it is unique, up to isometries.

- (5) It contains isometric complemented copies of all separable Banach spaces with 1-FDD. As a by-product, it contains isometric copies of all separable Banach spaces.
- (6) It is not an \mathcal{L}_{∞} -space.

In the second part of the paper we introduce and study the notions of space of universal complemented disposition (u.c.d.) and of space of universal complemented disposition with respect to separable spaces (ω -u.c.d.), their existence (Every Banach space can be isometrically embedded as a 1-complemented subspace of a space of (ω -)universal complemented disposition – cf. Propositions 6.2 and 7.2), universality and uniqueness properties.

The case of *p*-Banach spaces, 0 has been treated in a separate paper [8] with entirely different techniques.

2. Almost universal complemented disposition


All required technical results, definitions and constructions have been gathered in the Appendix section 9 at the end of the paper. The key notions for this paper are those of double-arrow and almost double arrow, that we state now. A $(1 + \varepsilon)$ -isometry is a linear continuous operator $f : A \to B$ such that for every $x \in A$ verifies

$$(1+\varepsilon)^{-1} ||x|| \le ||f(x)|| \le (1+\varepsilon) ||x||.$$

We will say that f is contractive if it verifies $(1 + \varepsilon)^{-1} ||x|| \le ||f(x)|| \le ||x||$.

Given $\alpha, \gamma > 1$ and $\beta \ge 0$ a (contractive) (α, β, γ) -arrow is a pair (f, \overline{f}) of linear continuous operators, $f : A \to B$ and $\overline{f} : B \to A$ in which f is a (contractive) α isometry, $\|\overline{f}\| \le \gamma$ and $\|\overline{f}f - 1_A\| < \beta$. To simplify some notation, (1, 0, 1)-arrows will be called double arrows, and pairs (f, \overline{f}) which are (α, β, γ) -arrows for suitable α, β, γ will be called almost double arrows and depicted as $(f, \overline{f}) : A \leftrightarrows B$.


Given three almost double arrows $(i_1, \overline{i_1}) : A \hookrightarrow C$, $(i_2, \overline{i_2}) : A \hookrightarrow B$ and $(i_3, \overline{i_3}) : B \hookrightarrow C$ we will say that the diagram they form



 ε -commutes if $||i_3i_2 - i_1|| \le \varepsilon$ and $||\overline{i}_2\overline{i}_3 - \overline{i}_1|| \le \varepsilon$. We will say it almost commutes if there exists $\varepsilon > 0$ such that the diagram ε -commutes. And we will say that it commutes if $i_3i_2 = i_1$ and $\overline{i}_2\overline{i}_3 = \overline{i}_1$.

Definition 2.1. A Banach space E will be called of almost universal complemented disposition (a.u.c.d., in short) if for every double arrow $(i, \overline{i}) : F \leftrightarrows G$ between finite dimensional spaces, every double arrow $(j, \overline{j}) : F \leftrightarrows E$ and every $\varepsilon > 0$ there exists a $(1 + \varepsilon, \varepsilon, 1)$ -double arrow $(J, \overline{J}) : G \leftrightarrows E$ making a commutative diagram

5



By Lemma 9.5, the condition is equivalent to the existence of a $(1+\varepsilon, \varepsilon, 1+\varepsilon)$ -double arrow $(J, \overline{J}) : G \leftrightarrows E$ making the diagram ε -commutative. This property essentially corresponds to property [E] of Garbulinska [13], although in that paper only the almost commutativity of injections is mentioned; the almost commutativity of projections is however used.

Our immediate purpose is to establish a key Approximation Lemma that will explain the structure of spaces of almost universal complemented disposition. To perturbate projections we will use Lemma 9.13, which is modeled upon [26, Thm. 1.a.9]. In order to give an estimate for the distance between projections, a proof is included there. Let us recall that we call a skeleton for E to a sequence (E_n) of finite-dimensional subspaces so that each E_n is 1-complemented in E_{n+1} and $E = \bigcup_n E_n$. See section 9.2 for details.

Lemma 2.2 (Approximation Lemma). Let $\varepsilon < 1/3$. If E is Banach space of almost universal complemented disposition admitting a skeleton then every $(1+\varepsilon,\varepsilon,1+\varepsilon)$ -arrow $(f,\overline{f}): F \hookrightarrow E$ with F finite-dimensional admits a (1,0,1)-arrow $(\phi,\overline{\phi}): F \hookrightarrow E$ at distance at most 72ε .

Proof. If (f, \overline{f}) is a $(1 + \nu, \nu, 1 + \nu)$ -arrow then, according to the estimate (3) in Lemma 9.6, $(f/(1 + \nu), \overline{f}/(1 + \nu))$ would be a contractive $(1 + \varepsilon, \varepsilon, 1)$ -arrow with $\varepsilon = 3\nu$. Thus, there is no loss of generality assuming that (f, \overline{f}) is a contractive $(1 + \varepsilon, \varepsilon, 1)$ -arrow.


i) Perturbation step. Since E has skeleton, it also has a sequence (E_n) of finite dimensional 1-complemented subspaces so that $E = \bigcup E_n$. Let $i_n : E_n \to E$ be the isometric embedding with 1-projection $\overline{i_n} : E \to E_n$. The perturbation arguments in Lemma 9.13 show that if one sets ε' so that $\varepsilon' \leq \varepsilon \frac{(1+\varepsilon)^2}{1-\varepsilon}$ and $1+\varepsilon' \leq \frac{1-\varepsilon^2}{1-3\varepsilon}$ then it is possible to find for $\varepsilon < 1/3$ and n large enough a $(1+\varepsilon')$ -isometry $\tau_{\varepsilon'} : f(F) \to F' \subset E_n$ so that

$$\|f - \tau_{\varepsilon'} f\| \le \varepsilon'$$

and a projection $p_{\varepsilon'}: E \to F'$ having norm at most $||p_{\varepsilon'}|| \leq 1 + \varepsilon'$ such that

(1)
$$\|p_{\varepsilon'} - \tau_{\varepsilon'} f \, \overline{f}\| \le \varepsilon'.$$

A diagram will help to understand the situation



It is clear that $\|\overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'}\| \leq \frac{1+\varepsilon'}{1-\varepsilon'} \leq 1+3\varepsilon' < 1+\varepsilon$ and it follows from the estimate (1) in Lemma 9.6 that $\tau_{\varepsilon'} f$ is a $(1+3\varepsilon)$ -isometry. Thus, $(\tau_{\varepsilon'} f, \overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'} i_n) : F \leftrightarrows E_n$ is a $(1+3\varepsilon, \varepsilon, 1+\varepsilon)$ -arrow and it follows from the estimate (3) in Lemma 9.6 that

$$(f_1, \overline{f_1}) = \left(\frac{\tau_{\varepsilon'} f}{1 + 3\varepsilon}, \frac{\overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'} i_n}{1 + 3\varepsilon'}\right)$$


is a contractive $(1 + 6\varepsilon, 6\varepsilon, 1)$ -arrow. Moreover, $||f - f_1|| \le 1 - \frac{1+\varepsilon'}{1+3\varepsilon} \le \frac{3\varepsilon+\varepsilon'}{1+3\varepsilon} \le 4\varepsilon$ and, taking into account the estimate (1) above, one gets

$$\|\overline{f} - \overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'}\| \le \|\overline{f} - \overline{f} \ \tau_{\varepsilon'}^{-1} (\tau_{\varepsilon'} f \ \overline{f} + p_{\varepsilon'} - \tau_{\varepsilon'} f \ \overline{f})\| = \|\overline{f} \ \tau_{\varepsilon'}^{-1} (p_{\varepsilon'} - \tau_{\varepsilon'} f \ \overline{f})\| + \varepsilon \le \frac{\varepsilon}{1 - \varepsilon'}$$

which gives

$$\begin{aligned} \|\overline{f} - \overline{f_1}\| &= \left(1 - \frac{1}{1 + 3\varepsilon'}\right) + \frac{1}{1 + 3\varepsilon'} \|\overline{f} - \overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'}\| \\ &\leq 3\varepsilon' + (1 + \varepsilon') \|\overline{f} - \overline{f} \ \tau_{\varepsilon'}^{-1} p_{\varepsilon'}\| \\ &\leq 3\varepsilon' + \frac{(1 + \varepsilon')\varepsilon}{1 - \varepsilon'} \\ &< 4\varepsilon. \end{aligned}$$


ii) Correction step. Apply the correction Lemma 9.12 to $(f_1, \overline{f_1}) : F \leftrightarrows E_n$ to get a 6ε -commutative diagram



in which (i_1, π_1) and (i_2, π_2) are (1, 0, 1)-arrows and moreover $\pi_1 i_2 = \overline{f_1}$ and $\pi_2 i_1 = f_1$.

7

iii) Almost universal complemented disposition step. Use now that E is of a.u.c.d. to get an ε' -commutative diagram



in which $(g_1, \overline{g_1}) : G_1 \hookrightarrow E$ is a $(1 + \varepsilon', \varepsilon', 1)$ -arrow that extends $(\iota_n, \overline{\iota_n})$. Thus, $(g_1 \iota_1, \pi_1 \overline{g_1}) : F \hookrightarrow E$ is a $(1 + \varepsilon', \varepsilon', 1)$ -arrow. Moreover,

 $\|g_1 i_1 - f\| \le \|g_1 i_1 - g_1 i_2 f_1\| + \|g_1 i_2 f_1 - f\| \le (1 + \varepsilon') 6\varepsilon + \varepsilon' + \|\imath_n f_1 - f\| \le 8\varepsilon + 4\varepsilon = 12\varepsilon$ And

 $\|\pi_1 \ \overline{g_1} - \overline{f}\| \le \|\pi_1 \ \overline{g_1} - \overline{f_1} \ \pi_2 \ \overline{g_1}\| + \|\overline{f_1} \ \pi_2 \ \overline{g_1} - \overline{f}\| \le 6\varepsilon + \|\overline{f_1} \ \pi_2 \ \overline{g_1} - \overline{f}\| \le 6\varepsilon + \varepsilon' + \|\overline{f_1} \ \overline{\imath_n} - \overline{f}\|.$ From where we get $\|(\pi_1 \overline{g_1} - \overline{f})|_{E_n}\| \le 12\varepsilon.$

We have thus obtained that each $(1 + \varepsilon, \varepsilon, 1 + \varepsilon)$ -arrow $F \cong E$ can be 36ε approximated by a $(1 + \varepsilon', \varepsilon', 1 + \varepsilon')$ -arrow for any $\varepsilon' \in (0, \varepsilon/3)$ on E_n for n large
enough.

iv) Ultraperturbation and iteration. Assume without loss of generality that $E_n = E_1$ in the first step, $E_n = E_2$ in the second step and so on. We have thus obtained a sequence $(f_n, \overline{f_n})$ of contractive $(1 + \varepsilon_n, \varepsilon_n, 1 + \varepsilon_n)$ -arrows such that $||(f_n, \overline{f_n}) - (f_{n+1}, \overline{f_{n+1}})|| \leq 36\varepsilon_n$ on E_n . Pick the sequence of ε_n monotone decreasing so that $\sum \varepsilon_n = \varepsilon$.

We use now a the ultraperturbation argument explained in Lemma 9.5, with a slight variation since this particular case is simpler. Pick a countably incomplete ultrafilter \mathcal{U} on \mathbb{N} . It is clear that $F_{\mathcal{U}} \cong (E_n)_{\mathcal{U}}$ is a (1,0,1)-arrow at distance 36ε of (f,\overline{f}) on the whole canonical copy of E inside $E_{\mathcal{U}}$. The point is that its image likely does not lie in E. We can use then principle of local reflexivity we can push-down this arrow back to E using the argument in Lemma : given the finite dimensional subspace $[f_n](F)$ pick an ε' -isometry $T_{\varepsilon'}: [f_n](F) \to E$ which is almost a projection and replace the embedding $[f_n]$ by $T_{\varepsilon'}[f_n]$ (see Lemma for details). The projection thus remains as it was while the inclusion $[f_n]$ is slightly perturbed with the ε' one prefers so that it takes values in E. Call $[f_n]'$ this perturbed inclusion to simplify. The new arrow $(u_n, \overline{u_n}) = ([f_n]', [\overline{f_n}]_{|E}) : F \cong E$ is a $(1 + \varepsilon', \varepsilon', 1)$ -arrow at distance at most $36\varepsilon + \varepsilon'$ of the original (f, \overline{f}) and $||(u_n, \overline{u_n}) - (u_{n+1}, \overline{u_{n+1}})|| \leq 36\varepsilon_n$ on E. Then, both (u_n) and $(\overline{u_n})$ are Cauchy sequences and thus they converge to a (1, 0, 1) arrow $(\phi, \overline{\phi})$ at distance 72ε from (f, \overline{f}) .

An immediate corollary from the Approximation Lemma is:

Proposition 2.3. A Banach space of almost universal complemented disposition with skeleton contains isometric 1-complemented copies of every finite-dimensional Banach space.

It may seem strange, but we do not know if this result can be obtained without the skeleton assumption. Observe that another reading of Approximation Lemma 2.2 is

that every finite dimensional subspace of a space of almost universal complemented disposition is contained in a finite dimensional 1-complemented subspace. Thus, the space has property π_1 . We obtain now one of the fundamental structural results:

Theorem 2.4. Every space of almost universal complemented disposition with skeleton contains isometric 1-complemented copies of every space with a skeleton.

Proof. Assume E is a space of almost universal complemented disposition with a 1-FDD (E_n) having canonical (1, 0, 1)-arrows $(i_n, \overline{i_n}) : E_n \leftrightarrows E$, and let Y be a space with a skeleton defined by the sequence of (1, 0, 1)-arrows $(\delta_n, \overline{\delta_n}) : Y_n \leftrightarrows Y_{n+1}$. Assuming without loss of generality that both Y_0 and E_0 are of dimension 1, pick a (1, 0, 1)-arrow $(f_0, \overline{f_0}) : Y_0 \leftrightarrows E_0$. Fix $\varepsilon = \sum \varepsilon_n$ with $(1 + \varepsilon_{n+1})^2 \leq 1 + \varepsilon_n$.

• Form first the push-out diagram as in Lemma 9.10:


which yields (1, 0, 1)-arrows $(\delta'_0, \overline{\delta'_0})$ and $(f'_0, \overline{f'_0})$ making the diagram ε_1 commutative (in fact, it is commutative in both directions; i.e., $\delta'_0 f_0 = f'_0 \delta_0$ and $\overline{\delta}_0 \overline{f'_0} = \overline{f}_0 \overline{\delta'_0}$).

• Inductive step. Assume that one has obtained an ε_n -commutative diagram

$$(f_n,\overline{f_n}) \bigvee_{E_n} \underbrace{ \overset{(\delta_n,\overline{\delta_n})}{\underbrace{\longleftarrow}} Y_{n+1}}_{(\delta'_n,\overline{\delta'_n})} P_{n+1}$$

in which $(f_n, \overline{f_n})$ is a contractive $(1 + \varepsilon_n, 0, 1 + \varepsilon_n)$ -arrow, $(\delta_n, \overline{\delta_n})$ and $(\delta'_n, \overline{\delta'_n})$ are (1, 0, 1)-arrows and $(f'_n, \overline{f'_n})$ is a contractive $(1 + \varepsilon_{n+1}, 0, 1 + \varepsilon_{n+1})$ -arrow

The a.u.c.d. disposition character of E yields a $(1 + \varepsilon_{n+1}, \varepsilon_{n+1}, 1)$ -arrow $(j_n, \overline{j_n}) : P_{n+1} \hookrightarrow E$ making a commutative diagram



Thus, $(j_n f'_n, \overline{f'_n}, \overline{j_n})$ is a $((1 + \varepsilon_{n+1})^2, \varepsilon_{n+1}, 1 + \varepsilon_n)$ -arrow. The approximation lemma 2.2 yields a (1, 0, 1)- arrow $(g_{n+1}, \overline{g_{n+1}}) : Y_{n+1} \hookrightarrow E$ at distance $72\varepsilon_n$ of $(j_n f'_n, \overline{f'_n}, \overline{j_n})$.

A small perturbation τ_{n+1} , as in Lemma 9.13, of $(g_{n+1}, \overline{g_{n+1}})$ yields a $(1 + \varepsilon_{n+2}, 0, 1 + \varepsilon_{n+2})$ arrow $(f_{n+1}, \overline{f_{n+1}}) : Y_{n+1} \hookrightarrow E_{n+1}$ (in which we set $f_{n+1} =$

(2)

 $\tau_{n+1}g_{n+1}$ and assume that the large E_{α} is E_{n+1} of course) at a distance ε_{n+2} . Form the push-out to get a commutative diagram

$$\begin{array}{cccc}
 & (\delta_{n+1},\overline{\delta_{n+1}}) \\
Y_{n+1} & & Y_{n+2} \\
(f_{n+1},\overline{f_{n+1}}) & & & & \downarrow \uparrow (f'_{n+1},\overline{f'_{n+1}}) \\
 & E_{n+1} & \underbrace{\longleftrightarrow}_{(\delta'_{n+1},\overline{\delta'_{n+1}})} P_{n+2} \\
\end{array}$$

According to diagram (7) one gets that $(\delta'_{n+1}, \overline{\delta'_{n+1}})$ and $(f'_{n+1}, \overline{f_n n + 1'})$ are both contractive $(1 + \varepsilon_{n+2}, 0, 1 + \varepsilon_{n+2})$ arrows. Induction will be over if the left downward arrow would be contractive. To make it so, we replace $(f_{n+1}, \overline{f_{n+1}})$ by $(\frac{1}{1+\varepsilon_{n+2}}f_{n+1}, (1+\varepsilon_{n+2})\overline{f_{n+1}})$, which is a contractive $((1+\varepsilon_{n+2})^2, 0, (1+\varepsilon_{n+2})^2)$ arrow (in particular a contractive $(1 + \varepsilon_{n+1}, 0, 1 + \varepsilon_{n+1})$ -arrow) and yields an ε_{n+1} -commutative diagram

This concludes the induction.

Relabel the left downwards arrow as $(f_{n+1}, \overline{f_{n+1}})$ to simplify notation. After this relabeling, we define now the (1, 0, 1) arrow $(f, \overline{f}) : Y \cong E$ we are looking for. Given $y \in Y$ so that $y = \lim_{n \to \infty} y_n$ with $y_n \in Y_n$ and $\sum ||y_{n+1} - y_n|| < +\infty$ then we set

$$f(y) = \lim_{n} f_n(y_n).$$

Since f_n is a $(1 + \varepsilon_n)$ -isometric embedding, whenever $\lim y_n = 0$ then $\lim f_n(y_n) = 0$ and thus f(y) does not depend on the choice of the sequence. To check that f is well defined observe that

$$\begin{aligned} \|f_{n|Y_{n-1}} - f_{n-1}\| &= \|\tau_n g_{n|Y_{n-1}} - f_{n-1}\| \\ &\leq \|(\tau_n g_n - g_n)|_{Y_{n-1}}\| + \|g_{n|Y_{n-1}} - f_{n-1}\| \\ &\leq \varepsilon_{n+1} + 72\varepsilon_{n-1} + \|f'_{n-1|Y_{n-1}} - f_{n-1}\| \\ &\leq \varepsilon_{n+1} + 72\varepsilon_{n-1} + \varepsilon_{n-1} \end{aligned}$$

and thus, with the proper choice of (ε_n) the sequence $(f_n(y_n))$ is Cauchy

$$\begin{aligned} \|f_{n+1}y_{n+1} - f_ny_n\| &= \|f_{n+1}y_{n+1} - f_{n+1}y_n + f_{n+1}y_n - f_ny_n\| \\ &\leq \|y_{n+1} - y_n\| + (\varepsilon_{n+1} + 73\varepsilon_{n-1})\|y_n\|. \end{aligned}$$

The map f is quite clearly an isometric embedding. We define the projection \overline{f} as follows

$$\overline{f}(e) = \lim_{n} f_n \overline{f}_n \overline{\iota}_n(e).$$

The operator \overline{f} is well defined: if $e = \lim e_n$ with $e_n \in E_n$ and $\sum ||e_{n+1} - e_n|| < +\infty$ then $\overline{f}(e) = \lim_n f_n \overline{f}_n(e_n)$. Observe that (with a slight abuse of notation)

$$\begin{aligned} \|\overline{f}_{n+1}\|_{E_n} - \overline{f}_n\| &\leq \|\overline{f}_{n+1} - \overline{g}_{n+1}\| + \|\overline{g}_{n+1} - \overline{f}'_n\| + \|\overline{f}'_n - \overline{f}_n\| \\ &\leq \varepsilon_{n+2} + 72\varepsilon_{n+1} + 2\varepsilon_n \end{aligned}$$

and thus one gets

$$\begin{aligned} \|f_{n+1}\overline{f}_{n+1}(e_{n+1}) - f_{n}\overline{f}_{n}(e_{n})\| &\leq \|f_{n+1}\overline{f}_{n+1}(e_{n+1}) - f_{n+1}\overline{f}_{n+1}(e_{n})\| \\ &+ \|f_{n+1}\overline{f}_{n+1}e_{n} - f_{n}\overline{f}_{n}e_{n}\| \\ &\leq (1 + \varepsilon_{n+1})^{2}\|e_{n+1} - e_{n}\| + \\ &+ \|f_{n+1}\|\|\overline{f}_{n+1}(e_{n}) - \overline{f}_{n}(e_{n})\| \\ &+ \|f_{n+1}|_{E_{n}} - f_{n}\|\|\overline{f}_{n}\|\|e_{n}\| \\ &\leq (1 + \varepsilon_{n+1})^{2}\|e_{n+1} - e_{n}\| \\ &+ (1 + \varepsilon_{n+1})\left(\varepsilon_{n+2} + 72\varepsilon_{n+1} + 2\varepsilon_{n}\right)\|e_{n}\| \\ &+ (\varepsilon_{n+1} + 72\varepsilon_{n} + 2\varepsilon_{n-1})\left(1 + \varepsilon_{n}\right)\|e_{n}\| \end{aligned}$$

and thus $(f_n \overline{f}_n(e_n))$ is a Cauchy sequence.

It remains to prove that for $y = \lim y_m \in Y$

$$\overline{f}(f(y)) = \lim_{m} f_n \overline{f}_n \overline{i}_n (\lim_{m} f_m y_m)) = \lim_{m} f_m y_m$$

as it immediately follows form the estimate:

$$\begin{aligned} \|\overline{f}_{n}(f_{m}(y_{m})) - y_{m}\| &\leq \|\overline{f}_{n}(f_{m}(y_{m})) - \overline{f}_{n}(f_{n}(y_{n}))\| + \|y_{n} - y_{m}\| \\ &\leq (1 + \varepsilon_{n})\|f_{m}y_{m} - f_{n}y_{n}\| + \|y_{n} - y_{m}\|. \end{aligned}$$


It is then clear that \overline{f} is a norm one projection.

In addition to the statement of Theorem 2.4, since C[0, 1] contains isometric copies of every separable Banach space and has skeleton, one gets:

Corollary 2.5. Every space of almost universal complemented disposition with skeleton contains isometric copies of every separable Banach space.

3. DIGRESSION ON BANACH SPACES OF ALMOST UNIVERSAL DISPOSITION

Recall (see e.g., [1, 3]) that a Banach space E is said to be of *almost universal* disposition if for every into isometry $i: F \to G$ between finite dimensional spaces, every into isometry $j: F \to E$ and every $\varepsilon > 0$ there exists a $(1 + \varepsilon)$ -isometry $J: G \to E$ making a commutative diagram



Let us show that the approximation lemma remains true in this context.

Proof. Let $f: F \to E$ be contractive a $(1 + \varepsilon)$ -isometry from a finite-dimensional space F into a space E of almost universal disposition. We apply the Correction Lemma 9.12 to the couple $f: F \to f(F)$ to find another space G = G(f, F, f(F)) and two isometries $i_F: F \to G$ and $j_F: f(F) \to G$ such that $||j_F f - i_F|| \leq \varepsilon$. If we call $\delta: f(F) \to E$ the canonical inclusion, the almost universal disposition property of E provides a $(1 + \varepsilon')$ -isometry $f_G: G \to E$ such that $f_G j_F = \delta$. Hence $f_G i_F: F \to E$ is a $(1 + \varepsilon')$ -isometry such that

$$\|f_G i_F - \delta f\| = \|f_G (i_F - j_F f + j_F f) - \delta f\| \le \varepsilon \|f_G\| + \|f_G j_F f - \delta f\| = \varepsilon (1 + \varepsilon').$$

Thus, every $(1 + \varepsilon)$ -isometry f admits a $(1 + \varepsilon/2)$ -isometry f_1 2ε -close. So f_1 admits a $(1 + \varepsilon/4)$ -isometry f_2 at distance at most ε and we thus obtain a sequence f_n of $(1 + \varepsilon/2^n)$ -isometries so that $||f_n - f_{n-1}|| \leq 3\varepsilon/2^{n-1}$. In particular, $(f_n(x))$ is a Cauchy sequence for every $x \in F$. The map $\phi : F \to E$ given by

$$\phi(x) = \lim f_n(x)$$

is an into isometry and $||f - \phi|| \leq 3\varepsilon$.

F. Cabello suggested to us that Lemma 3.1 could be true. An immediate (and well-known) consequence of Lemma 3.1 is:

Corollary 3.2. Every space of almost universal disposition contains isometric copies of all separable spaces.

Proof. Let X be a separable space, which we write as the closure of the union $X = \bigcup_{n=0}^{\infty} X_n$ of a sequence of finite-dimensional spaces X_n . Assume that X_0 is one dimensional. Let E be a space of almost universal disposition and thus, fixing $\varepsilon > 0$, any isometric embedding $f_0 : X_0 \to E$ can be extended to a $(1 + \varepsilon)$ -isometry $f'_0 : X_1 \to E$ which, by Lemma 3.1, admits an isometric embedding $f_1 : X_1 \to E$ at distance 2ε . Now it is f_1 which admits a $(1 + \varepsilon/2)$ -isometric extension $f'_1 : X_2 \to E$ which, by the lemma, admits an isometric embedding $f_2 : X_2 \to E$ at distance ε . Continue in this way and define $f : \bigcup_n X_n \to E$ as $f(x) = \lim f_n(x)$. This is an isometric embedding that extends to an isometric embedding $X \to E$ as desired.

The assertion in Corollary 3.2 was proved by Gurariy [15] for his space and by Gevorkyan [14] in full generality.

4. Construction of separable spaces of almost universal complemented disposition

We show now that the basic construction device as presented in [1, 3, 11], and used in [7], that provided a unified method to construct spaces of (almost) universal disposition, such as the Gurariy, Kubiś or the \mathcal{L}_{∞} -envelopes, can be adapted to construct separable spaces of almost universal complemented disposition.

Theorem 4.1. Every Banach space X with separable dual can be isometrically embedded as a 1-complemented subspace of a separable space $\mathcal{K}(X)$ of almost universal complemented disposition.

Proof. Let $\mathfrak{U} = \{(u, \overline{u}) : F_u \leftrightarrows G_u\}$ be a countable set of double arrows between finite dimensional spaces as in Lemma 9.15. The space F_u will be called the domain of uand G_u its codomain. We will call $dom\mathfrak{U}$ the set of the domains of the elements uso that $(u, \overline{u}) \in \mathfrak{U}$. For fixed $F_u \in dom\mathfrak{U}$ and X with separable dual, any subspace of $\mathfrak{L}(F_u, X) \oplus \mathfrak{L}(X, F_u)$ is separable. Thus, for $m \in \mathbb{N}$, let $L_m(F_u, X)$ be a countable dense subset of the space of all contractive $(1+2^{-m}, 0, 1+2^{-m})$ -arrows $F_u \leftrightarrows X$. Form now the countable set

$$D(E) = \bigcup_{u \in \mathfrak{U}} \bigcup_{m \in \mathbb{N}} L_m(F_u, E).$$

We start fixing an enumeration $\{d_{0,j}, j \in \mathbb{N}\}$ of D(X). To avoid ambiguities, the map $F_{d_{0,1}} \to G_{d_{0,1}}$, which should be called $d_{0,1}$ by the general convention above, will be called v_0 . Our first step is then to form the push-out

in which u_0 is an isometric embedding by Lemma 9.9. Observe that $P_1/X = G_{d_{0,1}}/F_{d_{0,1}}$ is finite-dimensional.

Assume now that spaces P_1, \ldots, P_n having separable dual have already been obtained together with into isometries $u_k : P_k \to P_{k+1}$ so that one can assume that P_k is a subspace of P_{k+1} and enumerations $\{d_{k,j}, j \in \mathbb{N}\}$ of $D(P_k)$ have also been fixed. Let us call $I_{n+1} = \{d_{i,j} : i+j \le n+1\}$ and form the push-out

$$\begin{array}{ccc} \ell_1(I_{n+1}, F_u) & \xrightarrow{\oplus u} & \ell_1(I_{n+1}, G_u) \\ \Sigma d & & \downarrow \\ P_n & \xrightarrow{u_n} & P_{n+1}. \end{array}$$

in which $\oplus u$ is the natural (into isometry) amalgamation of the maps $F_u \to G_u$ that appear involved in I_{n+1} and $\sum d$ is the (contractive) operator sum of the operators in I_{n+1} . This, again by Lemma 9.9, makes u_n an into isometry. Since $P_{n+1}/P_n = \ell_1(I_{n+1}, G_u)/\ell_1(I_n, F_u)$ is finite-dimensional, as well as P_1/X and X has separable dual, all P_n have separable dual and the process can be actually performed and the space $\mathcal{K}(X) = \overline{\cup}P_n$ is separable. Let us show that:

- $\mathcal{K}(X)$ contains an isometric 1-complemented copy of X.
- $\mathcal{K}(X)$ is a space of almost universal complemented disposition.

The first part follows from the "Moreover" part of lemma 9.10, which says that the $u_n : P_n \to P_{n+1}$ maps are actually part of certain (1, 0, 1)-arrows (u_n, U_n) ; which means that each P_n is 1-complemented in P_{n+1} and therefore X is 1-complemented in $\mathcal{K}(X)$.

To prove the almost universal complemented disposition of $\mathcal{K}(X)$, fix $0 < \varepsilon < 1/3$ and consider a double arrow $(\delta, \overline{\delta}) : F \leftrightarrows G$ between two finite dimensional spaces and a double arrow $(f, \overline{f}) : F \leftrightarrows \mathcal{K}(X)$. Assume that the subspace A = f(F) of $\mathcal{K}(X)$ is *n*-dimensional. A is complemented by a projection $f\overline{f}$ of norm C = 1. Pick $\{a_1, \ldots, a_n\}$ a basis for A so that $\operatorname{dist}(A, \ell_1^n)^{-1} \sum |\lambda_i| \leq \|\sum \lambda_i a_i\| \leq \sum |\lambda_i|$. Fix k large enough so that $2^{-k} < \varepsilon \operatorname{dist}(F, \ell_1^n)^{-1}$, there exist $x_i \in P_{n_k}, i \in \{1, \cdots, n\}$, such that $\|x_i - a_i\| \leq 2^{-k} \leq \frac{\varepsilon}{\operatorname{dist}(F, \ell_1^n)}$ and assume without loss of generality $P_k = P_{n_k}$.

By Lemma 9.13(1), the map $\tau a_i = x_i$ is a $(1 + \varepsilon)$ -isometry $A \to [x_i]$ and thus $\tau f: F \to P_k$ is a $(1+\varepsilon)$ -isometry with range $[x_i]$ and complemented via some projection p' of norm $\frac{1-\varepsilon^2}{1-3\varepsilon}$ for which $\|p' - \tau f\overline{f}\| \leq \frac{\varepsilon(1+\varepsilon)^2}{1-\varepsilon}$, besides $\|f - \tau f\| \leq \varepsilon$. Set $f_{\varepsilon} = \frac{\tau f}{1+\varepsilon}$ and $\overline{f_{\varepsilon}} = (1+\varepsilon)\overline{f}p'$ so that, again by Lemma 9.13(2), $(f_{\varepsilon}, \overline{f_{\varepsilon}}): F \leftrightarrows P_k$ is a contractive $(1+3\varepsilon, 0, \frac{1+3\varepsilon}{1-3\varepsilon})$ -arrow, $\|f - f_{\varepsilon}\| \leq 2\varepsilon$ and

$$\|\overline{f_{\varepsilon}} - f_{\varepsilon}\overline{f}\| \le 3\varepsilon$$

Using Lemma 9.15 we pick then $(u, \overline{u}) : F_u \simeq G_u$ in \mathfrak{U} for which there exist surjective contractive $(1 + 2^{-k})$ -isometries α, β so that the square

$$\begin{array}{ccc} F_u & \stackrel{u}{\longrightarrow} & G_u \\ \downarrow^{\alpha} & & \downarrow^{\beta} \\ F & \stackrel{}{\longrightarrow} & G \end{array}$$

is commutative in both directions, i.e., $\delta \alpha = \beta u$ and $\alpha \overline{u} = \overline{\delta} \beta$. Thus, $(f_{\varepsilon} \alpha, \alpha^{-1} \overline{f_{\varepsilon}})$: $F_u \leftrightarrows P_k$ is a contractive $(1+3\varepsilon, 0, \frac{1+3\varepsilon}{1-3\varepsilon})$ -arrow. Thus, some contractive $(1+3\varepsilon, 0, \frac{1+3\varepsilon}{1-3\varepsilon})$ arrow $(f', \overline{f'}) : F_u \leftrightarrows P_k$ at distance ε' (to be chosen) must exist in some set $L_m(F_u, P_k)$, which means that $(f', \overline{f'}) = d_{k,s}$ for some s. Since $d_{k,s} \in I_{k+s}$ we have that $(f', \overline{f'})$ is one of the elements forming the operator $\sum d$ that appears in the push-out diagram

According to Lemma 9.11, $(f', \overline{f'})$ admits a contractive $(1 + 3\varepsilon, 0, \frac{1+3\varepsilon}{1-3\varepsilon})$ -arrow extension $(f'', \overline{f''}) : G_u \cong P_{k+s+1}$. The composition $(f''\beta^{-1}, \beta \overline{f''})$ is a, say, $(1 + 7\varepsilon, 0, \frac{1+7\varepsilon}{1-7\varepsilon})$ -arrow such that

$$\|f''\beta^{-1}\delta - f\| = \|f''u\alpha^{-1} - f\|$$

$$= \|f'\alpha^{-1} - f\|$$

$$= \|(f' - f_{\varepsilon}\alpha + f_{\varepsilon}\alpha)\alpha^{-1} - f\|$$

$$\leq \varepsilon' + \|f_{\varepsilon} - f\|$$

$$\leq \varepsilon' + 2\varepsilon$$

and also

$$\| \left(\overline{\delta} \beta \ \overline{f''} - \overline{f} \right)_{|P_{s+k}} \| = \| \left(\alpha \ \overline{u} \ \overline{f''} - \overline{f} \right)_{|P_{s+k}} \| \\ = \| \left(\alpha \ \overline{f'} \overline{u_{k+s}} - \overline{f} \right)_{|P_{s+k}} \| \\ = \| \left(\alpha \left(\overline{f'} - \alpha^{-1} \overline{f_{\varepsilon}} + \alpha^{-1} \overline{f_{\varepsilon}} \right) \overline{u_{k+s}} - \overline{f} \right)_{|P_{s+k}} \| \\ \leq \varepsilon' + \| \left(\overline{f_{\varepsilon}} - \overline{f} \right)_{|P_{s+k+1}} \| \\ \leq \varepsilon' + 3\varepsilon.$$

We have thus obtained that

 $\forall \varepsilon > 0 \quad \exists n \in \mathbb{N} \ \exists (f_n, \overline{f}_n) : G \leftrightarrows P_n : \|f_n \delta - f\| \le 4\varepsilon; \quad \text{and} \quad \|\overline{\delta f}_n - \overline{f}_{|P_{n-1}}\| \le 4\varepsilon$

and $(f_n, \overline{f_n})$ is a $((1+7\varepsilon), 0, \frac{1+7\varepsilon}{1-7\varepsilon})$ -arrow. An ultraperturbation argument we sketch now is sufficient to conclude that for each ε there is a $((1+\varepsilon), 0, (1+\varepsilon))$ -arrow $G \leftrightarrows \mathcal{K}(X)$ making the diagram ε -commutative, which is condition i) in Lemma 9.5, and therefore $\mathcal{K}(X)$ is a space of almost universal complemented disposition.

Ultraperturbation argument: Observe that the problem lies in that the projection $\overline{f_m}$ behave well only on P_m . Inclusions behave well in the sense that once some f_n has been obtained then one can set $f_m = u_{m-1} \dots u_n f_n$. To get a good projection defined on the whole $\mathcal{K}(X)$ just define $[\overline{f_n}] : (\Pi P_n)_{\mathfrak{U}} \leftrightarrows G_{\mathfrak{U}} = G$ and compose with the diagonal canonical embedding $\mathcal{K}(X) \to \mathcal{K}(X)_{\mathfrak{U}}$.

5. Uniqueness

Definition 5.1. Given a Banach space X with separable dual we will denote $\mathcal{K}(X)$ the space constructed in Theorem 4.1.

We need a simple observation:

Lemma 5.2. If X has separable dual and skeleton then $\mathcal{K}(X)$ has skeleton.

Proof. Using the enumeration of Theorem 4.1 one gets that X is 1-complemented and has finite codimension in P_1 , and then P_n is 1-complemented and has finite codimension in P_{n+1} . Let us write $P_{n+1} = P_n \oplus C_n$ and $P_1 = X \oplus F$ with F finite-dimensional. If $(X_n)_n$ is a skeleton of X then $(X_n \oplus C_n)$ is a skeleton for $\mathcal{K}(X)$.

Thus, contrarily to what occurs with Gurariy space:

Proposition 5.3. There are non-isomorphic separable spaces of almost universal complemented disposition.

Proof. When X has not the BAP the space $\mathcal{K}(X)$ cannot have skeleton (it cannot have BAP) and thus it cannot be isomorphic to any space $\mathcal{K}(Y)$ constructed over a space Y with skeleton by virtue of the previous lemma

This marks a neat difference with the situation for separable spaces of almost universal disposition. Still, there is only one space of almost universal complemented disposition with skeleton, up to isomorphism: on one side the class of separable spaces

with BAP is closed under c_0 -sums, which means by Lemma 9.1 that there is only one complementably universal member, up to isomorphisms; since all spaces of complementably universal disposition with skeleton are complementably universal for the class of separable spaces with BAP, by Theorem 2.4, the assertion follows. Let us show that the space is unique, up to isometries

ameno

Theorem 5.4. Let U, V be two spaces of almost universal complemented disposition having a skeleton. let $i : A \to B$ be an isometry between two finite-dimensional 1complemented subspaces $A \subset U$ and $B \subset V$. For every $\varepsilon > 0$ there exists an isometry $\tau : U \to V$ such that $\|\tau_{|A} - i\| \leq \varepsilon$. In particular, all spaces of almost universal complemented disposition with skeleton are isometric.

Proof. The proof is a simple back-and-forth argument combining the Approximation Lemma 2.2 and the Perturbation argument of Lemma 9.13: let (U_n) (resp. (V_n)) be a skeleton for U (resp. V), so that $(u_n, \overline{u_n}) : U_n \hookrightarrow U$ and $(v_n, \overline{v_n}) : V_n \hookrightarrow V$ are double arrows.


Let $A \subset U$ be a 1-complemented subspace with embedding and projection $(\iota_A, \overline{\iota_A})$; and, analogously, $B \subset U$ be a 1-complemented subspace with embedding and projection $(\iota_B, \overline{\iota_B})$. Let $\iota : A \to B$ be an isometry between them. Set $\varepsilon = \sum \varepsilon_n$. After some ε' -perturbation of $(\iota_A, \overline{\iota_A})$ using Lemma 9.13 —which we do not relabel— we can assume that $[\iota_A(A) + U_1] \subset U_2$ (actually some n_2 , but again we do not relabel). This small perturbation we ignore by using the Approximation Lemma 2.2, so we still assume that that $(\iota, \iota^{-1}\overline{\iota_B})$ is a double arrow $A \leftrightarrows V$. By the a.u.c.d character of V this double arrow extends to some ε_1 -arrow $U_2 \leftrightarrows V$ that can therefore be ε_1 approximated by a double arrow: $(\iota_1, \overline{\iota_1}) : U_2 \leftrightarrows V$. Now repeat the argument back: we work with $(j_1, \overline{j_1}) = (\iota_1^{-1}, \iota_1\overline{u_2}) : \iota_1(U_2) \leftrightarrows U$. After some ε_2 -perturbation we assume that $\iota_1(U_2) \subset V_2$, use the Approximation Lemma 2.2 to not relabel, so that $(j_1, \overline{j_1})$ is a double arrow that the a.u.c.d. character of U allows one to extend to an ε_2 -arrow $V_2 \leftrightarrows U$ that can therefore be ε_2 -approximated by a double arrow: $(j_2, \overline{j_2}) : V_2 \leftrightarrows U$. And forth again. Iterate the argument.

This result should be compared to [13, Thm. 7.3]. We will (improperly) call Kadec space to $\mathcal{K}(\mathbb{R})$, the only (up to isometries) separable space of almost universal complemented disposition having skeleton. Which is of course complementably universal for all separable spaces with BAP. We say "improperly" because we cannot prove that the Kadec space \mathcal{K} constructed in [17] is of almost universal complemented disposition, although we know that it is isomorphic to $\mathcal{K}(\mathbb{R})$.

6. Spaces of universal complemented disposition

Spaces of universal disposition (i.e., the case $\varepsilon = 0$) were studied in [15, 1, 3]. In the same spirit, we have:

Definition 6.1. A Banach space E will be called of universal complemented disposition if given a double arrow $(i, \overline{i}) : F \leftrightarrows G$ between finite dimensional spaces and a double arrow $(j, \overline{j}) : F \leftrightarrows E$ there exists a double arrow $(J, \overline{J}) : G \leftrightarrows E$ making a commutative diagram.



The additional hypothesis of having separable dual is no longer required, and one gets:

Proposition 6.2. Every Banach space can be isometrically embedded as a 1-complemented subspace of a space of universal complemented disposition.

Proof. We will use the same device as for the construction of $\mathcal{K}(X)$, although everything is much simpler now since no correction lemmata or countable dense sets are required. The construction has now ω_1 steps. At step α , assuming P_{α} has been obtained, we get $P_{\alpha+1}$ as the push out in the diagram

(4)
$$\begin{array}{ccc} \ell_1(I_{\alpha}, F_u) & \xrightarrow{\oplus u} & \ell_1(I_{\alpha}, G_u) \\ \Sigma^d \downarrow & & \downarrow \\ P_{\alpha} & \xrightarrow{u_{\alpha}} & P_{\alpha+1}. \end{array}$$

Here I_{α} represents an index set containing: all (1, 0, 1)-arrows $d : F_u \cong P_{\alpha}$ from a finite dimensional space F_u into P_{α} each of them repeated as many times as (1, 0, 1)-arrows $u : F_u \cong G_u$ between finite dimensional spaces. The operator $\oplus u$ is the vector sum of all operators u and $\sum d$ the sum of all operators d. For $\alpha = 0$ set $P_{\alpha} = X$. If α is a limit ordinal then $P_{\alpha} = \overline{\bigcup_{\beta < \alpha} P_{\beta}}$.

The resulting space $\mathcal{K}_{\omega_1}(X)$ is of universal complemented disposition. Indeed, consider a double arrow $(\delta, \overline{\delta}) : F \leftrightarrows G$ between two finite dimensional spaces and a double arrow $(f, \overline{f}) : F \leftrightarrows \mathcal{K}_{\omega_1}(X)$. We choose $\alpha < \omega_1$ in such a way that f(F) is actually contained in P_{α} . So, $(f, \overline{f}_{|P_{\alpha}}) : F \leftrightarrows P_{\alpha}$ is one of the arrows d appearing in diagram (4) and can therefore be extended through any double arrow $F \leftrightarrows G$, in particular $(\delta, \overline{\delta})$ to a double arrow $G \leftrightarrows P_{\alpha+1}$. We have obtained now that

$$\exists \beta \ \forall \ \alpha \geq \beta \quad \exists (f_{\alpha}, \overline{f_{\alpha}}) \ : G \leftrightarrows P_{\alpha+1} : f_{\alpha}\delta = f \quad \text{and} \quad \overline{\delta} \ \overline{f_{\alpha}} = \overline{f_{P_{\alpha}}}$$

which a simple ultraperturbation argument transforms into a double arrow $G \cong \mathcal{K}_{\omega_1}(X)$ extending (f, \overline{f}) .


Definition 6.3. Given a Banach space X we will call $\mathcal{K}_{\omega_1}(X)$ the Banach space constructed in Theorem 6.2.

Proposition 6.4.


- (1) Under CH, if X has an ω -skeleton then the space $\mathcal{K}_{\omega_1}(X)$ has an ω -skeleton.
- (2) Under CH, if X has skeleton then $\mathcal{K}_{\omega_1}(X)$ has the BAP.

Proof. Under CH ($\mathfrak{c} = \aleph_1$) and thus a set of size \mathfrak{c} can be written as an increasing union of ω_1 countable sets. Now, the set we are considering is that of double arrows between two separable spaces A, B, which has the size of $\mathfrak{L}(A, B) \oplus \mathfrak{L}(B, A)$, namely $\mathfrak{c}^{\aleph_0} = \mathfrak{c}$. Let

 $(X_{\alpha})_{\alpha < \omega_1}$ be the ω -skeleton of X. Proceed as in the proof of Proposition 6.2 except that now we can arrange things so that all P_{α} are separable. This is done by representing each size \mathfrak{c} set I_{α} (since X_{α} is countable) as an increasing union $I_{\alpha} = \bigcup_{\mu < \omega_1} \Gamma_{\alpha,\mu}$ of countable sets and performing successive "diagonalizations" of those sets so that each push-out is done using only with a countable number of operators, so that the resulting push-out space is separable. Indeed, start with $P_0 = X_0$ and decompose the first set $I_0 = \bigcup_{\mu < \omega_1} \Gamma_{0,\mu}$ as an increasing union of countable sets. Make the first push-out only with the elements of $\Gamma_{0,1}$. The space P'_1 is thus a separable superspace of X_0 . Make a new push-out



to obtain a new separable enlargement of X_1 . Decompose now the set $I_1 = \bigcup_{\mu < \omega_1} \Gamma_{1,\mu}$ as an increasing union of countable sets $\Gamma_{1,\mu}$ and make now push-out only with the elements of $\Gamma_{0,2} \cup \Gamma_{1,1}$. The new space thus obtained P'_2 is a separable enlargement of P_1 . Assume now that a separable P'_{α} has already been obtained, make the new push-out



and write now $I_{\alpha} = \bigcup_{\mu < \omega_1} \Gamma_{\alpha,\mu}$ as the increasing union of countable sets $\Gamma_{\alpha,\mu}$ and make push-out only with the elements of $\bigcup_{i+j \le \alpha+1} \Gamma_{i,j}$. This yields a separable $P'_{\alpha+1}$. The skeleton of $\mathcal{K}_{\omega_1}(X)$ are the spaces $(P_{\alpha})_{\alpha < \omega_1}$.

To prove (2) we will actually show that the construction can be modified so that for each scountable α the push-out space P_{α} has a skeleton. Let us simplify the notation assuming that the space P_{α} has been obtained making push-out with the countable set I_{α} . Decompose I_{α} into an increasing sequence of finite sets $I_{\alpha} = \bigcup_m F_m$ and observe that P_{α} could have been obtained making just a sequence of iterated push outs starting with X: at step m make push-out with only the elements of F_m . Next, observe that the real content of Lemma 5.2 is that when X has skeleton then so does the space P_{ω} . Thus, P_{α} has skeleton. This immediately implies that $\mathcal{K}_{\omega_1}(X)$ has the BAP since any of its finite dimensional subspaces is contained into some P_{α} , and a Banach space such that any finite dimensional subspace is contained into a λ -complemented subspace with the λ -BAP must have the λ -BAP.

The general version of Theorem 2.4 becomes:

Proposition 6.5. A Banach space of universal complemented disposition that has a ω -skeleton formed by spaces with skeleton contains isometric 1-complemented copies of every Banach space with a skeleton.

Sketch of proof. Let Y be a space with skeleton and let E be a space of universal complemented disposition having a ω -skeleton of spaces E_{α} so that each E_{α} admits a skeleton $E_{\alpha,n}$. The uncountable cofinality of ω_1 will make he image of Y obtained be

lying in some of the separable spaces E_{α} ; and since these are 1-complemented in E the point is to obtain the copy of Y complemented in some E_{α} .

Let us proceed as in the proof of Theorem 2.4. A close examination of that proof reveals that the major part of the difficulties and the hard work in that proof was to get the image of Y complemented, something that could be done because E had skeleton: so one just needed to inductively add one by one the pieces of the skeleton to finally get a projection defined on all of E.

What has to be done now is to inductively add, one step each, finite dimensional pieces E_{α_n,m_n} with $E_{\alpha_i,m_i} \subset E_{\alpha_j,m_j}$ for i < j in such a way that $E_{\sup \alpha_n} = \overline{\bigcup_n E_{\alpha_n,m_n}}$. This would provide Y complemented in $E_{\sup \alpha_n}$ and the argument is complete.

The assumption "having a ω -skeleton" is necessary. To show this, let us consider a different way to obtain spaces of universal complemented disposition: Let Z be a space of almost universal complemented disposition and let \mathcal{U} be a countably incomplete ultrafilter on \mathbb{N} . The ultrapower $Z_{\mathcal{U}}$ is quite obviously a space of universal complemented disposition. In particular, one thus has:

Proposition 6.6. Let X be a dual separable Banach space. The space $\mathcal{K}(X)_{\mathfrak{U}}$ is a space of universal complemented disposition.

Recall that a Banach space X is said to have the Uniform Approximation Property (UAP) when every ultrapower of X has the BAP. It is clear that the UAP exactly means that X has the λ -AP and there exists a "control function" $f : \mathbb{N} \to \mathbb{N}$ so that, given F and $\lambda' > \lambda$, one can choose T such that $\operatorname{rank}(T) \leq f(\dim F)$ and Tf = f for all $f \in F$, with $||T|| \leq \lambda'$. Since X^{**} is complemented in some ultrapower of X, when X has the UAP then all even duals have the UAP. And since approximation properties pass from the dual to the space, when X has the UAP all its duals have the UAP. See [9, Section 7] for details. Therefore, Banach spaces with the BAP but whose duals do not have the BAP (see [9, Section 7]) admit ultrapowers without the BAP.

Regarding uniqueness, there are at least two (three under CH) non isomorphic spaces of universal complemented disposition:

Proposition 6.7.

- (1) The spaces $\mathfrak{K}(c_0)_{\mathfrak{U}}$ and $\mathfrak{K}_{\omega_1}(c_0)$ are not isomorphic.
- (2) Under CH, the spaces $\mathcal{K}(\mathbb{R})_{\mathfrak{U}}$, $\mathcal{K}_{\omega_1}(\mathbb{R})$ and $\mathcal{K}_{\omega_1}(X)$ for X a separable Banach space without BAP are not isomorphic

Proof. In [2] it was proved that infinite dimensional ultrapowers never contain complemented copies of c_0 , and thus $\mathcal{K}(X)_{\mathfrak{U}}$ cannot contain c_0 complemented. Since any copy of c_0 must be complemented in any space with ω -skeleton, thanks to Sobczyk's theorem, assertion (1) is clear. The space $\mathcal{K}(\mathbb{R})_{\mathfrak{U}}$ cannot have ω -skeleton nor the BAP: otherwise, every ultrapower $X_{\mathfrak{U}}$ of a separable space with BAP should have the BAP, which is false. The space $\mathcal{K}_{\omega_1}(\mathbb{R})$ has the BAP and ω -skeleton; and the space $\mathcal{K}_{\omega_1}(X)$ has ω -skeleton but not BAP. All this proves (2).

Regarding universality results, observe that

Corollary 6.8. No Banach space with ω -skeleton can be universal for the class of spaces with density character \aleph_1 .


Proof. As it has been said, every copy of c_0 must be complemented in a space with ω -skeleton; and thus, spaces with density character \aleph_1 but containing uncomplemented copies of c_0 cannot embed in a space with ω -skeleton.

Thus, there is no point in asking if a space of universal complemented disposition contains isometric copies of all spaces with density character at most \aleph_1 (since one must exclude those with ω -skeleton). It is quite curious that the spaces $K_{\mathfrak{U}}$ with K of almost universal complemented disposition contain, at least under CH, isometric copies of all spaces with density character \aleph_1 : indeed, K contains C[0, 1], hence $K_{\mathfrak{U}}$ contains $C[0, 1]_{\mathfrak{U}}$ which is, under CH, isomorphic to ℓ_{∞}/c_0 by [4, Proposition 2.4.1]; and this last space is universal for all spaces with density character \aleph_1 by Parovičenko's theorem ([6], [36, p.81]). A different thing is to ask if space of universal complemented disposition must contain isometric copies of all spaces with ω -skeleton. See Proposition 7.4 and Problem (6).

7. Spaces of universal complemented disposition for separable spaces

In the same way that the notion of space of universal disposition can be extended to "space of universal disposition with respect to the class of separable spaces", we can define:

Definition 7.1. A Banach space E will be called of ω -universal complemented disposition if given a double arrow $(i, \overline{i}) : S_1 \leftrightarrows S_2$ between separable spaces and a double arrow $(j, \overline{j}) : S_1 \leftrightarrows E$ there exists a double arrow $(J, \overline{J}) : S_2 \leftrightarrows E$ making a commutative diagram



One has:

Proposition 7.2. Every Banach space can be isometrically embedded as a 1-complemented subspace of a space of ω -universal complemented disposition.

The construction is immediate after that in Proposition 6.2 just replacing "finite dimensional" by "separable". Let us call $\mathcal{K}_{\omega_1}^{\mathfrak{S}}(X)$ the resulting space. It is of ω -universal complemented disposition exactly as in the proof of Proposition 6.2, which remains valid since no countable set is cofinal in ω_1 , and thus any operator from a separable space into $\mathcal{K}_{\omega_1}^{\mathfrak{S}}(X)$ actually has its image contained in some space P_{α} for some $\alpha < \omega_1$. The ω -version of Theorem 5.4 is:

Theorem 7.3. Let U, V be two spaces of ω -universal complemented disposition having a ω -skeleton. Let $i : A \to B$ be an isometry between two separable 1-complemented subspaces $A \subset U$ and $B \subset V$. There exists an isometry $\tau : U \to V$ such that $\tau_{|A} = i$. In particular, all spaces of ω -universal complemented disposition with ω -skeleton are isometric. *Proof.* The proof is much simpler than that of Theorem 5.4 since no approximation of perturbation is required: just a straightforward back-and-forth argument. \Box

Under CH, the space $\mathcal{K}_{\omega_1}^{\mathfrak{S}}(X)$ has an ω -skeleton when X has an ω -skeleton; and thus all the spaces of ω -universal complemented disposition with ω -skeleton are isometric to $\mathcal{K}_{\omega_1}^{\mathfrak{S}}(\mathbb{R})$. Let us call this unique space $\mathcal{K}_{\omega_1}^{\mathfrak{S}}$ from now on. Since X is 1-complemented in $K_{\omega_1}^{\mathfrak{S}}(X)$, under CH, $\mathcal{K}_{\omega_1}^{\mathfrak{S}}$ contains isometric 1-complemented copies of every Banach space with ω -skeleton; i.e.,

Proposition 7.4. Under CH, a space of ω -universal complemented disposition with ω -skeleton contains isometric 1-complemented copies of all spaces with ω -skeleton.

Observe that, even outside CH, spaces of ω -universal complemented disposition contain isometric 1-complemented copies of all separable spaces. It therefore follows from the Johnson-Szankowski theorem [16] that spaces of ω -universal complemented disposition must have density character at least \aleph_1 . Spaces of universal complemented disposition need not be of ω -universal complemented disposition: indeed, ultrapowers of spaces of almost universal disposition are of universal complemented disposition, although they cannot be of ω -universal complemented disposition since they cannot contain complemented copies of c_0 .

8. Open ends

We leave open a few questions which appeared during the course of this paper.

- (1) Is the Kadec space \mathcal{K} of [17] of almost universal complemented disposition. Equivalently, is it isometric to $\mathcal{K}(\mathbb{R})$?
- (2) Does a space of almost universal complemented disposition contain isometric 1-complemented copies of all finite-dimensional Banach spaces?
- (3) Does a separable space of almost universal complemented disposition contain isometric 1-complemented copies of all separable spaces with 1-FDD? Without separability assumption the answer is no. On the other hand, the spaces $\mathcal{K}(X)$ are 1-complementably universal for spaces with 1-FDD, regardless of whether they have or not skeleton, since $\mathcal{K}(X)$ contains a 1-complemented copy of $\mathcal{K}(\mathbb{R})$.
- (4) Do separable spaces of universal complemented disposition exist?
- (5) Is there a continuum of non-isomorphic spaces of universal complemented disposition? The corresponding question of the existence of many different spaces of universal disposition has been treated, although not completely solved, in [1] and [10].
- (6) Does a space of universal complemented disposition contain isometric copies of all spaces with ω -skeleton? Observe that a space of ω -universal complemented disposition contains isometric copies of all spaces with ω -skeleton.
- (7) Prus shows in [35, Thm. 2.1] that there is a reflexive separable space with basis that is complementably universal for all separable super-reflexive spaces with BAP. This suggest the possibility of obtaining other spaces of "almost universal complemented disposition with respect to certain subclasses of finite dimensional spaces" (see also [10, Prop. 3.1]). However, we cannot see how the methods in this paper could cover the reflexive case. See also [30]

9. Appendix: Basic constructions

We begin with the observation in the Introduction that the spaces of Pełcyński, Kadec and Wojtaszczyk are isomorphic. This is consequence of Pełczyński decomposition method.

Lemma 9.1. Let \mathcal{M} be a class of quasi-Banach spaces such that for some $0 it is closed under <math>\ell_p$ -sums. There is only one complementably universal member for \mathcal{M} , up to isomorphisms.

Proof. Let \mathcal{U} be a complementably universal member. The spaces \mathcal{U} and $\ell_p(\mathcal{U})$ contain complemented copies of each other, and $\ell_p(\ell_p(\mathcal{U})) \simeq \ell_p(\mathcal{U})$, hence $\mathcal{U} \simeq \ell_p(\mathcal{U})$. In particular, $\mathcal{U} \simeq \mathcal{U} \oplus \mathcal{U}$. Now, if A, B are two complementably universal members of \mathcal{M} , each of them contains a complemented copy of the other and both are isomorphic to their squares, so they are isomorphic. \Box

9.1. Arrows and double arrows.


Definition 9.2. We say that $f: A \to B$ is a $(1 + \varepsilon)$ -isometry if it is a linear continuous operator $f: A \to B$ such that for every $x \in A$ verifies $(1 + \varepsilon)^{-1} ||x|| \leq ||f(x)|| \leq (1+\varepsilon)||x||$. We will say that f is a contractive $(1+\varepsilon)$ -isometry if it is a linear continuous operator $f: A \to B$ such that for every $x \in A$ verifies $(1 + \varepsilon)^{-1} ||x|| \leq ||f(x)|| \leq ||x||$.

We define now "double arrows" (f, \overline{f}) between Banach spaces.

Definition 9.3. Given $\alpha, \gamma > 1$ and $\beta \ge 0$ a (contractive) (α, β, γ) -arrow is a pair (f, \overline{f}) of linear continuous operators, $f : A \to B$ and $\overline{f} : B \to A$ in which f is a (contractive) α -isometry, $\|\overline{f}\| \le \gamma$ and $\|\overline{f}f - 1_A\| < \beta$.

Throughout the paper, (1, 0, 1)-arrows have been called double arrows, and pairs (f, \overline{f}) which are (α, β, γ) -arrows for suitable α, β, γ have been called almost double arrows and depicted as $(f, \overline{f}) : A \simeq B$. The composition of two almost double arrows is $(f, \overline{f})(g, \overline{g}) = (fg, \overline{gf})$. The operator $\overline{f} : B \to A$ can be considered as a kind of "projection". This means that if one has a true projection $\pi : B \to B$ with range f(A) we will understand that the $\overline{f} : B \to A$ is $f^{-1}\pi$. When no confusion arises, given an (α, β, γ) -arrow (f, \overline{f}) we will simply say that \overline{f} is a β -projection along f of norm at most γ . To measure the commutativity of diagrams we will need a fourth parameter.

Definition 9.4. Let $(i_1, \overline{i_1}) : A \hookrightarrow C$, $(i_2, \overline{i_2}) : A \hookrightarrow B$ and $(i_3, \overline{i_3}) : B \hookrightarrow C$ be almost double arrows. We will say that the diagram they form



i) ε -commutes if $||i_3i_2 - i_1|| \le \varepsilon$ and $||\overline{i_2i_3} - \overline{i_1}|| \le \varepsilon$. ii) Almost commutes if there exists $\varepsilon > 0$ such that the diagram ε -commutes. iii) Commutes if $i_3i_2 = i_1$ and $\overline{i_2i_3} = \overline{i_1}$.

We present now a technique that allows one to pass from almost-commutative diagrams with bad projections to commutative diagrams with good projections. **Lemma 9.5** (Ultraperturbation lemma). Let X be a Banach space. Given a double arrow $(i, \overline{i}) : F \leftrightarrows G$ between finite dimensional spaces and a double arrow $(j, \overline{j}) : F \leftrightarrows X$, the following properties are equivalent:

- i) For every $\varepsilon > 0$ there exists a $(1 + \varepsilon, \varepsilon, 1 + \varepsilon)$ -double arrow $(J, \overline{J}) : G \leftrightarrows X$ making the diagram ε -commute.
- ii) For every $\varepsilon > 0$ there exists a $(1 + \varepsilon, \varepsilon, 1)$ -double arrow $(J, \overline{J}) : G \leftrightarrows X$ making the diagram commute.

Proof. It is clear that $ii \Rightarrow i$, so we only need to prove that $i \Rightarrow ii$. Consider a positive sequence (ε_n) with $\lim \varepsilon_n = 0$ and, by i), $(1 + \varepsilon_n, \varepsilon_n, 1 + \varepsilon_n)$ -double arrows $(J_n, \overline{J_n}) : G \leftrightarrows X$ making the diagram ε_n -commute. Take a non-trivial ultrafilter \mathcal{U} on \mathbb{N} and form the operators $[J_n] : G_{\mathcal{U}} \to X_{\mathcal{U}}$ and $[\overline{J_n}] : X_{\mathcal{U}} \to G_{\mathcal{U}}$. It turns out that $[J_n]$ is an into isometry and $[\overline{J_n}]$ a norm 1 projection through $[J_n]$, so $([J_n], [\overline{J_n}]) : G_{\mathcal{U}} \leftrightarrows X_{\mathcal{U}}$ is a double arrow. And if $([i], [\overline{i}]) : F_{\mathcal{U}} \leftrightarrows G_{\mathcal{U}}$ and $([j], [\overline{j}]) : F_{\mathcal{U}} \hookrightarrow X_{\mathcal{U}}$ are the natural double arrows, the diagram is commutative since $[J_n][i] = [j]$ and $[\overline{i}][\overline{J_n}] = [\overline{j}]$.

Since F and G are finite dimensional spaces then $F = F_{\mathcal{U}}$ and $G = G_{\mathcal{U}}$. So $[J_n](G_{\mathcal{U}})$ is finite dimensional and we can choose a $\varepsilon/2$ -net $f_1, ..., f_N$ in the dual unit ball $B_{[J_n](G)^*}$, which we can assume to be in the dual unit ball of $(X_{\mathcal{U}})^*$, such that for every $g \in G$,

$$|\langle [J_n]g, f_k \rangle| \ge (1+\varepsilon)^{-1} ||[J_n]g||, \text{ for some } k.$$

Now, observe that the key feature behind the Principle of Local Reflexivity of Lindenstrauss and Rosenthal [25] is the notion of local complementation identified by Kalton [18], as it appears implicitly in [29] and explicitly in [31]. We will consider from now on a Banach space X isometrically embedded into its ultrapower $X_{\mathcal{U}}$ via the map $x \to [x]$. In this form, X is locally complemented in $X_{\mathcal{U}}$. Thus, once the functionals f_1, \ldots, f_N are set, given $\varepsilon > 0$ there is an operator $T_{\varepsilon} : [J_n](G) \to X$ such that

(1) $||T_{\varepsilon}|| \leq 1 + \varepsilon$

(2)
$$(T_{\varepsilon})_{|[J_n](G)\cup X} = 1_{[J_n](G)\cap X}$$

(3)
$$(T_{\varepsilon}^* f_k)_{|[J_n](G)} = (f_k)_{|[J_n](G)}$$

Therefore, the map $T_{\varepsilon}[J_n]$ is a $1 + \varepsilon$ -isometry since $||T_{\varepsilon}[J_n](g)|| \leq (1 + \varepsilon)||g||$ and

$$|T_{\varepsilon}[J_n](g)|| \geq | < T_{\varepsilon}[J_n](g), f_k > |$$

$$= | < [J_n](g), T_{\varepsilon}^* f_k > |$$

$$= | < [J_n](g), f_k > |$$

$$\geq (1 + \varepsilon)^{-1} || [J_n](g) ||$$

On the other hand, the norm 1 projection we need is $[\overline{J_n}]_{|X}$. The couple $(T_{\varepsilon}[J_n], [\overline{J_n}]_{|X}) : G \leftrightarrows X$ is a $(1 + \varepsilon, \varepsilon, 1)$ -double arrow since, for suitably chosen f_k one has

$$\begin{aligned} \|[\overline{J_n}]T_{\varepsilon}[J_n][g] - [g]\| &= \|[\overline{J_n}]T_{\varepsilon}[J_n][g] - [\overline{J_n}J_ng]\| \\ &= \|[\overline{J_n}T_{\varepsilon}[J_n][g]] - [\overline{J_n}J_ng]\| \\ &= \|[T_{\varepsilon}[J_n][g]] - [J_ng]\| \\ &\leq \langle [T_{\varepsilon}[J_n][g]] - [J_ng], f_k \rangle + \varepsilon \\ &= \varepsilon. \end{aligned}$$

The diagram commutes since $T_{\varepsilon}[J_n][i] = T_{\varepsilon}[j] = j$ and $\overline{i}[\overline{J_n}]|_X = [\overline{iJ_n}]|_X = \overline{j}|_X = \overline{j}$. \Box

Observe that adding "contractive" to the hypothesis does not improve the results. We conclude this section with a set of elementary estimates that will be useful later.

Lemma 9.6.

- (1) If f is a (contractive) $(1+\varepsilon)$ isometry and τ is a (contractive) $(1+\varepsilon')$ -isometry then τf is a (contractive) $(1+\varepsilon)(1+\varepsilon')$ -isometry.
- (2) If f is a $(1+\varepsilon)$ -isometry then $\frac{1}{1+\varepsilon}f$ is a contractive $(1+\varepsilon)^2$ -isometry.
- (3) If (f,\overline{f}) is an (α,β,γ) -arrow then $(\frac{1}{\alpha}f,\frac{1}{\gamma}\overline{f})$ is a contractive $(\alpha^2,\frac{\beta+\gamma\alpha-1}{\gamma\alpha},1)$ arrow.
- (4) If (f, \overline{f}) is a (contractive) $(1+\varepsilon, \varepsilon, 1)$ -arrow then $(f, (\overline{f} \ f)^{-1}\overline{f})$ is a (contractive) $(1+\varepsilon, 0, 1+\frac{\varepsilon}{1-\varepsilon})$ arrow. Moreover, $\|\overline{f} (\overline{f} \ f)^{-1}\overline{f}\| \leq \frac{\varepsilon}{1-\varepsilon}$

Proof. Probably only assertion (4) requires some explanation. Since $||1 - \overline{f} f|| \le \varepsilon < 1$ then $1 - (1 - \overline{f} f) = \overline{f} f$ is invertible and its inverse has norm at most $1 + \varepsilon + \varepsilon^2 + \cdots = \frac{1}{1-\varepsilon} = 1 + \frac{\varepsilon}{1-\varepsilon}$. Then $(\overline{f} f)^{-1}$ exists and $(\overline{f} f)^{-1}p$ is a true projection along f since $(\overline{f} f)^{-1}\overline{f} f = 1$. Finally

$$\|\overline{f} - (\overline{f} f)^{-1}\overline{f}\| = \|\overline{f} f f^{-1} - (\overline{f} f)^{-1}\overline{f} f f^{-1}\| \le \|\overline{f} f - 1\|\|f^{-1}\| \le \frac{\varepsilon}{1-\varepsilon}.$$

9.2. Skeletons. Different approximation notions are essential in the theory of spaces of complemented disposition. A Banach space X is said to have the λ -approximation property (λ -BAP in short) if for each finite dimensional subspace $F \subset X$ and every $\lambda' > \lambda$ there is a finite-rank operator $T: X \to X$ such that $||T|| \leq \lambda'$ and T(f) = ffor each $f \in F$. This is not the standard definition, but it is an equivalent formulation (see [9, Theorem 3.3]). The space is said to have the Bounded Approximation Property (BAP in short) if it enjoys the λ -BAP for some λ . A μ -complemented subspace of a space with the λ -BAP has the $\lambda\mu$ -BAP. When X is separable, the λ -BAP is equivalent to the existence of a sequence $B_n: X \to X$ of linear finite-dimensional operators with norms $||B_n|| \leq \lambda$ that is pointwise convergent to the identity. This sequence of operators can be asked to verify $B_m B_n = B_n$ for m > n. The sequence is called a a Finite Dimensional Decomposition (FDD, in short) if, moreover, for every $m, n \in \mathbb{N}$, $B_n B_m = B_{\min\{m,n\}}$. By a well-known result of Pełczyński [32], spaces complementably universal for spaces with FDD are also complementably universal for separable spaces with the BAP.

An essential part in our arguments and in the classification of spaces of (almost) universal complemented disposition is played by the notion of *skeleton* which, as we will show next, coincides with that of 1-Finite Dimensional Decomposition, although the skeleton formulation is more adapted to the problems treated in this paper:

Definition 9.7. We say that a Banach space E admits a *skeleton* if there exists a sequence (E_n) of finite-dimensional subspaces and of double arrows $(\delta_n, \overline{\delta_n}) : E_n \cong E_{n+1}$ so that $E = \overline{\bigcup_n E_n}$. We will say that E admits a ω -skeleton if there is a continuous

chain $(E_{\alpha})_{\alpha < \omega_1}$ of separable subspaces and double arrows $(\delta_{\alpha}, \overline{\delta_{\alpha}}) : E_{\alpha} \leftrightarrows E_{\alpha+1}$. Here *continuous* means that for every limit ordinal β one has $E_{\beta} = \overline{\bigcup_{\alpha < \beta} E_{\alpha}}$.

In each case we shall say that $(\delta_{\alpha}, \overline{\delta_{\alpha}})$ is the family of double arrows defining the (ω) skeleton. Of course that spaces admitting a skeleton must be separable and spaces admitting a ω -skeleton must have density character at most \aleph_1 .

Lemma 9.8.

- A Banach space has a skeleton if and only if it has a 1-FDD.
- A Banach space has an ω-skeleton if and only if it is a 1-Plichko space with density character at most ℵ₁.

Proof. Assume that a Banach space E has a skeleton $(\delta_n, \overline{\delta_n}) : E_n \leftrightarrows E_{n+1}$. The spaces E_k are 1-complemented in E since one can define norm one projections $P_k : E \to E_k$ as follows: if $x \in \bigcup E_n$ and $x \in E_{n+1}$ then set $P_k(x) = \overline{\delta_k} \dots \overline{\delta_{n-1}\delta_n}(x)$ and extend P_k to E by density. Notice that if n + 1 < k, then $P_k(x) = x$. These projections verify $\lim P_k(x) = x$. Thus, spaces with skeleton have the π_1 -property [9, Def.5.1]; i.e., there is a net of finite rank norm one projections pointwise convergent to the identity. The π_1 property in a separable space implies 1-FDD [9, Prop.5.4]. It is clear that spaces with 1-FDD have a skeleton. The second part can be found in [20, Section 6].

 ω -skeletons will only appear in the final Sections 8 and 9, where we will maintain the name by coherence with the rest of the paper and because statements are shorter this way. The first assertion in Lemma 9.8 appears used in [13]. Kubiś [21] and other authors have given more general notions of *projectional skeleton* by considering a partially ordered index space.

9.3. Push-out constructions.

9.3.1. The push-out. Given operators $i: Y \to A$ and $j: Y \to B$, the associated push-out diagram is

(5)
$$\begin{array}{cccc} Y & \stackrel{i}{\longrightarrow} & A \\ j & & \downarrow j' \\ B & \stackrel{i'}{\longrightarrow} & PO \end{array}$$

Here, the push-out space PO = PO(i, j) is the quotient of the direct sum $A \oplus_1 B$, the product space endowed with the sum norm, by the closure of the subspace $\Delta = \{(iy, -jy) : y \in Y\}$. We will call $Q : A \oplus_1 B \to (A \oplus_1 B)/\overline{\Delta}$, the natural quotient map. The map i' is given by the inclusion of B into $A \oplus_1 B$ followed by Q, so that $i'(b) = (0, b) + \overline{\Delta}$ and, analogously, $j'(a) = (a, 0) + \overline{\Delta}$.

The diagram (5) is commutative: j'i = i'j. Moreover, it is 'minimal' in the sense of having the following universal property: if $j'': A \to C$ and $i'': B \to C$ are operators such that j''i = i''j, then there is a unique operator $\gamma: PO \to C$ such that $i'' = \gamma i'$ and $j'' = \gamma j'$. Clearly, $\gamma((a, b) + \overline{\Delta}) = j''(a) + i''(b)$ and one has $\|\gamma\| \leq \max\{\|i''\|, \|j''\|\}$. Regarding the behaviour of the maps in diagram (5) one has (see [3, Lemma A.19] for details):

Lemma 9.9.

- (a) If i is an isomorphic embedding, then Δ is closed.
- (b) The norm of the operators i' and j' is less than or equal to one.
- (c) If i is an isometric embedding and $||j|| \leq 1$ then i' is an isometric embedding.
- (d) If i is an isomorphic embedding then i' is an isomorphic embedding.
- (e) If $||j|| \leq 1$ and i is an isomorphism then i' is an isomorphism and

$$||(i')^{-1}|| \le \max\{1, ||i^{-1}||\}.$$

9.3.2. The almost-complemented push-out. We establish now that the push-out construction can be adapted to cover the case of ε -projections.

Lemma 9.10. Given almost double arrows $(i, \overline{i}) : A \cong B$ and $(j, \overline{j}) : A \cong X$ there is a commutative diagram

(6)

$$A \xrightarrow{(i,i)} B$$

$$(j,\overline{j}) \bigwedge \bigwedge \bigwedge (j',\overline{j'})$$

$$X \xrightarrow{(i',\overline{i'})} PO$$

so that if (i,\overline{i}) is an $(\alpha,0,\gamma)$ -arrow and (j,\overline{j}) is a (u,v,w)-arrow then $(i',\overline{i'})$ is a contractive $(\alpha u, 0, u\gamma)$ -arrow and $(j',\overline{j'})$ is a contractive $(u\alpha, \alpha v\gamma, \max\{\alpha w, 1 + \alpha v\gamma\})$ -arrow. Moreover (compare with Lemma 9.9 (c) above) if (i,\overline{i}) is a (1,0,1)-arrow, and (j,\overline{j}) is a contractive (u,v,w)-arrow then $(i',\overline{i'})$ is a (1,0,1)-arrow and $(j',\overline{j'})$ is a contractive $(u,v,\max\{w, 1+v\})$ -arrow.

All this can be depicted for mnemonical reasons as

(7)
$$(u,v,w) \bigvee_{\cdot \leftarrow \leftarrow \leftarrow \leftarrow}^{(\alpha,0,\gamma)} \cdot (u,v,w) \bigvee_{\cdot \leftarrow \leftarrow \leftarrow \leftarrow}^{(\alpha,0,\gamma)} \cdot (u\alpha, \alpha v\gamma, \max\{w\alpha, 1+\alpha v\gamma\})$$

and

(8)
$$\begin{array}{c} \underbrace{(1,0,1)}_{\text{contractive }(u,v,w)} \downarrow \uparrow \underbrace{(1,0,1)}_{\text{contractive }(u,v,\max\{w,1+v\})} \\ \cdot \underbrace{(1,0,1)}_{\text{contractive }(u,v,\max\{w,1+v\})} \end{array}$$

Proof. To obtain $\overline{j'}$ observe that the diagram

(9)
$$\begin{array}{ccc} A & \stackrel{i}{\longrightarrow} & B \\ i \downarrow & & \downarrow^{1_B + i(\overline{j}j - 1_A)\overline{i}} \\ X & \stackrel{i}{\longrightarrow} & B \end{array}$$

is commutative, and thus the universal property of the push-out yields the existence of a unique operator $\overline{j'}: \text{PO} \to B$ such that

(3.a) $\underline{j'} i' = i \, \overline{j};$ (3.b) $\underline{j'} j' = 1_B + i(\overline{j} \, j - 1_A) \, \overline{i};$ (3.c) $\|\overline{j'}\| \le \max\{\|i\overline{j}\|, \|1_B + i(\overline{j} \, j - 1_A) \, \overline{i}\|\}.$

Notice that by properties of the push-out construction, $||i'|| \leq 1$ and $||j'|| \leq 1$ independently of the norms of *i* and *j*. To estimate the norm of their inverse maps observe that for every $x \in X$,

$$\begin{aligned} \|x\| &\leq \inf_{a \in A} \{ \|x - ja\| + \|ja\| \} \\ &\leq \inf_{a \in A} \{ \|x - ja\| + u\alpha \|ia\| \} \\ &\leq u\alpha \|i'(x)\|_{PO}; \end{aligned}$$

thus $(u\alpha)^{-1}||x|| \le ||i'(x)|| \le ||x||$. Except when *i* is an into isometry and $||j|| \le 1$, in which case

$$||x|| = \inf_{a \in A} \{ ||x - ja|| + ||ja|| \}$$

$$\leq \inf_{a \in A} \{ ||x - ja|| + ||ia|| \}$$

$$= ||i'(x)||_{PO},$$

and thus $||x|| = ||i'(x)||_{PO}$. In the same way, for every $b \in B$,

$$\begin{aligned} \|b\| &\leq \inf_{a \in A} \{ \|b + ia\| + \|ia\| \} \\ &\leq \inf_{a \in A} \{ \|b + ia\| + \alpha u \|ja\| \}, \end{aligned}$$

and thus $(u\alpha)^{-1} ||b|| \le ||j'(b)||_{PO} \le ||b||$. To obtain $\overline{i'}$, since the diagram

(10)
$$\begin{array}{ccc} A & \stackrel{i}{\longrightarrow} & B \\ j \downarrow & & \downarrow j\overline{i} \\ X & \stackrel{i}{\longrightarrow} & X, \end{array}$$

is commutative, the universal property of the push-out yields a unique operator $\overline{i'}:\mathrm{PO}\to X$ such that

(4.a) $\overline{i'} i' = 1_X;$ (4.b) $\overline{i'} j' = j \overline{i};$ (4.c) $\|\overline{i'}\| \le \max\{\|1_X\|, \|j \overline{i}\|\}.$

Let us check that the just defined projection $\overline{i'}$ and ε -projection $\overline{j'}$ make commutative the original diagram (6). To this end, it is enough to observe that since diagram

(11)
$$\begin{array}{ccc} A & & & \\ & \downarrow^{i} & & \downarrow^{\overline{j}j\overline{i}} \\ X & & & \\ & & & \overline{j} & A. \end{array}$$

is commutative, the universal property of the push-out yields a unique operator $\gamma:\mathrm{PO}\to A$ such that

- (5.a) $\gamma i' = \overline{j}$ (5.b) $\gamma j' = \overline{j} j \overline{i}$.
- (5.c) $\|\gamma\| \le \max\{\|\overline{j}\|, \|\overline{j}\ j\ \overline{i}\|\}.$

Since $\overline{j} \ \overline{i'}i' = \overline{j}$ and $\overline{j} \ \overline{i'} \ j' = \overline{j}j\overline{i}$ (by (4.b)), the uniqueness (see (5.a) and (5.b)) yields $\gamma = \overline{j} \ \overline{i'}$. On the other hand, also $\gamma = \overline{i} \ \overline{j'}$ since $\overline{i} \ \overline{j'} \ i' = \overline{j}$ (by (3.a)) and $\overline{i} \ \overline{j'}j' = \overline{i}(1_B + i(\overline{j}j - 1_A)\overline{i}) = \overline{i} + (\overline{j} \ j - 1_A)\overline{i} = \overline{j} \ \overline{j} \ \overline{i}$.

Modifying the proof above in an obvious way we obtain the result of Kubis [22, Section 5] (see also [13, Lemma 4.1] and the comments before the lemma) that in a push-out diagram

$$\begin{array}{ccc} A & & & \\ & & i \end{array} & & & \\ i & & & & \downarrow i' \\ C & & & & PO \end{array}$$

in which both i, j have complemented ranges via projections p, q then also i', j' have complemented ranges via projections p', q' yielding a diagram

$$\begin{array}{ccc} A & \stackrel{\stackrel{p}{\longrightarrow}}{\longrightarrow} & B \\ j & \downarrow \uparrow q & j' & \downarrow \uparrow q' \\ C & \stackrel{\stackrel{p'}{\longrightarrow}}{\longrightarrow} & \mathrm{PO} \end{array}$$

commutative in both directions i.e., pq' = qp' and, moreover, such that jp = p'j' and iq = q'i'. One has to proceed just as the proof of Lemma 9.10 but, in diagram (9), take 1_B instead of $i(qj - 1_A)p + 1_B$ and, in diagram (11), take p instead of qjp.

9.3.3. The complementation feature of multiple push-out. Let us check now that almost complementation is preserved in almost complemented push-out with several factors:

Lemma 9.11. Let $(i_1, \overline{i_1}) : A_1 \to B_1$ and $(i_2, \overline{i_2}) : A_2 \to B_2$ be (1, 0, 1)-arrows. Let $(j_1, \overline{j_1}) : A_1 \to X$ be a (u, v, w)-arrow and let $j_2 : A_2 \to X$ be an operator. Consider the push-out diagram

(12)
$$\begin{array}{cccc} A_1 \oplus_1 A_2 & \xrightarrow{i_1 \oplus i_2} & B_1 \oplus_1 B_2 \\ & & & \downarrow^{j_1+j_2} \downarrow & & \downarrow^{J} \\ & X & \xrightarrow{(i_1 \oplus i_2)'} & \text{PO} \,. \end{array}$$

The restriction $J_{|B_1}$ admits an arrow $\overline{J_{|B_1}}$: PO $\rightarrow B_1$ so that $(J_{|B_1}, \overline{J_{|B_1}})$ is a contractive tive $(u, v, \max\{w, 1+v\}u||j_2||)$ -arrow. In particular, if $(j_1, \overline{j_1})$ is a contractive (u, v, w) arrow then $(J_{|B_1}, \overline{J_{|B_1}})$ is a contractive $(u, v, \max\{w, 1+v\}||j_2||)$ -arrow.

Moreover, $\overline{i_1}\overline{J_{|B_1}} = \overline{j_1}(i_1 \oplus i_2)'$.

Proof. Perform first the almost-complemented push-out as in the diagram (6) in Lemma 9.10 to get

(13)

$$A_{1} \xrightarrow{(i_{1},\overline{i_{1}})} B_{1}$$

$$(j_{1},\overline{j_{1}}) \bigwedge \bigwedge \bigwedge (j'_{1},\overline{j'_{1}}) \bigwedge (j'_{1},\overline{j'_{1}})$$

$$X \xleftarrow{(i'_{1},\overline{i'_{1}})} P_{1}$$

in which $(i'_1, \overline{i'_1})$ is a contractive (u, 0, u)-arrow and $(j'_1, \overline{j'_1})$ is a contractive $(u, v, \max\{w, 1+v\})$ -arrow. Now make the push-out of the arrows i_2 and $i'_1 j_2$

(14)

$$A_{2} \xrightarrow{(i_{2},\overline{i_{2}})} B_{2}$$

$$i_{1}'_{j_{2}} \bigvee \qquad \qquad \downarrow \qquad \qquad \downarrow (i_{1}'_{j_{2}})'$$

$$P_{1} \xrightarrow{(i_{2},\overline{i_{2}'})} P_{2}$$

The map $\overline{i'_2}$ is obtained according to diagram (10) in the proof of Lemma 9.10, in such a way that $\overline{i'_2}i'_2 = 1_{P_1}$ and $\|\overline{i'_2}\| \leq max\{1, \|i'_1j_2\overline{i_2}\|\}$. On the other hand, since the following square is commutative

(15)
$$\begin{array}{cccc} A_1 \oplus_1 A_2 & \xrightarrow{i_1 \oplus i_2} & B_1 \oplus_1 B_2 \\ & & & & \downarrow i'_2 j'_1 + (i'_1 j_2) \\ & X & \xrightarrow{i'_2 i'_1} & P_2. \end{array}$$

there must be a unique operator arrow $\tau : PO \to P_2$ such that

(1) $\tau(i_1 \oplus i_2)' = i'_2 i'_1$ (2) $\tau J = i'_2 j'_1 + (i'_1 j_2)'$ (3) $\|\tau\| \le \max\{\|i'_2 i'_1\|, \|i'_2 j'_1 + (i'_1 j_2)'\|\} \le \max\{\|i'_2 i'_1\|, \|i'_2 j'_1 + (i'_1 j_2)'\|\} = 1.$

The almost projection is going to be $\overline{J_{|B_1}} = \overline{j'_1} \ \overline{i'_2}\tau$: PO $\rightarrow B_1$, where $\overline{j'_1}$ has been obtained in diagram (13) while $\overline{i'_2}$ has been obtained in diagram (14). To check this observe that


$$\begin{aligned} \|\overline{j'_1} \ \overline{i'_2} \ \tau \ J_{|B_1} - \mathbf{1}_{B_1} \| &= \|\overline{j'_1} \ \overline{i'_2} \ (i'_2 j'_1 + (i'_1 j_2)')_{|B_1} - \mathbf{1}_{B_1} \| \\ &= \|\overline{j'_1} \ \overline{i'_2} \ i'_2 j'_1 - \mathbf{1}_{B_1} \| \\ &= \|\overline{j'_1} \ j'_1 - \mathbf{1}_{B_1} \| \\ &\leq v. \end{aligned}$$

Since $\|\overline{j'_1} \ \overline{i'_2}\tau\| \leq \max\{w, 1+v\}u\|j_2\|$ it turns out that $(J_{|B_1}, \overline{J_{|B_1}})$ is a contractive $(u, v, \max\{w, 1+v\}u\|j_2\|)$ -arrow. If $\|j_1\| \leq 1$ then $(J_{|B_1}, \overline{J_{|B_1}})$ is a contractive $(u, v, \max\{w, 1+v\}\|j_2\|)$ -arrow. Finally, according again to diagram (10) in the proof of Lemma 9.10, there exists and operator $\overline{(i_1 \oplus i_2)'}$: $PO \to X$ such that $\overline{(i_1 \oplus i_2)'(i_1 \oplus i_2)'} = 1_X$, and the "moreover" part is clear.

9.3.4. The almost-push-out. Garbulinska introduces in [13, Lemma 3.1] a useful correction lemma. Let us show that it can be understood as an "almost" push-out construction, which moreover admits an extension to cover the case of almost double arrows.

Lemma 9.12 (Correction lemma).

- Given a contractive $1 + \varepsilon$ -isometry $f : X \to Y$ between Banach spaces, there exists a space E(f, X, Y) and isometries $i_f : X \to E(f, X, Y)$, $j_f : Y \to E(f, X, Y)$ such that $\|j_f f i_f\| \leq \varepsilon$ with the following universal property: for any couple of arrows $k : X \to V$ and $l : Y \to V$ such that $\|lf k\| \leq \varepsilon$ there exists a unique arrow $\gamma : E(f, X, Y) \to V$ such that $\gamma i_f = k$ and $\gamma j_f = l$.
- Given a contractive $(1 + \varepsilon, \varepsilon, 1)$ -arrow $(f, \overline{f}) : X \leftrightarrows Y$ there exist a space E = E(f, X, Y) and double arrows $(i, \overline{i}) : X \leftrightarrows E, (j, \overline{j}) : Y \leftrightarrows E$ making the diagram



 ε -commutative and verifying also $\overline{ij} = \overline{f}$ and $\overline{ji} = f$.

Proof. Let us first see that there exists a push-out diagram which partially corrects the almost-isometry f. To this purpose, consider the isometric (for $\varepsilon < 1$) embedding operator $\delta_{\varepsilon} : X \to X \oplus_{\infty} X$, $\delta_{\varepsilon}(x) = (x, \varepsilon x)$ and make the push-out square

$$\begin{array}{ccc} X & \stackrel{\delta_{\varepsilon}}{\longrightarrow} & X \oplus_{\infty} X \\ f & & & \downarrow f' \\ Y & \stackrel{\delta'}{\longrightarrow} & \text{PO} \,. \end{array}$$

By the general properties of the push-out, f' is a $(1 + \varepsilon)$ -isometry and δ' is an into isometry. Recall that PO is the quotient of $(X \oplus_{\infty} X) \oplus_1 Y$ via the natural quotient map $Q: (X \oplus_{\infty} X) \oplus_1 Y \to \text{PO}$ with kernel X that defines the push-out. We form a subspace of PO where X and Y embed isometrically at the cost of loosing commutativity by taking

$$E(f, X, Y) = Q(X \oplus_{\infty} 0 \oplus_{1} Y)$$

and define the map $s: X \oplus_{\infty} X \to E(f, X, Y)$ by s(x, z) = f'(x, 0). The map δ' is already well defined as a map $Y \to E(f, X, Y)$. The resulting square

$$\begin{array}{cccc} X & \stackrel{\delta_{\varepsilon}}{\longrightarrow} & X \oplus_{\infty} X \\ f & & & \downarrow^{s} \\ Y & \stackrel{\delta'}{\longrightarrow} & E(f, X, Y) \end{array}$$

is ε -commutative:

 $\|\delta' f(x) - s\delta_{\varepsilon}(x)\| = \|\overline{(x, 0, -f(x))}\| \le \inf_{\omega \in X} \|(x - \omega, \varepsilon\omega, f(\omega) - f(x)\| \le \varepsilon \|x\|.$ Moreover $s\delta_{\varepsilon} : X \to E(f, X, Y)$ is an into isometry:


$$\begin{aligned} \|x\| &\leq \|x\| - \|\omega\| + \|\omega\| - \|f(\omega)\| + \|f(\omega)\|, \ \forall \omega \in X \\ &\leq \|x - \omega\| + \varepsilon \|\omega\| + \|f(\omega)\| \\ &\leq \|\overline{(x, 0, 0)}\|_{\mathrm{PO}} = \|s\delta_{\varepsilon}(x)\|_{E}. \end{aligned}$$

We must therefore set: $i_f = s\delta_{\varepsilon}$ and $j_f = \delta'$.

We prove now the universal property mentioned above: let $k : X \to V$ and $l : Y \to V$ be operators such that $||lf - k|| \leq \varepsilon$. The map $t : X \oplus X \to V$ defined by $t(x, z) = k(x) + \varepsilon^{-1}(lf - k)(z)$ verifies $t\delta_{\varepsilon} = lf$. By the universal property of the push-out there exists a unique arrow $\gamma : \text{PO} \to V$ such that $\gamma f' = t$ and $\gamma \delta' = l$. And the recontractive on of γ to E(f, X, Y) yields $\gamma i_f(x) = \gamma f'(x, 0) = t(x, 0) = k(x)$; while $\gamma j_f = \gamma \delta' = l$.

The complemented version of the Correction lemma will follow from the universal property of the "almost push-out" applied first to the arrows 1_X and $\overline{f}: Y \to X$, so we get $\overline{i}: E \to X$ such that $\overline{ij} = \overline{f}$ and $\overline{ii} = 1_X$; and then to f and 1_Y , obtaining $\overline{j}: E \to Y$ such that $\overline{jj} = 1_Y$ and $\overline{ji} = f$. In addition, $\|\overline{f}\| \leq \|\overline{i}\|$ and $\|f\| \leq \|\overline{j}\|$.

Now, when one has a push-out diagram



and two arrows $\gamma : \text{PO} \to Z$ and $h : \text{PO} \to Z$ so that $\gamma\beta' = h\beta'$ and $\|\gamma\alpha' - h\alpha'\| \leq \varepsilon$ then $\|\gamma - h\| \leq \varepsilon$: indeed, for given $(c, b) + \Delta \in \text{PO}$ with $\|(c, b) + \Delta\| \leq 1$ pick a representative $(c_1, b_1) + \Delta$ so that $\|c_1\| \leq 1$. Since $(c_1, b_1) + \Delta = \alpha'(c_1) + \beta'(b_1)$ one has

$$\|(\gamma - h)((c, b) + \Delta)\| = \|(\gamma - h)((c_1, b_1) + \Delta)\| = \|\gamma \alpha'(c_1) - h\alpha'(c_1)\| \le \varepsilon.$$

Thus, since $\overline{f} \ \overline{j} \ j = \overline{f}$ and $\overline{f} \ \overline{j} \ i = \overline{f} f$ and $\overline{i} \ j = \overline{f}$ and $\|\overline{i} \ i - \overline{f} \ f\| \le \varepsilon$, it turns out that $\|\overline{f} \ \overline{j} - \overline{i}\| \le \varepsilon$.

The condition $\|\overline{f} \ \overline{j} - \overline{i}\| \leq \varepsilon$ that we have obtained does not appear in either [13] or [7], where the authors only consider the almost commutativity condition $\|jf-i\| \leq \varepsilon$ for embeddings. Observe that the almost commutativity for embeddings and projections implies $\|\overline{i} \ j - \overline{f}\| \leq \varepsilon$ and $\|\overline{j} \ i - f\| \leq \varepsilon$ (but there is no equality).

9.4. Perturbation of projections.

Lemma 9.13. Let A be an n-dimensional subspace of E which is complemented by some projection p of norm C. Let $\delta = \operatorname{dist}(A, \ell_1^n)$. Let $\{a_1, \ldots, a_n\}$ be a basis for A so that $\delta^{-1} \sum |\lambda_i| \leq \|\sum \lambda_i a_i\| \leq \sum |\lambda_i|$. Given $0 < \varepsilon < 1/3$, if $\|x_i - a_i\| \leq \frac{\varepsilon}{\delta C}$ then the map $\tau a_i = x_i$ is a $(1 + \varepsilon)$ -isometry and the space $X = [x_1, \ldots, x_n]$ is:

(1) complemented via some projection p' of norm at most $C\frac{1-\varepsilon^2}{1-3\varepsilon}$ for which

$$\|p' - \tau p\| \le \varepsilon \frac{(1+\varepsilon)^2}{1-\varepsilon}C.$$

(2) In particular, $\frac{1}{1+\varepsilon}\tau$ is a contractive $(1+\varepsilon)^2$ -isometry with projection $(1+\varepsilon)p'$ having norm at most $C\frac{(1+\varepsilon)(1-\varepsilon^2)}{1-3\varepsilon}$ and so that

$$\|(1+\varepsilon)p' - \frac{1}{1+\varepsilon}\tau p\| \le C3\varepsilon.$$

Proof. The operator $\tau : A \to X$ that sends $\tau(a_i) = x_i$ is a $(1 + \varepsilon)$ -isometry. And if $p : E \to A$ is a norm-one projection, on every $x = \sum \lambda_i x_i \in X$ one has

$$\|\tau px - x\| = \|\tau p(\sum \lambda_i x_i) - \sum \lambda_i \tau pa_i\| \le (1 + \varepsilon) \|\sum \lambda_i (x_i - a_i)\| \le \varepsilon \frac{1 + \varepsilon}{1 - \varepsilon} \|x\|.$$

The estimates now are as in Lemma 9.6 (4). We call $\mu = \varepsilon \frac{(1+\varepsilon)}{1-\varepsilon}$. Since $(1_E - \tau p)_{|X|}$ has norm $\mu < 1$ for $\varepsilon < 1/3$, then $\tau p_{|X|} = 1_E - (1_E - \tau p_{|X|})$ is invertible and its inverse has norm at most $1 + \mu + \mu^2 + \cdots = \frac{1}{1-\mu}$. So, $(\tau p)_{|X|}$ is an isomorphism and $\|(\tau p)_{|X|}^{-1}\| \leq \frac{1}{1-\mu}$. It turns out that $p' = (\tau p)_{|X|}^{-1} \tau p$ is a projection onto X since

$$p^{\prime 2} = (\tau p)_{|X}{}^{-1}\tau p(\tau p)_{|X}{}^{-1}\tau p = (\tau p)_{|X}{}^{-1}(\tau p)_{|X}(\tau p)_{|X}{}^{-1}\tau p = (\tau p)_{|X}{}^{-1}\tau p$$

with norm at most $\frac{1+\varepsilon}{1-\mu}$. Moreover,

$$\begin{aligned} \|p' - \tau p\| &= \|(\tau p)_{|X}^{-1} \tau p - \tau p\| \\ &\leq \|(\tau p)_{|X}^{-1} - 1_X\| \|\tau p\| \\ &\leq \|(\tau p)_{|X}^{-1}\| \|1_X - (\tau p)_{|X}\| \|\tau\| \\ &\leq \frac{1}{1 - \mu} \mu (1 + \varepsilon) \\ &\leq \frac{\varepsilon (1 + \varepsilon)^2}{1 - 3\varepsilon} \\ &\leq \frac{4\varepsilon}{1 - 3\varepsilon}. \end{aligned}$$

The "in particular" estimate easily follows:

$$\begin{split} \|(1+\varepsilon)p' - \frac{1}{1+\varepsilon}\tau p\| &= (1+\varepsilon)\|p' - \frac{1}{(1+\varepsilon)^2}\tau p\| \\ &\leq (1+\varepsilon)\left(\|p' - \tau p\| + \left(1 - \frac{1}{(1+\varepsilon)^2}\right)\|\tau p\|\right) \\ &\leq (1+\varepsilon)\left(\|p' - \tau p\| + \left(1 - \frac{1}{(1+\varepsilon)^2}\right)\|\tau p\|\right) \\ &\leq (1+\varepsilon)\left(\frac{\varepsilon(1+\varepsilon)^2}{1-3\varepsilon}C + \frac{(1+\varepsilon)^2 - 1}{(1+\varepsilon)^2}(1+\varepsilon)C\right) \\ &\leq C\left(\frac{\varepsilon(1+\varepsilon)^3}{1-3\varepsilon} + 2\varepsilon + \varepsilon^2\right) \\ &\leq C(3\varepsilon - \varepsilon^2) \\ &\leq C3\varepsilon. \end{split}$$

9.5. Countable dense sets of double arrows between finite-dimensional spaces. To produce a separable space as output a basic ingredient is to have a countable set of double arrows between finite dimensional spaces that is "dense". To this end, consider for fixed $n \leq k$ the set of double arrows

$$\mathcal{U}_{n,k} = \{ (f, \overline{f}) : A \leftrightarrows B \quad \dim A = n; \quad \dim B = k \}$$

in which elements are identified as: $(f, \overline{f}) \sim (g, \overline{g})$ when there are surjective isometries $a : A \to A'$ and $b : B \to B'$ such that bf = ga and $a\overline{f} = \overline{g}b$. We call $\mathcal{U}(n, k)$ the quotient space endowed with the metric induced by

 $d((f,\overline{f}),(g,\overline{g})) = \inf\{\log(1+\varepsilon) > 0 : \exists a, b \ (1+\varepsilon) - \text{onto isometries} : bf = ga \text{ and } a\overline{f} = \overline{g}b\}.$ One has:

Lemma 9.14. The space $\mathcal{U}(m, l)$ is a compact metric space for all m, l.

Proof. Let $(A_k, B_k, f_k, \overline{f_k})$ be a sequence. In the Banach-Mazur distance –for spaces– and the operator norm -for operators- there is a subsequence (no need to relabel) so that $\lim A_k = A$, $\lim B_k = B$, $\lim f_k = f$ and $\lim \overline{f_k} = \overline{f}$. There is no loss of generality assuming that the almost isometries that yield the Banach-Mazur distance are the identity. Which in particular means that if one fixes a basis in each A_k and e_j^k is the jth element in A_k then $e_j^k \to a_j$, the elements a_j form a basis for A and $f_k(a_j) \to b_j$ form a basis for f(A) in B, which we complete with as many $b'_i s$ as necessary. Let \mathcal{U} be a free ultrafilter on N. One has $A = [A_1, A_2, \ldots, A_n, \ldots]_{\mathcal{U}}$ and $B = [B_1, B_2, \ldots, B_n, \ldots]_{\mathcal{U}}$. The map $f = [f_k]$ is thus an isometry between them and $\overline{f} = [\overline{f_k}]$ a 1-projection. Moreover, given a finite dimensional space F one has $F = [F]_{\mathcal{U}}$ and thus one can identify A_k with the its ultrapower $[A_k]_{\mathfrak{U}}$ and B_k with $[B_k]_{\mathfrak{U}}$. In this way, the formal identity 1_k : $A_k = [A_k]_{\mathcal{U}} \to [A_n]_{\mathcal{U}} = A$ is a $1 + \text{dist}(A_k, A)$ -isometry. To check that $(f,\overline{f}):A \cong B$ is the limit of $(f_k,\overline{f_k})$ we set 1_k on the left and do as follows on the right: given k, we call $t_k : B_k \to B$ the $1 + \varepsilon$ -isometry that fixes all b_i while sending $f_k(a_i)$ to b_i (of course that ε depends on k, but goes to 0 when k goes to infinity). Form $[t_k]$ and observe that the diagram

$$\begin{array}{ccc} A_k & \xrightarrow{\overline{f_k}} & B_k \\ & & & & \downarrow^{[t_k]} \\ 1_k \downarrow & & & \downarrow^{[t_k]} \\ & & & \downarrow^{[t_k]} \\ & & & & [A_n]_{\mathfrak{U}} & \xrightarrow{[\overline{f_k}]} & [B_n]_{\mathfrak{U}} \end{array}$$

is commutative in both directions.

Now, observe that there is no loss of generality in assuming that the two $(1 + \varepsilon)$ isometries a, b in the definition of the distance $d(\cdot, \cdot)$ at the beginning of section 9.5 are contractive $(1 + \varepsilon)$ -isometries: indeed, given a, b so that bf = ga and $a\overline{f} = \overline{g}b$ one can set $a' = \frac{1}{1+\varepsilon}a$ and $b' = \frac{1}{1+\varepsilon}b$, who still satisfy b'f = ga' and $a'\overline{f} = \overline{g}b'$. For the same reason, one can also make a^{-1}, b^{-1} contractive $(1 + \varepsilon)$ -isometries. Thus, since metrizable compacta are separable we get:

Lemma 9.15. There is a countable set \mathfrak{U} of double arrows between finite dimensional spaces with the following property: given a double arrow $(w, \overline{w}) : A \cong B$ between finite dimensional spaces and $\varepsilon > 0$, there is $(u, \overline{u}) : A_u \cong B_u$ in \mathfrak{U} , and surjective contractive $(1 + \varepsilon)$ -isometries $a : A_u \to A$ and $b : B_u \to B$ making the square

$$\begin{array}{cccc}
A_u & \stackrel{u}{\longrightarrow} & B_u \\
a \downarrow & & \downarrow_b \\
A & \stackrel{w}{\longrightarrow} & B
\end{array}$$

commutative both directions; i.e., wa = bu and $a\overline{u} = \overline{w}b$.

9.6. Distances between double arrows and the role of dual separable spaces. Almost double arrows $A \cong B$ form a subset of $\mathfrak{L}(A, B) \oplus \mathfrak{L}(B, A)$, and thus the distance between two almost double arrows $(j, \overline{j}) : A \cong B$ and $(i, \overline{i}) : A \cong B$ is defined as $\max\{\|j-i\|, \|\overline{j}-\overline{i}\|\}$. The following lemma is here to justify the additional hypothesis in Theorem 4.1.

Lemma 9.16. Let F be a finite dimensional Banach space. There is a countable set of double arrows $F \to X$ which is dense in the set of all double arrows $F \to X$ if and only if X^* is separable.

Proof. (Necessity) Set $F = \mathbb{R}$, without loss of generality. Every double arrow $(f, p) : \mathbb{R} \leftrightarrows X$ is an isometric embedding f and a 1-projection onto $f(\mathbb{R})$. Or, which is the same, a norm one element $u \in X$ and a norm one functional $\phi \in X^*$ so that $\phi(u) = 1$. The projection is $p(x) = \phi(x)u$. Assume there is a countable set of (f, p) so that for every (g, q) there is one of them for which $||f - g|| + ||p - q|| \leq \varepsilon$. Let ψ be a norm one element of X^* . Find norm one $v \in X$ for which $\psi(v) = 1 - \varepsilon$ and then form the isometric embedding g(1) = v with projection $q(x) = (1 - \varepsilon)^{-1}\psi(x)v$. Find one of those countable elements (f, p) close to (g, q). If $p(x) = \phi(x)u$ with f(1) = u then for ||x|| = 1 one has

$$\begin{aligned} |\phi(x) &- (1-\varepsilon)^{-1}\psi(x)| &= \|\phi(x)u - (1-\varepsilon)^{-1}\psi(x)u\| \\ &\leq \|\phi(x)u - (1-\varepsilon)^{-1}\psi(x)v\| + \|(1-\varepsilon)^{-1}\psi(x)v - (1-\varepsilon)^{-1}\psi(x)u\| \\ &\leq \|p-q\| + (1-\varepsilon)^{-1}\|v-u\| \\ &= \|p-q\| + (1-\varepsilon)^{-1}\|f(1) - g(1)\| \\ &\leq \|p-q\| + (1-\varepsilon)^{-1}\|f - g\| \\ &\leq 2\varepsilon + \frac{2\varepsilon}{1-\varepsilon}. \end{aligned}$$

(Sufficiency) The set of double arrows so it is separable when both $\mathfrak{L}(F,X)$ and $\mathfrak{L}(X,F)$ are separable; that is, when X^* is separable.

References

- A. Avilés, F. Cabello, J.M.F. Castillo, M. González, and Y. Moreno, Banach spaces of universal disposition, J. Funct. Anal. 261 (2011) 2347-2361.
- [2] A. Avilés, F. Cabello, J.M.F. Castillo, M. González, and Y. Moreno, On ultraproducts of Banach space of type \mathcal{L}_{∞} , Fundamenta Math. 222 (2013) 195-212.
- [3] A. Avilés, F. Cabello, J.M.F. Castillo, M. González, and Y. Moreno, Separably injective Banach spaces, Lecture Notes in Mathematics 2132, Springer 2016.
- [4] P. Bankston, Reduced coproducts of compact Hausdorff spaces, J. Symbolic Logic 52 (1987), 404–424.
- [5] C. Bargetz, K. Kąkol, W. Kubiś, A separable Fréchet space of almost universal disposition, J. Funct. Anal. 272 (2017) 1876–1891.
- [6] A. Błaszyk and A. Szymański, Concerning Parovičenko's theorem, Bull. Acad. Polon. Sci. Math. 28 (1980) 311–314.
- F. Cabello, J. Garbulinska, W. Kubis, Quasi-Banach spaces of almost universal disposition, J. Funct. Anal. 267 (2014), 744–771.
- [8] F. Cabello, JMF. Castillo, Y. Moreno, On the bounded approximation property on subspaces of ℓ_p when 0 and related issues, Forum Math. (in press).
- [9] P.G. Casazza, Approximation properties. Handbook of the geometry of Banach spaces, Vol. I; pp. 271–316. North-Holland, Amsterdam, 2001.
- [10] Jesús M. F. Castillo M. A. Simoes, On Banach spaces of universal disposition, New York J. Math. 22 (2016) 605-613.
- [11] JMF Castillo and J. Suárez, On L_∞-envelopes of Banach spaces, J. Math. Anal. Appl. 394 (2012) 152–158.
- [12] P. Dodos, Banach spaces and descriptive set theory: selected topics, Lecture Nothes in Mathematics 10993, Springer 2010.
- [13] J. Garbulinska, Isometric uniqueness of a complementably universal Banach space for Schauder decompositions, Banach J. Math. Anal. 8 (2014), 211-220.
- [14] Yu L. Gevorkyan, The universality of spaces of almost universal displacement, (Russian) Funkcional. Anal. i Priložen 8 (1974), no. 2, 72.
- [15] V.I. Gurariy, Spaces of universal placement, isotropic spaces and a problem of Mazur on rotations of Banach spaces (Russian). Sibirsk. Mat. Z. 7 (1966), 1002–1013.
- [16] W.B. Johnson, A. Szankowski, Complementably universal Banach spaces, Studia Math. 58 (1976) 91–97.
- [17] M. I. Kadec, On complementably universal Banach spaces, Studia Math. 40 (1971) 85–89.
- [18] N.J. Kalton, Locally complemented subspaces and \mathcal{L}_p -spaces for 0 , Math. Nachr. 115 (1984) 71û97.

BANACH SPACES OF ALMOST UNIVERSAL

- [19] N. Kalton, Universal spaces and universal bases in metric linear spaces, Studia Math. 61 (1977), 161–191
- [20] W. Kubiś, Linearly ordered compacta and Banach spaces with a projectional resolution of the identity, Topology and Appl. 154 (2007) 749–757.
- [21] W. Kubiś, Banach spaces with projectional skeletons, J. Math. Anal. Appl. 350 (2009) 758–776.
- [22] W. Kubiś, Fraïssé sequences: category-theoretic approach to universal homogeneous structures. Ann. Pure Appl. Logic 165 (2014), no. 11, 1755–1811.
- [23] O. Kurka, Amalgamations of classes of Banach spaces with a monotone basis, Studia Math. 3496 (2016), 121 – 148
- [24] A.J. Lazar and J. Lindenstrauss, Banach spaces whose duals are L₁-spaces and their representing matrices. Acta Math. 126 (1971) 165–193
- [25] J. Lindenstrauss, H.P. Rosenthal, The \mathcal{L}_p spaces, Israel J. Math. 7 (1969), 325–349.
- [26] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces I, sequence spaces, Ergebnisse der Mathematik und ihrer Grenzgebiete 92, Springer-Verlag 1977.
- [27] W. Lusky, The Gurarii spaces are unique. Arch. Math. 27 (1976) 627-635.
- [28] W. Lusky, Separable Lindenstrauss spaces. Functional Analysis: Surveys and Recent Results. North-Holland 1977, pp. 15–28.
- [29] A. Martínez-Abejón, An elementary proof of the principle of local reflexivity, Proc. Amer. Math. Soc. 127 (1999) 1397–1398.
- [30] E. Odell, T. Schlumprecht, Embedding into Banach spaces with finite dimensional decompositions, Rev. R. Acad. Cien. Mat. 100 (2006) 295–323.
- [31] E. Oja and M. Põldvere, Principle of local reflexivity revisited, Proc. Amer. Math. Soc. 135 (2007) 1081–1088.
- [32] A. Pełczyński, Any separable Banach space with the bounded approximation property is a complemented subspace of a Banach space with a basis, Studia Math. 40 (1971) 239–243.
- [33] A. Pełczyński, Universal bases, Studia Math. 32 (1969) 247–268.
- [34] A. Pełczyński and P. Wojtaszczyk, Banach spaces with finite dimensional expansions of identity and universal bases of finite dimensional spaces. Studia Math. XL (1971) 91–108.
- [35] S. Prus, Finite Dimensional Decompositions with p-Estimates and Universal Banach Spaces, Bull. Polish. Acad. Sci 31 (1983) 281-288.
- [36] R. C. Walker, The Stone-Cech compactification, Ergeb. Math. und ihrer Grenzgebiete 83, Springer-Verlag 1974.
- [37] P. Wojtaszczyk, Some remarks on the Gurarij space, Studia Math. XLI (1972), 207–210.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS, 06071-BADAJOZ, SPAIN

E-mail address: castillo@unex.es

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS, 06071-BADAJOZ, SPAIN

E-mail address: ymoreno@unex.es