# Gaps between divisible terms in $a^2(a^2+1)$

Tsz Ho Chan

#### Abstract

Suppose  $a^2(a^2 + 1)$  divides  $b^2(b^2 + 1)$  with b > a. In this paper, we improve a previous result and prove a gap principle, without any additional assumptions, namely  $b \gg a(\log a)^{1/8}/(\log \log a)^{12}$ . We also obtain  $b \gg_{\epsilon} a^{15/14-\epsilon}$  under the abc conjecture.

# 1 Introduction and Main Results

We are interested in increasing sequence of positive integers  $(a_n)_{n\geq 0}$  with each term dividing the next one (i.e.  $a_n|a_{n+1}$ ). A simple example is  $(2^n)_{n\geq 0}$ . Another example is  $(n!)_{n\geq 1}$ . These are simple recursively defined sequences. It is more interesting and challenging to require each term of the sequence to have a special form. For example,  $(3^n)_{n\geq 0}$  has all terms odd. For another special example, let us consider the Fibonacci numbers

$$F_1 = 1, F_2 = 1, F_{n+1} = F_n + F_{n-1}$$
 for  $n \ge 2$ .

By the well-known fact that  $F_m|F_n$  if and only if m|n, the sequence  $(F_{2^n})_{n\geq 0}$ has all terms of Fibonacci-type and each term dividing the next one. One may restrict the sequence to numbers of the form  $n^2+1$  or other polynomials, and we are interested in the growth of such sequences. In this paper, we shall focus on numbers of the form  $n^2(n^2+1)$  and study the following

**Question 1** Suppose  $a^2(a^2+1)$  divides  $b^2(b^2+1)$ . Must it be true that there is some gap between a and b? More preciously, is it true that  $b > a^{1+\epsilon}$  for some small  $\epsilon > 0$ ?

Key words and phrases: divisibility, gap principle, hyper-elliptic curve, abc conjecture AMS 2010 Mathematics Subject Classifications: 11B83

In [1], the author studied the above question with some additional restrictions on a and b. In this paper, we remove all these restrictions and prove

**Theorem 2** Let a and b be positive integers with a < b. Suppose  $a^2(a^2 + 1)$  divides  $b^2(b^2 + 1)$ . Then

$$b \gg \frac{a(\log a)^{1/8}}{(\log \max(e, \log a))^{12}}.$$

Recently, Stephen Choi, Peter Lam and the author [2] defined gap principle of order n for polynomials with integer coefficients as follows:

**Definition 3** Let n be a positive integer and f(x) be a polynomial with integral coefficients. Consider the set of all positive integers  $a_0 < a_1 < a_2 < \ldots < a_n$  such that  $f(a_i)$  divides  $f(a_{i+1})$  for  $0 \le i \le n-1$ . We say that f(x) satisfies the gap principle of order n if  $\lim a_n/a_0 = \infty$  as  $a_0 \to \infty$ .

Hence, Theorem 2 implies that the polynomial  $f(x) = x^2(x^2 + 1)$  satisfies the gap principle of order 1.

Assuming the abc conjecture, we can obtain a better gap result than that in [1], without any extra assumptions.

**Theorem 4** Let a and b be positive integers with a < b. Suppose  $a^2(a^2 + 1)$  divides  $b^2(b^2 + 1)$ . Then, under the abc conjecture with any small  $\epsilon > 0$ ,

 $b \gg_{\epsilon} a^{15/14-\epsilon}$ 

Thus, this answers question 1 in the affirmative under the abc conjecture. We hope this article would inspire readers to study questions of similar nature.

**Some Notations** The symbol a|b means that a divides b. The notations  $f(x) \ll g(x)$  and  $g(x) \gg f(x)$  are all equivalent to  $|f(x)| \leq Cg(x)$  for some constant C > 0. Finally  $f(x) = O_{\lambda}(g(x)), f(x) \ll_{\lambda} g(x)$  or  $g(x) \gg_{\lambda} f(x)$  mean that the implicit constant C may depend on  $\lambda$ .

#### 2 Proof of Theorem 2

Since  $a^{2}(a^{2}+1)$  divides  $b^{2}(b^{2}+1)$ , say

$$ta^2(a^2+1) = b^2(b^2+1)$$

for some integer t > 1. We may assume  $t \le \log a$ , for otherwise the theorem is true automatically. Let D be the greatest common divisor of a and b. Suppose a = Dx and b = Dy with (x, y) = 1, and let  $T = (D^2x^2+1, D^2y^2+1)$ . Then

$$tx^2 \frac{D^2 x^2 + 1}{T} = y^2 \frac{D^2 y^2 + 1}{T}.$$
(1)

Since (x, y) = 1,  $x^2$  must divide  $(D^2y^2 + 1)/T$ . Say  $(D^2y^2 + 1)/T = mx^2$  for some integer m. Then  $t(D^2x^2 + 1)/T = my^2$ , and we have

$$t(D^2x^2 + 1) = mTy^2$$
(2)

and

$$D^2 y^2 + 1 = mTx^2. (3)$$

Multiplying (2) by  $D^2$ , (3) by mT and combining, we have

$$[(mT)^2 - tD^4]x^2 = tD^2 + mT.$$
(4)

Similarly, multiplying (2) by mT, (3) by  $tD^2$  and combining, we have

$$[(mT)^{2} - tD^{4}]y^{2} = tD^{2} + tmT.$$
(5)

Subtracting (4) from (5), we get

$$s(y^2 - x^2) = (t - 1)mT$$
(6)

with

$$s = (mT)^2 - tD^4.$$
 (7)

From (3), we have (mT, D) = 1. Hence,

$$(s, mT) = ((mT)^2 - tD^4, mT) = (tD^4, mT) = (t, mT).$$

Therefore, by combining this with (6), we have s|(t-1)t and (7) gives us an hyperelliptic curve

$$Y^2 = tX^4 + s$$

by setting Y = mT and X = D. Let  $\lambda := (t - 1)t$ . By Theorem 1 of Voutier [3] on the study of integral solutions to hyperelliptic curves using transcendental number theory, we have

$$\max(X, Y) < e^{C_1 \lambda (\max(1, \log \lambda))^{96}}$$

for some constant  $C_1 \ge 1$ . Suppose  $\lambda \le \frac{1}{C_1} \frac{\log D}{(\log \max(e, \log D))^{96}}$ . Then

$$D = X < e^{C_1 \frac{1}{C_1} \frac{\log D}{(\log \max(e, \log D))^{96}} (\log \max(e, \log D))^{96}} = D$$

which is a contradiction. Therefore,  $(t-1)t > \frac{1}{C_1} \frac{\log D}{(\log \max(e, \log D))^{96}}$  and

$$t \gg \frac{\sqrt{\log D}}{(\log \max(e, \log D))^{48}}.$$
(8)

From (2), we have  $2tD^2x^2 \ge mTy^2$  and  $tD^2 \gg mT$ . This together with (4) gives  $tD^2 \gg x^2$ . Hence, as  $t \le \log a$  and a = Dx, we have  $\log D \gg \log a$ . Therefore, (8) gives

$$t \gg \frac{\sqrt{\log a}}{(\log \max(e, \log a))^{48}}$$

and we have Theorem 2 as  $b^4/a^4 \gg t$ .

### 3 Proof of Theorem 4

Firstly, for any integer n, let  $R(n) := \prod_{p|n} p$  be the radical or kernel of an integer n. The abc-conjecture is

**Conjecture 5** For every  $\epsilon > 0$ , there exists a constant  $C_{\epsilon}$  such that for all triples (a, b, c) of coprime positive integers, with a + b = c, then

$$c < C_{\epsilon} R(abc)^{1+\epsilon}.$$

Secondly, let us state a lemma which follows easily from the unique prime factorization of numbers.

**Lemma 6** Suppose  $a|A^2$  and  $a = a_1a_2^2$  with  $a_1$  squarefree. Then  $a_1a_2|A$ .

Basically, we follow the proof of Theorem 2. Using the same notation as in section 2,

$$a = Dx$$
,  $b = Dy$ , and  $s + tD^4 = (mT)^2$ 

with (mT, D) = 1. Let

 $(t, (mT)^2) = d$  and  $d = d_1 d_2^2$  with  $d_1$  squarefree.

By Lemma 6, we have  $mT = d_1 d_2 S$ . Then

$$\frac{s}{d} + \frac{t}{d}D^4 = d_1 S^2. \tag{9}$$

Here, the three terms are integers and pairwise relatively prime. We can apply the abc conjecture and obtain

$$\frac{t}{d}D^4 < C_\epsilon \left(\frac{(t-1)t}{d}Dd_1S\right)^{1+\epsilon} \tag{10}$$

since s|(t-1)t. Suppose  $t \leq 10D^4$ . As  $d_1 \leq d$ , (9) gives

$$(d_1S)^2 \ll tD^4$$
 or  $d_1S \ll t^{1/2}D^2$ .

Putting this into (10), we have

$$\frac{t}{d}D^4 \ll_{\epsilon} \left(\frac{t^{5/2}}{d}D^3\right)^{1+\epsilon}.$$

Hence,

$$D \ll_{\epsilon} t^{3/2+8\epsilon}. \tag{11}$$

As  $s + tD^4 = (mT)^2$ , s|(t-1)t and  $t \leq D^4$ , we have  $mT \ll t^{1/2}D^2$ . This together with (4) gives

$$x^2 \ll \frac{tD^2}{s}$$
 or  $x \ll t^{1/2}D$ .

Therefore, by (11),

$$a = Dx \ll t^{1/2} D^2 \ll_{\epsilon} t^{7/2 + 16\epsilon} \text{ or } t \gg_{\epsilon} a^{2/7 - 2\epsilon}$$

which gives Theorem 4 as  $b^4/a^4 \gg t$ .

We are left with the case  $t > 10D^4$ . Suppose  $y/x = b/a \le a^{1/5} = (Dx)^{1/5}$ . From (1), we have  $t^{1/4} \le \sqrt[4]{2}y/x$ . Hence,

$$10D^4 < t \le 2(Dx)^{4/5}.$$
 (12)

This gives

$$x > \frac{10}{2^{5/4}} D^4. \tag{13}$$

On the other hand, (3) and (1) give

$$mT \le 2D^2 y^2 / x^2 \le 2\sqrt{2}t^{1/2}D^2$$

Putting this into (4), we have

$$x^{2} \leq (1+2\sqrt{2})tD^{2} \text{ or } \frac{1}{1+2\sqrt{2}} \left(\frac{x}{D}\right)^{2} \leq t.$$
 (14)

Combining (12) and (14), we obtain

$$\frac{1}{1+2\sqrt{2}} \left(\frac{x}{D}\right)^2 \le 2(Dx)^{4/5} \text{ or } x \le [2(1+\sqrt{2})]^{5/6} D^{7/3}$$

which contradicts (13). Therefore, we must have  $b/a > a^{1/5}$  which also gives Theorem 4.

# References

- T.H. Chan, Common factors among pairs of consecutive integers, Int. J. Number Theory 14 (2018), no. 3, 871–880.
- [2] Tsz Ho Chan, Stephen Choi and Peter Cho-Ho Lam, Divisibility on the sequence of perfect squares minus one: The gap principle, J. Number Theory 184 (2018), 473–484.
- [3] P. Voutier, An Upper Bound for the Size of Integral Solutions to  $Y^m = f(X)$ , J. Number Theory 53 (1995), 247–271.

Department of Mathematical Sciences University of Memphis Memphis, TN 38152 U.S.A. thchan6174@gmail.com