

BURGERS EQUATION IN THE ADHESION MODEL

YURI GLIKLIKH AND EVELINA SHAMAROVA

ABSTRACT. We prove the existence and uniqueness of a classical solution to a multidimensional non-potential stochastic Burgers equation with Hölder continuous initial data. Our motivation is the adhesion model in the theory of formation of the large-scale structure of the universe. Importantly, we drop the assumption on the potentiality of the velocity flow that has been questioned in physics literature.

Keywords: Multidimensional non-potential stochastic Burgers equation, adhesion model

AMS subject classifications: 60H15, 60H10, 35K59

1. Introduction

In this work, we are concerned with a multidimensional stochastic Burgers equation of the form

$$(1) \quad y(t, x) = \varphi(x) + \int_0^t [\nu \Delta y(s, x) - (y, \partial_x)y(s, x)] ds + \eta(t, x)$$

with a view of its application to the adhesion model in cosmology. Here, $(t, x) \in [0, T] \times \mathbb{R}^d$, φ is a Hölder continuous bounded initial function, and $\eta(t, x)$ is a noise, rough in time and smooth in space. The presence of the noise, however, is introduced for the sake of generality since in our main application $\eta = 0$.

The adhesion model was introduced by Gurbatov and Saichev [9] as a generalization of Zel'dovich's approximation [19] aiming to represent the effect of gravitational sticking in the formation of the large-scale structure of the universe. Zel'dovich's approximation is based on an inviscid potential Burgers equation, and is only valid in the early linear regime of the universe expansion. It is essential that in this part of the theory, the potentiality of the velocity field is used in the derivation of the basic equation of the model. To formulate the adhesion model, the term $\nu \Delta y$ was artificially added to the (inviscid) Burgers equation, and then the model was tested numerically showing a qualitative and a quantitative agreement with a gravitational N -body simulation [6] (a numerical solution of the equations of motion for N gravitationally interacting particles). It turned out that this agreement holds not only in the early linear regime but also in the subsequent strongly non-linear regime, described as follows: the matter, previously concentrated in Zel'dovich's pancakes, moves towards the faces, the edges, and the vertices of the emerging mosaic structure, and then the latter is deformed due to the gravitational interaction [11]. However, according to what was discussed in [5], edges and vertices, representing high-density regions of the structure (galaxy clusters and superclusters), are associated with strong vortex flows. This means that the velocity flow is no longer potential. Remark that the assumption on its potentiality in the derivation of Zel'dovich's approximation is only valid in the early linear regime while in the adhesion model the initial condition is still assumed potential. This latter fact is

our main motivation for studying a non-potential Burgers equation which agrees with the adhesion model otherwise. By the latter, we mean that the initial function should be stationary against translations and scale-invariant [18]. The above properties are satisfied by a fractional Brownian sheet $W^H(x)$. Since we require the initial data to be bounded, we consider $\varphi(x) = \zeta(x)W^H(x)$, where $\zeta(x)$ is a C^∞ -cutting function, i.e., a mollified indicator function, of a bounded domain where the expansion of the universe takes place according to the adhesion model, and whose diameter is compatible with the scale of validity of the latter [11].

Thus, in this work, we are concerned with the existence and uniqueness of a classical solution to equation (1) when the initial function $\varphi(x)$ possesses just a Hölder regularity. According to the results of [17] (Theorem 6.2), the aforementioned property is satisfied by a fractional Brownian sheet. Thus, the above-described choice of the initial data is suitable for the adhesion model.

A stochastic Burgers equation of form (1) has been extensively studied in the literature over the past two decades (see, e.g., [1, 2, 3, 7, 13, 8, 10, 11, 14]); however, in most cases this study was restricted to the one-dimensional or the potential case. The interest to the potential case is mainly based on the existence of an exact solution by means of the Cole-Hopf transformation. In the non-potential case, it is only known that a multidimensional stochastic Burgers equation possesses a solution in an L_p -space [4]. However, an L_p -solution is not a classical, and not even a continuous solution, and, therefore, is not suitable for the adhesion model. Indeed, the evolution of the large-scale structure of the universe is regarded as a continuous process of transport of the matter. Thus, apart of our main important application, a classical solution to equation (1) in the non-potential case represents an interesting mathematical question (see also the discussion in [2] on this topic).

Our pathway to obtaining a classical solution to (1) is as follows. Equation (1) is first reduced to a PDE with random coefficients. The sequence of classical solutions y_m corresponding to a sequence φ_m of mollified initial functions is shown to be bounded and Hölder continuous uniformly in m which implies the existence of a converging subsequence. The limit function $y^{(1)}$ is then plugged into the non-linear term in (1), i.e., the non-linear term becomes $(y^{(1)}, \partial_x)y$, reducing (1) to a linear equation. For a linear PDE, in turn, it is well known (see, e.g., [12]) that a Hölder regularity of the initial data is sufficient for the existence of a classical solution.

2. Preliminaries

2.1 Existence of solution to a deterministic Burgers-type equation

In this section, we obtain the existence of solution to the Burgers-type equation

$$(2) \quad \begin{cases} \partial_t y(t, x) = \nu \Delta y(t, x) - (g(t, x, y), \partial_x)y(t, x) + f(t, x, y), \\ y(0, x) = \varphi(x), \end{cases}$$

where the force f is smooth in time and space variables and φ is of class $C_b^{2+\beta}(\mathbb{R}^d)$, $\beta \in (0, 1)$. By the latter, we understand the class of twice differentiable functions possessing a Hölder continuous second derivative. This result will be required in the next section.

Recall that the Hölder constants $[\Phi]_{\frac{\beta}{2}}^t$ and $[\Phi]_{\beta}^x$ are defined as follows:

$$[\Phi]_{\frac{\beta}{2}}^t = \sup_{\substack{t, t' \in [0, T], \\ t \neq t'}} \frac{|\Phi(t, x, y) - \Phi(t', x, y)|}{|t - t'|^{\frac{\beta}{2}}}; \quad [\Phi]_{\beta}^x = \sup_{\substack{x, x' \in \mathbb{R}^d, \\ 0 < |x - x'| < 1}} \frac{|\Phi(t, x, y) - \Phi(t, x', y)|}{|x - x'|^{\beta}};$$

$[\Phi]_{\beta}^y$ is defined likewise.

Theorem 2.1 below (the existence theorem for equation (2)) and its proof are included here for the sake of completeness since in [15] both are given only for the case one equation.

THEOREM 2.1. *Let the functions f , g , and φ satisfy assumptions (i) – (iii) below*

- (i) $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}^d$ belongs to $C_b^{2+\beta}(\mathbb{R}^d)$, $\beta \in (0, 1)$;
- (ii) the partial derivatives $\partial_t f$, $\partial_x f$, $\partial_y f$, $\partial_t g$, $\partial_x g$, $\partial_y g$ are continuous;
- (iii) the following estimates hold on $[0, T] \times \mathbb{R}^d \times \mathbb{R}^d$:

$$\begin{aligned} (f(t, x, y), y) &< C(1 + |y|^2), \\ |g(t, x, y)| + |\partial_x g(t, x, y)| + |\partial_y g(t, x, y)| + [g]_{\frac{\beta}{2}}^t &\leq \Lambda_1(|y|), \\ |f(t, x, y)| + |\partial_x f(t, x, y)| + |\partial_y f(t, x, y)| + [f]_{\frac{\beta}{2}}^t &\leq \Lambda_2(|y|), \end{aligned}$$

where C is a constant and $\Lambda_1(\cdot)$ and $\Lambda_2(\cdot)$ are positive non-decreasing functions.

Then, there exists a $C_b^{1,2}$ -solution $y(t, x)$ to problem (4). Further, the bound for $|y(t, x)|$ depends only on \mathcal{C} , T , and $\sup_{x \in \mathbb{R}^d} |\varphi(x)|$.

Proof. Consider the initial-boundary value problem

$$(3) \quad \begin{cases} \partial_t y(t, x) = \nu \Delta y(t, x) - (g(t, x, y), \nabla) y(t, x) + \zeta(x) f(t, x, y), \\ y(0, x) = \varphi(x) \zeta(x), \quad y(t, x) \Big|_{\partial B_r} = 0, \end{cases}$$

where B_r is an open ball of radius $r > 1$, ∂B_r is its boundary, and $\zeta(x)$ is a smooth cutting function of the ball B_{r-1} with the properties $\zeta(x) = 1$ if $x \in B_{r-1}$, $\zeta(x) = 0$ if $x \notin B_r$, $0 \leq \zeta(x) \leq 1$, $\zeta(x)$ is bounded together with its derivatives of all orders. Consider, for example, a mollified indicator function of the ball $B_{r-\frac{1}{2}}$: $\zeta = \mathbb{1}_{B_{r-\frac{1}{2}}} * \rho_{\frac{1}{2}}$, where $\rho_{\frac{1}{2}}$ is the standard mollifier supported on the ball of radius $\frac{1}{2}$. The class of initial-boundary value problems for systems of quasilinear parabolic PDEs, which includes problem (3), was considered by Ladyzhenskaya et al. in [15] (Theorem 7.1, p. 596). By Theorem 7.1 of [15], problem (3) possesses a unique solution on $[0, T] \times \overline{B}_r$ which belongs to the Hölder space $C^{1+\frac{\beta}{2}, 2+\beta}([0, T] \times \overline{B}_r)$ with the norm

$$\begin{aligned} \|y\|_{C^{1+\frac{\beta}{2}, 2+\beta}([0, T] \times \overline{B}_r)} &= \|y\|_{C^{1,2}([0, T] \times \overline{B}_r)} + \sup_{t \in [0, T]} [\partial_t y]_{\beta}^x + \sup_{t \in [0, T]} [\partial_{xx}^2 y]_{\beta}^x \\ &\quad + \sup_{x \in \overline{B}_r} [\partial_t y]_{\frac{\beta}{2}}^t + \sup_{x \in \overline{B}_r} [\partial_x y]_{\frac{1+\beta}{2}}^t + \sup_{x \in \overline{B}_r} [\partial_{xx}^2 y]_{\frac{\beta}{2}}^t. \end{aligned}$$

First of all, we note that by Theorem 6.1 (p. 592) from [15], there exists a bound for the gradient $\partial_x y(t, x)$ that only depends on ν , \mathcal{C} , T , and the bounds for $|\varphi(x)|$ and $|\nabla \varphi(x)|$. Also, according to the results of §7 from [15], the bound for $y(t, x)$ depends only on \mathcal{C} , T , and the bound for $|\varphi(x)|$.

Furthermore, by Theorem 5.1 from [15] (Chapter VII), the $C^{1+\frac{\beta}{2}, 2+\beta}([0, T] \times \overline{B_{r-1}})$ -norm of the solution $y(t, x)$ possesses a bound that depends only on \mathcal{C} , ν , T , and on the Hölder norm $\|\varphi\|_{C_b^{2+\beta}(\mathbb{R}^d)}$. Note that the bound for this norm does not depend on the radius r of B_r .

To prove the existence of solution to (4), we employ the diagonalization argument similar to the one presented in [15] (p. 493) for the case of one equation. Take a closed ball $\overline{B_R}$ of radius R . Let $y_r(t, x)$ be the solution to problem (3) in the ball B_{r+1} with $r > R$. Since the Hölder norms $\|y_r\|_{C^{1+\frac{\beta}{2}, 2+\beta}([0, T] \times \overline{B_r})}$ possess a bound not depending on r , then, by Arzelà-Ascoli's theorem, the family of functions $y_r(t, x)$, parametrized by r , is relatively compact in $C^{1,2}([0, T] \times \overline{B_R})$. Therefore, we can find a sequence $\{y_{r_n}\}$ which converges in $C^{1,2}([0, T] \times \overline{B_R})$. We can also find a further subsequence $\{y_{r_n}^{(1)}\}$ that converges in $C^{1,2}([0, T] \times \overline{B_{R+1}})$. Proceeding this way, we find a subsequence $\{y_{r_n}^{(k)}\}$ that converges in $C^{2,1}([0, T] \times \overline{B_{R+k}})$. It remains to note that the diagonal subsequence $\{y_{r_n}^{(n)}\}$ converges at each point of $[0, T] \times \mathbb{R}^d$ to a function $y(t, x)$, while its derivatives $\partial_t y_{r_n}^{(n)}$, $\partial_x y_{r_n}^{(n)}$, and $\partial_{xx}^2 y_{r_n}^{(n)}$ converge to the corresponding derivatives of $y(t, x)$. Clearly, $y(t, x)$ is a solution to (4). Since the $C^{1,2}$ -norm of each function $y_{r_n}^{(n)}(t, x)$ has the same bound, the solution $y(t, x)$ belongs to $C_b^{1,2}([0, T] \times \mathbb{R}^d)$. Moreover, the bound for $y(t, x)$ depends only on \mathcal{C} , T , and the bound for $|\varphi(x)|$; the bound for $\partial_x y(t, x)$ depends on ν , \mathcal{C} , T , and the bounds for $|\varphi(x)|$ and $|\nabla\varphi(x)|$. □

2.2 Examples of the noise process

Example 1: Stochastic integral. $\eta(t, x) = \int_0^t g(s, x) dB_s = \sum_{i=1}^l \int_0^t g_i(s, x) dB_s^i$, where $g_i(t, x) \in C^{0, 2+\beta}([0, T] \times \mathbb{R}^d)$, $\beta \in (0, 1)$, is such that $g_i(t, x) = g_i(t, x)\zeta(x)$ with $\zeta(x)$ being a C^∞ -cutting function of the ball $\mathbb{B}_R = \{x \in \mathbb{R}^d : |x| < R\}$ described in the previous subsection; further, B_t^i are independent real-valued Brownian motions, and the stochastic integral is defined for each $x \in \mathbb{R}^d$.

LEMMA 2.1. *There is a version of the stochastic integral $\int_0^t g(s, x) dB_s$ which belongs to $C^{0,2}([0, T] \times \mathbb{R}^d)$.*

The proof of Lemma 2.1 follows from Kolmogorov's continuity theorem.

Example 2: x -regularized space-time white noise. Let $\dot{W}^i(t, x)$, $i = 1, \dots, d$, be independent space-time white noises, and let $\dot{W}_\varepsilon^i(t, x)$ be a regularization in x of $\dot{W}^i(t, x)$, that is, $\dot{W}_\varepsilon^i(t, x) = (\dot{W}^i(t, \cdot) * \rho_\varepsilon)(x)$, where ρ_ε is a standard mollifier supported on the ball of radius ε . Alternatively, one can write $\dot{W}_\varepsilon^i(t, x) = (W^i(t, \cdot) * \partial_{x_1 \dots x_n}^n \rho_\varepsilon)(x)$, where $W^i(t, x)$ is an $(n+1)$ -parameter Brownian sheet on \mathbb{R}^d . Since we are interested in noises of class $C_b^{0,2}(\mathbb{R}^n)$, define $\dot{\eta}^i(t, x)$ as $\dot{W}_\varepsilon^i(t, x)\zeta(x)$, where $\zeta(x)$ is a C^∞ -cutting function of a bounded domain $D \subset \mathbb{R}^d$.

3. Classical solution to equation (1)

In this section, we will prove the existence of a classical solution to equation (1) under assumptions (A1) and (A2):

- (A1) $\eta(t, x)$ is of class $C_b^{0,2}([0, T] \times \mathbb{R}^d)$;
- (A2) the initial data $\varphi(x)$ is of class $C_b^\beta(\mathbb{R}^d)$, $\beta \in (0, 1)$.

THEOREM 3.1. *Let (A1) and (A2) be fulfilled. Then equation (1) has a solution.*

Proof. Step 1. Transformation to a Burgers-type PDE. The substitution $\hat{y}(t, x) = y(t, x) - \eta(t, x)$ transforms (1) to the following Burgers-type equation

$$(4) \quad \begin{cases} \partial_t \hat{y}(t, x) = \nu \Delta \hat{y}(t, x) - (\hat{y} + \eta, \partial_x) \hat{y}(t, x) + f(t, x, \hat{y}), \\ \hat{y}(0, x) = \varphi(x), \end{cases}$$

where $f(t, x, y) = \nu \Delta \eta(t, x) - \partial_x \eta(t, x) \eta(t, x) - (y, \partial_x) \eta(t, x)$.

Step 2. Mollification. Let φ_m be a mollification of φ , and $\eta_m(t, x)$ be a mollification of $\eta(t, x)$ in (t, x) , both by means of standard mollifiers. Consider the problem

$$(5) \quad \begin{cases} \partial_t \hat{y}(t, x) = \nu \Delta \hat{y}(t, x) - (\hat{y} + \eta_m, \partial_x) \hat{y}(t, x) + f_m(t, x, \hat{y}), \\ \hat{y}(0, x) = \varphi_m(x), \end{cases}$$

where $f_m(t, x, y) = \nu \Delta \eta_m(t, x) - \partial_x \eta_m(t, x) \eta_m(t, x) - (y, \partial_x) \eta_m(t, x)$. By Theorem 2.1, the above problem has a unique $C_b^{1,2}([0, T] \times \mathbb{R}^m)$ -solution $y_m(t, x)$.

Step 3. Uniform boundedness of $y_m(t, x)$. Note that $\bar{y}_m(t, x) = y_m(T - t, x)$ is the unique $C_b^{2,1}([0, T], \mathbb{R}^d)$ -solution to

$$(6) \quad y(t, x) = \varphi_m(x) + \int_t^T [\nu \Delta y(s, x) - (y + \bar{\eta}_m, \partial_x) y(s, x) + \bar{f}_m(s, x, y)] ds,$$

where $\bar{f}_m(t, x, y) = f_m(T - t, x, y)$, $\bar{\eta}_m(t, x) = \eta_m(T - t, x)$.

Consider the forward-backward SDE (FBSDE) associated to (6):

$$(7) \quad \begin{cases} X_t^{\tau, x} = x + \int_\tau^t (Y_s^{\tau, x} + \bar{\eta}_m(s, X_s^{\tau, x})) ds + \sqrt{2\nu} (W_t - W_\tau), \\ Y_t^{\tau, x} = \varphi_m(X_T^{\tau, x}) + \int_t^T \bar{f}_m(s, X_s^{\tau, x}, Y_s^{\tau, x}) ds - \sqrt{2\nu} \int_t^T Z_s^{\tau, x} dW_s. \end{cases}$$

Let $X_t^{\tau, x}$ be a solution to

$$X_t^{\tau, x} = x + \int_\tau^t (\bar{y}_m(s, X_s^{\tau, x}) + \bar{\eta}_m(s, X_s^{\tau, x})) ds + \sqrt{2\nu} (W_t - W_\tau).$$

It is known that (see, e.g. [16])

$$(8) \quad Y_t^{\tau, x, m} = \bar{y}_m(t, X_t^{\tau, x}), \quad Z_t^{\tau, x, m} = \partial_x \bar{y}_m(t, X_t^{\tau, x})$$

is a solution to (7). In what follows, for any function $\phi(t, x)$, $\nabla \phi(t, x)$ will denote $\partial_x \phi(t, x)$ not to confuse with the partial derivative w.r.t. the upper index x in $(X_t^{\tau, x}, Y_t^{\tau, x}, Z_t^{\tau, x})$. Also, for simplicity of notation, sometimes we skip the upper index τ, x .

Itô's formula, applied to the BSDE in (7), implies

$$(9) \quad \mathbb{E}|Y_t^m|^2 \leq \mathbb{E}|\varphi_m(X_T)|^2 + \int_t^T \mathbb{E}|\nu \Delta \bar{\eta}_m(t, X_s) - \nabla \bar{\eta}_m(t, X_s) \bar{\eta}_m(t, X_s)|^2 ds \\ + \int_t^T \mathbb{E}(1 + 2|\nabla \bar{\eta}_m(s, X_s)|) |Y_s^m|^2 ds.$$

Since $\bar{\eta}_m, \nabla \bar{\eta}_m, \Delta \bar{\eta}_m$, and φ_m are bounded uniformly in m , by Gronwall's inequality, $\mathbb{E}|Y_t^{\tau, x, m}|^2$, and, consequently, $\bar{y}_m(\tau, x)$, is bounded over $[0, T] \times \mathbb{R}^d$ uniformly in m .

Step 4. $y_m(t, x)$ contains a pointwise converging subsequence. By Theorem 1.1 (Chapter V) in [15], under (A1) and (A2), $y_m(t, x)$ is of class $C^{\frac{\alpha}{2}, \alpha}([0, T] \times \mathbb{B}_R)$

for any ball $\mathbb{B}_R = \{|x| < R\}$, where the exponent α and the Hölder constant does not depend on m . By the Arzelá-Ascoli theorem, the family $y_m(t, x)$ is relatively compact in $C([0, T], \overline{\mathbb{B}}_R)$. Therefore, there exists a subsequence $y_{m_k}(t, x)$ and a function $y^{(1)}(t, x) \in C([0, T], \overline{\mathbb{B}}_R)$ which is the uniform limit of $y_{m_k}(t, x)$ on $[0, T] \times \overline{\mathbb{B}}_R$. Choosing a further subsequence of $y_{m_k}(t, x)$, we prove that there exists its uniform limit (which we again denote by $y^{(1)}(t, x)$) on $[0, T] \times \overline{\mathbb{B}}_{R+1}$. For every $N \in \mathbb{N}$, in a finite number of steps, we find a subsequence of y_m converging uniformly on $[0, T] \times \overline{\mathbb{B}}_{R+N}$. Its limit function, again, will be denoted by $y^{(1)}(t, x)$. Passing to the diagonal subsequence, we obtain a subsequence converging to $y^{(1)}(t, x)$ pointwise on $[0, T] \times \mathbb{R}^d$ and uniformly on each closed ball \mathbb{B}_R centered at the origin. Remark that since y_m is of class $C^{\frac{\alpha}{2}, \alpha}([0, T] \times \mathbb{B}_R)$, so is $y^{(1)}(t, x)$. Also remark that if R is fixed, by aforementioned Theorem 1.1 (Chapter V) in [15], due to the global boundedness of η_m , $\partial_x \eta_m$, and f_m (uniform in m), the Hölder constant for y_m in a ball $K_R = \{x : |x - a| < R\}$ does not depend on the center a of the ball. This implies that this Hölder constant is uniform in m over \mathbb{R}^d . Therefore, $y^{(1)} \in C^{\frac{\alpha}{2}, \alpha}([0, T] \times \mathbb{R}^d)$.

The pointwise converging (diagonal) subsequence constructed above, we again denote by $\{y_m\}$, i.e., $y^{(1)}(t, x) = \lim_{m \rightarrow \infty} y_m(t, x)$.

Step 5. The limit of y_m is a solution to equation (4). Consider a linear PDE with respect to v :

$$(10) \quad \begin{cases} \partial_t v(t, x) = \nu \Delta v(t, x) - (y^{(1)} + \eta, \partial_x) v(t, x) + f(t, x, y^{(1)}(t, x)), \\ v(0, x) = \varphi(x). \end{cases}$$

By Theorem 12 (Chapter 1, Paragraph §7) in [12],

$$(11) \quad y^{(2)}(t, x) = \int_{\mathbb{R}^d} G(t, x; 0, z) \varphi(z) dz + \int_0^t \int_{\mathbb{R}^d} G(t, x; s, z) f(t, z, y^{(1)}(t, z)) dz ds$$

is a solution to problem (10) which is continuous on $[0, T] \times \mathbb{R}^d$ and is of class $C^{1,2}$ on $(0, T] \times \mathbb{R}^d$, where $G(t, x; s, z)$ is the fundamental solution for the differential operator $\nu \Delta - (y^{(1)} + \eta, \partial_x)$.

We use the associated BSDE to show that $y^{(2)}$ is also a limit of y_m . This will prove that $\hat{y}(t, x) = y^{(2)}(t, x) = y^{(1)}(t, x)$ is a solution to problem (4), and, therefore, $y(t, x) = \hat{y}(t, x) + \eta(t, x)$ is a solution to original equation (1).

Define $B_t^{\tau, x} = x + \sqrt{2\nu}(W_t - W_\tau)$, $\bar{y}^{(1)}(t, x) = y^{(1)}(T - t, x)$, and note (see, e.g., [16]) that

$$Y_t^{\tau, x} = y^{(2)}(T - t, B_t^{\tau, x}), \quad Z_t^{\tau, x} = \partial_x y^{(2)}(T - t, B_t^{\tau, x})$$

satisfies the BSDE

$$(12) \quad \begin{aligned} Y_t^{\tau, x} = y^{(2)}(\varepsilon, B_T^{\tau, x}) &+ \int_t^{T-\varepsilon} [\bar{f}(s, B_s^{\tau, x}, \bar{y}^{(1)}(s, B_s^{\tau, x})) \\ &- Z_s^{\tau, x}(\bar{y}^{(1)}(s, B_s^{\tau, x}) + \bar{\eta}(s, B_s^{\tau, x}))] ds - \sqrt{2\nu} \int_t^{T-\varepsilon} Z_s^{\tau, x} dW_s, \end{aligned}$$

where $\varepsilon > 0$ is a sufficiently small number that appears due to the fact that $y^{(2)}$ may not be differentiable in t at zero. Further, since f and φ in (11) are bounded, then $y^{(2)}$ is also bounded. Indeed, the fundamental solution $G(t, x; s, z)$ possesses estimates by Gaussian densities (see, e.g., [12], Chapter 1, formula (6.12)). This

implies the boundedness of $Y_t^{\tau,x}$, and, therefore, by Itô's formula, there exists a constant $k > 0$, independent of ε , such that

$$\mathbb{E} \int_t^{T-\varepsilon} |Z_s|^2 ds \leq k.$$

Note that the above identity also holds for $\varepsilon = 0$ which allows us to consider (12) for $\varepsilon = 0$. On the other hand,

$$(13) \quad Y_t^{\tau,x,m} = \bar{y}_m(t, B_t^{\tau,x}), \quad Z_t^{\tau,x,m} = \partial_x \bar{y}_m(t, B_t^{\tau,x})$$

is a solution to

$$(14) \quad Y_t^{\tau,x,m} = \varphi_m(B_T^{\tau,x}) + \int_t^T [\bar{f}_m(s, B_s^{\tau,x}, Y_s^{\tau,x,m}) - Z_s^{\tau,x,m}(Y_s^{\tau,x,m} + \bar{\eta}_m(s, B_s^{\tau,x}))] ds - \sqrt{2\nu} \int_t^T Z_s^{\tau,x,m} dW_s,$$

where \bar{y}_m is the solution to (6). Remark that (13) is a different process than (8), introduced earlier. However, we use the same symbols to simplify notation. Since $Y_t^{\tau,x,m}$, $\bar{\eta}_m(t, B_t^{\tau,x})$, and $\bar{y}^{(1)}(t, B_t^{\tau,x})$ are globally bounded by a constant that does not depend on m , by the standard argument involving Itô's formula, from (12) and (14) we obtain

$$(15) \quad \begin{aligned} \mathbb{E}|Y_t^m - Y_t|^2 + 2\nu \mathbb{E} \int_t^T |Z_s^m - Z_s|^2 ds &\leq \mathbb{E}|\varphi(B_T^{\tau,x}) - \varphi_m(B_T^{\tau,x})|^2 \\ &+ \varepsilon \mathbb{E} \int_t^T |Z_s^m - Z_s|^2 ds + K \mathbb{E} \int_t^T |Y_s^m - Y_s|^2 ds \\ &+ \mathbb{E} \int_t^T |Z_s|^2 (|\bar{y}^{(1)} - \bar{y}_m|^2 + |\bar{\eta} - \bar{\eta}_m|^2)(s, B_s^{\tau,x}) ds \\ &+ \mathbb{E} \int_t^T (|\bar{y}^{(1)} - \bar{y}_m|^2 + |\nabla \bar{\eta}_m - \nabla \bar{\eta}|^2 + |\bar{\eta}_m - \bar{\eta}|^2 + |\Delta \eta_m - \Delta \eta|)(s, B_s^{\tau,x}) ds, \end{aligned}$$

where $\varepsilon > 0$ is a number smaller than 2ν and $K > 0$ is a constant. Above, we skipped the upper index τ, x in $Y_t^{m,\tau,x}$, $Z_t^{m,\tau,x}$, $Y_t^{\tau,x}$, and $Z_t^{\tau,x}$ to simplify notation. Note that the first term and the terms in the last two lines on the right-hand side go to zero by Lebesgue's dominated convergence theorem. Therefore, by Gronwall's inequality, $\mathbb{E}|Y_t^m - Y_t|^2 \rightarrow 0$ as $m \rightarrow \infty$, and hence, $\lim_{m \rightarrow \infty} y_m = y^{(2)}$ pointwise on $[0, T] \times \mathbb{R}^d$. This proves that $y^{(1)}(t, x) = y^{(2)}(t, x)$.

Step 6. Uniqueness. Suppose there are two solutions $y_1, y_2 \in C_b^{0,2}((0, T] \times \mathbb{R}^d)$ to problem (1), and let Y_t^1 and Y_t^2 be the solutions to the associated BSDEs of form (12) (with $\varepsilon = 0$):

$$Y_t^i = \varphi(B_T^{\tau,x}) + \int_t^T [\bar{f}(s, B_s^{\tau,x}, Y_s^i) - Z_s^i(Y_s^i + \bar{\eta}(s, B_s^{\tau,x}))] ds - \sqrt{2\nu} \int_t^T Z_s^i dW_s,$$

$i = 1, 2$. The same argument as in Step 3 implies the boundedness of y_1 and y_2 by the same constant. Since $\int_0^T |Z_s^i|^2 ds \leq k$ for some constant k (which follows from Itô's formula), then an estimate similar to (15) implies that $\mathbb{E}|Y_t^1 - Y_t^2|^2 = 0$ by Gronwall's inequality. Hence, $y_1 = y_2$. The theorem is proved. \square

Finally, we prove that the solution to (1) is adapted to the same filtration as $\eta(t, x)$.

THEOREM 3.2. *Let \mathcal{G}_t be a filtration and let $\eta(t, x)$ be \mathcal{G}_t -adapted for each $x \in \mathbb{R}^d$. Then, the solution $y(t, x)$ to (1) is also \mathcal{G}_t -adapted.*

Proof. First, let us prove that if for each (x, y) , $\eta(t, x)$ and $f(t, x, y)$ are \mathcal{G}_t -adapted, so is the solution to (4). We can prove the theorem for a smooth initial data and a smooth noise since according to the proof of Theorem 3.1, the solution to (4) can be approximated pointwise by solutions to (5).

By [15], the central argument in the proof of existence for initial-boundary value problem (3) is the Leray-Schauder theorem. According to the latter, the map $\Gamma : v \mapsto y$ defined by

$$(16) \quad \begin{cases} \partial_t y(t, x) = \nu \Delta y(t, x) - (v(t, x) + \eta(t, x), \nabla) v(t, x) + \zeta(x) f(t, x, v), \\ y(0, x) = \varphi(x) \zeta(x), \quad y(t, x) \Big|_{\partial B_r} = 0, \end{cases}$$

has a fixed point. Here, the cutting function $\zeta(x)$ is defined as in the proof of Theorem 2.1. Hence,

$$(17) \quad y(t, x) = \int_{B_r} G(x, z, t) \varphi(z) \zeta(z) dz + \int_0^t \int_{B_r} G(x, z, t-s) (\zeta(z) f(s, z, v(s, z)) - (v(s, z) + \eta(s, z), \partial_z) v(s, z)) dz ds,$$

where $G(x, z, t)$ is the Green function for the heat equation $\partial_t y(t, x) = \nu \Delta y(t, x)$ on the ball B_r (with zero boundary condition). If $v(t, x)$ is a $C^{0,1}([0, T] \times B_r)$ deterministic function, then $y(t, x) = \Gamma(v)(t, x)$ is clearly \mathcal{G}_t -adapted. Therefore, the fixed point of Γ , which is the solution to (16), is \mathcal{G}_t -adapted. The solution to Cauchy problem (4) is then \mathcal{G}_t -adapted as a pointwise limit (by construction) of solutions to initial-boundary value problems (16).

Observe that since $\eta(t, x)$ is \mathcal{G}_t -adapted for each x , $f(t, x, y)$, defined via $\eta(t, x)$ in Step 1 of the proof of Theorem 3.1, is also \mathcal{G}_t -adapted. The connection between the solutions to problems (4) and (1) implies that the latter is \mathcal{G}_t -adapted. \square

Acknowledgements

The research of the first-named author was partially supported by RFBR Grant N 18-01-00048.

References

- [1] A. BORITCHEV, *Multidimensional Potential Burgers Turbulence*, Communications in Mathematical Physics, vol. 342, issue 2, (2016) pp. 441–489.
- [2] J. BEC AND K. KHANIN, *Burgers turbulence*, Phys. Rep, 447, (2007), pp. 1–66.
- [3] L. BERTINI, N. CANCRINI, G. JONA-LASINIO, *The stochastic Burgers equation*, Comm. Math. Phys. 165 (1994) 211–232.
- [4] Z. BRZEZNIAK, B. GOLDYS, M. NEKLYUDOV, *Multidimensional stochastic Burgers equation*, SIAM JOURNAL of Mathematical Analysis, vol. 46, no. 1, (2014), pp 871–889.
- [5] T. BUCHERT, *Lagrangian theory of gravitational instability of Friedman-Lemaitre cosmologies and the “Zel’dovich approximation”*, MNRAS 254, (1992), pp. 729–737.
- [6] P. COLES, A.L. MELOTT, S. SHANDARIN, *Testing approximations for non-linear gravitational clustering*, Mon. Not. R. Astron. Soc. 260 (1993), pp. 765–766.
- [7] N. ENGLEZOS, N. E. FRANGOS, X-I KARTALA, AND A N. YANNAKOPOULOS, *Stochastic Burgers PDEs with random coefficients and a generalization of the Cole-Hopf transformation*, Stoch. Proc. and their Appl. 123, 8, (2013), pp. 3239–3272

- [8] T. GOTOH, R. H. KRAICHNAN. *Burgers turbulence with large scale forcing*, Phys. Fluids A 10, (1998), pp. 2859–2866.
- [9] S.N. GURBATOV, A.I. SAICHEV, *Probability distribution and spectra of potential hydrodynamic turbulence*, Radiophysics and Quantum Electronics 27(4), (1984), pp. 303–313.
- [10] S. GURBATOV, A. MOSHKOV, A. NOULLEZ. *Evolution of anisotropic structures and turbulence in the multidimensional Burgers equation*, Phys. Rev. E 81(4), (2010), 046312.
- [11] S. GURBATOV, A. SAICHEV, S. SHANDARIN, *Large-scale structure of the Universe. The Zel'dovich approximation and the adhesion model*, Physics-Uspekhi, vol 55, 3, (2012), pp. 223–249.
- [12] A. FRIEDMAN, *Partial differential equations of parabolic type*, Robert E. Krieger publishing company, 1983.
- [13] U. FRISCH, J. BEC *Burgulence*, New trends in turbulence Turbulence: nouveaux aspects, vol. 74 of the series Les Houches - Ecole d'Ete de Physique Theorique, Springer Berlin Heidelberg, (2002), pp. 341–383.
- [14] R. ITURRIAGA, K. KHANIN. *Burgers turbulence and random Lagrangian systems*, Comm. Math. Phys., vol. 232, issue 3, (2003), pp. 377–428.
- [15] O.A. LADYŽENSKAYA; V.A. SOLONNIKOV; N.N. URAL'CEVA, *Linear and quasi-linear equations of parabolic type*, Translations of Mathematical Monographs 23, Providence, RI: American Mathematical Society, pp. XI+648, 1968.
- [16] J. MA, P. PROTTER, J. YONG, *Solving forward backward stochastic differential equations explicitly: a four step scheme*, Probability Theory and Related Fields, 98 (1994), pp. 339–359.
- [17] M. MEERSCHAERT, W. WANG, Y. XIAO, *Fernique-type inequalities and moduli of continuity for anisotropic Gaussian random fields*, Trans Am Math Soc., vol 365, (2012), pp. 1081–1107.
- [18] M. VERGASSOLA, B. DUBRULLE, U. FRISCH, A. NOULLEZ, *Burgers' equation, Devil's staircases and the mass distribution for large-scale structures*, Astro. Astrophys. 289, (1994), pp. 325–356.
- [19] YA. B. ZEL'DOVICH, *Gravitational instability: An approximate theory for large density perturbations*, Astro. Astrophys. 5, (1970), pp. 84–89.

VORONEZH STATE UNIVERSITY, UNIVERSITETSKAYA PL. 1, VORONEZH, RUSSIA
E-mail address: yeg@math.vsu.ru

MATHEMATICS DEPARTMENT, FEDERAL UNIVERSITY OF PARAÍBA, JOÃO PESSOA, BRAZIL
E-mail address: evelina@mat.ufpb.br