

ESSENTIAL DIMENSION, SYMBOL LENGTH AND p -RANK

ADAM CHAPMAN AND KELLY MCKINNIE

ABSTRACT. We prove that the essential dimension of central simple algebras of degree $p^{\ell m}$ and exponent p^m over fields F containing a base-field k of characteristic p is at least $\ell + 1$ when k is perfect. We do this by observing that the p -rank of F bounds the symbol length in $\mathrm{Br}_{p^m}(F)$ and that there exist indecomposable p -algebras of degree $p^{\ell m}$ and exponent p^m . We also prove that the symbol length of the Milne-Kato cohomology group $H_{p^m}^{n+1}(F)$ is bounded from above by $\binom{r}{n}$ where r is the p -rank of the field, and provide upper and lower bounds for the essential dimension of Brauer classes of a given symbol length.

1. INTRODUCTION

Given a field k and a covariant functor $\mathcal{F} : \text{Fields}/k \rightarrow \text{Sets}$, the essential dimension of an object $x \in \mathcal{F}(F)$, denoted $\mathrm{ed}_{\mathcal{F}}(x)$, where F is a field containing k , is the minimal transcendence degree of a field E with $k \subseteq E \subseteq F$ for which there exists $x_0 \in \mathcal{F}(E)$ such that $x = x_0 \otimes_E F$. The essential dimension of the functor, denoted $\mathrm{ed}(\mathcal{F})$, is the supremum on the essential dimension of all the objects $x \in \mathcal{F}(F)$ for all fields $F \supseteq k$. The essential p -dimension of an object $x \in \mathcal{F}$, denoted $\mathrm{ed}_{\mathcal{F}}(x; p)$, is defined to be the minimal $\mathrm{ed}_{\mathcal{F}}(x \otimes L)$ where L ranges over all prime to p field extensions of F . The essential p -dimension of $\mathcal{F}$, denoted $\mathrm{ed}(\mathcal{F}; p)$, is defined to be the supremum on the essential p -dimension of all objects $x \in \mathcal{F}(F)$ for all fields $F \supseteq k$. Note that $\mathrm{ed}_{\mathcal{F}}(x; p) \leq \mathrm{ed}_{\mathcal{F}}(x)$ and $\mathrm{ed}(\mathcal{F}; p) \leq \mathrm{ed}(\mathcal{F})$. See [Mer13] for a comprehensive discussion on these definitions and associated open problems.

Given a prime number p , a field k of $\mathrm{char}(k) = p$ and integers m, n with $n \geq m$, let Alg_{p^n, p^m} denote the functor mapping every field F containing k to the set of isomorphism classes of central simple algebras of degree p^n and exponent dividing p^m over F . The computation of $\mathrm{ed}(\mathrm{Alg}_{p^n, p^m})$ is largely open (see [BM09, McK17, Mer13] for example). It is known from [Bae11] that for fields k of $\mathrm{char}(k) = 2$ we have $\mathrm{ed}(\mathrm{Alg}_{8,2}) \leq 10$, and when $k \supseteq \mathbb{F}_4$ we also have $\mathrm{ed}(\mathrm{Alg}_{4,2}) = 3$. In the same paper it was also proven that for any k , $\mathrm{ed}(\mathrm{Alg}_{p^n, p^m}; p) \geq 3$ when $n > m$. In [McK17] it was shown that when k is algebraically closed, $\ell + 1 \leq \mathrm{ed}(\mathrm{Alg}_{p^{\ell}, p}; p)$ for any ℓ , improving the previous lower bound of 3 when $\ell \geq 2$.

The goal of this paper (Theorem 5.4) is to prove that $\ell + 1 \leq \mathrm{ed}(\mathrm{Alg}_{p^{\ell m}, p^m}; p)$ when k is perfect for any ℓ and m , recovering and extending the result from [McK17]. The techniques in this paper are simpler than [McK17], relying on symbol length bounds and indecomposable division algebras which already exist in the literature, instead of analyzing sums of generic p -symbols. We also provide upper and lower bounds for the essential dimension of Brauer classes (and other Kato-Milne cohomology groups) of a given symbol length.

2010 *Mathematics Subject Classification.* 16K20 (primary); 13A35, 19D45, 20G10 (secondary) .

Key words and phrases. Essential Dimension, Symbol Length, p -rank, Fields of Positive Characteristic, Brauer Group, Central Simple Algebras, Kato-Milne Cohomology.

2. PRELIMINARIES

For a field F of $\text{char}(F) = p$ and positive integers m and n , the Kato-Milne cohomology group $H_{p^m}^{n+1}(F)$ is the additive group $W_m(F) \otimes \underbrace{F^\times \otimes \cdots \otimes F^\times}_{n \text{ times}}$ modulo the relations

- $(\omega^p - \omega) \otimes b_1 \otimes \cdots \otimes b_n = 0$,
- $(0 \dots 0, a, 0, \dots, 0) \otimes a \otimes b_2 \otimes \cdots \otimes b_n = 0$, and
- $\omega \otimes b_1 \otimes \cdots \otimes b_n = 0$ where $b_i = b_j$ for some $i \neq j$,

where $W_m(F)$ is the ring of truncated Witt vectors of length m over F , and for each $\omega = (\omega_1, \dots, \omega_m)$, ω^p stands for $(\omega_1^p, \dots, \omega_m^p)$. For a comprehensive reference on these groups see [AJO18]. The generators $\omega \otimes b_1 \otimes \cdots \otimes b_n$ are called “ (p^m) -symbols”. For $n = 1$, these groups describe the p^m -torsion of the Brauer group, i.e., $H_{p^m}^2(F) \cong \text{Br}_{p^m}(F)$ with the isomorphism given by $\omega \otimes b \mapsto [\omega, b]_F$, where $[\omega, b]_F$ stands for the cyclic algebra generated by $\theta_1, \dots, \theta_m$ and y satisfying

$$\vec{\theta}^p - \vec{\theta} = \omega, \quad y^{p^m} = b, \quad \text{and} \quad y \vec{\theta} y^{-1} = \vec{\theta} + \vec{1}$$

where $\vec{\theta} = (\theta_1, \theta_2, \dots, \theta_m)$ is a truncated Witt vector, $\vec{\theta}^p = (\theta_1^p, \theta_2^p, \dots, \theta_m^p)$, and $\vec{1} = (1, 0, \dots, 0)$ (see [MM91] for reference). The symbol length of a class in $H_{p^m}^{n+1}(F)$ is the minimal t for which the class can be written as the sum of t symbols. In the special case of symbol p -algebras of exponent dividing p^m , (i.e., $H_{p^m}^2(F)$) the symbol length translates into the minimal t for which the algebra is Brauer equivalent to a tensor product of t cyclic algebras of degree p^m . We denote the symbol length of such a Brauer class $[A]$ by $\text{sl}_{p^m}([A])$.

For any $t < m$, there is a shift map from the group $H_{p^t}^{n+1}(F)$ to $H_{p^m}^{n+1}(F)$ given by $(a_1, \dots, a_t) \otimes b_1 \otimes \cdots \otimes b_n \mapsto (0, \dots, 0, a_1, \dots, a_t) \otimes b_1 \otimes \cdots \otimes b_n$. Since the map taking each symbol $\pi \in H_{p^m}^{n+1}(F)$ to $\underbrace{\pi + \cdots + \pi}_{p^{m-1} \text{ times}}$ takes each $(a_1, \dots) \otimes b_1 \otimes \cdots \otimes b_n$ to $(0, \dots, 0, a_1^{p^{m-1}}) \otimes b_1 \otimes \cdots \otimes b_n$ (which is equal to $(0, \dots, 0, a_1) \otimes b_1 \otimes \cdots \otimes b_n$ in this group), it gives rise to another map $\text{Exp} : H_{p^m}^{n+1}(F) \rightarrow H_p^{n+1}(F)$.

Theorem 2.1 ([AJO18, Theorem 2.31]; see also [Izh00, Theorem 1] and [Izh96, Lemma 6.2]). *The following sequence is exact*

$$0 \longrightarrow H_{p^{m-1}}^{n+1}(F) \xrightarrow{\text{Shift}} H_{p^m}^{n+1}(F) \xrightarrow{\text{Exp}} H_p^{n+1}(F) \longrightarrow 0.$$

3. SYMBOL LENGTH AND p -RANK

Proposition 3.1. *Let p be a prime integer, k a field of $\text{char}(k) = p$, and m and ℓ positive integers with $\ell \geq 2$. Exclude the case of $p = \ell = 2$ and $m = 1$. Then there exists a p -algebra A of degree $p^{\ell m}$ and exponent p^m over a field F containing k with $\text{sl}_{p^m}([A_L]) \geq \ell + 1$ for all prime to p field extensions L of F .*

Proof. If the symbol length of A is at most ℓ then A decomposes (as an algebra) as a tensor product of ℓ cyclic algebras of degree p^m . Hence we need only find a field $F \supset k$ and a p -algebra A of degree $p^{\ell m}$ and exponent p^m over F which does not decompose as a tensor product of ℓ cyclic algebras of degree p^m and retains this quality after any prime to p extension. Such an algebra exists by [Kar95] (see the introduction of that paper for explanation for $m > 1$ and Sections 2 and 3 for $m = 1$; the papers [Kar98] and [McK08] also discuss such examples). $\square$

Every field F of $\text{char}(F) = p$ is a vector space over F^p , and if $[F : F^p]$ is finite then it is p^r for some nonnegative integer r , called the “ p -rank of F ”, and denoted $\text{rank}_p(F)$.

Proposition 3.2 (cf. [CM19, Remark 3.1]). *Given a prime integer p , a field F of $\text{char}(F) = p$ and $\text{rank}_p(F) = r < \infty$ and a p -algebra A of exponent p^m over F , we have $\text{sl}_{p^m}([A]) \leq r$.*

Proof. Let $\alpha_1, \dots, \alpha_r$ be a p -basis for F , i.e., F is spanned by $\{\alpha_1^{d_1} \dots \alpha_r^{d_r} : 0 \leq d_1, \dots, d_r \leq p-1\}$ over F^p . Since F is isomorphic to F^p , by induction we get that F is spanned by $\{\alpha_1^{d_1} \dots \alpha_r^{d_r} : 0 \leq d_1, \dots, d_r \leq p^m-1\}$ over F^{p^m} . Since p^m -symbols are split by $K = F[\sqrt[p^m]{\alpha_1}, \dots, \sqrt[p^m]{\alpha_r}]$ (because $K^{p^m} = F$) and $\text{Br}_{p^m}(F)$ is generated by p^m -symbols, every class in $\text{Br}_{p^m}(F)$ is split by restriction to K as well. By [Alb68, Theorem 28] (for a more modern reference see [GS17, Thm. 9.1.1]), each class in $\text{Br}_{p^m}(F)$ decomposes as a tensor product $C_1 \otimes \dots \otimes C_r$ of p^m -symbols where each C_i contains $F[\sqrt[p^m]{\alpha_i}]$, and therefore the symbol length of every class in $\text{Br}_{p^m}(F)$ is at most r . $\square$

In the remainder of the section, we provide another proof for Proposition 3.2, and generalize it to higher cohomology groups.

Lemma 3.3. *Given a p -basis $\alpha_1, \dots, \alpha_r$ for a field F of $\text{char}(F) = p$, $a \in W_m(F)$ and $b_1, \dots, b_n \in F^\times$, there exist $w_i \in W_m(F)$ so that the following equality holds true in $H_{p^m}^{n+1}(F)$:*

$$a \otimes b_1 \otimes \dots \otimes b_n = \sum_{\substack{i=(i_1, \dots, i_n) \\ 1 \leq i_1 < \dots < i_n \leq r}} w_i \otimes \alpha_{i_1} \otimes \dots \otimes \alpha_{i_n}.$$

Proof. We prove it by induction on m . We know it holds for $m = 1$ (see [CM19, Remark 3.1]). Suppose it holds for all positive integers smaller than m . Take now a class π in $H_{p^m}^{n+1}(F)$. The class $\text{Exp}(\pi)$ lives in $H_p^{n+1}(F)$, and so the statement holds true for it, i.e.

$$\text{Exp}(\pi) = \sum_{\substack{i=(i_1, \dots, i_n) \\ 1 \leq i_1 < \dots < i_n \leq r}} c_i \otimes \alpha_{i_1} \otimes \dots \otimes \alpha_{i_n}$$

for some $c_i \in F$. Then by Theorem 2.1 the class π differs from

$$\sum_{\substack{i=(i_1, \dots, i_n) \\ 1 \leq i_1 < \dots < i_n \leq r}} (c_i, 0, \dots, 0) \otimes \alpha_{i_1} \otimes \dots \otimes \alpha_{i_n}$$

by a class π_0 from the embedding of $H_{p^{m-1}}^{n+1}(F)$ into $H_{p^m}^{n+1}(F)$, and so the statement holds true also for π_0 . Then π is a sum of two classes for which the statements holds true, and by adding the Witt vector coefficients we see that it holds true also for π . $\square$

Corollary 3.4. *Let F be a field of characteristic p and finite p -rank r . Then the symbol length of a class in $H_{p^m}^{n+1}(F)$ is at most $\binom{r}{n}$, and in particular the symbol length in $\text{Br}_{p^m}(F)$ is at most r .*

4. OTHER UPPER BOUNDS ON THE SYMBOL LENGTH

As we saw, the p -rank provides a useful bound on the symbol length of classes in $H_{p^m}^{n+1}(F)$. However, in certain cases the p -rank can be infinite, and still one can obtain a reasonable finite upper bound on the symbol length.

Proposition 4.1. *Given finite symbol lengths t of $H_{p^{m+1}}^{n+1}(F)$ and s of $H_p^{n+1}(F)$, the symbol length of $H_{p^{m+1}}^{n+1}(F)$ is at most $t + s$.*

Proof. Consider a class π in $H_{p^{m+1}}^{n+1}(F)$. The class $\text{Exp}(\pi)$ lives in $H_p^{n+1}(F)$, and therefore its symbol length is at most s , i.e. $\text{Exp}(\pi) = a_1 \otimes b_1 + \dots + a_s \otimes b_s$ for some $a_1, \dots, b_s \in F$. Then by Theorem 2.1 π differs from $(a_1, 0, \dots, 0) \otimes b_1 + \dots + (a_s, 0, \dots, 0) \otimes b_s$ by a class

whose exponent divides p^m , so it belongs to the embedding of $H_{p^m}^{n+1}(F)$ into $H_{p^{m+1}}^{n+1}(F)$, and therefore its symbol length is at most t . Altogether, the symbol length of π is at most $t + s$. $\square$

Corollary 4.2. *Given a finite symbol length s of $H_p^{n+1}(F)$, the symbol length of $H_{p^m}^{n+1}(F)$ is at most $m \cdot s$.*

Proof. By induction on m : if the symbol length of $H_{p^m}^{n+1}(F)$ is bounded from above by $m \cdot s$, then by the previous proposition the symbol length of $H_{p^{m+1}}^{n+1}(F)$ is bounded from above by $m \cdot s + s = (m + 1) \cdot s$. $\square$

As proven in [CM19], $\prod_{i=1}^n (\frac{u(F)}{2} - 2^i + 1)$ is an upper bound for the symbol length of $H_2^{n+1}(F)$ when $u(F) < \infty$. Recall that $u(F)$ is the maximal dimension of a nonsingular anisotropic quadratic form over F , and it can be finite even when the 2-rank is infinite (see [MTW91]). As a result, we obtain an upper bound of $m \cdot \prod_{i=1}^n (\frac{u(F)}{2} - 2^i + 1)$ for the symbol length of $H_{2^m}^{n+1}(F)$. Similarly, over the $\widetilde{C}_{p,r}$ -fields F studied in [CM18], for which the symbol length of $H_p^2(F)$ was bounded from above by $p^{r-1} - 1$, the symbol length of $H_{p^m}^2(F)$ is therefore bounded from above by $m \cdot (p^{r-1} - 1)$.

5. THE Br_{p^m} FUNCTOR

Let Br_{p^m} denote the functor mapping each field F containing k to the group $\text{Br}_{p^m}(F)$ of Brauer classes of central simple algebras of exponent dividing p^m over F . For any central simple algebra A , let $[A]$ denote its Brauer class.

Remark 5.1. For a given central simple algebra A of degree p^n and exponent p^m over a field F containing k , we have $\text{ed}_{\text{Br}_{p^m}}([A]) \leq \text{ed}_{\text{Alg}_{p^n, p^m}}(A)$ and $\text{ed}_{\text{Br}_{p^m}}([A]; p) \leq \text{ed}_{\text{Alg}_{p^n, p^m}}(A; p)$. The reason is that for any algebra A_0 of degree p^n and exponent p^m over a field F_0 satisfying $k \subseteq F_0 \subseteq F$ and $A_0 \otimes F = A$, $[A]$ descends to $[A_0]$. It is not necessarily an equality, because $[A]$ may descend to a p^m -torsion Brauer class $[B]$ where B is of degree than p^n .

Lemma 5.3 below forms the basis for our lower bound on the essential dimension.

Lemma 5.2 ([Bou90, Chapter V, Section 16.6, Corollary 3]). *Suppose F is a finitely generated field extension of transcendence degree t of a field k of positive characteristic p and $\text{rank}_p(k) = r$. Then $\text{rank}_p(F) = r + t$.*

Lemma 5.3. *Let k be a field of $\text{char}(k) = p > 0$ and $\text{rank}_p(k) = r$, and A a p -algebra over a field F containing k of exponent p^m , $m \geq 1$. Then $\text{ed}_{\text{Br}_{p^m}}([A]) + r \geq \text{sl}_{p^m}([A])$.*

Proof. Suppose the symbol length is s and the essential dimension is t where $r + t < s$. Then $[A]$ descends to $[A_0] \in \text{Br}_{p^m}(E)$ with $k \subseteq E \subseteq F$ and E finitely generated of finite transcendence degree t over k . By Lemma 5.2 the p -rank of E is $r + t$. By Proposition 3.2 the symbol length of A_0 is at most $r + t$, and so is the symbol length of A , contradiction. $\square$

Theorem 5.4. *Let k be a field of $\text{char}(k) = p > 0$ and $\text{rank}_p(k) = r$, and m and ℓ be positive integers with $\ell \geq 2$. Then $\text{ed}(\text{Alg}_{p^{\ell m}, p^m}; p) \geq \ell + 1 - r$. In particular, when k is perfect, we have $\text{ed}(\text{Alg}_{p^{\ell m}, p^m}; p) \geq \ell + 1$.*

Proof. The case of $p = n = 2$ and $m = 1$ is covered by [Bae11, Corollary 2.2], so we exclude this case in the following discussion. Suppose to the contrary that the essential p -dimension of this functor is at most $\ell - r$. Let A be a central simple algebra of degree $p^{\ell m}$ and exponent p^m over a field F containing k . Then there exists a prime to p extension L/F such that the essential dimension of $[A \otimes L]$ is $\leq \ell - r$, and therefore by Lemma 5.3 the

symbol length of $[A \otimes L]$ is at most ℓ . However, by Proposition 3.1 there exists an algebra A in this category whose symbol length is at least $\ell + 1$ under restriction to any prime to p field extension L of F , contradiction. Therefore $\text{ed}_{\text{Br}_{p^m}}([A]; p) \geq \ell + 1 - r$, which means $\text{ed}_{\text{Alg}_{p^{\ell m}, p^m}}(A; p) \geq \ell + 1 - r$, and as a result $\text{ed}(\text{Alg}_{p^{\ell m}, p^m}; p) \geq \ell + 1 - r$. $\square$

Note that the special case of Theorem 5.4 for $m = 1$ and k algebraically closed was proven in [McK17]. The main two advantages of our approach is that (1) it is much simpler, and (2) it holds true for any field k of finite p -rank, not just algebraically closed fields. If the essential dimension of Alg_{p^r, p^m} is under discussion, the lower bound of $\ell + 1 - r$ is obtained from Theorem 5.4 by taking $\ell = \lfloor \frac{r}{m} \rfloor$ as a result of $\text{ed}(\text{Alg}_{p^r, p^m}) \geq \text{ed}(\text{Alg}_{p^{\ell m}, p^m})$.

6. THE H_p^{n+1} FUNCTOR

Given a field k of $\text{char}(k) = p$, consider the functor H_p^{n+1} mapping each field F containing k to the group $H_p^{n+1}(F)$.

For $1 \leq i \leq \ell$ let $x_{i,1}, \dots, x_{i,m}, y_{i,1}, \dots, y_{i,n}$ be independent indeterminates over k and set $F_{\ell,m,n} = k(x_{1,1}, \dots, y_{\ell,n})$ the rational function field over k in $(m+n)\ell$ indeterminates. Set $A_{\ell,m,n} = \sum_{i=1}^{\ell} (x_{i,1}, \dots, x_{i,m}) \otimes y_{i,1} \otimes \dots \otimes y_{i,n}$. This class is “the generic sum of ℓ symbols in $H_p^{n+1}(F_{\ell,m,n})$ ”. Note that it depends on the choice of k . The following theorem gives a lower bound for the essential dimension of this generic sum of symbols:

Theorem 6.1. *Given a prime integer p , an algebraically closed field k of $\text{char}(k) = p$ and integers $m, n, \ell \geq 1$, the generic sum $A_{\ell,m,n} = \sum_{i=1}^{\ell} (x_{i,1}, \dots, x_{i,m}) \otimes y_{i,1} \otimes \dots \otimes y_{i,n}$ of ℓ symbols in $H_p^{n+1}(F_{\ell,m,n})$ is of $\text{ed}_{H_p^{n+1}}(A_{\ell,m,n}; p) \geq \ell + n$.*

Proof. By [McK17, Theorem 5.8] the generic sum $A_{\ell,1,n}$ of ℓ symbols in $H_p^{n+1}(F_{\ell,1,n})$ has $\text{ed}_{H_p^{n+1}}(A_{\ell,1,n}; p) \geq \ell + n$. Write $B = A_{\ell,1,n} \otimes F_{\ell,m,n}$. By taking k' to be the algebraic closure of $k(x_{i,j} : 1 \leq i \leq \ell, 2 \leq j \leq m)$, the class $B' = B \otimes k'(x_{1,1}, \dots, x_{\ell,1}, y_{1,1}, \dots, y_{\ell,n})$ is the generic sum of ℓ symbols over k' . Note that [McK17, Theorem 5.8] only requires k to be algebraically closed, so one can consider the same functors over k' as well (because it is also algebraically closed), and over k' we have $\text{ed}_{H_p^{n+1}}(B'; p) \geq \ell + n$, and therefore over k we have $\text{ed}_{H_p^{n+1}}(B; p) \geq \ell + n$. Now, the generic sum $A_{\ell,m,n}$ of ℓ symbols in $H_p^{n+1}(F_{\ell,m,n})$ is a pre-image of $A_{\ell,1,n} \otimes F_{\ell,m,n}$ under Exp . If for some prime to p field extension L of $F_{\ell,m,n}$, $A_{\ell,m,n} \otimes L$ descends to π in $H_p^{n+1}(E)$ for a field E of transcendence degree less than $\ell + n$ over k , then $B \otimes L$ descends to $\text{Exp}(\pi)$ which is in $H_p^{n+1}(E)$, contradiction. $\square$

Remark 6.2. The special case of Theorem 6.1 for $n = 1$ coincides with Theorem 5.4 for k algebraically closed. Note that the general case of Theorem 6.1 with $n \geq 1$ requires the use of [McK17] rather than the existence of indecomposable algebras as in Theorem 5.4.

In the rest of the section, we present an upper bound for the symbol length of sums of ℓ symbols in $H_p^{n+1}(F)$ (which can also be the generic sum). For this we recall the definition of $\text{ed}(G)$, for G a finite group, to be the essential dimension of the functor $\mathcal{F} : \text{Fields}/k \rightarrow \text{Sets}$ sending each field F containing k to the set of Galois field extensions K/F with Galois group $\text{Gal}(K/F) = G$. In the following example of Ledet, we have changed exponent p^m to exponent dividing p^m , a trivial generalization of the original statement.

Lemma 6.3 ([Led04, Example, pg. 162]). *Let m and ℓ be positive integers, k a field of $\text{char}(k) = p$ with $|k| \geq p^\ell$ and G a p -group of exponent dividing p^m minimally generated by ℓ elements. Then $\text{ed}(G) \leq m$.*

Remark 6.4. There is a correspondence between finite subgroups of $W_m(F)/\wp(W_m(F))$ and Galois field extensions K/F with a Galois p -group G of exponent dividing p^m : Each class in $W_m(F)/\wp(W_m(F))$ of exponent p^t ($t \leq m$) is represented by a class ω coming from the embedding of $W_t(F)/\wp(W_t(F))$ into $W_m(F)/\wp(W_m(F))$. Written as an element $\omega = (\omega_1, \dots, \omega_t)$ in $W_t(F)/\wp(W_t(F))$, the corresponding field extension is C/F where C is generated over F by $\theta_1, \dots, \theta_t$ subject to the relations

$$(\theta_1, \dots, \theta_t)^p - (\theta_1, \dots, \theta_t) = \omega$$

thinking about $(\theta_1, \dots, \theta_t)$ as a Witt vector in $W_t(C)$ and $(\theta_1, \dots, \theta_t)^p = (\theta_1^p, \dots, \theta_t^p)$ (see the Preliminaries section, and also [Jac89, Chapter 8] and [MM91]). The extension K/F corresponding to G is thus the field compositum of the field extensions corresponding to the classes G consists of. By Lemma 6.3 we thus conclude that given any finite subgroup $G' = \langle \vec{x}_1, \dots, \vec{x}_\ell \rangle$ of $W_m(F)/\wp(W_m(F))$, there exists an intermediate field $k \subseteq E \subseteq F$ of $\text{tr.deg}_k(E) \leq m$ and a subgroup of $W_m(E)/\wp(W_m(E))$ to which G' descends, and in particular, there exist classes $\vec{z}_1, \dots, \vec{z}_\ell$ in $W_m(E)/\wp(W_m(E))$ for which $\vec{z}_i \otimes F \equiv \vec{x}_i \pmod{\wp(W_m(F))}$ for each $i \in \{1, \dots, \ell\}$.

Proposition 6.5. *Let k be a field of $\text{char}(k) = p$ with $|k| \geq p^\ell$, F a field containing k and let π be the sum of ℓ symbols in $H_{p^m}^{n+1}(F)$. Then $\text{ed}_{H_{p^m}^{n+1}}(\pi) \leq m + \ell n$.*

Proof. Let $\pi = \sum_{i=1}^\ell (x_{i,1}, \dots, x_{i,m}) \otimes y_{i,1} \otimes \dots \otimes y_{i,n}$. By Remark 6.4 there exists an intermediate field $k \subseteq E \subseteq F$ and $z_{i,j} \in E$ so that $\vec{x}_i = (x_{i,1}, \dots, x_{i,m}) \in W_m(F)$ satisfies $\vec{x}_i \equiv \vec{z}_i \otimes F \pmod{\wp(W_m(F))}$ and $\text{tr.deg}_k(E) \leq m$. Therefore, π is defined over $E(y_{i,j} : 1 \leq i \leq \ell, 1 \leq j \leq n)$. This field has transcendence degree at most $m + \ell n$. $\square$

Corollary 6.6. *Let k be a field of $\text{char}(k) = p$ with $|k| \geq p^\ell$, F a field containing k and let π be the sum of ℓ symbols in $\text{Br}_{p^m}(F)$. Then $\text{ed}_{\text{Br}_{p^m}}(\pi) \leq m + \ell$.*

Proof. This is the case of $n = 1$ in Proposition 6.5. $\square$

Combining what we have obtained so far, we can outline the bounds as follows:

Corollary 6.7. *For any positive integer m and a class $[A]$ in $\text{Br}_{p^m}(F)$ where $F \supseteq k$ and k is an infinite perfect field of $\text{char}(k) = p$, we have*

$$\text{sl}_{p^m}([A]) \leq \text{ed}_{\text{Br}_{p^m}}([A]) \leq \text{sl}_{p^m}([A]) + m.$$

And in particular, we get the following elegant upper bounds for the essential dimension when k is an infinite perfect field:

Corollary 6.8.

- If A is a p -algebra of degree p^n and exponent p^m over F , then $\text{ed}_{\text{Br}_{p^m}}([A]) \leq p^n + m - 1$.
- If $p = 2$ and A is of degree 8 and exponent 2 over F , then $\text{ed}_{\text{Br}_2}([A]) \leq 5$.

Proof. By [Flo13], the symbol length of A of degree p^n and exponent p is at most $p^n - 1$. By [Row84], the symbol length of A of degree 8 and exponent 2 is at most 4. $\square$

Note that 5 is much better than the upper bound of 10 for $\text{ed}_{\text{Alg}_{8,2}}(A)$ obtained in [Bae11], but one should note that $\text{ed}_{\text{Alg}_{8,2}}(A) \geq \text{ed}_{\text{Br}_2}([A])$ and it is not necessarily an equality.

Remark 6.9. We believe that when k is algebraically closed, the lower bound in Corollary 6.7 can be improved to $\text{sl}_{p^m}([A]) + 1 \leq \text{ed}_{\text{Br}_{p^m}}([A])$. If this is indeed true, that would mean that over fields of transcendence degree d over k , every algebra A of exponent p^m has index $\text{ind}(A) \leq p^{\text{sl}([A])} \leq p^{\text{ed}([A]) - 1} \leq p^{d-1}$, proving an important well-known conjecture on the period-index problem for this particular case. Unfortunately, we do not have a proof yet.

APPENDIX – STANDARD CASE

We conclude with some comments on the case of an algebraically closed field k of characteristic prime to p . Given a field E of transcendence degree r over k , E is a C_r field, and so the symbol length of a central simple algebra A in $\text{Br}_{p^m}(E)$ of degree $p^{\ell m}$ is at most $m(p^{r-1} - 1)$ by [Mat16, Theorem 8.2]. Therefore, if we start with a central simple algebra A of degree $p^{\ell m}$ and exponent p^m over a field F containing k whose essential dimension is r , then $\text{sl}_{p^m}([A]) \leq m(p^{r-1} - 1)$ (a formula which was already implicitly obtained in [Mat16, Section 5]). By solving for r , we obtain the formula

$$\text{ed}_{\text{Alg}_{p^{\ell m}, p^m}}(A) \geq \text{ed}_{\text{Br}_{p^m}}([A]) \geq 1 + \log_p \left(\frac{\text{sl}_{p^m}([A])}{m} + 1 \right).$$

Excluding the case of $p = m = \ell = 2$, the existence of indecomposable algebras A of degree $p^{\ell m}$ and exponent p^m provides algebras A of symbol length at least $\ell + 1$, which gives the bound

$$\text{ed}(\text{Alg}_{p^{\ell m}, p^m}) \geq 1 + \log_p \left(\frac{\ell + 1}{m} + 1 \right).$$

This lower bound is by no means as good as the known bounds in the literature (see [BM12]), but it is possible that this technique could lead to a better bound if we found a way of constructing indecomposable algebras of sufficiently large prescribed symbol length, or if the upper bound for the symbol length for C_r -fields from [Mat16, Theorem 8.2] could be improved (for example to the conjectured bound of r).

ACKNOWLEDGEMENTS

We thank Skip Garibaldi, Alexander Merkurjev and Jean-Pierre Tignol for their comments on the manuscript. We also thank the anonymous referee for the careful reading of the manuscript and the helpful comments.

REFERENCES

- [AJO18] Roberto Aravire, Bill Jacob, and Manuel O’Ryan. The de Rham Witt complex, cohomological kernels and p^m -extensions in characteristic p . *J. Pure Appl. Algebra*, 222(12):3891–3945, 2018.
- [Alb68] A. Albert. *Structure of Algebras*, volume 24 of *Colloquium Publications*. American Math. Soc., 1968.
- [Bae11] Sanghoon Baek. Essential dimension of simple algebras in positive characteristic. *C. R. Math. Acad. Sci. Paris*, 349(7-8):375–378, 2011.
- [BM09] Sanghoon Baek and Alexander Merkurjev. Invariants of simple algebras. *Manuscripta Math.*, 129(4):409–421, 2009.
- [BM12] Sanghoon Baek and Alexander S. Merkurjev. Essential dimension of central simple algebras. *Acta Math.*, 209(1):1–27, 2012.
- [Bou90] N. Bourbaki. *Algebra. II. Chapters 4–7*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1990. Translated from the French by P. M. Cohn and J. Howie.
- [CM18] Adam Chapman and Kelly McKinnie. Kato-Milne cohomology and polynomial forms. *J. Pure Appl. Algebra*, 222(11):3547–3559, 2018.
- [CM19] Adam Chapman and Kelly McKinnie. The u^n -invariant and the symbol length of $H_2^n(F)$. *Proc. Amer. Math. Soc.*, 147(2):513–521, 2019.
- [Flo13] Mathieu Florence. On the symbol length of p -algebras. *Compos. Math.*, 149(8):1353–1363, 2013.
- [GS17] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*, volume 165 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2017. Second edition of [MR2266528].
- [Izh96] O. T. Izhboldin. On the cohomology groups of the field of rational functions. In *Mathematics in St. Petersburg*, volume 174 of *Amer. Math. Soc. Transl. Ser. 2*, pages 21–44. Amer. Math. Soc., Providence, RI, 1996.

- [Izh00] Oleg Izhboldin. p -primary part of the Milnor K -groups and Galois cohomologies of fields of characteristic p . In *Invitation to higher local fields (Münster, 1999)*, volume 3 of *Geom. Topol. Monogr.*, pages 19–41. Geom. Topol. Publ., Coventry, 2000. With an appendix by Masato Kurihara and Ivan Fesenko.
- [Jac89] Nathan Jacobson. *Basic algebra. II*. W. H. Freeman and Company, New York, second edition, 1989.
- [Kar95] Nikita A. Karpenko. Torsion in CH^2 of Severi-Brauer varieties and indecomposability of generic algebras. *Manuscripta Math.*, 88(1):109–117, 1995.
- [Kar98] Nikita A. Karpenko. Codimension 2 cycles on Severi-Brauer varieties. *K-Theory*, 13(4):305–330, 1998.
- [Led04] Arne Ledet. On the essential dimension of p -groups. In *Galois theory and modular forms*, volume 11 of *Dev. Math.*, pages 159–172. Kluwer Acad. Publ., Boston, MA, 2004.
- [Mat16] Eliyahu Matzri. Symbol length in the Brauer group of a field. *Trans. Amer. Math. Soc.*, 368(1):413–427, 2016.
- [McK08] Kelly McKinnie. Indecomposable p -algebras and Galois subfields in generic abelian crossed products. *J. Algebra*, 320(5):1887–1907, 2008.
- [McK17] Kelly McKinnie. Essential dimension of generic symbols in characteristic p . *Forum Math. Sigma*, 5:e14, 30, 2017.
- [Mer13] Alexander S. Merkurjev. Essential dimension: a survey. *Transform. Groups*, 18(2):415–481, 2013.
- [MM91] P. Mammone and A. Merkurjev. On the corestriction of p^n -symbol. *Israel J. Math.*, 76(1-2):73–79, 1991.
- [MTW91] P. Mammone, J.-P. Tignol, and A. Wadsworth. Fields of characteristic 2 with prescribed u -invariants. *Math. Ann.*, 290(1):109–128, 1991.
- [Row84] Louis H. Rowen. Division algebras of exponent 2 and characteristic 2. *J. Algebra*, 90(1):71–83, 1984.

ADAM CHAPMAN, DEPARTMENT OF COMPUTER SCIENCE, TEL-HAI COLLEGE, UPPER GALILEE, 12208 ISRAEL
E-mail address: adam1chapman@yahoo.com

KELLY MCKINNIE DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MONTANA, MISSOULA, MT 59812, USA
E-mail address: kelly.mckinnie@mso.umt.edu