Piecewise autoregression for general integer-valued time series

November 5, 2019

Mamadou Lamine DIOPa, b 1 and William KENGNEa 2

THEMA, Université de Cergy-Pontoise, 33 Boulevard du Port, 95011 Cergy-Pontoise Cedex, France
 LERSTAD, Université Gaston Berger, BP 234 Saint-Louis, Sénégal
 E-mail: mamadou-lamine.diop@u-cergy.fr; william.kengne@u-cergy.fr

Abstract: This paper proposes a piecewise autoregression for general integer-valued time series. The conditional mean of the process depends on a parameter which is piecewise constant over time. We derive an inference procedure based on a penalized contrast that is constructed from the Poisson quasi-maximum likelihood of the model. The consistency of the proposed estimator is established. From practical applications, we derive a data-driven procedure based on the slope heuristic to calibrate the penalty term of the contrast; and the implementation is carried out through the dynamic programming algorithm, which leads to a procedure of $\mathcal{O}(n^2)$ time complexity. Some simulation results are provided, as well as the applications to the US recession data and the number of trades in the stock of Technofirst.

Keywords: Multiple change-points, model selection, integer-valued time series, Poisson quasi-maximum likelihood, penalized quasi-likelihood, slope heuristic.

1 Introduction

We consider a \mathbb{N}_0 -valued ($\mathbb{N}_0 = \mathbb{N} \cup \{0\}$) process $Y = \{Y_t, t \in \mathbb{Z}\}$ where the conditional mean

$$\lambda_t = \lambda_t(\theta_t^*) = \mathbb{E}(Y_t | \mathcal{F}_{t-1}) \tag{1.1}$$

is a function (see below) of the whole information \mathcal{F}_{t-1} up to time t-1 and of an unknown parameter θ_t^* belongs to a compact subset $\Theta \subset \mathbb{R}^d$ $(d \in \mathbb{N})$. The inference in the cases where $\theta_t^* = \theta^*$ is constant or the distribution of $Y_t | \mathcal{F}_{t-1}$ is known have been studied by many authors in several directions; see for instance Fokianos et al. (2009), Fokianos and Tjøstheim (2011, 2012), Douc et al. (2017) among others, for some recent works. We consider here a more general setting where θ_t^* is piecewise constant and the distribution of $Y_t | \mathcal{F}_{t-1}$ is unknown.

¹Supported by the Institute for advanced studies - IAS (Université de Cergy-Pontoise, France), the MME-DII center of excellence (ANR-11-LABEX-0023-01) and by the CEA-MITIC (Université Gaston Berger, Sénégal)

²Developed within the ANR BREAKRISK: ANR-17-CE26-0001-01

We consider the observations Y_1, \dots, Y_n generated as in model (1.1) and assume that the parameter θ_t^* is piecewise constant. Assume that $\exists K^* \in \mathbb{N}, \underline{\theta}^* = (\theta_1^*, \dots, \theta_{K^*}^*) \in \Theta^{K^*}$ and $0 < t_1^* < \dots < t_{K^*-1}^* < n$ such that, $\{Y_t, t_{j-1}^* < t \le t_j^*\}$ is a trajectory of the process $\{Y_{t,j}, t \in \mathbb{Z}\}$ (see Section 2) satisfying:

$$\mathbb{E}(Y_{t,j}|\mathcal{F}_{t-1}) = f(Y_{t-1,j}, Y_{t-2,j}, \dots; \theta_i^*), \ \forall \ t_{j-1}^* < t \le t_i^*$$
(1.2)

where $\mathcal{F}_t = \sigma(Y_{s,j}, s \leq t, j = 1, \cdots, K^* - 1)$ is the σ -field generated by the whole information up to time t and f a measurable non-negative function assumed to be known up to the parameter θ_t^* . K^* is the number of segments (or regimes) of the model; the jth segment corresponds to $\{t_{i-1}^*+1,t_{i-1}^*+2,\cdots,t_i^*\}$ and depends on the parameter θ_i^* . $t_1^*, \dots, t_{K^*-1}^*$ are the change-point locations; by convention, $t_0^* = -\infty$ and $t_{K^*}^* = \infty$. To ensure identifiability of the change-point locations, it is reasonable to assume that $\theta_j^* \neq \theta_{j+1}^*$ for $j=1,\cdots,K^*-1$. The case $K^*=1$ corresponds to the model without change. In the sequel, we assume that the random variables Y_t , $t \in \mathbb{Z}$ have the same (up to the parameter θ_t^*) distribution P and denote by $P(\lambda_t(\theta_t^*))$ the distribution of $Y_t|\mathcal{F}_{t-1}$. Our main focus of interest is the estimation of the unknown parameters $(K^*, (t_i^*)_{1 \le j \le K^*-1}, (\theta_i^*)_{1 \le j \le K^*})$ in the model (1.2). This can be viewed as a classical model selection problem. Assume that the observations Y_1, \dots, Y_n are generated from (1.2). Let K_{max} be the upper bound of the number of segments (note that $K_{\text{max}} < n$). Denote by \mathcal{M}_n the set of partitions of [1, n] into at most K_{max} contiguous segments. Set $m = \{T_1, \dots, T_K\}$ a generic element of K segments in \mathcal{M}_n . Consider the collection $\{S_m, m \in \mathcal{M}_n\}$ where, for a given $m \in \mathcal{M}_n$, S_m is the families of sequence (θ_t) which are piecewise constant on the partition m. Any $\theta = (\theta_t) \in \mathcal{S}_m$ depends on the parameter $\underline{\theta} = (\theta_1, \dots, \theta_K)$ which is the piecewise values of θ_t on each segment. Set $\mathcal{S} = \bigcup_{m \in \mathcal{M}_n} \mathcal{S}_m$. Denote by θ a generic element of \mathcal{S} , with partition m and parameter $\underline{\theta}$. $|\underline{\theta}| = K$ denote the number of the piecewise segment, also called the dimension of ϑ . The true model ϑ^* with dimension K^* , depends on a partition m^* and the parameter θ^* .

For any $\vartheta \in \mathcal{S}$, set $\lambda_t^{\vartheta} = \sum_{k=1}^K \lambda_t(\theta_k) \mathbb{1}_{t \in T_k}$ and denote by $P(\lambda_t^{\vartheta})$ the distribution of $Y_t | \mathcal{F}_{t-1}, \vartheta$; let $p(\cdot | \mathcal{F}_{t-1}, \vartheta) = p(\cdot; \lambda_t^{\vartheta})$ be the probability density function of this distribution. For $\vartheta \in \mathcal{S}$, let $P_{n,\vartheta}$ be the conditional distribution of $(Y_1, \dots, Y_n) | \mathcal{F}_{n-1}, \vartheta$. We consider the log-likelihood contrast: $\forall \vartheta \in \mathcal{S}$,

$$\gamma_n(\vartheta) := \gamma_n(P_{n,\vartheta}) = -\log P_{n,\vartheta}(Y_1, \dots, Y_n) = -\sum_{t=1}^n \log p(Y_t | \mathcal{F}_{t-1}, \vartheta) = -\sum_{t=1}^n \log p(Y_t; \lambda_t^{\vartheta}).$$

Thus, the minimal contrast estimator $\widehat{\vartheta}_m$ of ϑ^* on the collection \mathcal{S}_m is obtained by minimizing the contrast $\gamma_n(\vartheta)$ over $\vartheta \in \mathcal{S}_m$; that is $\widehat{\vartheta}_m = \underset{\vartheta \in \mathcal{S}_m}{\operatorname{argmin}} \ \gamma_n(\vartheta)$. The main approaches of the model selection procedures take into account the model complexity and select the estimator $\widehat{\vartheta}_{m_n}$ such that, m_n minimizes the penalized criterion

$$\operatorname{crit}_n(m) = \gamma_n(\widehat{\vartheta}_m) + \operatorname{pen}_n(m), \text{ for all } m \in \mathcal{M}_n$$
 (1.3)

where $pen_n: \mathcal{M}_n \to \mathbb{R}_+$ is a penalty function, possibly data-dependent. We now address the following issues.

(i) **Semi-parametric setting**. Cleynen and Lebarbier (2014 and 2017) recently consider the change-point type problem (1.2) with i.i.d. observations; in their works, the distribution P is assumed to be known and could be Poisson, Negative binomial or belongs to the exponential family distribution. From the practical viewpoint, we consider the case where P is unknown and deal with the Poisson quasi-likelihood (see for instance Ahmad and Francq (2016)). So in the sequel, γ_n is the Poisson quasi-likelihood contrast and $\widehat{\vartheta}_m$ is the Poisson quasi-maximum likelihood estimator (PQMLE).

(ii) Multiple change-point problem from a non-asymptotic point of view. This question is tacked by model selection approach. Numerous works have been devoted to this issue; see among others, Lebarbier (2005), Arlot and Massart (2009), Cleynen and Lebarbier (2014 and 2017), Arlot and Celisse (2016). In this (quasi)log-likelihood framework, it is more usual to consider the Kullback-Leibler risk. For any $\vartheta \in \mathcal{S}$, the Kullback-Leibler divergence between P_{n,ϑ^*} and $P_{n,\vartheta}$ is

$$KL(\vartheta^*, \vartheta) := KL(P_{n,\vartheta^*}, P_{n,\vartheta}) = \mathbb{E}\Big[\log \frac{P_{n,\vartheta^*}(Y_1, \dots, Y_n)}{P_{n,\vartheta}(Y_1, \dots, Y_n)}\Big] = \sum_{t=1}^n \mathbb{E}\Big[\log \frac{p(Y_t | \mathcal{F}_{t-1}, \vartheta^*)}{p(Y_t | \mathcal{F}_{t-1}, \vartheta)}\Big]$$
$$= \sum_{t=1}^n \mathbb{E}\Big[\log p(Y_t; \lambda_t^{\vartheta^*})\Big] - \sum_{t=1}^n \mathbb{E}\Big[\log p(Y_t; \lambda_t^{\vartheta})\Big]$$

where \mathbb{E} denotes the expectation with respect to the true distribution of the observations. In the case where γ_n is the likelihood contrast, we get $KL(\vartheta^*,\vartheta) = \mathbb{E}[\gamma_n(\vartheta) - \gamma_n(\vartheta^*)]$. The "ideal" partition $m(\vartheta^*)$ (the one whose estimator is closest to ϑ^* according to the Kullback-Leibler risk) satisfying:

$$m(\vartheta^*) = \underset{m \in \mathcal{M}_n}{\operatorname{argmin}} \mathbb{E}[KL(\vartheta^*, \widehat{\vartheta}_m)].$$

The corresponding estimator, $\widehat{\vartheta}_{m(\vartheta^*)}$ called the *oracle*, depends on the true sample distribution, and cannot be computed in practice. The goal is to calibrate the penalty term, such that the segmentation \widehat{m} provides an estimator $\widehat{\vartheta}_{\widehat{m}}$ where the risk of $\widehat{\vartheta}_{\widehat{m}}$ is close as possible to the risk of the *oracle*, namely such that

$$\mathbb{E}[KL(\vartheta^*, \widehat{\vartheta}_{\widehat{m}})] \le C \, \mathbb{E}[KL(\vartheta^*, \widehat{\vartheta}_{m(\vartheta^*)})] \tag{1.4}$$

for a nonnegative constant C, expected close to 1. This issue is addressed in the above mentioned papers, and the results obtained are heavily relied on the independence of the observations. In our setting here, it seems to be a more difficult task. But, we believe that the coupling method can be used as in Lerasle (2011) to overcome this difficulty. We leave this question as the topic of a different research project.

(iii) Multiple change-point problem from an asymptotic point of view. The aim here is to consistently estimate the parameters of the change-point model. This issue has been addressed by several authors using the classical contrast/criteria optimization or binary/sequential segmentation/estimaion; see for instance Bai and Perron (1998), Davis et al. (2008), Harchaoui and Lévy-Leduc (2010), Bardet et al. (2012), Davis and Yau (2013), Davis et al. (2016), Yau and Zhao (2016), Inclan and Tiao (1994), Bai (1997), Fryzlewicz and Subba Rao (2014), Fryzlewicz (2014), among others, for some advanced towards this issue. These works and many other papers in the literature on the asymptotic study of multiple change-point problem are often focussed on continuous valued time series; moreover, the case of a large class of semi-parametric model for discrete-valued time series (such as those discussed earlier) have not yet addressed.

We consider (1.2) and derive a penalized contrast of type (1.3). We assume that there exists a partition $\underline{\tau}^*$ of [0,1] such that $[\underline{\tau}^*n] = m^*$, where $[\underline{\tau}^*n]$ is the corresponding partition of $[\![1,n]\!]$ obtained from $\underline{\tau}^*$. We provide sufficient conditions on the penalty pen_n, for which the estimators \widehat{m} and $\widehat{\vartheta}_{\widehat{m}}$ are consistent; that is:

$$\left(|\widehat{m}|, \frac{\widehat{m}}{n}, \widehat{\vartheta}_{\widehat{m}}\right) \xrightarrow[n \to \infty]{\mathcal{P}} \left(K^*, \underline{\tau}^*, \vartheta^*\right)$$

where $\frac{\widehat{m}}{n}$ is the corresponding partition of [0, 1] obtained from \widehat{m} .

The paper is organized as follows. In Section 2, we set some notations, assumptions and define the Poisson QMLE. In Section 3, we derive the estimation procedure and provide the main results. Some simulations results are displayed in Section 4 whereas Section 5 focus on applications on the US recession data and the daily number of trades in the stock of Technofirst. Section 6 provides the proofs of the main results.

2 Notations and Poisson QMLE

We set the following classical Lipschitz-type condition on the function f.

Assumption A_i(Θ) (i=0,1,2): For any $y \in \mathbb{N}_0^{\mathbb{N}}$, the function $\theta \mapsto f(y;\theta)$ is i times continuously differentiable on Θ and there exists a sequence of non-negative real numbers $(\alpha_k^{(i)})_{k\geq 1}$ satisfying $\sum_{k=1}^{\infty} \alpha_k^{(0)} < 1$ (or $\sum_{k=1}^{\infty} \alpha_k^{(i)} < \infty$ for i=1,2); such that for any $y,y' \in \mathbb{N}_0^{\mathbb{N}}$,

$$\sup_{\theta \in \Theta} \left| \frac{\partial^i f(y; \theta)}{\theta^i} - \frac{\partial^i f(y'; \theta)}{\theta^i} \right| \le \sum_{k=1}^{\infty} \alpha_k^{(i)} |y_k - y_k'|.$$

In the whole paper, it is assumed that for $j=1,\cdots,K^*$, there exists a stationary and ergodic process $\{Y_{t,j},t\in\mathbb{Z}\}$ satisfying

$$\mathbb{E}(Y_{t,j}|\mathcal{F}_{t-1,j}) = f(Y_{t-1,j}, Y_{t-2,j}, \dots; \theta_i^*), \ \forall \ t \in \mathbb{Z}$$

where $\mathcal{F}_{t,j} = \sigma(Y_{s,j}, s \leq t)$ is the σ -field generated by $\{Y_{s,j}, s \leq t\}$; and

$$\exists C > 0, \epsilon > 1$$
, such that $\forall t \in \mathbb{Z}$, $\mathbb{E}Y_{t,i}^{1+\epsilon} < C$. (2.2)

 $\{Y_{t,j}, t \in \mathbb{Z}\}$ is a stationary solution of the jth regime. The focus process $Y = \{Y_t, t \in \mathbb{Z}\}$ is modelled by these stationary regimes; that is, for any $j = 1, \dots, K^*$, $\{Y_t, t_{j-1}^* < t \le t_j^*\}$ is a trajectory of the process $\{Y_{t,j}, t \in \mathbb{Z}\}$.

Ahmad and Francq [1] (Section 3) have discussed about the stationarity and ergodicity issues. In many classical integer-valued time series, the assumption $\mathbf{A}_0(\Theta)$ is enough to enable the existence of a stationary and ergodic process satisfying (2.1).

2.1 Notations

Assume that a trajectory (Y_1, \dots, Y_n) of Y is observed; with $0 < t_1^* < \dots < t_{K^*-1}^* < n$. By convention $t_0^* = -\infty$ and $t_{K^*}^* = \infty$. We will use the following notations.

- For any finite set A, |A| denote the cardinality of A.
- For $a, b \in \mathbb{R}$ (with $a \leq b$), $[a, b] = \mathbb{N} \cap [a, b]$ is the set of integers between a and b.
- For $K \in \mathbb{N}$, $\mathcal{M}_n(K) = \{\underline{t} = (t_1, \dots, t_{K-1}) ; 0 < t_1 < \dots < t_{K-1} < n\}$; in particular, $\underline{t}^* = (t_1^*, \dots, t_{K^*-1}^*) \in \mathcal{M}_n(K^*)$ is the true vector of the locations of breaks. When K = 1, $\mathcal{M}_n(1)$ corresponds to the model with no break.

In the sequel, any configuration $\underline{t} = (t_1, \dots, t_{K-1}) \in \mathcal{M}_n(K)$ is also used as a partition $\{T_1, T_2, \dots, T_K\}$ of [1, n] into K contiguous segments, where $T_1 = \{1, \dots, t_1\}$, $T_j = \{t_{j-1} + 1, \dots, t_j\}$ for $j = 2, \dots, K-1$, $T_K = \{t_{K-1} + 1, \dots, n\}$. In particular, $T_1^* = \{1, \dots, t_1^*\}$, $T_j^* = \{t_{j-1}^* + 1, \dots, t_j^*\}$ for $j = 2, \dots, K^* - 1$,

 $T_{K^*} = \{t_{K^*-1} + 1, \dots, n\}$. $\mathcal{M}_n(K)$ corresponds to the set of partitions of $[\![1, n]\!]$ into K contiguous segments.

- For $K \in \mathbb{N}^*$ and $\underline{t} \in \mathcal{M}_n(K)$ fixed, we set $n_k = |T_k|$ for $1 \le k \le K$. In particular $n_j^* = |T_j^*|$ for $1 \le j \le K^*$. For $1 \le k \le K$ and $1 \le j \le K^*$, let $n_{k,j} = |T_j^* \cap T_k|$.
- Let $\underline{\theta}^* = (\theta_1^*, \dots, \theta_{K^*}^*) \in \Theta^{K^*}$ be the vector of the true parameters of the model (1.2).

Throughout the sequel, the following norms will be used:

- $||x|| := \sqrt{\sum_{i=1}^p |x_i|^2}$ for any $x \in \mathbb{R}^p$;
- $||f||_{\Theta} := \sup_{\theta \in \Theta} (||f(\theta)||)$ for any function $f : \Theta \longrightarrow \mathbb{R}^{d'}$;
- for $x = (x_1, \dots, x_K) \in \mathbb{R}^K$, $||x||_m = \max_{1 \le i \le K} |x_i|$;
- if Y is a random vector with finite r-order moments, we set $\|Y_t\|_r = \mathbb{E}(\|Y\|^r)^{1/r}$.

2.2 Poisson QMLE

Let (Y_1, \ldots, Y_n) be a trajectory generated from the model (1.2). Since the conditional distribution is assumed to be unknown, the likelihood of the model is unknown. The estimation procedure of the parameters θ_j^* is based on the Poisson quasi-maximum likelihood introduced by Ahmad and Francq (2016). The conditional Poisson (quasi)log-likelihood of the model (1.2) computed on a segment $T \subset \{1, \ldots, n\}$ is given (up to a constant) by

$$\widehat{L}_n(T,\theta) := \sum_{t \in T} (Y_t \log \widehat{\lambda}_t(\theta) - \widehat{\lambda}_t(\theta)) = \sum_{t \in T} \widehat{\ell}_t(\theta) \text{ with } \widehat{\ell}_t(\theta) = Y_t \log \widehat{\lambda}_t(\theta) - \widehat{\lambda}_t(\theta)$$
 (2.3)

where $\widehat{\lambda}_t(\theta) = \widehat{f}_t^{\theta} = f(Y_{t-1}, \dots, Y_1, 0, \dots, 0; \theta).$

According to (2.3), the Poisson quasi-likelihood estimator (PQMLE) of θ_j^* computed on T is defined by

$$\widehat{\theta}_n(T) := \underset{\theta \in \Theta}{\operatorname{argmax}} (\widehat{L}_n(T, \theta)). \tag{2.4}$$

Now, for $j = 1, \dots, K^*$, define the Poisson (quasi)log-likelihood of the jth regime by

$$L_{n,j}(T_j^*, \theta) := \sum_{t \in T_j^*} (Y_{t,j} \log \lambda_{t,j}(\theta) - \lambda_{t,j}(\theta)) = \sum_{t \in T_j^*} \ell_{t,j}(\theta) \text{ with } \ell_{t,j}(\theta) = Y_{t,j} \log \lambda_{t,j}(\theta) - \lambda_{t,j}(\theta)$$

where $\lambda_{t,j}(\theta) = f_{t,j}^{\theta} = f(Y_{t-1,j}, Y_{t-2,j}, \dots; \theta)$. It can be approximated by

$$\widehat{L}_{n,j}(T_j^*, \theta) := \sum_{t \in T_j^*} (Y_{t,j} \log \widehat{\lambda}_{t,j}(\theta) - \widehat{\lambda}_{t,j}(\theta)) = \sum_{t \in T_j^*} \widehat{\ell}_{t,j}(\theta) \quad \text{with } \widehat{\ell}_{t,j}(\theta) = Y_{t,j} \log \widehat{\lambda}_{t,j}(\theta) - \widehat{\lambda}_{t,j}(\theta)$$
 (2.5)

where $\widehat{\lambda}_{t,j}(\theta) = \widehat{f}_{t,j}^{\theta} = f(Y_{t-1,j}, \dots, Y_{t_{j-1}^*+1,j}, 0 \dots 0; \theta).$

According to (2.5), the PQMLE of θ_i^* computed on T_i^* is defined by

$$\widetilde{\theta}_n(T_j^*) := \underset{\theta \in \Theta}{\operatorname{argmax}}(\widehat{L}_{n,j}(T_j^*, \theta)).$$
 (2.6)

To avoid problems of parameter identifiability and to study asymptotic normality of the PQMLE, we shall assume:

 $\begin{aligned} & (\mathbf{A0}) \colon \text{ for all } (\theta,\theta') \in \Theta^2, \ \left(f(Y_{t-1},Y_{t-2},\cdots;\theta) = f(Y_{t-1},Y_{t-2},\cdots;\theta') \text{ a.s. for some } t \in \mathbb{N} \right) \Rightarrow \ \theta = \theta' \ ; \\ & \text{moreover, } \exists \underline{c} > 0 \text{ such that } \inf_{\theta \in \Theta} f(y_1,y_2,\cdots;\theta) \geq \underline{c}, \text{ for all } y \in \mathbb{N}_0^{\mathbb{N}}. \end{aligned}$

In order to ensure the consistency and asymptotic normality of the PQMLE, we set the following assumptions for each segment $j = 1, \dots, K^*$ (see also [1]):

(A1): θ_i^* is an interior point of $\Theta \subset \mathbb{R}^d$;

(**A2**):
$$a_{t,j} \longrightarrow 0$$
 and $Y_{t,j}a_{t,j} \longrightarrow 0$ as $t \to \infty$, where $a_{t,j} = \sup_{\theta \in \Theta} \left| \widehat{\lambda}_{t,j}(\theta) - \lambda_{t,j}(\theta) \right|$;

$$(\mathbf{A3}): \ J_j = \mathbb{E}\left[\frac{1}{\lambda_{t,j}(\theta_j^*)} \frac{\partial \lambda_{t,j}(\theta_j^*)}{\partial \theta} \frac{\partial \lambda_{t,j}(\theta_j^*)}{\partial \theta'}\right] < \infty \ \text{ and } \ I_j = \mathbb{E}\left[\frac{\operatorname{Var}(Y_{t,j}|\mathcal{F}_{t-1})}{\lambda_{t,j}^2(\theta_j^*)} \frac{\partial \lambda_{t,j}(\theta_j^*)}{\partial \theta} \frac{\partial \lambda_{t,j}(\theta_j^*)}{\partial \theta'}\right] < \infty;$$

(**A4**): for all
$$c' \in \mathbb{R}$$
, $c' \frac{\partial \lambda_{t,j}(\theta_j^*)}{\partial \theta} = 0$ a.s $\Rightarrow c' = 0$;

(A5): there exists a neighborhood $V(\theta_j^*)$ of θ_j^* such that: for all $i, k \in \{1, \dots, d\}$,

$$\mathbb{E}\left[\sup_{\theta\in V(\theta_j^*)}\left|\frac{\partial^2}{\partial\theta_i\partial\theta_k}\ell_{t,j}(\theta)\right|\right]<\infty;$$

(A6): $b_{t,j}$, $b_{t,j}Y_{t,j}$ and $a_{t,j}d_{t,j}Y_{t,j}$ are of order $O(t^{-h})$ for some h > 1/2, where

$$b_{t,j} = \sup_{\theta \in \Theta} \left\{ \mathbb{E} \left[\left\| \frac{\partial \widehat{\lambda}_{t,j}(\theta)}{\partial \theta} - \frac{\partial \lambda_{t,j}(\theta)}{\partial \theta} \right\| \right] \right\} \text{ and } d_{t,j} = \sup_{\theta \in \Theta} \max \left\{ \mathbb{E} \left[\left\| \frac{1}{\widehat{\lambda}_{t,j}(\theta)} \frac{\partial \widehat{\lambda}_{t,j}(\theta)}{\partial \theta} \right\| \right], \mathbb{E} \left[\left\| \frac{1}{\lambda_{t,j}(\theta)} \frac{\partial \lambda_{t,j}(\theta)}{\partial \theta} \right\| \right] \right\}.$$

These aforementioned assumptions hold for many classical models, see Ahmad and Francq [1]. These authors have established that the estimator $\widetilde{\theta}_n(T_j^*)$ is strongly consistent, for each regime $j \in \{1, \dots, K^*\}$; that is,

$$\widetilde{\theta}_n(T_j^*) \xrightarrow[n \to \infty]{a.s.} \theta_j^*.$$

They have also proved the asymptotic normality of $\theta_n(T_i^*)$; that is,

$$\sqrt{n_j^*}(\widetilde{\theta}_n(T_j^*) - \theta_j^*) \xrightarrow[n \to \infty]{\mathcal{D}} \mathcal{N}(0, \Sigma_j), \ \forall j = 1, \cdots, K^*,$$

where $\Sigma_j := J_j^{-1} I_j J_j^{-1}$. Under the above assumptions, for any $j = 1, \dots, K^*$, the matrix Σ_j can be consistently estimated by (see [1])

$$\widehat{\Sigma}_{j} = \widehat{J}_{j}^{-1} \widehat{I}_{j} \widehat{J}_{j}^{-1}, \text{ where}$$

$$\widehat{J}_{j} = \frac{1}{n} \sum_{t=1}^{n} \frac{1}{\widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{j}^{*}))} \frac{\partial \widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{j}^{*}))}{\partial \theta} \frac{\partial \widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{j}^{*}))}{\partial \theta'},$$

$$\widehat{I}_{j} = \frac{1}{n} \sum_{t=1}^{n} \left(\frac{Y_{t}}{\widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{i}^{*}))} - 1 \right)^{2} \frac{\partial \widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{j}^{*}))}{\partial \theta} \frac{\partial \widehat{\lambda}_{t,j}(\widetilde{\theta}_{n}(T_{j}^{*}))}{\partial \theta'}.$$
(2.7)

If we consider the process $\{Y_t, t \in \mathbb{Z}\}$, these properties are also verified on the segment T_1^* since it is easy to see that $\{(Y_t, \lambda_t), t \in T_1^*\}$ is a stationary process while $\{(Y_t, \lambda_t), t > t_1^*\}$ is not.

The following proposition establishes the consistency of the estimator $\widehat{\theta}_n(T_j^*)$, for any $j \in \{1, \dots K^*\}$.

Proposition 2.1 Assume that (A0)-(A2) and $(A_0(\Theta))$ hold. Then

$$\widehat{\theta}_n(T_j^*) \xrightarrow[n \to \infty]{a.s.} \theta_j^*, \ \forall j = 1, \cdots, K^*.$$

The results of this Proposition have been obtained by Ahmad and Francq (2016) when (Y_t, λ_t) is strictly stationary.

3 Estimation procedure and main results

In this section, we carry out the estimation of the number of breaks $K^* - 1$ and the instants of breaks \underline{t}^* by using a penalized contrast. Some asymptotic studies are also reported.

3.1 Penalized Poisson quasi-likelihood estimator

For any configuration of periods $K \ge 1$, $\underline{t} \in \mathcal{M}_n(K)$ and $\underline{\theta} = (\theta_1, \dots, \theta_K) \in \Theta^{K^*}$, we define the contrast

$$(QLIK) \qquad \widehat{J}_n(K, \underline{t}, \underline{\theta}) := -2 \sum_{k=1}^K \widehat{L}_n(T_k, \theta_k). \tag{3.1}$$

According to the proprieties of the PQMLE (see [1]), when K^* is known, a natural estimator of $(\underline{t}^*, \underline{\theta}^*) = ((t_j^*)_{1 \leq j \leq K^*-1}, (\theta_j^*)_{1 \leq j \leq K^*})$ for the model (1.2) is therefore the PQMLE on every interval $[t_j + 1, \dots, t_{j+1}]$ and every parameters θ_j for $1 \leq j \leq K^*$. But, since K^* is assumed to be unknown, we cannot directly use such method. To take into account the estimation of K^* , the most classical solution is to penalize the contrast by an additional term $\kappa_n K$, where κ_n represents a regularization parameter.

Now, define the penalized contrast QLIK, called penQLIK, by

$$(penQLIK) \qquad \widetilde{J}_n(K, t, \theta) := \widehat{J}_n(K, t, \theta) + \kappa_n K, \tag{3.2}$$

with $\kappa_n \leq n$ and $\kappa_n \underset{n \to \infty}{\longrightarrow} + \infty$.

The estimator of $(K^*, \underline{t}^*, \underline{\theta}^*)$ is defined as one of the minimizers of the penalized contrast:

$$\left(\widehat{K}_{n}, \underline{\widehat{t}}_{n}, \underline{\widehat{\theta}}_{n}\right) \in \underset{1 \leq K \leq K_{\max}}{\operatorname{argmin}} \underset{(\underline{t}, \underline{\theta}) \in \mathcal{M}_{n}(K) \times \Theta^{K}}{\operatorname{argmin}} \left(\widetilde{J}_{n}(K, \underline{t}, \underline{\theta})\right) \text{ and } \underline{\widehat{\tau}}_{n} = \frac{\underline{\widehat{t}}_{n}}{n}.$$
(3.3)

We will adapt the slope heuristic procedure to calibrate the penalty term from data (see Baudry et al. (2010)). In this procedure, the criteria QLIK is a linear transformation of the penalty (here the number of periods K) for the most complex models (with K close to K_{max}). This slope should be close to $-\kappa_n/2$. The slope estimation procedure considers only the linear part of -QLIK(K) with $1 \le K \le K_{\text{max}}$. Note that, in practice, a numerical algorithm can be used to compute the estimator on each segment; therefore, a minimum size is needed for the numerical computation of the criteria. Thus, we consider only the periods of length larger than some u_n and we can a priori fix K_{max} smaller than $[n/u_n]$. The complete procedure can be summarized as follows:

- 1. For each $1 \le K \le K_{\max}$, draw $(K, -\min_{\underline{t},\underline{\theta}} QLIK(K))$. Then compute the slope of the linear part: this slope is $\widehat{\kappa}_n/2$.
- 2. Using $\kappa_n = \widehat{\kappa}_n$, draw $(K, -\min_{\underline{t},\underline{\theta}} penQLIK(K))_{1 \leq K \leq K_{\text{max}}}$. This curve has a global minimum at \widehat{K}_n .

3.2 Asymptotic behavior

Under some assumptions, we will establish the asymptotic behavior of the estimator $(\widehat{K}_n, \underline{\hat{t}}_n, \underline{\widehat{\theta}}_n)$. Throughout this article, we set the following classical assumptions in the problem of break detection:

Assumption B. $\min_{1 \leq j \leq K^*-1} \left\| \theta_{j+1}^* - \theta_j^* \right\| > 0$. Also, there exists a vector $\underline{\tau}^* = (\tau_1^*, \dots, \tau_{K-1}^*)$ with $0 < \tau_1^* < \dots < \tau_{K-1}^* < 1$, called the vector of breaks such that $t_j^* = [n\tau_j^*]$, for $j = 1, \dots, K$ (where $[\cdot]$ is the integer part).

The following theorem gives the consistency of the estimator $(\widehat{K}_n, \widehat{t}_n, \widehat{\underline{\theta}}_n)$.

Theorem 3.1 Assume $K_{\text{max}} > K^*$ and $(\mathbf{A0})$ - $(\mathbf{A2})$, \mathbf{B} . If $\mathbf{A}_0(\Theta)$ holds and (κ_{ℓ}) satisfies

$$\sum_{\ell \ge 1} \frac{1}{\kappa_{\ell}} \sum_{k \ge \ell} \alpha_k^{(0)} < \infty, \tag{3.4}$$

then

$$(\widehat{K}_n, \widehat{\underline{\tau}}_n, \widehat{\underline{\theta}}_n) \xrightarrow[n \to \infty]{\mathcal{P}} (K^*, \underline{\tau}^*, \underline{\theta}^*).$$

By convention, throughout the sequel, if the vectors $\underline{\hat{t}}_n$ and \underline{t}^* do not have the same length, complete the shorter of the two vectors with 0 before computing the norm $\left\|\underline{\hat{t}}_n - \underline{t}^*\right\|_m$. The following theorem establishes the rates of convergence of the estimators $\underline{\hat{\tau}}_n$.

Theorem 3.2 Assume $K_{\max} > K^*$ and $(\mathbf{A0})$ - $(\mathbf{A2})$, \mathbf{B} . If $\mathbf{A}_i(\Theta)$ (i = 0, 1, 2), (2.2) (with $\epsilon > 2$) hold and (κ_{ℓ}) satisfies

$$\sum_{\ell \ge 1} \frac{1}{\kappa_{\ell}} \sum_{k \ge \ell} \alpha_k^{(i)} < \infty, \tag{3.5}$$

then the sequence $\left(\left\|\widehat{\underline{t}}_n - \underline{t}^*\right\|_m\right)_{n>1}$ is uniformly tight in probability, that is,

$$\lim_{\delta \to \infty} \lim_{n \to \infty} \mathbb{P}\left(\left\| \underline{\hat{t}}_n - \underline{t}^* \right\|_m > \delta \right) = 0.$$

Now, we give the convergence in distribution of the estimator of $\widehat{\underline{\theta}}_n$. By convention, if $\widehat{K}_n < K^*$, set $\widehat{T}_j = \widehat{T}_{\widehat{K}_n}$, for $j \in \{\widehat{K}_n, \dots, K^*\}$. The following theorem establishes the asymptotic normality of $\widehat{\theta}_n(\widehat{T}_j)$.

Theorem 3.3 Assume $K_{\text{max}} > K^*$ and $(\mathbf{A0})$ - $(\mathbf{A6})$ and \mathbf{B} . If (2.2) (with $\epsilon > 2$), $\mathbf{A}_i(\Theta)$ (i = 0, 1, 2) hold, such that (κ_{ℓ}) satisfies

$$\sum_{\ell>1} \max\left(\frac{1}{\kappa_{\ell}}, \frac{1}{\sqrt{\ell}}\right) \sum_{k>\ell} \alpha_k^{(i)} < \infty, \tag{3.6}$$

then

$$\sqrt{n_j^*}\left(\widehat{\theta}_n(\widehat{T}_j) - \theta_j^*\right) \xrightarrow[n \to \infty]{\mathcal{D}} \mathcal{N}_d\left(0, \Sigma_j\right), \quad \forall j = 1, \cdots, K^*,$$

where $\Sigma_j := J_j^{-1}(\theta_j^*)I_j(\theta_j^*)J_j^{-1}(\theta_j^*)$ with

$$J_{j}(\theta_{j}^{*}) = \mathbb{E}\left[\frac{1}{\lambda_{t,j}(\theta_{j}^{*})} \frac{\partial \lambda_{t,j}(\theta_{j}^{*})}{\partial \theta} \frac{\partial \lambda_{t,j}(\theta_{j}^{*})}{\partial \theta'}\right] \quad and \quad I_{j}(\theta_{j}^{*}) = \mathbb{E}\left[\frac{\operatorname{Var}(Y_{t,j}|\mathcal{F}_{t-1})}{\lambda_{t,j}^{2}(\theta_{j}^{*})} \frac{\partial \lambda_{t,j}(\theta_{j}^{*})}{\partial \theta} \frac{\partial \lambda_{t,j}(\theta_{j}^{*})}{\partial \theta'}\right].$$

Remark 3.4 The conditions on the regularization parameters $(\kappa_n)_{n\in\mathbb{N}}$ can be obtained if the Lipschitzian coefficients of $f(\cdot;\theta)$ and its derivatives are bounded by a geometric or Riemanian sequence:

- 1. the geometric case: if $\alpha_k^{(i)} = O(a^k)$ (i = 0, 1, 2) with $0 \le a < 1$, then any choice of $(\kappa_n)_{n \in \mathbb{N}}$ such that $\kappa_n \le n$ and $\kappa_n \to \infty$ satisfies (3.4) and (3.6) (for instance κ_n of order $\log n$ as in the BIC approach).
- 2. the Riemanian case: if $\alpha_k^{(i)} = O(k^{-\gamma})$ (i = 0, 1, 2) with $\gamma > 3/2$,
 - if $\gamma > 2$, then the conditions (3.4) and (3.6) hold for any choice of $(\kappa_n)_{n \in \mathbb{N}}$ such that $\kappa_n \leq n$ and $\kappa_n \to \infty$.
 - if $3/2 < \gamma \le 2$, then one can choose any sequence such that $\kappa_n = O(n^{\delta})$ with $\delta > 2 \gamma$ or $\kappa_n = O(n^{2-\gamma}(\log n)^{\delta})$ with $\delta > 1$.

4 Some simulations results

In this section, we implement the procedure on the R software (developed by the CRAN project). We will restrict our attention to the estimation of the vector (K^*, \underline{t}^*) ; *i.e* the number of segments K^* and the instants of breaks \underline{t}^* . For the performances of the estimator of the parameter $\underline{\theta}^*$, we refer to the works of Ahmad and Francq (2016). For each process, we generate 100 replications following the scenarios considered. The estimated number of segments is computed by using QLIK criteria penalized with $\kappa_n = \widehat{\kappa}_n$, $\kappa_n = \log n$ and $\kappa_n = \log n^{1/3}$. The value of the estimator $\widehat{\kappa}_n$ is calibrated by using the slope estimation procedure (see Baudry et al. (2010)) as described above. Once the regularization parameter κ_n obtained, the dynamic programming algorithm is used to minimize the criteria. With this algorithm, the complexity of the procedure declines from $\mathcal{O}(n^{K_{\text{max}}})$ to $\mathcal{O}(n^2)$.

4.1 Implementation procedure

We give the steps of the dynamic programming algorithm for computing the number of segments \widehat{K}_n and the optimal configuration of the breaks \widehat{t}_n . This algorithm is such that if (t_1, \dots, t_{K-1}, t) represents the optimal configuration of Y_1, \dots, Y_t into K segments, then (t_1, \dots, t_{K-1}) is the optimal configuration of $Y_1, \dots, Y_{t_{K-1}}$ into K-1 segments. Assume that the regularization parameter κ_n is known and let ML be the upper triangular matrix of dimension $n \times n$ with $ML_{i,l} = \widehat{L}(T_{i,l}, \widehat{\theta}_n(T_{i,l}))$, where $T_{i,l} = \{i, i+1, \dots, l\}$, for $1 \le i \le l \le n$. We summarize the implementation of the procedure as follows:

- The number of segments \widehat{K}_n : Let C be an upper triangular matrix of dimension $K_{\max} \times n$. For $1 \leq K \leq K_{\max}$ and $K \leq t \leq n$, $C_{K,t}$ will be the minimum penalized criteria of Y_1, \dots, Y_t into K segments. For $t = 1, \dots, n$, $C_{1,t} = -2ML_{1,t} + \kappa_n$ and the relation $C_{K+1,t} = \min_{K \leq l \leq t-1} (C_{K,l} 2ML_{l+1,t} + \kappa_n)$ is satisfied. Hence, $\widehat{K}_n = \operatorname{argmin}_{1 \leq K \leq K_{\max}} (C_{K,n})$.
- The change-point locations $\widehat{\underline{t}}_n$: Let Z be an upper triangular matrix of dimension $(K_{\max}-1)\times n$. For $1\leq K\leq (K_{\max}-1)$ and $K+1\leq t\leq n$, $Z_{K,t}$ will be the Kth potential break-point of Y_1,\cdots,Y_t . Therefore, the relation $Z_{K,t}=\min_{K\leq l\leq t-1}\left(C_{K,l}-2ML_{l+1,t}+\kappa_n\right)$ is satisfied for $K=1,\cdots,(K_{\max}-1)$. The break-points are obtained as follow: set $\widehat{t}_{\widehat{K}_n}=n$ and for $K=\widehat{K}_n-1,\cdots,1$, $\widehat{t}_K=Z_{K,\widehat{t}_{K+1}}$.

4.2 Results of simulations

4.2.1 Poisson-INARCH models

We consider the problem (1.2) for a Poisson-INARCH(1), i.e. (Y_1, \dots, Y_n) is a trajectory of the process $Y = \{Y_t, t \in \mathbb{Z}\}$ satisfying:

$$Y_t | \mathcal{F}_{t-1} \sim \mathcal{P}(\lambda_t) \; ; \; \lambda_t = f(Y_{t-1}, Y_{t-2}, \dots; \theta_j^*) = \alpha_0^{(j)} + \alpha_0^{(j)} Y_{t-1}, \quad \forall \; t \in T_j^*, \; \forall \; j \in \{1, \dots, K^*\} \,.$$
 (4.1)

The parameter vector is $\theta_j^* = (\alpha_0^{(j)}, \alpha^{(j)})$, for all $j \in \{1, \dots, K^*\}$.

For n = 500 and n = 1000, we generate a sample (Y_1, \dots, Y_n) in the following situations:

- scenario IA₀: $\theta_1^* = (0.5, 0.6)$ is constant $(K^* = 1)$;
- scenario IA₁: $\theta_1^* = (0.5, 0.6)$ changes to $\theta_2^* = (1.0, 0.6)$ at $t^* = 0.5n$ $(K^* = 2)$;
- scenario IA₂: $\theta_1^* = (0.5, 0.6)$ changes to $\theta_2^* = (1.0, 0.6)$ at $t_1^* = 0.3n$ which changes to $\theta_3^* = (1.0, 0.25)$ at $t_2^* = 0.7n$ ($K^* = 3$).

For scenario \mathbf{IA}_2 , Figure 1 shows the slope of the linear part of the -QLIK criteria minimized in $(\underline{t},\underline{\theta})$. We obtain $\widehat{\kappa}_n \approx 4.6$ for n=500 and $\widehat{\kappa}_n \approx 5.9$ for n=1000. Using these above values for κ_n (i.e. $\kappa_n = \widehat{\kappa}_n$), we minimize the penQLIK in $(K,\underline{t},\underline{\theta})$, with $1 \leq K \leq K_{\text{max}}$. Figure 2 displays the points $(K,\min_{\underline{t},\underline{\theta}}penQLIK(K))$ for $1 \leq K \leq K_{\text{max}} = 15$. One can see that the estimated number of segments is $\widehat{K}_n = 3$ for n=500 and n=1000. The estimated instants of breaks are $\widehat{\underline{t}}_n = (157,349)$ ($\underline{t}^* = (150,350)$) for n=500 and $\widehat{\underline{t}}_n = (291,702)$ ($\underline{t}^* = (300,700)$) for n=1000 (see Figure 3).

Now, we are going to generate 100 replications of a Poisson-INGARCH(1,1) process following the scenarios \mathbf{IA}_0 - \mathbf{IA}_2 . Table 1 indicates the frequencies of number of replications where $\widehat{K}_n = K^*$, $\widehat{K}_n < K^*$ and $\widehat{K}_n > K^*$, for the regularization parameter $\kappa_n = \widehat{\kappa}_n$, $\log n$, $n^{1/3}$. For the scenarios \mathbf{IA}_1 and \mathbf{IA}_2 , we also consider the replications where the true number of breaks is achieved (i.e. $\widehat{K}_n = K^*$) and we present some elementary statistics of the estimated instants of breaks (see Table 1).




Figure 1: The curve of $-\min_{\underline{t},\underline{\theta}} QLIK(K)$, for $1 \le K \le K_{\text{max}}$ for a Poisson-INARCH(1) process in scenario \mathbf{IA}_2 . The solid line represents the linear part of this curve with slope $\widehat{\kappa}_n/2 = 2.307$ when n = 500 and $\widehat{\kappa}_n/2 = 2.928$ when n = 1000.




Figure 2: The graph $(K, \min_{\underline{t},\underline{\theta}} penQLIK(K))$, for $1 \leq K \leq K_{max}$ for a Poisson-INARCH(1) process in scenario \mathbf{IA}_2 .




Figure 3: The estimated of breakpoints for a trajectory of a Poisson-INARCH(1) process in scenario \mathbf{IA}_2 . The solid lines represent the estimated instants of breaks and the dotted lines represent the true ones.

The results of Table 1 show that for the penalties considered, the performance increase with n in all scenarios. In accordance with Theorem 3.1, the consistency of the penalties $\log n$ and $n^{1/3}$ is numerically convincing. Moreover, the $n^{1/3}$ penalty outperforms the other procedures when n=1000.

Table 1: Breaks estimated after 100 replications for Poisson-INARCH(1) process following scenarios \mathbf{IA}_0 - \mathbf{IA}_2 . The first three columns show the frequencies of the estimation of the true, low and high number of breaks. The last three columns give some elementary statistics of the change-point locations when the true number of breaks is achieved.

Scenarios			Frequencies $\widehat{K}_n = K^* \qquad \widehat{K}_n < K^* \qquad \widehat{K}_n > K^*$		Mean \pm s.d. $\widehat{\tau}_1$ $\widehat{\tau}_2$		Mean $\left\ \widehat{\underline{\tau}}_n - \underline{\tau}^* \right\ $	
			16			. 1	· <u>z</u>	-n -
\mathbf{IA}_0	n = 500	$\kappa_n = \kappa_n$	0.72	0.00	0.28			
$(K^* = 1)$		$\kappa_n = \log n$	0.76	0.00	0.24			
		$\kappa_n = n^{1/3}$	0.94	0.00	0.06			
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.94	0.00	0.06			
		$\kappa_n = \log n$	0.90	0.00	0.10			
		$\kappa_n = n^{1/3}$	1.00	0.00	0.00			
IA_1	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.76	0.03	0.21	0.497 ± 0.064		0.038
$(K^* = 2)$		$\kappa_n = \log n$	0.83	0.03	0.14	0.495 ± 0.066		0.040
		$\kappa_n = n^{1/3}$	0.87	0.09	0.04	0.495 ± 0.064		0.038
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.89	0.00	0.11	0.507 ± 0.033		0.019
		$\kappa_n = \log n$	0.87	0.00	0.13	0.507 ± 0.034		0.020
		$\kappa_n = n^{1/3}$	0.98	0.00	0.02	0.506 ± 0.032		0.019
IA_2	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.62	0.13	0.25	0.311 ± 0.071	0.689 ± 0.060	0.061
$(K^* = 3)$		$\kappa_n = \log n$	0.73	0.12	0.15	0.317 ± 0.073	0.690 ± 0.072	0.067
		$\kappa_n = n^{1/3}$	0.64	0.33	0.03	0.310 ± 0.058	0.685 ± 0.070	0.061
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.87	0.00	0.13	0.300 ± 0.034	0.693 ± 0.030	0.034
		$\kappa_n = \log n$	0.84	0.00	0.16	0.302 ± 0.043	0.692 ± 0.030	0.038
		$\kappa_n = n^{1/3}$	0.93	0.05	0.02	0.300 ± 0.051	0.694 ± 0.028	0.037

4.2.2 Poisson-INGARCH models

We consider the problem (1.2) for a Poisson-INGARCH(1,1), i.e. (Y_1, \dots, Y_n) is a trajectory of the process $Y = \{Y_t, t \in \mathbb{Z}\}$ satisfying:

$$Y_t | \mathcal{F}_{t-1} \sim \mathcal{P}(\lambda_t) ; \ \lambda_t = \alpha_0^{(j)} + \alpha_0^{(j)} Y_{t-1} + \beta_0^{(j)} \lambda_{t-1}, \ \forall \ t \in T_j^*, \ \forall \ j \in \{1, \dots, K^*\}.$$
 (4.2)

The parameter vector is $\theta_j^* = (\alpha_0^{(j)}, \alpha^{(j)}, \beta^{(j)})$, for all $j \in \{1, \dots, K^*\}$.

For n = 500 and n = 1000, we generate 100 replications of the model (4.2) in the following situations:

- scenario IG₀: $\theta_1^* = (1.0, 0.2, 0.15)$ is constant $(K^* = 1)$;
- scenario IG₁: $\theta_1^* = (1.0, 0.2, 0.15)$ changes to $\theta_2^* = (1.0, 0.45, 0.15)$ at $t^* = 0.5n$ $(K^* = 2)$;
- scenario IG₂: $\theta_1^* = (0.1, 0.3, 0.6)$ changes to $\theta_2^* = (0.5, 0.3, 0.6)$ at $t_1^* = 0.3n$ which changes to $\theta_3^* = (0.5, 0.3, 0.2)$ at $t_2^* = 0.7n$ $(K^* = 3)$.

Table 2 indicates the frequencies of the true number of breaks estimated and some elementary statistics of the estimators of the change-point locations. It appears that the results of the $n^{1/3}$ -penalty and the slope procedure are quite satisfactory except for the case of two breaks. In this later case, the $n^{1/3}$ -penalty and the slope procedure over-penalizes the number of breaks, while the log n-penalty under-penalizes. But, overall, the performances of the proposed procedures increase with n and the estimation of the breakpoints locations is well achieved.

Table 2: Breaks estimated after 100 replications for Poisson-INGARCH(1,1) process following scenarios \mathbf{IG}_0 - \mathbf{IG}_2 . The first three columns show the frequencies of the estimation of the true, low and high number of breaks.

The last three columns give some elementary statistics of the change-point locations when the true number of breaks is achieved.

			Frequencies		Mean \pm s.d.		Mean	
Scenarios			$\widehat{K}_n = K^*$	$\widehat{K}_n < K^*$	$\widehat{K}_n > K^*$	$\widehat{ au_1}$	$\widehat{ au_2}$	$\left\ \widehat{\underline{\tau}}_n - \underline{\tau}^* \right\ $
IG_0	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.86	0.00	0.14			
$(K^* = 1)$		$\kappa_n = \log n$	0.96	0.00	0.04			
		$\kappa_n = n^{1/3}$	1.00	0.00	0.00			
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.92	0.00	0.08			
		$\kappa_n = \log n$	0.96	0.00	0.04			
		$\kappa_n = n^{1/3}$	1.00	0.00	0.00			
IG_1	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.75	0.05	0.20	0.515 ± 0.066		0.038
$(K^* = 2)$		$\kappa_n = \log n$	0.70	0.03	0.27	0.514 ± 0.073		0.040
		$\kappa_n = n^{1/3}$	0.78	0.06	0.16	0.512 ± 0.066		0.038
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.75	0.05	0.20	0.507 ± 0.031		0.019
		$\kappa_n = \log n$	0.58	0.00	0.42	0.508 ± 0.034		0.021
		$\kappa_n = n^{1/3}$	0.83	0.03	0.13	0.501 ± 0.048		0.022
IG_2	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.53	0.41	0.06	0.299 ± 0.078	0.691 ± 0.073	0.053
$(K^* = 3)$		$\kappa_n = \log n$	0.58	0.23	0.19	0.299 ± 0.074	0.693 ± 0.070	0.049
		$\kappa_n = n^{1/3}$	0.37	0.49	0.14	0.300 ± 0.076	0.697 ± 0.015	0.047
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.62	0.26	0.12	0.293 ± 0.050	0.702 ± 0.010	0.025
		$\kappa_n = \log n$	0.60	0.06	0.34	0.293 ± 0.051	0.702 ± 0.010	0.026
		$\kappa_n = n^{1/3}$	0.56	0.29	0.15	0.301 ± 0.029	0.699 ± 0.011	0.016

4.2.3 Negative binomial INGARCH models

We consider the problem (1.2) for a negative binomial INGARCH(1,1) (NB-INGARCH(1,1)), i.e. (Y_1, \dots, Y_n) is a trajectory of the process $Y = \{Y_t, t \in \mathbb{Z}\}$ satisfying:

$$Y_{t}|\mathcal{F}_{t-1} \sim NB(r, p_{t}) \; ; \; r\frac{(1-p_{t})}{p_{t}} = \lambda_{t} = \alpha_{0}^{(j)} + \alpha^{(j)}Y_{t-1} + \beta^{(j)}\lambda_{t-1}, \; \forall \; t \in T_{j}^{*}, \; \forall \; j \in \{1, \dots, K^{*}\} \; ; \quad (4.3)$$

where the parameter vector is $\theta_j^* = (\alpha_0^{(j)}, \alpha^{(j)}, \beta^{(j)})$, for all $j \in \{1, \dots, K^*\}$ and NB(r, p) denotes the negative binomial distribution with parameters r and p.

For r = 14 (used for transaction data, see [13]), n = 500 and n = 1000, we generate a sample (Y_1, \dots, Y_n) in the following situations:

- scenario NB-IG₀: $\theta_1^* = (1.0, 0.2, 0.15)$ is constant $(K^* = 1)$;
- scenario NB-IG₁: $\theta_1^* = (1, 0.2, 0.15)$ changes to $\theta_2^* = (1, 0.45, 0.15)$ at $t^* = 0.5n$ $(K^* = 2)$;
- scenario NB-IG₂: $\theta_1^* = (0.1, 0.3, 0.6)$ changes to $\theta_2^* = (0.5, 0.3, 0.6)$ at $t_1^* = 0.3n$ which changes to $\theta_3^* = (0.5, 0.3, 0.2)$ at $t_2^* = 0.7n$ ($K^* = 3$).

Once again, it appears in Table 3 that the performances of the proposed procedures increase with n and the estimation of the breakpoints locations remain satisfactory even in this case where the Poisson quasi-likelihood used is quite different from the true distribution of the observations.

Table 3: Breaks estimated after 100 replications for NB-INGARCH(1,1) process following scenarios $\mathbf{NB} - \mathbf{IG}_0 - \mathbf{NB} - \mathbf{IG}_2$. The first three columns show the frequencies of the estimation of the true, low and high number of breaks. The last three columns give some elementary statistics of the change-point locations when the true number of breaks is achieved.

Camaniaa			Frequencies $\widehat{K}_n = K^* \widehat{K}_n < K^* \widehat{K}_n > K^*$		$^{\wedge}$ Mean \pm s.d. $^{\wedge}$		Mean ^*	
Scenarios			$K_n = K$	$K_n < K$	$K_n > K$	$ au_1$	$ au_2$	$\left\ \underline{\tau}_{n} - \underline{\tau}^{*} \right\ $
NB-IG ₀	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.90	0.00	0.10			
$(K^* = 1)$		$\kappa_n = \log n$	0.95	0.00	0.05			
		$\kappa_n = n^{1/3}$	0.98	0.00	0.02			
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.92	0.00	0.08			
		$\kappa_n = \log n$	0.94	0.00	0.06			
		$\kappa_n = n^{1/3}$	0.98	0.00	0.02			
$NB-IG_1$	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.60	0.09	0.31	0.512 ± 0.122		0.072
$(K^* = 2)$		$\kappa_n = \log n$	0.55	0.04	0.41	0.514 ± 0.110		0.063
		$\kappa_n = n^{1/3}$	0.65	0.12	0.23	0.519 ± 0.106		0.060
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.69	0.02	0.29	0.507 ± 0.065		0.037
		$\kappa_n = \log n$	0.56	0.00	0.44	0.500 ± 0.054		0.030
		$\kappa_n = n^{1/3}$	0.83	0.02	0.15	0.505 ± 0.061		0.037
$\mathbf{NB}\text{-}\mathbf{IG}_2$	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.45	0.51	0.04	0.330 ± 0.084	0.696 ± 0.040	0.057
$(K^* = 3)$		$\kappa_n = \log n$	0.41	0.13	0.46	0.319 ± 0.080	0.700 ± 0.026	0.057
		$\kappa_n = n^{1/3}$	0.43	0.28	0.29	0.328 ± 0.061	0.685 ± 0.066	0.055
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.70	0.25	0.05	0.304 ± 0.060	0.699 ± 0.020	0.033
		$\kappa_n = \log n$	0.54	0.02	0.44	0.299 ± 0.066	0.699 ± 0.014	0.033
		$\kappa_n = n^{1/3}$	0.68	0.19	0.13	0.325 ± 0.090	0.697 ± 0.018	0.043

4.2.4 Binary Time Series

Consider the problem (1.2) for a binary INARCH(1) (BIN-INARCH(1)) time series model, i.e. (Y_1, \dots, Y_n) is a trajectory of the process $Y = \{Y_t, t \in \mathbb{Z}\}$ satisfying:

$$Y_t | \mathcal{F}_{t-1} \sim \mathcal{B}(p_t) \; ; \; p_t = \lambda_t = \alpha_0^{(j)} + \alpha^{(j)} Y_{t-1}, \; \forall \; t \in T_i^*, \; \forall \; j \in \{1, \dots, K^*\} \; ;$$
 (4.4)

where the parameter vector is $\theta_j^* = (\alpha_0^{(j)}, \alpha^{(j)})$ (with $0 < \alpha_0^{(j)} + \alpha^{(j)} < 1$), for all $j \in \{1, \dots, K^*\}$ and $\mathcal{B}(p)$ denotes the Bernoulli distribution with parameter p.

For n = 500 and n = 1000, we generate a sample (Y_1, \dots, Y_n) in the following situations:

- scenario BIN-IA₀: $\theta_1^* = (0.15, 0.75)$ is constant $(K^* = 1)$;
- scenario BIN-IA₁: $\theta_1^* = (0.15, 0.75)$ changes to $\theta_2^* = (0.04, 0.60)$ at $t^* = 0.5n$ $(K^* = 2)$;
- scenario BIN-IA₂: $\theta_1^* = (0.15, 0.75)$ changes to $\theta_2^* = (0.04, 0.60)$ at $t_1^* = 0.3n$ which changes to $\theta_3^* = (0.25, 0.35)$ at $t_2^* = 0.7n$ ($K^* = 3$).

The scenario $BIN-IA_1$ is related and close to the real data example (see below).

Table 4 shows that the procedure provides satisfactory results with BIN-INARCH(1) model, except that the $n^{1/3}$ -penalty in the case of two breaks. But, the performances of these procedures increase with n and the breakpoints locations are overall well estimated.

Table 4: Breaks estimated after 100 replications for BIN-INARCH(1) process following scenarios $BIN - IA_0$ - $BIN - IA_2$. The first three columns show the frequencies of the estimation of the true, low and high number of breaks. The last three columns give some elementary statistics of the change-point locations when the true number of breaks is achieved.

Scenarios			$\widehat{K}_n = K^*$	Frequencies $\widehat{K}_n < K^*$	$\widehat{K}_n > K^*$	$\widehat{ au}_1$ Mean	\pm s.d. $\widehat{\tau}_2$	Mean $\left\ \widehat{\underline{\tau}}_n - \underline{\tau}^* \right\ $
BIN-IA ₀	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.84	0.00	0.16			
$(K^* = 1)$		$\kappa_n = \log n$	0.98	0.00	0.02			
		$\kappa_n = n^{1/3}$	0.98	0.00	0.02			
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.86	0.00	0.14			
		$\kappa_n = \log n$	1.00	0.00	0.00			
		$\kappa_n = n^{1/3}$	1.00	0.00	0.00			
BIN-IA ₁	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.69	0.10	0.21	0.499 ± 0.091		0.055
$(K^* = 2)$		$\kappa_n = \log n$	0.72	0.27	0.01	0.491 ± 0.087		0.051
		$\kappa_n = n^{1/3}$	0.52	0.48	0.00	0.484 ± 0.091		0.054
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.89	0.00	0.11	0.499 ± 0.036		0.020
		$\kappa_n = \log n$	0.96	0.01	0.03	0.500 ± 0.035		0.019
		$\kappa_n = n^{1/3}$	0.85	0.15	0.00	0.484 ± 0.091		0.054
$BIN-IA_2$	n = 500	$\kappa_n = \widehat{\kappa}_n$	0.75	0.18	0.07	0.324 ± 0.094	0.695 ± 0.044	0.060
$(K^* = 3)$		$\kappa_n = \log n$	0.43	0.57	0.00	0.312 ± 0.044	0.694 ± 0.030	0.035
		$\kappa_n = n^{1/3}$	0.23	0.77	0.00	0.306 ± 0.033	0.702 ± 0.017	0.026
	n = 1000	$\kappa_n = \widehat{\kappa}_n$	0.95	0.03	0.02	0.303 ± 0.046	0.696 ± 0.019	0.028
		$\kappa_n = \log n$	0.90	0.10	0.00	0.299 ± 0.037	0.697 ± 0.017	0.025
		$\kappa_n = n^{1/3}$	0.52	0.48	0.00	0.296 ± 0.021	0.697 ± 0.019	0.020

4.2.5 INARCH(∞) models

Now, consider a Poisson-INARCH(∞), *i.e.* (Y_1, \dots, Y_n) is a trajectory of the process $Y = \{Y_t, t \in \mathbb{Z}\}$ satisfying:

$$Y_t | \mathcal{F}_{t-1} \sim \mathcal{P}(\lambda_t) \; ; \; \lambda_t = \alpha_0^{(j)} + \sum_{k=1}^{\infty} \alpha_k Y_{t-k}, \quad \forall \; t \in T_j^*, \; \forall \; j \in \{1, \dots, K^*\},$$
 (4.5)

where $\alpha_0^{(j)} > 0$, $\alpha_k \ge 0$ (for all $k \ge 1$ and $j = 1, \dots, K^*$) and $\sum_{k \ge 1} \alpha_k < 1$; that is, we focus on the change in the parameter $\alpha_0^{(j)}$. This process corresponds to a particular case of the problem (1.2) with $f(y_1, y_2, \dots, \alpha_0^{(j)}) = \alpha_0^{(j)} + \sum_{k=1}^{\infty} \alpha_k y_k$ for each regime $j \in \{1, \dots, K^*\}$. We deal with a scenario where the consistency of the BIC procedure is not ensured. Therefore, we consider the Riemanian case with $\alpha_k = O(k^{-1.7})$ (in the scenario detailed below). More precisely, we consider the model (4.5) with

$$\lambda_t = \alpha_0^{(j)} + \frac{1}{2.2} \sum_{k=1}^{\infty} \frac{1}{k^{1.7}} Y_{t-k}.$$

The number 1/2.2 is obtained from the values of the Riemann zeta function, and allows the condition $\frac{1}{2.2} \sum_{k=1}^{\infty} \frac{1}{k^{1.7}} < 1$. According to Theorem 3.1 and Remark 3.4, if the regularization parameter verifies $\kappa_n = O(n^{\delta})$ with $\delta > 0.3$, then the consistency holds. Thus, the consistency of the BIC penalty is not ensured.

Now, for n = 500 and n = 100, we generate a trajectory (Y_1, \dots, Y_n) of the model (4.5) in the following scenarios:

- scenario IA-INF₀: $\alpha_0^{(1)} = 0.5$ is constant $(K^* = 1)$;
- scenario IA-INF₁: $\alpha_0^{(1)} = 0.5$ changes to $\alpha_0^{(2)} = 0.1$ at $t^* = 0.5n$ $(K^* = 2)$.

Table 5: Breaks estimated after 100 replications for INARCH(∞) process following scenarios $\mathbf{IA} - \mathbf{INF}_0$ and $\mathbf{IA} - \mathbf{INF}_1$. The first three columns show the frequencies of the estimation of the true, low and high number of breaks. The last two columns give some elementary statistics of the change-point locations when the true number of breaks is achieved.

Scenarios			$\widehat{K}_n = K^*$	Frequencies $\widehat{K}_n < K^*$	$\widehat{K}_n > K^*$	Mean \pm s.d. $\widehat{\tau}_1$	
$ \begin{array}{c} \mathbf{IA-INF}_0 \\ (K^* = 1) \end{array} $	n = 500	$\kappa_n = \widehat{\kappa}_n$ $\kappa_n = \log n$ $\kappa_n = n^{1/3}$	0.15 0.56 0.84	0.00 0.00 0.00	0.85 0.44 0.16		
	n = 1000	$\kappa_n = \kappa_n$ $\kappa_n = \log n$ $\kappa_n = n^{1/3}$	0.17 0.57 0.95	0.00 0.00 0.00	0.83 0.43 0.05		
$ \begin{array}{c} \mathbf{IA\text{-}INF}_1 \\ (K^* = 2) \end{array} $	n = 500	$\kappa_n = \widehat{\kappa}_n$ $\kappa_n = \log n$ $\kappa_n = n^{1/3}$	0.66 0.49 0.82	0.04 0.02 0.03	0.34 0.49 0.15	0.498 ± 0.0057 0.498 ± 0.0013 0.497 ± 0.0062	0.003 0.002 0.003
	n = 1000	$\kappa_n = \widehat{\kappa}_n$ $\kappa_n = \log n$ $\kappa_n = n^{1/3}$	0.79 0.43 0.93	0.00 0.00 0.00	0.21 0.57 0.07	0.499 ± 0.0004 0.499 ± 0.0006 0.499 ± 0.0005	0.001 0.001 0.001

In Table 5, on can see that the $n^{1/3}$ -penalty uniformly outperforms the other two procedures. Moreover, the performances of the proposed procedures increase with n, except the log n-penalty whose the performances decrease with n. Hence, the consistency of the BIC procedure is quite questionable in this case.

5 Real data application

We apply our change-point procedure to two examples of real data series. To compute the estimator \widehat{K}_n , the $\widehat{\kappa}_n$ penalty is used with $u_n = \lceil (\log(n))^{\delta} \rceil$ (where $3/2 \le \delta \le 2$) and $K_{\max} = 15$.

5.1 The US recession data

Firstly, we consider the series of the quarterly recession data from the USA for the period 1855-2013 (see Figure 5). This series (Y_t) represents a binary variable that is equal to 1 if there is a recession in at least 1 month in the quarter and 0 otherwise. There are 636 quarterly observations obtained from The National Bureau of Economic Research. These data have already been analyzed by several authors. Hudecová (2013) has applied a change-point procedure based on a normalized cumulative sums of residuals and has found a break in the first quarter of 1933. Recently, Diop and Kengne (2017) have applied a change-point test based on the maximum likelihood estimator of the model's parameter and have detected a break in the last quarter of 1932.

We consider the INARCH(1) representation and apply the penQLIK contrast procedure. This choice is motivated by the fact that the estimation of the last component of θ (i.e. the parameter β) is not significant in the INGARCH(1,1) representation (see Diop and Kengne [13]). The test of nullity of one coefficient (TNOC) proposed by Ahmad and Francq [1], applied a posteriori (after change-point detection) also confirms these results. As noted in the implementation of the dynamic programming algorithm, we begin by the calibration of the regularization parameter κ_n . The slope estimation procedure applied with $u_n = [(\log n)^2]$ returns the values $\hat{\kappa}_n \approx 3.21$ and the estimation of the number of segments is $\hat{K}_n = 2$, i.e. one break is detected (see Figure 4). The location of the breakpoint estimated is $\hat{t} = 313$. The change detected at $\hat{t} = 313$ corresponds

to the first quarter of 1933 (see Figure 5). These results are in concordance with those obtained by Diop and Kengne (2017) and Hudecová (2013). The estimated model with one breakpoint is

$$\mathbb{E}(Y_t|\mathcal{F}_{t-1}) = \lambda_t = \begin{cases} 0.120 + 0.749Y_{t-1}, & \text{for } t \le 313\\ (0.028) & (0.215) \end{cases}$$

$$0.047 + 0.681Y_{t-1}, & \text{for } t > 313,\\ (0.013) & (0.230) \end{cases}$$

where in parentheses are the standard errors of the estimators obtained from the robust sandwich matrix $\widehat{H}_j^{-1}\widehat{\Sigma}_j^{-1}\widehat{H}_j^{-1}$ computed on each regime j, where $\widehat{\Sigma}_j$ is given by (2.7) and $\widehat{H}_j = \frac{1}{n}\sum_{t=1}^n \frac{\partial^2 \widehat{\ell}_{t,j}(\widehat{\theta}_n(\widehat{T}_j))}{\partial \theta \partial \theta'}$. These parameters estimation display a distortion in term of standard errors; it can be explained by the fact that the true distribution of the observations (which is binary), is quite different from the Poisson quasi-likelihood used.




Figure 4: The curve of $-\min_{\underline{t},\underline{\theta}} QLIK(K)$ and the graph $(K,\min_{\underline{t},\underline{\theta}} penQLIK(K))$ for the US recession data with a INARCH(1) model. The solid line represents the linear part of this curve with slope $\widehat{\kappa}_n/2 = 1.605$.




Figure 5: The US recession data with the estimated location of the breakpoint \hat{t} .

5.2 Number of trades in the stock of Technofirst

Secondly, we apply our change-point detection procedure to a financial time series data. We consider the daily number of trades in the stock of Technofirst listed in the NYSE Euronext group. It is a series of

1000 observations from 04 January 2010 to 20 April 2016 (see Figure 7). The data are available online at the website "https://www.euronext.com/en/products/equities/FR0011651819-ALXP". These data have been analyzed by Ahmad and Francq [1] with the PQMLE, and have concluded that the INGARCH(1,3) is more appropriate. We carry out an INGARCH(1,1) with the possibility of change in the observations.

The slope estimation procedure obtained with $u_n = [(\log(n))^2]$ returns $\hat{\kappa}_n \approx 23.04$ and the estimation of the number of segments is $\widehat{K}_n = 3$, *i.e.* two changes are detected (see Figure 6). The locations of the breakpoints estimated are $\hat{t}_1 = 230$ (06 April 2011) and $\hat{t}_2 = 311$ (06 September 2011), see also Figure 7.

The estimated model with change-points is

$$\mathbb{E}(Y_t|\mathcal{F}_{t-1}) = \lambda_t = \begin{cases} 2.436 + 0.368Y_{t-1}, & \text{for } t \le 230 \\ (0.126) & (0.032) \end{cases}$$

$$\frac{4.643 + 0.607Y_{t-1} + 0.032\lambda_{t-1}, & \text{for } 230 < t \le 311}{(0.649) & (0.022) & (0.033)}$$

$$\frac{1.113 + 0.166Y_{t-1} + 0.531\lambda_{t-1}, & \text{for } t > 311,}{(0.226) & (0.016) & (0.071)}$$

where in parentheses are the robust standard errors of the estimators. Let us note that, we have applied the TNOC, which fund that the INARCH(1) representation is the most appropriate for first regime.





Figure 6: The curve of $-\min_{\underline{t},\underline{\theta}} QLIK(K)$ and the graph $(K,\min_{\underline{t},\underline{\theta}} penQLIK(K))$ for the daily number of trades in the stock of Technofirst. The solid line represents the linear part of this curve with slope $\widehat{\kappa}_n/2 = 11.52$.

6 Proofs of the main results

In the sequel, C denotes a positive constant whom value may differ from an inequality to another and we set $v_n = n/\kappa_n$ for all $n \ge 1$.

6.1 Proof of Proposition 2.1

Remark that since $\{Y_{t,j}, t \in \mathbb{Z}\}$ is stationary and ergodic, the process $\{\ell_{t,j}(\theta), t \in \mathbb{Z}\}$ is also a stationary and ergodic sequence, for any $\theta \in \Theta$. Then, the proof can be divided into two parts. For $j = 1, \dots, K^*$, we






Figure 7: The daily number of trades in the stock of the Technofirst with the estimated locations of the breakpoints \hat{t}_1 and \hat{t}_2 .

will first (1.) show that $\left\|\frac{1}{n_j^*}\widehat{L}_n(T_j^*,\theta) - \mathbb{E}(\ell_{1,j}(\theta))\right\|_{\Theta} \xrightarrow[n\to\infty]{a.s.} 0$; secondly (2.), we will show that the function $\theta \mapsto \mathbb{E}(\ell_{1,j}(\theta))$ has a unique maximum in θ_j^* .

(1.) Recall that $\ell_{t,j}(\theta) = Y_{t,j} \log f_{t,j}^{\theta} - f_{t,j}^{\theta}$, for any $\theta \in \Theta$. We have

$$\begin{split} |\ell_{t,j}| &\leq Y_{t,j} |\log f_{t,j}^{\theta}| + |f_{t,j}^{\theta}| \\ &\leq Y_{t,j} \Big| \log \Big(\frac{f_{t,j}^{\theta}}{\underline{c}} \times \underline{c} \Big) \Big| + |f_{t,j}^{\theta}| \\ &\leq Y_{t,j} \Big(\Big| \frac{f_{t,j}^{\theta}}{\underline{c}} - 1 \Big| + |\log \underline{c}| \Big) + |f_{t,j}^{\theta}| \quad \text{(because for } x > 1, \ |\log x| \leq |x - 1|) \\ &\leq Y_{t,j} \Big(\Big| \frac{f_{t,j}^{\theta}}{\underline{c}} \Big| + 1 + |\log \underline{c}| \Big) + |f_{t,j}^{\theta}|. \end{split}$$

Hence,

$$\|\ell_{t,j}\|_{\Theta} \le Y_{t,j} \left(\frac{1}{c} \|f_{t,j}^{\theta}\|_{\Theta} + 1 + |\log \underline{c}|\right) + \|f_{t,j}^{\theta}\|_{\Theta}.$$
 (6.1)

We will show that $\mathbb{E}[\|\ell_{t,j}\|_{\Theta}] < \infty$. According to (6.1), we have

$$\mathbb{E}[\|\ell_{t,j}\|_{\Theta}] \leq \mathbb{E}\left[Y_{t,j}\left(\frac{1}{c}\|f_{t,j}^{\theta}\|_{\Theta} + 1 + |\log \underline{c}|\right) + \|f_{t,j}^{\theta}\|_{\Theta}\right] \leq C\mathbb{E}\left[\left(\frac{Y_{t,j}}{\underline{c}} + 1\right)\|f_{t,j}^{\theta}\|_{\Theta}\right]$$
$$\leq C\left(\mathbb{E}\left[\left(\frac{Y_{t,j}}{c} + 1\right)^{2}\right]\right)^{1/2} (\mathbb{E}\|f_{t,j}^{\theta}\|_{\Theta}^{2})^{1/2} \leq C(\mathbb{E}\|f_{t,j}^{\theta}\|_{\Theta}^{2})^{1/2}.$$

Since $(\mathbf{A}_0(\Theta))$ holds, we have

$$||f_{t,j}^{\theta}||_{\Theta} \leq ||f_{t,j}^{\theta} - f^{\theta}(0, \dots, 0)|_{\Theta} + ||f^{\theta}(0, \dots, 0)||_{\Theta} \leq \sum_{k \geq 1} \alpha_k^{(0)} |Y_{t-k,j}| + ||f^{\theta}(0, \dots, 0)||_{\Theta}.$$

Therefore,

$$\mathbb{E}[\|\ell_{t,j}\|_{\Theta}] \leq C \left[\mathbb{E} \left(\sum_{k \geq 1} \alpha_k^{(0)} |Y_{t-k,j}| + \|f^{\theta}(0, \dots, 0)\|_{\Theta} \right)^2 \right]^{1/2}$$

$$\leq C \sum_{k \geq 1} \alpha_k^{(0)} (\mathbb{E} |Y_{t-k,j}|^2)^{1/2} + (\mathbb{E} \|f^{\theta}(0, \dots, 0)\|_{\Theta}^2)^{1/2}$$

$$\leq C \sum_{k \geq 1} \alpha_k^{(0)} + (\mathbb{E} \|f^{\theta}(0, \dots, 0)\|_{\Theta}^2)^{1/2} < \infty.$$

By the uniform strong law of large number applied on the process $\{\ell_{t,j}(\theta), t \in \mathbb{Z}\}\$, it holds that

$$\left\| \frac{1}{n_j^*} L_{n,j}(T_j^*, \theta) - \mathbb{E}(\ell_{1,j}(\theta)) \right\|_{\Theta} = \left\| \frac{1}{n_j^*} \sum_{t \in T_j^*} \ell_{t,j}(\theta) - \mathbb{E}(\ell_{1,j}(\theta)) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$
 (6.2)

According to Ahmad and Francq [1], we have

$$\frac{1}{n_j^*} \|\widehat{L}_{n,j}(T_j^*, \theta) - L_{n,j}(T_j^*, \theta)\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0, \text{ for any } j = 1, \cdots, K^*.$$

$$(6.3)$$

From (6.2) and (6.3), we deduce that

$$\left\| \frac{1}{n_j^*} \widehat{L}_{n,j}(T_j^*, \theta) - \mathbb{E}(\ell_{1,j}(\theta)) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$
 (6.4)

The following lemma is needed to complete the proof of (1.).

Lemma 6.1 Under the assumptions of Theorem 3.1, for any $j = 1, \dots, K^*$,

$$\frac{v_{n_j^*}}{n_j^*} \| \widehat{L}_n(T_j^*, \theta) - \widehat{L}_{n,j}(T_j^*, \theta) \|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$

From (6.4) and Lemma 6.1, we have

$$\left\| \frac{1}{n_j^*} \widehat{L}_n(T_j^*, \theta) - \mathbb{E}(\ell_{1,j}(\theta)) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$
 (6.5)

(2.) Now, we show that the function $\theta \mapsto \mathbb{E}(\ell_{1,j}(\theta))$ has a unique maximum in θ_j^* . We will proceed as in Doukhan and Kengne (2015). For any $\theta \in \Theta$, define $L^{(j)}(\theta) := \mathbb{E}[\ell_{1,j}(\theta)]$. Let $\theta \in \Theta$, with $\theta \neq \theta_j^*$. We have

$$\begin{split} L^{(j)}(\theta_{j}^{*}) - L^{(j)}(\theta) &= \mathbb{E}[\ell_{1,j}(\theta_{j}^{*})] - \mathbb{E}[\ell_{1,j}(\theta)] \\ &= \mathbb{E}[Y_{1,j} \log f_{1,j}^{\theta_{j}^{*}} - f_{1,j}^{\theta_{j}^{*}}] - \mathbb{E}[Y_{1,j} \log f_{1,j}^{\theta} - f_{1,j}^{\theta}] \\ &= \mathbb{E}[f_{1,j}^{\theta_{j}^{*}} (\log f_{1,j}^{\theta_{j}^{*}} - \log f_{1,j}^{\theta})] - \mathbb{E}[f_{1,j}^{\theta_{j}^{*}} - f_{1,j}^{\theta}]. \end{split}$$

We apply the mean value theorem at the function $x \mapsto \log x$ defined in $[\underline{c}, +\infty[$. There exists ξ between $f_{1,j}^{\theta_j^*}$ and $f_{1,j}^{\theta}$ such that

$$\log f_{1,j}^{\theta_j^*} - \log f_{1,j}^{\theta} = \frac{1}{\xi} (f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta}).$$

Hence,

$$\begin{split} L^{(j)}(\theta_j^*) - L^{(j)}(\theta) &= \mathbb{E}\Big[\frac{f_{1,j}^{\theta_j^*}}{\xi}(f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta})\Big] - \mathbb{E}\Big[f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta}\Big] \\ &= \mathbb{E}\Big[\Big(\frac{f_{1,j}^{\theta_j^*}}{\xi} - 1\Big)\left(f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta}\right)\Big] = \mathbb{E}\Big[\frac{1}{\xi}(f_{1,j}^{\theta_j^*} - \xi)(f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta})\Big]. \end{split}$$

From assumption (**A0**), it follows that $\frac{1}{\xi}(f_{1,j}^{\theta_j^*} - \xi)(f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta}) \neq 0$ a.s, since $\theta \neq \theta_j^*$. Moreover,

- if $f_{1,j}^{\theta_j^*} < f_{1,j}^{\theta}$, then $f_{1,j}^{\theta_j^*} < \xi < f_{1,j}^{\theta}$ and hence $\frac{1}{\xi} (f_{1,j}^{\theta_j^*} \xi) (f_{1,j}^{\theta_j^*} f_{1,j}^{\theta}) > 0$;
- if $f_{1,j}^{\theta_j^*} > f_{1,j}^{\theta}$, then $f_{1,j}^{\theta} < \xi < f_{1,j}^{\theta_j^*}$ and hence $\frac{1}{\xi} (f_{1,j}^{\theta_j^*} \xi) (f_{1,j}^{\theta_j^*} f_{1,j}^{\theta}) > 0$.

We deduce that $\frac{1}{\xi}(f_{1,j}^{\theta_j^*} - \xi)(f_{1,j}^{\theta_j^*} - f_{1,j}^{\theta}) > 0$ a.s. Hence, $L^{(j)}(\theta_j^*) - L^{(j)}(\theta) > 0$ and the function $\theta \mapsto \mathbb{E}(\ell_{1,j}(\theta))$ has a unique maximum in θ_j^* .

(1.), (2.) and the standard arguments lead to the conclusion.

Proof of Lemma 6.1

For any $j = 1, \dots, K^*$, remark that

$$\begin{split} \|\widehat{L}_n(T_j^*,\theta) - \widehat{L}_{n,j}(T_j^*,\theta)\|_{\Theta} &\leq \sum_{t \in T_j^*} \|\widehat{\ell}_t(\theta) - \widehat{\ell}_{t,j}(\theta)\|_{\Theta} \\ &\leq \sum_{t \in T_j^*} \|Y_{t,j}\log\widehat{\lambda}_t(\theta) - \widehat{\lambda}_t(\theta) - Y_{t,j}\log\widehat{\lambda}_{t,j}(\theta) + \widehat{\lambda}_{t,j}(\theta)\|_{\Theta} \\ &\leq \sum_{t \in T_j^*} (Y_{t,j}\|\log\widehat{f}_t^{\theta} - \log\widehat{f}_{t,j}^{\theta}\|_{\Theta} + \|\widehat{f}_t^{\theta} - \widehat{f}_{t,j}^{\theta}\|_{\Theta}). \end{split}$$

According to the proprieties of the function $x \mapsto \log x$, we can show that $\|\log \widehat{f}_t^{\theta} - \log \widehat{f}_{t,j}^{\theta}\|_{\Theta} \leq \frac{1}{\underline{c}} \|\widehat{f}_t^{\theta} - \widehat{f}_{t,j}^{\theta}\|_{\Theta}$. Moreover, for $t \in T_j^*$, we have

$$\|\widehat{f}_{t}^{\theta} - \widehat{f}_{t,j}^{\theta}\|_{\Theta} = \|f(Y_{t-1}, \dots, Y_{1}, 0, \dots; \theta) - f(Y_{t-1,j}, \dots, Y_{t_{j-1}^{*}+1,j}, 0, \dots; \theta)\|_{\Theta}$$

$$\leq \sum_{k=t-t_{j-1}^{*}}^{t-1} \alpha_{k}^{(0)} |Y_{t-k}| \leq \sum_{k\geq t-t_{j-1}^{*}}^{t} \alpha_{k}^{(0)} Y_{t-k}.$$
(6.6)

Hence

$$\begin{split} \frac{v_{n_{j}^{*}}}{n_{j}^{*}} \| \widehat{L}_{n}(T_{j}^{*}, \theta) - \widehat{L}_{n,j}(T_{j}^{*}, \theta) \|_{\Theta} &\leq \frac{v_{n_{j}^{*}}}{n_{j}^{*}} \sum_{t \in T_{j}^{*}} \left[\left(\frac{Y_{t,j}}{\underline{c}} + 1 \right) \| \widehat{f}_{t}^{\theta} - \widehat{f}_{t,j}^{\theta} \|_{\Theta} \right] \\ &\leq \frac{v_{n_{j}^{*}}}{n_{j}^{*}} \sum_{t \in T_{j}^{*}} \sum_{k \geq t - t_{j-1}^{*}} \alpha_{k}^{(0)} \left[\left(\frac{Y_{t,j}}{\underline{c}} + 1 \right) Y_{t-k} \right] \\ &\leq \frac{v_{n_{j}^{*}}}{n_{j}^{*}} \sum_{\ell=1}^{n_{j}^{*}} \sum_{k > \ell} \alpha_{k}^{(0)} \left[\left(\frac{Y_{\ell+t_{j-1}^{*},j}}{\underline{c}} + 1 \right) Y_{\ell+t_{j-1}^{*} - k} \right]. \end{split}$$

By using Kounias and Weng (1969), it suffices to show that

$$\sum_{\ell>1} \frac{v_{\ell}}{\ell} \mathbb{E}\left[\sum_{k>\ell} \alpha_k^{(0)} \left(\frac{Y_{\ell+t_{j-1}^*,j}}{\underline{c}} + 1\right) Y_{\ell+t_{j-1}^*-k}\right] < \infty.$$

By using Hölder's inequality, for any $\ell \geq 1$, $k \geq \ell$, it holds that (see (2.2))

$$\mathbb{E}\Big[\Big(\frac{Y_{\ell+t^*_{j-1},j}}{c}+1\Big)Y_{\ell+t^*_{j-1}-k}\Big] \leq \Big(\mathbb{E}\Big[\Big(\frac{Y_{\ell+t^*_{j-1},j}}{c}+1\Big)^2\Big]\Big)^{1/2} \times (\mathbb{E}Y^2_{\ell+t^*_{j-1}-k})^{1/2} = C < \infty.$$

Hence,

$$\sum_{\ell \ge 1} \frac{v_{\ell}}{\ell} \mathbb{E} \Big[\sum_{k \ge \ell} \alpha_k^{(0)} \Big(\frac{Y_{\ell + t_{j-1}^*, j}}{\underline{c}} + 1 \Big) Y_{\ell + t_{j-1}^* - k} \Big] \le C \sum_{\ell \ge 1} \frac{1}{\kappa_{\ell}} \sum_{k \ge \ell} \alpha_k^{(0)} < \infty,$$

where the last equation follows from assumption (3.4) on the regularization parameter. Thus,

$$\frac{v_{n_j^*}}{n_j^*} \| \widehat{L}_n(T_j^*, \theta) - \widehat{L}_{n,j}(T_j^*, \theta) \|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$

6.2 Proof of Theorem 3.1

We will proceed as in Bardet *et al.* (2012). Firstly, we assume that K^* is known and we show $(\widehat{\underline{\tau}}_n, \widehat{\underline{\theta}}_n) \xrightarrow[n \to \infty]{\mathcal{P}} (\underline{\tau}^*, \underline{\theta}^*)$. Secondly, K^* is assumed to be unknown and we show $\widehat{K}_n \xrightarrow[n \to \infty]{\mathcal{P}} K^*$; which completes the proof of the theorem.

(1.) Assume that K^* is known and denote for any $\underline{t} \in \mathcal{M}_n(K^*)$:

$$\widehat{I}_n(\underline{t}) := \widehat{J}_n(K^*, \underline{t}, \widehat{\underline{\theta}}_n(\underline{t})) = -2 \sum_{k=1}^{K^*} \sum_{j=1}^{K^*} \widehat{L}_n(T_k \cap T_j^*, \widehat{\theta}_n(T_k)).$$

Hence, $\widehat{\underline{t}}_n = \underset{\underline{t} \in \mathcal{M}_n(K^*)}{\operatorname{argmin}} (\widehat{I}_n(\underline{t}))$. Let us show that $\widehat{\underline{\tau}}_n \xrightarrow{\mathcal{P}} \underline{\tau}^*$; which will implies that $\widehat{\theta}_n(\widehat{T}_{n,j}) \xrightarrow{\mathcal{P}} \widehat{\theta}_n(T_j^*)$ and from Proposition 2.1, $\widehat{\theta}_n(\widehat{T}_{n,j}) \xrightarrow{\mathcal{P}} \theta_j^*$ for all $j = 1, \cdots, K^*$. Without loss of generality, we assume that $K^* = 2$. Let t^* be the change-point location and $(u_n)_{n \geq 1}$ be a sequence of positive integers satisfying $u_n \to \infty$, $u_n/n \to 0$. For some $0 < \eta < 1$, define

$$V_{\eta,u_n} = \{ t \in \mathbb{Z} \ / \ |t - t^*| > \eta n \ ; \ u_n \le t \le n - u_n \},$$

$$W_{\eta,u_n} = \{ t \in \mathbb{Z} \ / \ |t - t^*| > \eta n \ ; \ 0 < t < u_n \text{ or } n - u_n < t \le n \}.$$

Remark that we have asymptotically $\mathbb{P}(\|\widehat{\underline{\tau}}_n - \underline{\tau}^*\|_m > \eta) \simeq \mathbb{P}(\|\widehat{t}_n - t^*\|_m > \eta n)$. But

$$\begin{split} \mathbb{P}(\|\widehat{t}_{n} - t^{*}\|_{m} > \eta n) &\leq \mathbb{P}(\widehat{t}_{n} \in V_{\eta, u_{n}}) + \mathbb{P}(\widehat{t}_{n} \in W_{\eta, u_{n}}) \\ &\leq \mathbb{P}(\min_{t \in V_{\eta, u_{n}}}(\widehat{I}_{n}(t) - \widehat{I}_{n}(t^{*})) \leq 0) + \mathbb{P}(\min_{t \in W_{\eta, u_{n}}}(\widehat{I}_{n}(t) - \widehat{I}_{n}(t^{*})) \leq 0). \end{split}$$

We will show that these two probabilities tend to 0. Let us show that $\mathbb{P}(\min_{t \in V_{\eta,u_n}}(\widehat{I}_n(t) - \widehat{I}_n(t^*)) \leq 0) \to 0$. Let $t \in V_{\eta,u_n}$ satisfying $t \geq t^*$ (without loss of generality). Then, we have $T_1 \cap T_1^* = T_1^*$, $T_2 \cap T_1^* = \emptyset$ and $T_2 \cap T_2^* = T_2$. We have the decomposition

$$\widehat{I}_{n}(t) - \widehat{I}_{n}(t^{*}) = 2 \left[\widehat{L}_{n}(T_{1}^{*}, \widehat{\theta}_{n}(T_{1}^{*})) - \widehat{L}_{n}(T_{1}^{*}, \widehat{\theta}_{n}(T_{1})) - \widehat{L}_{n}(T_{1} \cap T_{2}^{*}, \widehat{\theta}_{n}(T_{1})) + \widehat{L}_{n}(T_{2}, \widehat{\theta}_{n}(T_{2}^{*})) - \widehat{L}_{n}(T_{2}, \widehat{\theta}_{n}(T_{2})) + \widehat{L}_{n}(T_{1} \cap T_{2}^{*}, \widehat{\theta}_{n}(T_{2}^{*})) \right].$$

$$(6.7)$$

Since $|T_1^*| = t^*$, $|T_1 \cap T_2^*| = t - t^*$ and $|T_2| = n - t \ge u_n$, each term tends to ∞ with n. Recall that for $j = 1, 2, L^{(j)}(\theta) = \mathbb{E}(\ell_{1,j}(\theta))$. According to Proposition 2.1 and relation (6.5), we get the

Recall that for $j = 1, 2, L^{(j)}(\theta) = \mathbb{E}(\ell_{1,j}(\theta))$. According to Proposition 2.1 and relation (6.5), we get the following convergences, uniformly on V_{η,u_n} ,

$$\widehat{\theta}_n(T_1^*) \xrightarrow[n \to \infty]{a.s.} \theta_1^*, \quad \widehat{\theta}_n(T_2^*) \xrightarrow[n \to \infty]{a.s.} \theta_2^*, \quad \widehat{\theta}_n(T_2) \xrightarrow[n \to \infty]{a.s.} \theta_2^*, \quad \left\| \frac{1}{n} \widehat{L}_n(T_1^*, \theta) - \tau_1^* L^{(1)}(\theta) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0,$$

$$\left\| \frac{1}{t - t^*} \widehat{L}_n(T_1 \cap T_2^*, \theta) - L^{(2)}(\theta) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0 \text{ and } \left\| \frac{1}{n - t} \widehat{L}_n(T_2, \theta) - L^{(2)}(\theta) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$

Set $\eta_n = (t - t^*)/n$; clearly, $\varepsilon_n \in (\eta, 1)$. From (6.7), we get

$$\frac{1}{2n}(\widehat{I}_n(t) - \widehat{I}_n(t^*)) = \frac{1}{n}(\widehat{L}_n(T_1^*, \widehat{\theta}_n(T_1^*)) - \widehat{L}_n(T_1^*, \widehat{\theta}_n(T_1))) + \frac{1}{n}(\widehat{L}_n(T_2, \widehat{\theta}_n(T_2^*)) - \widehat{L}_n(T_2, \widehat{\theta}_n(T_2)))
+ \frac{1}{n}(\widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*)) - \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_1)))
= \tau_1^*(L^{(1)}(\theta_1^*) - L^{(1)}(\widehat{\theta}_n(T_1))) + o(1) + o(1) + \eta_n(L^{(2)}(\theta_2^*) - L^{(2)}(\widehat{\theta}_n(T_1))) + o(1)
= \tau_1^*(L^{(1)}(\theta_1^*) - L^{(1)}(\widehat{\theta}_n(T_1))) + \eta_n(L^{(2)}(\theta_2^*) - L^{(2)}(\widehat{\theta}_n(T_1))) + o(1).$$
(6.8)

Let V_1 and V_2 be two disjoint open neighborhoods of θ_1^* and θ_2^* respectively. For j=1,2, define

$$\delta_j = \inf_{\theta \in \mathcal{V}_i^c} \left(L^{(j)}(\theta_j^*) - L^{(j)}(\theta) \right).$$

Remark that $\delta_j > 0$ since the function $\theta \mapsto L^{(j)}(\theta)$ has a strict maximum in θ_j^* (see proof of Proposition 2.1). With $\varepsilon_n = \min(\tau_1^* \delta_1, \eta_n \delta_2)$ and $\varepsilon = \min(\tau_1^* \delta_1, \eta \delta_2)$, we have

- if $\widehat{\theta}_n(T_1) \in \mathcal{V}_1$, that is $\widehat{\theta}_n(T_1) \in \mathcal{V}_2^c$, then $\eta_n(L^{(2)}(\theta_2^*) L^{(2)}(\widehat{\theta}_n(T_1))) > \eta_n \delta_2$;
- if $\widehat{\theta}_n(T_1) \notin \mathcal{V}_1$, that is $\widehat{\theta}_n(T_1) \in \mathcal{V}_1^c$, then $\tau_1^* \left(L^{(1)}(\theta_1^*) L^{(1)}(\widehat{\theta}_n(T_1)) \right) > \tau_1^* \delta_1$.

In both cases, $\frac{1}{2n}(\widehat{I}_n(t) - \widehat{I}_n(t^*)) \ge \varepsilon_n + o(1) \ge \varepsilon + o(1)$, for any $t \in V_{\eta,u_n}$. This implies $\mathbb{P}\left(\min_{t \in V_{\eta,u_n}} \left(\widehat{I}_n(t) - \widehat{I}_n(t^*)\right) \le 0\right) \to 0$. By going along similar lines, one can prove that $\mathbb{P}\left(\min_{t \in W_{\eta,u_n}} \left(\widehat{I}_n(t) - \widehat{I}_n(t^*)\right) \le 0\right) \to 0$. Then, it follows that $\eta > 0$, $\mathbb{P}\left(\|\widehat{\underline{\tau}}_n - \underline{\tau}^*\|_m > \eta\right) \to 0$ as $n \to \infty$.

(2.) Now, assume that K^* is unknown. For $K \geq 2$, $x = (x_1, \dots, x_{K-1}) \in \mathbb{R}^{K-1}$, $y = (y_1, \dots, y_{K^*-1}) \in \mathbb{R}^{K^*-1}$, denote

$$||x - y||_{\infty} = \max_{1 \le j \le K^* - 1} \min_{1 \le k \le K - 1} |x_k - y_j|.$$

The following lemma will be useful in the sequel. It follows from (1.) and the definition of $\|\cdot\|_{\infty}$.

Lemma 6.2 Let $K \geq 1$, $(\underline{\hat{t}}_n, \underline{\hat{\theta}}_n)$ the estimator obtained by minimizing $\widehat{J}_n(K, \underline{t}, \underline{\theta})$ on $\mathcal{M}_n(K) \times \Theta^K$ and $\underline{\hat{\tau}}_n = \underline{\hat{t}}_n/n$. Under the assumptions of Theorem 3.1, if $K \geq K^*$, then $\|\underline{\hat{\tau}}_n - \underline{\tau}^*\|_{\infty} \xrightarrow[n \to \infty]{\mathcal{P}} 0$.

We will also use the following lemma, which the proof follows from the Lemma 3.3 of Lavielle and Ludena (2000) and the argument given in the proof of Lemma 6.5 in [6].

Lemma 6.3 Under the assumptions of Theorem 3.1, for any $K \geq 2$, there exists $C_K > 0$ such that:

$$\forall (\underline{t},\underline{\theta}) \in \mathcal{M}_n(K) \times \Theta^K, \quad e_n(\underline{t},\underline{\theta}) = 2\sum_{j=1}^{K^*} \sum_{k=1}^K \frac{n_{k,j}}{n} \left(L^{(j)}(\theta_j^*) - L^{(j)}(\theta_k) \right) \ge \frac{C_K}{n} \|\underline{t} - \underline{t}^*\|_{\infty}$$

where $L^{(j)}(\theta) = \mathbb{E}(\ell_{1,j}(\theta))$ for all $\theta \in \Theta$ and $j = 1, \dots, K^*$.

Now, let us use the Lemma 6.2 and 6.3 to show that $\widehat{K}_n \xrightarrow[n \to \infty]{\mathcal{P}} K^*$. To this end, we will show that $\mathbb{P}(\widehat{K}_n = K) \xrightarrow[n \to \infty]{} 0$, for $K < K^*$ and $K^* < K \le K_{\text{max}}$ separately. In any case, we have

$$\mathbb{P}(\widehat{K}_{n} = K) \leq \mathbb{P}\left(\inf_{(\underline{t},\underline{\theta}) \in \mathcal{M}_{n}(K) \times \Theta^{K}} (\widehat{J}_{n}(K,\underline{t},\underline{\theta})) \leq \widehat{J}_{n}(K^{*},\underline{t}^{*},\underline{\theta}^{*})\right) \\
\leq \mathbb{P}\left(\inf_{(\underline{t},\underline{\theta}) \in \mathcal{M}_{n}(K) \times \Theta^{K}} (\widehat{J}_{n}(K,\underline{t},\underline{\theta}) - \widehat{J}_{n}(K^{*},\underline{t}^{*},\underline{\theta}^{*})) \leq \frac{n}{v_{n}}(K^{*} - K)\right).$$
(6.9)

i-) For $K < K^*$, decompose $\widehat{J}_n(K, \underline{t}, \underline{\theta}) - \widehat{J}_n(K^*, \underline{t}^*, \underline{\theta}^*) = n \left(d_n(\underline{t}, \underline{\theta}) + e_n(\underline{t}, \underline{\theta}) \right)$, where $e_n(\underline{t}, \underline{\theta})$ is defined in Lemma 6.3 and

$$d_n(\underline{t},\underline{\theta}) = 2\Big[\sum_{i=1}^{K^*} \frac{n_j^*}{n} \Big(\frac{\widehat{L}_n\left(T_j^*,\theta_j^*\right)}{n_j^*} - L^{(j)}(\theta_j^*)\Big) + \sum_{k=1}^K \sum_{i=1}^{K^*} \frac{n_{k,j}}{n} \Big(L^{(j)}(\theta_k) - \frac{\widehat{L}_n\left(T_j^* \cap T_k,\theta_k\right)}{n_{k,j}}\Big)\Big].$$

Hence, from (6.9), we have

$$\mathbb{P}(\widehat{K}_n = K) \le \mathbb{P}\left(\inf_{(\underline{t}, \underline{\theta}) \in \mathcal{M}_n(K) \times \Theta^K} (d_n(\underline{t}, \underline{\theta}) + e_n(\underline{t}, \underline{\theta})) \le \frac{1}{v_n} (K^* - K)\right). \tag{6.10}$$

The equation (6.4) ensures that $d_n(\underline{t},\underline{\theta}) \to 0$ a.s. and uniformly on $\mathcal{M}_n(K) \times \Theta^K$. According to Lemma 6.3, there exists $C_K > 0$ such that $e_n(\underline{t},\underline{\theta}) \geq \frac{C_K}{n} \|\underline{t} - \underline{t}^*\|_{\infty}$, for all $(\underline{t},\underline{\theta}) \in \mathcal{M}_n(K) \times \Theta^K$. Since $K < K^*$, for any $\underline{t} \in \mathcal{M}_n(K)$, we have $\frac{1}{n} \|\underline{t} - \underline{t}^*\|_{\infty} = \|\underline{\tau} - \underline{\tau}^*\|_{\infty} \geq \min_{1 \leq j \leq K^*} (\tau_j^* - \tau_{j-1}^*)/2 > 0$. Then $e_n(\underline{t},\underline{\theta}) > 0$ for $(\underline{t},\underline{\theta}) \in \mathcal{M}_n(K) \times \Theta^K$ and since $\frac{1}{v_n} \xrightarrow[n \to \infty]{} 0$, we deduce from (6.10) that $\mathbb{P}(\widehat{K}_n = K) \xrightarrow[n \to \infty]{} 0$.

ii-) Now, assume that $K^* < K \le K_{\max}$. Denote $\widehat{\underline{t}}_n = (\widehat{t}_{n,1}, \dots, \widehat{t}_{n,K^*})$. From (6.9) and the Markov's inequality, we have:

$$\mathbb{P}(\widehat{K}_{n} = K) \leq \mathbb{P}(\widehat{J}_{n}(K, \underline{\widehat{t}}_{n}, \underline{\widehat{\theta}}_{n}) - \widehat{J}_{n}(K^{*}, \underline{t}^{*}, \underline{\theta}^{*}) + \frac{n}{v_{n}}(K - K^{*}) \leq 0)$$

$$\leq \mathbb{P}(|\widehat{J}_{n}(K, \underline{\widehat{t}}_{n}, \underline{\widehat{\theta}}_{n}) - \widehat{J}_{n}(K^{*}, \underline{t}^{*}, \underline{\theta}^{*})| > \frac{n}{v_{n}})$$

$$\leq \frac{v_{n}}{n} \mathbb{E}|\widehat{J}_{n}(K, \underline{\widehat{t}}_{n}, \underline{\widehat{\theta}}_{n}) - \widehat{J}_{n}(K^{*}, \underline{t}^{*}, \underline{\theta}^{*})|.$$
(6.11)

By Lemma 6.2, there exists some subset $\{k_j, 1 \leq j \leq K^* - 1\} \subset \{1, \dots, K - 1\}$ such that for any $j = 1, \dots, K^* - 1$, $\widehat{t}_{n,k_j}/n \to \tau_j^*$. Set $k_0 = 0$ and $k_{K^*} = K$. We have

$$\widehat{J}_{n}(K, \underline{\widehat{t}}_{n}, \underline{\widehat{\theta}}_{n}) - \widehat{J}_{n}(K^{*}, \underline{t}^{*}, \underline{\theta}^{*}) = 2\left(\sum_{j=1}^{K^{*}} \widehat{L}_{n}(T_{j}^{*}, \theta_{j}^{*}) - \sum_{k=1}^{K} \widehat{L}_{n}(\widehat{T}_{n,k}, \widehat{\theta}_{n,k})\right) \\
= 2\sum_{j=1}^{K^{*}} \left[\widehat{L}_{n}(T_{j}^{*}, \theta_{j}^{*}) - \sum_{k=k_{j-1}+1}^{k_{j}} \widehat{L}_{n}(\widehat{T}_{n,k}, \widehat{\theta}_{n,k})\right]$$

and from (6.11), it follows that

$$\mathbb{P}(\widehat{K}_{n} = K) \leq \frac{2v_{n}}{n} \sum_{j=1}^{K^{*}} \mathbb{E} |\widehat{L}_{n}(T_{j}^{*}, \theta_{j}^{*}) - \sum_{k=k_{j-1}+1}^{k_{j}} \widehat{L}_{n}(\widehat{T}_{n,k}, \widehat{\theta}_{n,k})| \\
\leq C \sum_{j=1}^{K^{*}} \frac{v_{n_{j}^{*}}}{n_{j}^{*}} \mathbb{E} |\widehat{L}_{n}(T_{j}^{*}, \theta_{j}^{*}) - \sum_{k=k_{j-1}+1}^{k_{j}} \widehat{L}_{n}(\widehat{T}_{n,k}, \widehat{\theta}_{n,k})|.$$

For any $j=1,\cdots,K^*$, one can easily get from the proof of Lemma 6.1 that

$$\frac{v_{n_j^*}}{n_j^*} \mathbb{E} |\widehat{L}_n(T_j^*, \theta_j^*) - \sum_{k=k_{j-1}+1}^{k_j} \widehat{L}_n(\widehat{T}_{n,k}, \widehat{\theta}_{n,k})| \underset{n \to \infty}{\longrightarrow} 0,$$

and thus $\mathbb{P}(\widehat{K}_n = K) \xrightarrow[n \to \infty]{} 0$.

6.3 Proof of Theorem 3.2

Also for this proof, We will proceed as in [6]. Without loss of generality, we can assume that $K^* = 2$. Let $(u_n)_{n\geq 1}$ be a sequence satisfying $u_n \xrightarrow[n\to\infty]{} \infty$, $\frac{u_n}{n} \xrightarrow[n\to\infty]{} 0$ and $\mathbb{P}(|\widehat{\underline{t}}_n - \underline{t}^*| > u_n) \xrightarrow[n\to\infty]{} 0$ (for instance $u_n = \sqrt{\max(\mathbb{E}|\widehat{\tau}_n - \tau^*|, n^{-1})}$). For any $\delta > 0$, since we have

$$\mathbb{P}(|\widehat{t}_n - t^*| > \delta) \le \mathbb{P}(\delta < |\widehat{t}_n - t^*| \le u_n) + \mathbb{P}(|\widehat{t}_n - t^*| > u_n),$$

it suffices to show that $\lim_{\delta \to \infty} \lim_{n \to \infty} \mathbb{P}(\delta < |\hat{t}_n - t^*| \le u_n) = 0.$

Denote $V_{\delta,u_n} = \{t \in \mathbb{Z}/\ \delta < |\widehat{t}_n - t^*| \le u_n\}$. Then

$$\mathbb{P}(\delta < |\widehat{t}_n - t^*| \le u_n) \le \mathbb{P}\left(\max_{t \in V_{\delta, u_n}} (\widehat{I}_n(t) - \widehat{I}_n(t^*)) \le 0\right).$$

Let $t \in V_{\delta,u_n}$ (for example $t \geq t^*$). With the notation of the proof of Theorem 3.1, we have $\widehat{L}_n(T_1^*,\widehat{\theta}_n(T_1^*)) \geq \widehat{L}_n(T_1^*,\widehat{\theta}_n(T_1))$ and from (6.7) we get

$$\widehat{I}_n(t) - \widehat{I}_n(t^*) \ge \widehat{L}_n\big(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*)\big) - \widehat{L}_n\big(T_1 \cap T_2^*, \widehat{\theta}_n(T_1)\big) + \widehat{L}_n\big(T_2, \widehat{\theta}_n(T_2^*)\big) - \widehat{L}_n\big(T_2, \widehat{\theta}_n(T_2)\big).$$

We consider the following two steps.

(1.) Let us show that $\widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*)) - \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_1)) > 0$, for n large enough.

For any $\theta \in \Theta$, we have $\frac{1}{n}\widehat{L}_n(T_1,\theta) = \frac{t^*}{n}\frac{\widehat{L}_n(T_1^*,\theta)}{t^*} + \frac{t-t^*}{n}\frac{\widehat{L}_n(T_1\cap T_2^*,\theta)}{t-t^*}$ and since $\frac{t-t^*}{n} \leq \frac{u_n}{n} \underset{n\to\infty}{\longrightarrow} 0$, it follows that

$$\widehat{\theta}_n(T_1) = \underset{\theta \in \Theta}{\operatorname{argmax}} \left(\frac{1}{n} \widehat{L}_n(T_1, \theta) \right) \xrightarrow[n, \delta \to \infty]{a.s} \theta_1^*.$$

Hence, $\frac{1}{t-t^*}(\widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*)) - \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_1)))$ converges a.s. and uniformly on V_{δ,u_n} to $L^{(2)}(\theta_2^*) - L^{(2)}(\theta_1^*) > 0$.

(2.) Let us show that $\frac{1}{t-t^*}(\widehat{L}_n(T_2,\widehat{\theta}_n(T_2^*)) - \widehat{L}_n(T_2,\widehat{\theta}_n(T_2))) \xrightarrow[n,\delta\to\infty]{a.s} 0$. For large value of n, remark that $\widehat{\theta}_n(T_2) \in \stackrel{\circ}{\Theta}$ so that $\partial \widehat{L}_n(T_2,\widehat{\theta}_n(T_2))/\partial \theta = 0$. From mean value theorem applied on $\partial \widehat{L}_n/\partial \theta_i$ for any $i = 1, \dots, d$, there exists $\widetilde{\theta}_{n,i} \in [\widehat{\theta}_n(T_2), \widehat{\theta}_n(T_2^*)]$ such that

$$0 = \frac{\partial \widehat{L}_n(T_2, \widehat{\theta}_n(T_2^*))}{\partial \theta_i} + \frac{\partial^2 \widehat{L}_n(T_2, \widetilde{\theta}_{n,i})}{\partial \theta \partial \theta_i} (\widehat{\theta}_n(T_2) - \widehat{\theta}_n(T_2^*))$$
(6.12)

where for $a, b \in \mathbb{R}^d$, $[a, b] = {\lambda a + (1 - \lambda)b; \lambda \in [0, 1]}.$

According the equalities $\widehat{L}_n(T_2^*,\theta) = \widehat{L}_n(T_1 \cap T_2^*,\theta) + \widehat{L}_n(T_2,\theta)$ and $\partial \widehat{L}_n(T_2^*,\widehat{\theta}_n(T_2^*))/\partial \theta = 0$, it comes from (6.12) that

$$\frac{\partial \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*))}{\partial \theta_i} = \frac{\partial^2 \widehat{L}_n(T_2, \widetilde{\theta}_{n,i})}{\partial \theta \partial \theta_i} (\widehat{\theta}_n(T_2) - \widehat{\theta}_n(T_2^*)), \text{ for any } i = 1, \dots, d,$$

and it follows that

$$\frac{1}{t-t^*} \frac{\partial \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*))}{\partial \theta} = \frac{n-t}{t-t^*} \widetilde{A}_n \cdot (\widehat{\theta}_n(T_2) - \widehat{\theta}_n(T_2^*))$$
(6.13)

where
$$\widetilde{A}_n := \left(\frac{1}{n-t} \frac{\partial^2 \widehat{L}_n(T_2, \widetilde{\theta}_{n,i})}{\partial \theta \partial \theta_i}\right)_{1 \leq i \leq d}$$
.

The following lemma will be useful in the sequel.

Lemma 6.4

• Assume that the conditions of Theorem 3.3 hold. Then, for any $j = 1, \dots, K^*$,

$$\frac{1}{\sqrt{n_j^*}} \left\| \frac{\partial^i \widehat{L}_n(T_j^*, \theta)}{\partial \theta^i} - \frac{\partial^i \widehat{L}_{n,j}(T_j^*, \theta)}{\partial \theta^i} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0; \tag{6.14}$$

• Assume that the conditions of Theorem 3.2 hold. Then, for any $j = 1, \dots, K^*$,

$$\left\| \frac{1}{n_j^*} \frac{\partial \widehat{L}_n(T_j^*, \theta)}{\partial \theta} - \frac{\partial L^{(j)}(\theta)}{\partial \theta} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0, \quad where \quad \frac{\partial L^{(j)}(\theta)}{\partial \theta} = \mathbb{E}\left(\frac{\partial \ell_{1,j}(\theta)}{\partial \theta}\right). \tag{6.15}$$

Hence, (6.15) gives
$$\underbrace{\frac{1}{t-t^*}} \frac{\partial \widehat{L}_n(T_1 \cap T_2^*, \widehat{\theta}_n(T_2^*))}{\partial \theta} \xrightarrow[n,\delta \to \infty]{a.s.} \underbrace{\frac{\partial L^{(2)}(\theta_2^*)}{\partial \theta}} = 0$$
 and $\widetilde{A}_n \xrightarrow[n,\delta \to \infty]{a.s.} \mathbb{E}\left(\frac{\partial^2 \ell_{1,2}(\theta_2^*)}{\partial \theta^2}\right)$.

Since $\mathbb{E}(\frac{\partial^2 \ell_{1,2}(\theta_2^*)}{\partial \theta^2})$ is a nonsingular matrix (see [1]), we deduce from (6.13) that

$$\frac{n-t}{t-t^*}(\widehat{\theta}_n(T_2) - \widehat{\theta}_n(T_2^*)) \xrightarrow[n.\delta \to \infty]{a.s} 0.$$
(6.16)

We conclude by the Taylor expansion on \widehat{L}_n that

$$\frac{1}{t-t^*}|\widehat{L}_n(T_2,\widehat{\theta}_n(T_2)) - \widehat{L}_n(T_2,\widehat{\theta}_n(T_2^*))| \le \frac{1}{2(t-t^*)} \|\widehat{\theta}_n(T_2) - \widehat{\theta}_n(T_2^*)\|^2 \sup_{\theta \in \Theta} \left\| \frac{\partial^2 \widehat{L}_n(T_2,\theta)}{\partial \theta^2} \right\| \to 0 \quad a.s.$$

Proof of Lemma 6.4

We detail the proof for the first order derivation; the proof for the second order derivation follows the same reasoning.

Let $j \in \{1, \dots, K^*\}$ and $l \in \{1, \dots, d\}$, we will show that

$$\frac{1}{\sqrt{n_j^*}} \left\| \frac{\partial \widehat{L}_n(T_j^*, \theta)}{\partial \theta_l} - \frac{\partial \widehat{L}_{n,j}(T_j^*, \theta)}{\partial \theta_l} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s} 0.$$
 (6.17)

Remark that

$$\begin{split} \frac{1}{\sqrt{n_{j}^{*}}} \left\| \frac{\partial \widehat{L}_{n}(T_{j}^{*}, \theta)}{\partial \theta_{l}} - \frac{\partial \widehat{L}_{n,j}(T_{j}^{*}, \theta)}{\partial \theta_{l}} \right\|_{\Theta} &\leq \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left\| \frac{\partial \widehat{\ell}_{t}(\theta)}{\partial \theta_{l}} - \frac{\partial \widehat{\ell}_{t,j}(\theta)}{\partial \theta_{l}} \right\|_{\Theta} \\ &\leq \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left[Y_{t,j} \left\| \frac{1}{\widehat{f}_{t}^{\theta}} \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{1}{\widehat{f}_{t,j}^{\theta}} \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} + \left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \right]. \end{split}$$

Thus, by using the inequality $|a_1b_1 - a_2b_2| \le |a_1||b_1 - b_2| + |b_2||a_1 - a_2| \ \forall a_1, a_2, b_1, b_2 \in \mathbb{R}$, we have

$$\frac{1}{\sqrt{n_{j}^{*}}} \left\| \frac{\partial \widehat{L}_{n}(T_{j}^{*}, \theta)}{\partial \theta_{l}} - \frac{\partial \widehat{L}_{n,j}(T_{j}^{*}, \theta)}{\partial \theta_{l}} \right\|_{\Theta} \leq \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left[Y_{t,j} \left(\left\| \frac{1}{\widehat{f}_{t}^{\theta}} \right\|_{\Theta} \left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} + \left\| \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \right\| \frac{1}{\widehat{f}_{t}^{\theta}} - \frac{1}{\widehat{f}_{t,j}^{\theta}} \right\|_{\Theta} \right) + \left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \right]$$

$$\leq \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left[\left(\frac{Y_{t,j}}{\underline{c}} + 1 \right) \left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} + \frac{1}{\underline{c}^{2}} Y_{t,j} \left\| \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \left\| \widehat{f}_{t}^{\theta} - \widehat{f}_{t,j}^{\theta} \right\|_{\Theta} \right] \\
\leq C \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left[\left(Y_{t,j} + 1 \right) \left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} + Y_{t,j} \left\| \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \left\| \widehat{f}_{t}^{\theta} - \widehat{f}_{t,j}^{\theta} \right\|_{\Theta} \right].$$

For $t \in T_i^*$, from $\mathbf{A}_1(\Theta)$, we have

$$\left\| \frac{\partial \widehat{f}_{t}^{\theta}}{\partial \theta_{l}} - \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} = \left\| \frac{\partial}{\partial \theta_{l}} f(Y_{t-1}, \dots, Y_{1}, 0, \dots; \theta) - \frac{\partial}{\partial \theta_{l}} f(Y_{t-1,j}, \dots, Y_{t_{j-1}^{*}+1,j}, 0, \dots; \theta) \right\|_{\Theta}$$

$$\leq \sum_{k=t-t_{j-1}^{*}}^{t-1} \alpha_{k}^{(1)} |Y_{t-k}| \leq \sum_{k\geq t-t_{j-1}^{*}}^{t} \alpha_{k}^{(1)} Y_{t-k}. \tag{6.18}$$

Hence, from (6.6) and (6.18), we get

$$\frac{1}{\sqrt{n_{j}^{*}}} \left\| \frac{\partial \widehat{L}_{n}(T_{j}^{*}, \theta)}{\partial \theta_{l}} - \frac{\partial \widehat{L}_{n,j}(T_{j}^{*}, \theta)}{\partial \theta_{l}} \right\|_{\Theta} \leq C \frac{1}{\sqrt{n_{j}^{*}}} \sum_{t \in T_{j}^{*}} \left[\left(Y_{t,j} + 1 \right) \sum_{k \geq t - t_{j-1}^{*}} \alpha_{k}^{(1)} Y_{t-k} + Y_{t,j} \left\| \frac{\partial \widehat{f}_{t,j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \sum_{k \geq t - t_{j-1}^{*}} \alpha_{k}^{(0)} Y_{t-k} \right].$$

$$\leq C \frac{1}{\sqrt{n_{j}^{*}}} \sum_{\ell=1}^{n_{j}^{*}} \left[\left(Y_{\ell+t_{j-1}^{*},j} + 1 \right) \sum_{k \geq \ell} \alpha_{k}^{(1)} Y_{\ell+t_{j-1}^{*}-k} + Y_{\ell+t_{j-1}^{*},j} \left\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^{*},j}^{\theta}}{\partial \theta_{l}} \right\|_{\Theta} \sum_{k \geq \ell} \alpha_{k}^{(0)} Y_{\ell+t_{j-1}^{*}-k} \right].$$

According to [24], (6.17) holds if

$$\sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \mathbb{E} \left[\left(Y_{\ell + t_{j-1}^*, j} + 1 \right) \sum_{k \geq \ell} \alpha_k^{(1)} Y_{\ell + t_{j-1}^* - k} + Y_{\ell + t_{j-1}^*, j} \left\| \frac{\partial \widehat{f}_{\ell + t_{j-1}^*, j}^{\theta}}{\partial \theta_l} \right\|_{\Theta} \sum_{k \geq \ell} \alpha_k^{(0)} Y_{\ell + t_{j-1}^* - k} \right] < \infty. \tag{6.19}$$

We have

$$\sum_{\ell \ge 1} \frac{1}{\sqrt{\ell}} \mathbb{E} \Big[\Big(Y_{\ell + t_{j-1}^*, j} + 1 \Big) \sum_{k \ge \ell} \alpha_k^{(1)} Y_{\ell + t_{j-1}^* - k} \Big] = C \sum_{\ell \ge 1} \frac{1}{\sqrt{\ell}} \sum_{k \ge \ell} \alpha_k^{(1)} < \infty.$$

Moreover,

$$\left\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^*,j}^{\theta}}{\partial \theta_l} \right\|_{\Theta} = \left\| \frac{\partial}{\partial \theta_l} f(0, \dots; \theta) \right\|_{\Theta} + \left\| \frac{\partial}{\partial \theta_l} f(Y_{\ell+t_{j-1}^*-1,j}, \dots; Y_{t_{j-1}^*+1,j}, 0, \dots; \theta) - \frac{\partial}{\partial \theta_l} f(0, \dots; \theta) \right\|_{\Theta}$$

$$\leq C + \sum_{k \geq 1} \alpha_k^{(1)} Y_{\ell+t_{j-1}^*-k,j}.$$

Hence, from Minkowski's inequality,

$$\left\| \left\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^*,j}^{\theta}}{\partial \theta_l} \right\|_{\Theta} \right\|_{3} = \left(\mathbb{E} \left[\left\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^*,j}^{\theta}}{\partial \theta_l} \right\|_{\Theta}^{3} \right] \right)^{1/3} \le C + \sum_{k \ge 1} \alpha_k^{(1)} \|Y_{\ell+t_{j-1}^*-k,j}\|_{3} < C(1 + \sum_{k \ge 1} \alpha_k^{(1)}) < C < \infty.$$

Thus, by using Hölder and Minkowski's inequalities, it comes that

$$\sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \mathbb{E} \Big[Y_{\ell+t_{j-1}^*,j} \Big\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^*,j}^{\theta}}{\partial \theta_l} \Big\|_{\Theta} \sum_{k \geq \ell} \alpha_k^{(0)} Y_{\ell+t_{j-1}^*-k} \Big] \leq \sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \| Y_{\ell+t_{j-1}^*,j} \|_{3} \Big\| \Big\| \frac{\partial \widehat{f}_{\ell+t_{j-1}^*,j}^{\theta}}{\partial \theta_l} \Big\|_{\Theta} \Big\|_{3} \Big\| \sum_{k \geq \ell} \alpha_k^{(0)} Y_{\ell+t_{j-1}^*-k} \Big\|_{3} \\
\leq C \sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \Big\| \sum_{k \geq \ell} \alpha_k^{(0)} Y_{\ell+t_{j-1}^*-k} \Big\|_{3} \\
\leq C \sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \sum_{k > \ell} \alpha_k^{(0)} \| Y_{\ell+t_{j-1}^*-k} \|_{3} \leq C \sum_{\ell \geq 1} \frac{1}{\sqrt{\ell}} \sum_{k \geq \ell} \alpha_k^{(0)} < \infty.$$

Hence, (6.19) is satisfied, and it holds that,

$$\frac{1}{\sqrt{n_i^*}} \left\| \frac{\partial \widehat{L}_n(T_j^*, \theta)}{\partial \theta_l} - \frac{\partial \widehat{L}_{n,j}(T_j^*, \theta)}{\partial \theta_l} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s} 0, \text{ for any } j = 1, \dots, K^* \text{ and } l \in \{1, \dots, d\}.$$

Thus, (6.14) follows.

Remark that, one can go along the same lines as above by replacing \hat{L}_n by L_n or $\hat{L}_{n,j}$; and obtain

$$\frac{1}{\sqrt{n_j^*}} \left\| \frac{\partial L_n(T_j^*, \theta)}{\partial \theta} - \frac{\partial \widehat{L}_{n,j}(T_j^*, \theta)}{\partial \theta} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s} 0, \text{ for any } j = 1, \dots, K^*.$$
 (6.20)

$$\frac{1}{\sqrt{n_i^*}} \left\| \frac{\partial \widehat{L}_n(T_j^*, \theta)}{\partial \theta} - \frac{\partial L_{n,j}(T_j^*, \theta)}{\partial \theta} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s} 0, \text{ for any } j = 1, \dots, K^*.$$
 (6.21)

Moreover, similar arguments as above can be easily applied to get

$$\forall t \in \mathbb{Z}, \ \mathbb{E} \left\| \frac{\partial \ell_{t,j}(\theta)}{\partial \theta} \right\|_{\Theta} = C < \infty \quad \text{for any } j = 1, \dots, K^*.$$

By the uniform strong law of large number applied on the process $\{\frac{\partial \ell_{t,j}(\theta)}{\partial \theta}, t \in \mathbb{Z}\}$, it holds that, for $j = 1, \dots, K^*$,

$$\left\| \frac{1}{n_j^*} \frac{\partial L_{n,j}(T_j^*, \theta)}{\partial \theta} - \frac{\partial L^{(j)}(\theta)}{\partial \theta} \right\|_{\Theta} = \left\| \frac{1}{n_j^*} \sum_{t \in T_j^*} \frac{\partial \ell_{t,j}(\theta)}{\partial \theta} - \mathbb{E}(\frac{\partial \ell_{1,j}(\theta)}{\partial \theta}) \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$
 (6.22)

Thus, (6.15) follows from (6.21) and (6.22). This achieves the proof of the lemma.

6.4 Proof of Theorem 3.3

Remark that for any $j \in \{1, \dots, K^*\}$ $(\widehat{\theta}_n(\widehat{T}_j) - \theta_j^*) = (\widehat{\theta}_n(\widehat{T}_j) - \widehat{\theta}_n(T_j^*)) + (\widehat{\theta}_n(T_j^*) - \theta_j^*)$. According to Theorem 3.2, it comes $(\widehat{t}_j - \widehat{t}_j^*) = o_P(\log(n))$. By relation (6.16), we obtain $(\widehat{\theta}_n(\widehat{T}_j) - \widehat{\theta}_n(T_j^*)) = o_P(\frac{\log(n)}{n})$. Therefore, $\sqrt{n_j^*}(\widehat{\theta}_n(\widehat{T}_j) - \widehat{\theta}_n(T_j^*)) \xrightarrow[n \to \infty]{\mathcal{P}} 0$ and it suffices to show that $\sqrt{n_j^*}(\widehat{\theta}_n(T_j^*) - \theta_j^*) \xrightarrow[n \to \infty]{\mathcal{P}} \mathcal{N}_d(0, \Sigma_j)$ to conclude.

Recall that for n large enough, $\widehat{\theta}_n(T_j^*) \in \overset{\circ}{\Theta}$. By the mean value theorem, there exists $(\widetilde{\theta}_{n,k})_{1 \leq k \leq d} \in [\widehat{\theta}_n(T_j^*), \theta_j^*]$ such that

$$\frac{\partial L_n(T_j^*, \widehat{\theta}_n(T_j^*))}{\partial \theta_k} = \frac{\partial L_n(T_j^*, \theta_j^*)}{\partial \theta_k} + \frac{\partial^2 L_n(T_j^*, \widetilde{\theta}_{n,k})}{\partial \theta \partial \theta_k} \left(\widehat{\theta}_n(T_j^*) - \theta_j^*\right). \tag{6.23}$$

Let $J_n = -\left(\frac{1}{n_j^*} \frac{\partial^2 L_n(T_j^*, \widetilde{\theta}_{n,k})}{\partial \theta \partial \theta_k}\right)_{1 \leq k \leq d}$. By relation (6.20), Lemma 6.4 and the proof of Theorem 2.2 of [1], we obtain $J_n \xrightarrow[n \to \infty]{} J_j(\theta_j^*)$, where J_j is the matrix defined in Theorem 3.3. But, by Assumption (**A4**), it is easy to see that $J_j(\theta_j^*)$ is a non singular matrix. Thus, for n large enough, J_n is invertible and (6.23) gives

$$\sqrt{n_j^*}(\widehat{\theta}_n(T_j^*) - \theta_j^*) = -J_n^{-1} \left[\frac{1}{\sqrt{n_j^*}} \left(\frac{\partial L_n(T_j^*, \widehat{\theta}_n(T_j^*))}{\partial \theta} - \frac{\partial L_n(T_j^*, \theta_j^*)}{\partial \theta} \right) \right]. \tag{6.24}$$

From the proof of Theorem 2.2 of [1], we get

$$\frac{1}{\sqrt{n_j^*}} \frac{\partial L_{n,j}(T_j^*, \theta_j^*)}{\partial \theta} \xrightarrow[n \to \infty]{\mathcal{D}} \mathcal{N}_d(0, I_j(\theta_j^*))$$
(6.25)

where I_j is given in Theorem 3.3. According to (6.14), (6.20) and (6.21), we have

$$\frac{1}{\sqrt{n_j^*}} \left| \frac{\partial L_n(T_j^*, \theta_j^*)}{\partial \theta} - \frac{\partial L_{n,j}(T_j^*, \theta_j^*)}{\partial \theta} \right| \xrightarrow[n \to \infty]{a.s.} 0.$$

Hence, (6.25) implies

$$\frac{1}{\sqrt{n_j^*}} \frac{\partial L_n(T_j^*, \theta_j^*)}{\partial \theta} \xrightarrow[n \to \infty]{\mathcal{D}} \mathcal{N}_d(0, I_j(\theta_j^*)). \tag{6.26}$$

Moreover, from (6.14) and (6.20), we have

$$\frac{1}{\sqrt{n_j^*}} \left\| \frac{\partial L_n(T_j^*, \theta)}{\partial \theta} - \frac{\partial \widehat{L}_n(T_j^*, \theta)}{\partial \theta} \right\|_{\Theta} \xrightarrow[n \to \infty]{a.s.} 0.$$

Therefore, since $\partial \widehat{L}_n(T_i^*, \widehat{\theta}_n(T_i^*))/\partial \theta = 0$, it follows that

$$\frac{1}{\sqrt{n_j^*}} \frac{\partial L_n(T_j^*, \widehat{\theta}_n(T_j^*))}{\partial \theta} = \frac{1}{\sqrt{n_j^*}} \left(\frac{\partial L_n(T_j^*, \widehat{\theta}_n(T_j^*))}{\partial \theta} - \frac{\partial \widehat{L}_n(T_j^*, \widehat{\theta}_n(T_j^*))}{\partial \theta} \right) \xrightarrow[n \to \infty]{a.s.} 0.$$

Thus, uses (6.24) and (6.26) to conclude.

References

- [1] Ahmad, A. and Franco, C. Poisson QMLE of count time series models. *Journal of Time Series Analysis* 37, (2016), 291-314.
- [2] Arlot, S., Celisse, A. and Celisse, Z. A kernel multiple change-point algorithm via model selection. arXiv:1202.3878v2: 2016.
- [3] ARLOT, S. AND MASSART, P. Data-driven Calibration of Penalties for Least-Squares Regression. *Journal of Machine Learning Research* 10, (2009), 245-279.
- [4] BAI, J. Estimating multiple breaks one at a time. Econometric theory 13 (3), (1997), 315-352.
- [5] BAI, J. AND PERRON, P. Estimating and testing linear models with multiple structural changes. *Econometrica* 66, (1998), 47-78.
- [6] BARDET, J. M., KENGNE, K. AND WINTENBERGER, O. Multiple breaks detection in general causal time series using penalized quasi-likelihood. *Electronic Journal of Statistics* 6, (2012), 435-477.
- [7] BAUDRY, J. P., AND MAUGIS, C. AND MICHEL, B. Slope Heuristics: overview and implementation. RR-INRIA no 7223, 2010.
- [8] CLEYNEN, A. AND LEBARBIER, E. Segmentation of the Poisson and negative binomial rate models: a penalized estimator. *ESAIM: Probability and Statistics* 18, (2014), 750-769.
- [9] CLEYNEN, A. AND LEBARBIER, E. Model selection for the segmentation of multiparameter exponential family distributions. *Electronic Journal of Statistics* 11, (2017), 800-842.
- [10] DAVIS, R. A., HANCOCK, S. A. AND YAO, Y.-C. On consistency of minimum description length model selection for piecewise autoregressions. *Journal of Econometrics* 194, (2016) 360-368.
- [11] DAVIS, R. A., LEE, T. C. M. AND RODRIGUEZ-YAM, G. A. Break detection for a class of nonlinear time series models. *Journal of Time Series Analysis* 29 (5), (2008), 834-867.
- [12] DAVIS, R. A. AND YAU, C. Y. Consistency of minimum description length model selection for piecewise stationary time series models. *Electronic Journal of Statistics* 7, (2013) 381-411.
- [13] DIOP, M.L. AND KENGNE, W. Testing parameter change in general integer-valued time series. J. Time Ser. Anal. 38, (2017) 880-894.
- [14] DOUC, R., FOKIANOS, K., AND MOULINES, E. Asymptotic properties of quasi-maximum likelihood estimators in observation-driven time series models. *Electronic Journal of Statistics* 11, (2017), 2707-2740.
- [15] DOUKHAN, P. AND KENGNE, W. Inference and testing for structural change in general poisson autoregressive models. *Electronic Journal of Statistics* 9, (2015), 1267-1314.
- [16] FOKIANOS, K., RAHBEK, A. AND TJØSTHEIM, D. Poisson autoregression. Journal of the American Statistical Association 104, (2009), 1430-1439.

- [17] FOKIANOS, K. AND TJØSTHEIM, D. Log-linear Poisson autoregression. *Journal of multivariate analysis* 102, (2011) 563-578.
- [18] FOKIANOS, K., AND TJØSTHEIM, D. Nonlinear Poisson autoregression. Ann. Inst. Stat. Math. 64, (2012) 1205-1225.
- [19] FRYZLEWICZ, P. Wild binary segmentation for multiple change-point detection. *The Annals of Statistics* 42 (6), (2014) 2243-228.
- [20] FRYZLEWICZ, P., AND SUBBA RAO, S. Multiple-change-point detection for auto-regressive conditional heteroscedastic processes. Journal of the Royal Statistical Society: series B 76 (5), (2014) 903-924.
- [21] HARCHAOUI, Z. AND LÉVY-LEDUC, C. Multiple Change-Point Estimation With a Total Variation Penalty. Journal of the American Statistical Association 105, (2010) 1480-1493.
- [22] Hudecová, Š. Structural changes in autoregressive models for binary time series. *Journal of Statistical Planning and Inference* 143, (2013) 1744-1752.
- [23] INCLÁN, C. AND TIAO, G. C. Use of cumulative sums of squares for retrospective detection of changes of variance. *Journal of the American Statistical Association* 89 (427), (1994), 913-923.
- [24] KOUNIAS, E.G. AND WENG, T.-S An inequality and almost sure convergence. *Annals of Mathematical Statistics* 33, (1969), 1091-1093.
- [25] LAVIELLE, M. AND LUDENA, C. The multiple change-points problem for the spectral distribution. Bernoulli 6, (2000), 845-869.
- [26] Lebarbier E. Detecting multiple change-points in the mean of Gaussian process by model selection. Signal Processing 85, (2005), 717-736.
- [27] LERASLE, M. Optimal model selection for density estimation of stationary data under various mixing conditions. *The Annals of Statistics 39 (4)*, (2003), 1852-1877.
- [28] Yau, C.Y. and Zhao, Z. Inference for multiple change points in time series via likelihood ratio scan statistics. *Journal of the Royal Statistical Society: Series B* 78, (2016), 895-916.