On certain linearized polynomials with high degree and kernel of small dimension

Olga Polverino, Giovanni Zini and Ferdinando Zullo*

April 23, 2020

Abstract

Let f be the \mathbb{F}_q -linear map over $\mathbb{F}_{q^{2n}}$ defined by $x \mapsto x + ax^{q^s} + bx^{q^{n+s}}$ with gcd(n,s) = 1. It is known that the kernel of f has dimension at most 2, as proved by Csajbók et al. in "A new family of MRD-codes" (2018). For n big enough, e.g. $n \ge 5$ when s = 1, we classify the values of b/a such that the kernel of f has dimension at most 1. To this aim, we translate the problem into the study of some algebraic curves of small degree with respect to the degree of f; this allows to use intersection theory and function field theory together with the Hasse-Weil bound. Our result implies a non-scatteredness result for certain high degree scattered binomials, and the asymptotic classification of a family of rank metric codes.

AMS subject classification: 11T06, 11G20, 51E20, 51E22

Keywords: linearized polynomial, algebraic curve, linear set, MRD code, Hasse-Weil bound

^{*}This research was supported by the project "VALERE: VAnviteLli pEr la RicErca" of the University of Campania "Luigi Vanvitelli", and by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

1 Introduction

Let q be a prime power and let m be a positive integer. A q-polynomial, or linearized polynomial, over \mathbb{F}_{q^m} is a polynomial of the form

$$f(x) = \sum_{i=0}^{t} a_i x^{q^i},$$

where $a_i \in \mathbb{F}_{q^m}$, t is a positive integer. If $a_t \neq 0$, we say that $t = \deg_q f(x)$ is the q-degree of f. We denote by $\mathcal{L}_{m,q}$ the set of all q-polynomials over \mathbb{F}_{q^m} and by $\tilde{\mathcal{L}}_{m,q}$ the following quotient $\mathcal{L}_{m,q}/(x^{q^m} - x)$. The \mathbb{F}_q -linear maps of \mathbb{F}_{q^m} can be identified with the polynomials in $\tilde{\mathcal{L}}_{m,q}$. This shows the relevance of linearized polynomials in the theory of finite fields and their algebraic and geometric applications. A fundamental problem in the theory of linearized polynomials is to characterize precisely the dimension of the kernel of the given polynomial in terms of its coefficients. Results in this direction are given in [6, 11, 18, 25, 28, 33, 34].

Let n, s be positive integers such that $s < 2n \operatorname{gcd}(s, n) = 1$. First in [8], and later in [28], the following polynomials are investigated

$$f_{a,b,s}(x) = x + ax^{q^s} + bx^{q^{s+n}} \in \tilde{\mathcal{L}}_{2n,q}.$$
(1)

The following results are known from [8] and [28]:

- if $N_{q^{2n}/q^n}(a) = N_{q^{2n}/q^n}(b)$, then $\dim_{\mathbb{F}_q} \ker f_{a,b,s}(x) \le 1$;
- if $N_{q^{2n}/q^n}(a) \neq N_{q^{2n}/q^n}(b)$, then $\dim_{\mathbb{F}_q} \ker f_{a,b,s}(x) \leq 2$;

where $N_{q^{2n}/q^n}(x) = x^{1+q^n}$.

Our main result is Theorem 1.1 and concerns the existence, for every $\delta \in \mathbb{F}_{q^{2n}}$ with $N_{q^{2n}/q^n}(\delta) \notin \{0, 1\}$, of an element $a \in \mathbb{F}_{q^{2n}}$ such that the kernel of $f_{a,\delta a,s}$ has dimension 2, providing n is large enough.

Theorem 1.1. Let q be a prime power and n, s be two relatively prime positive integers. Suppose that

$$n \ge \begin{cases} 4s+2 & \text{if } q=3 \text{ and } s>1, \text{ or } q=2 \text{ and } s>2;\\ 4s+1 & \text{otherwise.} \end{cases}$$

For every $\delta \in \mathbb{F}_{q^{2n}}^*$ with $N_{q^{2n}/q^n}(\delta) \neq 1$ there exists $a \in \mathbb{F}_{q^{2n}}^*$ such that $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2,$

where $b = \delta a$.

In Remark 3.2 we show that we can always suppose n > 2s, up to considering the adjoint polynomial.

The first step in the proof of Theorem 1.1 is to manipulate the shape of $f_{a,b,s}(x)$ to translate the condition on the dimension of the kernel into the existence of \mathbb{F}_{q^n} -rational points in the intersection of certain \mathbb{F}_{q^n} -rational hypersurfaces, which are described in Theorem 3.6. Then we prove that this intersection is described by means of an $\mathbb{F}_{q^{2n}}$ -rational curve \mathcal{X} . Using intersection theory and function field theory, the curve \mathcal{X} is shown to be absolutely irreducible of genus $q^{2s} - q^s - 1$; Theorem 1.1 now follows by Hasse-Weil bound.

Theorem 1.1 also has applications in the theory of scattered polynomials. A polynomial $f(x) \in \tilde{\mathcal{L}}_{m,q}$ is said to be *scattered* if

$$\dim_{\mathbb{F}_q} \ker(f(x) - \lambda x) \le 1, \quad \text{for all } \lambda \in \mathbb{F}_{q^m}.$$

Scattered polynomials have been widely investigated, especially after the paper [29], where Sheekey builds a bridge between scattered polynomials and rank metric codes. The family of linearized binomials $f_{\delta,s}(x) = x^{q^s} + \delta x^{q^{n+s}} \in$ $\mathcal{L}_{2n,q}$ with $\delta \neq 0$ contains a large number of scattered polynomials when n is 3 or 4, as proved in [8] and [28]. The question arises whether there exist other values of n, possibly infinitely many, for which $f_{\delta,s}(x)$ is scattered. Many authors have considered the problem of classifying exceptional scattered polynomials $f(x) \in \mathcal{L}_{m,q}$, i.e. scattered polynomials which remain scattered over infinitely many extensions $\mathbb{F}_{q^{\ell m}}$ of \mathbb{F}_{q^m} ; partial classification results have been provided by Bartoli and Zhou [4], Bartoli and Montanucci [2], Ferraguti and Micheli [15]. Their results rely on the fact that the order of $\mathbb{F}_{q^{\ell m}}$ is much larger than the degree of f(x); as a matter of fact, the key role in [4, 2] is played by the application of the Hasse-Weil bound to a curve whose degree has the same order of magnitude as deg f(x), and hence is small with respect to $q^{\ell m}$ (see [4, Lemma 2.1]). The aforementioned binomial $f_{\delta,s}(x)$ is not taken into account by their results, because deg $f_{\delta,s}(x) = q^{n+s}$ is high with respect to the order of $\mathbb{F}_{q^{2n}}$. As a byproduct of Theorem 1.1, we prove in Theorem 4.1 that $f_{\delta,s}(x)$ is not scattered when n is large enough with respect to s; for instance, when s = 1 it is enough to choose $n \ge 5$.

Finally, in Theorem 4.5 we use Theorem 1.1 to give an asymptotic classification of the family of rank-metric codes defined by the binomials $f_{\delta,s}(x)$.

The paper is organized as follows. Section 2 contains preliminary results about algebraic curves and function function fields which are used in Section 3. Section 3 is devoted to the proof of Theorem 1.1; the cases q odd and q even are studied separately, respectively in Section 3.1 and in Section 3.2. Section 4 provides the applications of Theorem 1.1; namely, Section 4.1 shows the applications to scattered polynomials and linear sets, while Section 4.2 shows the applications to rank metric codes.

2 Preliminaries on algebraic curves

Let \mathcal{C} be a projective, absolutely irreducible, algebraic curve over the algebraically closed field $\mathbb{K} = \overline{\mathbb{F}}_q$, embedded in a projective space $\mathrm{PG}(r, \mathbb{K})$ with homogeneous coordinates $(X_1: \ldots: X_{r+1})$ and not contained in the hyperplane at infinity $H_{\infty}: X_{r+1} = 0$. Let $I(\mathcal{C})$ be the ideal of \mathcal{C} . Denote by $\mathbb{K}(\mathcal{C})$ the field of (\mathbb{K} -)rational functions on \mathcal{C} , briefly the function field of \mathcal{C} . Clearly, $\mathbb{K}(\mathcal{C})$ is generated over \mathbb{K} by the coordinate functions x_1, \ldots, x_r with $x_i = \frac{X_i + I(\mathcal{C})}{X_{r+1} + I(\mathcal{C})}$, and $\mathbb{K}(\mathcal{C})$: \mathbb{K} is a field extension of transcendence degree 1. We denote by $\mathbb{P}(\mathcal{C})$ the set of places of \mathcal{C} , that is, the set of places of its function field $\mathbb{K}(\mathcal{C})$. For every $P \in \mathbb{P}(\mathcal{C})$ and every nonzero $z \in \mathbb{K}(\mathcal{C})$, we denote by $v_P(z) \in \mathbb{Z}$ the valuation of z at P; P is said to be a zero (resp. a pole) of z if $v_P(z) > 0$ (resp. $v_P(z) < 0$).

Suppose that \mathcal{C} is defined over \mathbb{F}_q , i.e. $I(\mathcal{C})$ is generated by polynomials over \mathbb{F}_q . Then $\mathbb{F}_q(\mathcal{C})$ denotes the \mathbb{F}_q -rational function field of \mathcal{C} , i.e. the field of \mathbb{F}_q -rational functions on \mathcal{C} . The \mathbb{F}_q -rational places of \mathcal{C} are those places $P \in \mathbb{P}(\mathcal{C})$ which are defined over \mathbb{F}_q ; that is, \mathbb{F}_q -rational places of \mathcal{C} are the places of degree 1 in $\mathbb{F}_q(\mathcal{C})$, which are exactly the restriction to $\mathbb{F}_q(\mathcal{C})$ of the places of $\mathbb{K}(\mathcal{C})$ in the constant field extension $\mathbb{K}(\mathcal{C}) \colon \mathbb{F}_q(\mathcal{C})$. The center of an \mathbb{F}_q -rational place is an \mathbb{F}_q -rational point of \mathcal{C} ; conversely, if P is a simple \mathbb{F}_q -rational point of \mathcal{C} , then the only place centered at P is \mathbb{F}_q -rational, and may be identified with P.

Let $\varphi : \mathcal{C}' \to \mathcal{C}$ be a covering of curves, i.e. a non-constant rational map from the curve \mathcal{C}' to the curve \mathcal{C} , of degree $\deg(\varphi) = [\mathbb{K}(\mathcal{C}'): \mathbb{K}(\mathcal{C})]$. We denote by φ also the induced map $\mathbb{P}(\mathcal{C}') \to \mathbb{P}(\mathcal{C})$; if φ is \mathbb{F}_q -rational, then φ maps \mathbb{F}_q -rational places of \mathcal{C}' to \mathbb{F}_q -rational places of \mathcal{C} . The pull-back of φ is denoted by $\varphi^* : \mathbb{K}(\mathcal{C}) \to \mathbb{K}(\mathcal{C}')$. When $P \in \mathbb{P}(\mathcal{C})$ and $P' \in \mathbb{P}(\mathcal{C}')$ satisfy $\varphi(P') = P$, we write P'|P and say that P' lies over P in φ . We denote by e(P'|P) the ramification index of P'|P, that is the unique positive integer such that $v_{P'}(\varphi^*(w)) = e(P'|P) \cdot v_P(w)$ for all $w \in \mathbb{K}(\mathcal{C})$; we have $\sum_{P':P'|P} e(P'|P) = \deg(\varphi)$. We say that P' is ramified over P if e(P'|P) > 1, and totally ramified if $e(P'|P) = \deg(\varphi)$; otherwise it is unramified. A ramified place P' is wildly ramified (resp. tamely ramified) if e(P'|P) is divisible (resp. not divisible) by p. We refer to [19, 32] for further details on algebraic curves and function fields.

Theorem 2.1. (Hurwitz genus formula, [32, Theorem 3.4.13]) Let $\mathcal{C}, \mathcal{C}'$ be two absolutely irreducible curves over $\mathbb{K} = \overline{\mathbb{F}}_q$ and $\varphi : \mathcal{C}' \to \mathcal{C}$ be a covering. For every place P of \mathcal{C} and every place P' of \mathcal{C}' lying over P in φ , let $t \in \mathbb{K}(\mathcal{C})$ be a local parameter at $P, t' \in \mathbb{K}(\mathcal{C}')$ be a local parameter at P', and $\varphi^*(t) \in \mathbb{K}(\mathcal{C}')$ be the pull-back of t with respect to φ . Then

$$2g(\mathcal{C}') - 2 = \deg(\varphi) \cdot (2g(\mathcal{C}) - 2) + \sum_{P' \in \mathbb{P}(\mathcal{C}')} v_{P'}\left(\frac{d\varphi^*(t)}{dt'}\right).$$

If P' is not wildly ramified, then $v_{P'}\left(\frac{d\varphi^*(t)}{dt'}\right) = e(P'|P) - 1$. We now recall two important types of coverings. The following results are the application of [32, Corollary 3.7.4] and [32, Theorem 3.7.10] in the case of an algebraically closed constant field \mathbb{K} .

Theorem 2.2. [32, Corollary 3.7.4] Let C: F(X, Y) = 0 be an absolutely irreducible plane curve defined over a finite field \mathbb{F}_q of characteristic p, and mbe a positive integer with gcd(m, p) = 1. Let $f(X, Y) \in \mathbb{F}_q[X, Y]$ be such that there exists an $\overline{\mathbb{F}}_q$ -rational place Q of C at which the valuation of the rational function f(x, y) is coprime with m, i.e. $gcd(v_Q(f(x, y)), m) = 1$. Let C' be the curve given by the two affine equations F(X, Y) = 0 and $Z^m = f(X, Y)$. Then the following holds.

- \mathcal{C}' is absolutely irreducible and defined over \mathbb{F}_q ; \mathcal{C}' is called a Kummer cover of \mathcal{C} .
- The \mathbb{F}_q -rational covering $\varphi : \mathcal{C}' \to \mathcal{C}, (X, Y, Z) \mapsto (X, Y)$, has degree m.
- For every place P of C and every place P' of C' lying over P in φ , we have $e(P'|P) = m/r_P$, where $r_P = \gcd(v_P(f(x, y)), m) > 0$.
- The Hurwitz genus formula reads

$$g(\mathcal{C}') = 1 + m(g(\mathcal{C}) - 1) + \frac{1}{2} \sum_{P \in \mathbb{P}(\mathcal{C})} (m - r_P).$$

If \mathcal{C}' is an absolutely irreducible curve over \mathbb{F}_q defined by the two affine equations F(X,Y) = 0 and L(Z) = f(X,Y), for some $f(X,Y), F(X,Y) \in$ $\mathbb{F}_q[X,Y]$ and some separable *p*-polynomial $L(T) \in \mathbb{F}_q[T]$, then \mathcal{C}' is said to be a generalized Artin-Schreier cover of the curve $\mathcal{C} : F(X,Y) = 0$, with generalized Artin-Schreier covering $\varphi : \mathcal{C}' \to \mathcal{C}, (X,Y,Z) \mapsto (X,Y)$.

Theorem 2.3. [32, Theorem 3.7.10] Let C : F(X,Y) = 0 be an absolutely irreducible plane curve defined over a finite field \mathbb{F}_q of charateristic p. Let $L(T) \in \mathbb{F}_q[T]$ be a separable p-polynomial of degree \bar{q} with all its roots in \mathbb{F}_q . Let $f(X,Y) \in \mathbb{F}_q[X,Y]$ be such that for every place $P \in \mathbb{P}(C)$ there exists a rational function ω on C (depending on P) satisfying either $v_P(f(x,y) - L(\omega)) \ge 0$ or $v_P(f(x,y) - L(\omega)) = -m$ with m > 0 and $p \nmid m$. Define $m_P =$ -1 in the former case and $m_P = m$ in the latter case. Let C' be the space curve given by the two affine equations F(X,Y) = 0 and L(Z) = f(X,Y). If there exists a place $Q \in \mathbb{P}(C)$ with $m_Q > 0$, then C' is a generalized Artin-Schreir cover of C, defined over \mathbb{F}_q .

With the above notation, the following holds for generalized Artin-Schreier curves.

- The \mathbb{F}_q -rational covering $\varphi : \mathcal{C}' \to \mathcal{C}, (X, Y, Z) \mapsto (X, Y)$, has degree \bar{q} .
- For every place P of C and every place P' of C' lying over P in φ , e(P'|P) is equal either to 1 or to \bar{q} according to $m_P = -1$ or $m_P > 0$, respectively.
- The Hurwitz genus formula reads

$$g(\mathcal{C}') = \bar{q} \cdot g(\mathcal{C}) + \frac{\bar{q} - 1}{2} \cdot \left(-2 + \sum_{P \in \mathbb{P}(\mathcal{C})} (m_P + 1) \right).$$

We now recall the well-known Hasse-Weil bound.

Theorem 2.4. [32, Theorem 5.2.3] (Hasse-Weil bound) Let C be an absolutely irreducible curve defined over \mathbb{F}_q and with genus g. Then the number N_q of \mathbb{F}_q -rational places of C satisfies

$$q+1-2g\sqrt{q} \le N_q \le q+1+2g\sqrt{q}.$$

3 Proof of Theorem 1.1

In this section we prove Theorem 1.1. First we determine necessary and sufficient conditions on a and b for $f_{a,b,s}(x)$ having kernel of dimension 2; cf. Theorem 3.6. Then we investigate such conditions by means of algebraic-geometric tools.

The first remark shows that different choices of a, b with the same norm of b/a over \mathbb{F}_{q^n} provide polynomials $f_{a,b,s}(x)$ with the same behaviour.

Remark 3.1. Assume that the linearized polynomial $f_{a,b,s}(x) = x + ax^{q^s} + bx^{q^{s+n}} \in \mathbb{F}_{q^{2n}}[x]$, with gcd(s, n) = 1 and $b = \delta a$, has kernel of dimension two. Clearly, for each $\lambda \in \mathbb{F}_{q^{2n}}^*$ we have

$$\dim_{\mathbb{F}_q} \ker(\lambda^{-1} f_{a,b,s}(\lambda x)) = 2,$$

where

$$\lambda^{-1} f_{a,b,s}(\lambda x) = x + a\lambda^{q^s - 1} x^{q^s} + a\lambda^{q^s - 1} \delta\lambda^{q^s(q^n - 1)} x^{q^{s+n}} = f_{a',b',s}(x),$$

with $a' = a\lambda^{q^s-1}$, $\delta' = \lambda^{q^s(q^n-1)}\delta$ and $b' = a'\delta'$. Note that for each element $\delta' \in \mathbb{F}_{q^{2n}}$ with $N_{q^{2n}/q^n}(\delta') = N_{q^{2n}/q^n}(\delta)$ there exists $\lambda \in \mathbb{F}_{q^{2n}}$ such that $\delta' = \delta\lambda^{q^s(q^n-1)}$. Therefore, if $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2$, with $b = \delta a$, then for each $\delta' \in \mathbb{F}_{q^{2n}}$ with $N_{q^{2n}/q^n}(\delta') = N_{q^{2n}/q^n}(\delta)$ there exists $a' \in \mathbb{F}_{q^{2n}}$ such that $\dim_{\mathbb{F}_q} \ker(f_{a',b',s}(x)) = 2$, with $b' = \delta'a'$.

The second remark shows that we may assume s < n/2.

Remark 3.2. The adjoint of a q-polynomial $f(x) = \sum_{i=0}^{n-1} a_i x^{q^i}$, with respect to the bilinear form $\langle x, y \rangle = \operatorname{Tr}_{q^n/q}(xy)$, is given by

$$\hat{f}(x) = \sum_{i=0}^{n-1} a_i^{q^{n-i}} x^{q^{n-i}}.$$

In particular, if f(x) is a q-polynomial of shape (1), then

$$f_{a,b,s}(x) = x + ax^{q^s} + bx^{q^{n+s}} \in \tilde{\mathcal{L}}_{2n,q},$$

with gcd(s, n) = 1 and its adjoint is

$$\hat{f}_{a,b,s}(x) = x + a^{q^{2n-s}} x^{q^{2n-s}} + b^{q^{n-s}} x^{q^{n-s}}.$$

Therefore, choosing s' = 2n - s, $a' = a^{q^{2n-s}}$, $b' = b^{q^{n-s}}$, we get

$$\hat{f}_{a,b,s}(x) = f_{a',b',s'}(x),$$

while choosing s'' = n - s, $a'' = b^{q^{n-s}}$, $b'' = a^{q^{2n-s}}$, we get

$$\hat{f}_{a,b,s}(x) = f_{a'',b'',s''}(x),$$

i.e. $f_{a,b,s}(x)$ is of shape (1). Therefore, the family of q-polynomials we are studying is closed by the adjoint operation. Furthermore, we underline that by [1, Lemma 2.6], the kernels of $f_{a,b,s}$ and $\hat{f}_{a,b,s}$ have the same dimension (see also [7, pages 407-408]). Thus, we can assume s < n/2.

We now prove that the shape of δ can be chosen as in (2).

Theorem 3.3. Let $f_{a,b,s}(x) \in \mathbb{F}_{q^{2n}}[x]$, with $b = a\delta$. Then $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2$ if and only if $\dim_{\mathbb{F}_q} \ker(f_{\overline{a},\overline{b},s}(x)) = 2$, with

$$\overline{\delta} = \frac{\xi^{q^{s+n}} - \xi^{q^n}}{\xi^{q^n} - \xi^{q^s}},\tag{2}$$

for some $\xi \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^n}$ and some $\overline{a} \in \mathbb{F}_{q^{2n}}$, $\overline{b} = \overline{\delta}\overline{a}$.

Proof. Assume that $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2$, i.e. there exist $x_0 \in \mathbb{F}_{q^{2n}}^*$ and $y_0 \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ such that $x_0/y_0 \notin \mathbb{F}_q$ and

$$\frac{x_0^{q^s} + \delta x_0^{q^{s+n}}}{x_0} = \frac{y_0^{q^s} + \delta y_0^{q^{s+n}}}{y_0},$$

which may be rewritten as follows

$$\delta(y_0 x_0^{q^{s+n}} - x_0 y_0^{q^{s+n}}) = x_0 y_0^{q^s} - y_0 x_0^{q^s}.$$

If $y_0 x_0^{q^{s+n}} - x_0 y_0^{q^{s+n}}$ would be zero, than $x_0/y_0 \in \mathbb{F}_{q^{2n}} \cap \mathbb{F}_{q^{s+n}} = \mathbb{F}_q$, a contradiction. Hence,

$$\delta = \frac{x_0 y_0^{q^s} - y_0 x_0^{q^s}}{y_0 x_0^{q^{s+n}} - x_0 y_0^{q^{s+n}}},$$

and, since $y_o = \xi x_0$ for some $\xi \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$, we have

$$\delta = \frac{1}{-x_0^{q^{s+n}-q^s}} \frac{\xi^{q^s} - \xi}{\xi^{q^{s+n}} - \xi}$$

By Remark 3.1, $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2$ if and only if there exists $\overline{a}, \overline{b}$ as in the claim such that $\dim_{\mathbb{F}_q} \ker(f_{\overline{a},\overline{b},s}(x)) = 2$. If $\xi \in \mathbb{F}_{q^n}$, then $\overline{\delta} = -1$, and hence $\dim_{\mathbb{F}_q} \ker(f_{\overline{a},\overline{b},s}(x)) \leq 1$. The claim follows.

As a consequence of Theorem 3.3 we get the following result.

Corollary 3.4. There exist $\delta \in \mathbb{F}_{q^{2n}}^*$ for which $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) \leq 1$, with $b = \delta a$, for each $a \in \mathbb{F}_{q^{2n}}^*$ if and only if

$$\left| \left\{ N_{q^{2n}/q^n} \left(\frac{\xi^{q^{n+s}} - \xi^{q^n}}{\xi^{q^n} - \xi^{q^s}} \right) : \xi \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^n} \right\} \right| < q^n - 1.$$

Since $\xi \notin \mathbb{F}_{q^n}$, we have that ξ is the root of an irreducible polynomial $X^2 - SX - T \in \mathbb{F}_{q^n}[X]$, where $N_{q^{2n}/q^n}(\xi) = -T$ and $\operatorname{Tr}_{q^{2n}/q^n}(\xi) = S$. Also, $\{1, \xi\}$ is an \mathbb{F}_{q^n} -basis of $\mathbb{F}_{q^{2n}}$ and so there exist $A, B \in \mathbb{F}_{q^n}$ such that $\xi^{q^s} = A + B\xi$. In the next we give some relations involving A, B, S and T.

Proposition 3.5. The following holds:

- 1. $S^{q^s} = 2A + BS;$
- 2. $-T^{q^s} = A^2 + B(AS BT).$

In particular, $\operatorname{Tr}_{q^{2n}/q^n}(\xi^{q^s+1}) = 2BT + AS + BS^2$ and $\operatorname{Tr}_{q^{2n}/q^n}(\xi^{q^s+q^n}) = AS - 2BT$.

Proof. As

$$\xi^{q^s+q^n} = (A+B\xi)(S-\xi) = AS - BT - A\xi$$

and

$$\xi^{1+q^{n+s}} = -BT + (S^{q^s} - A - BS)\xi,$$

we have that

$$\operatorname{Tr}_{q^{2n}/q^n}(\xi^{q^s+q^n}) = AS - 2BT + (S^{q^s} - 2A - BS)\xi$$

Since $\operatorname{Tr}_{q^{2n}/q^n}(\xi^{q^s+q^n}) \in \mathbb{F}_{q^n}$, we get the first relation. Also,

$$-T^{q^s} = N_{q^{2n}/q^n}(\xi^{q^s}) = A^2 + ABS - B^2T,$$

i.e. the second relation.

Let $\alpha \in \mathbb{F}_{q^n}^*$ with $\alpha \neq 1$. Then

$$N_{q^{2n}/q^n}\left(\frac{\xi^{q^{n+s}}-\xi^{q^n}}{\xi^{q^n}-\xi^{q^s}}\right) = \frac{\xi^{q^n+1}+\xi^{q^s+q^{n+s}}-(\xi^{1+q^{n+s}}+\xi^{q^s+q^n})}{\xi^{q^n+1}+\xi^{q^s+q^{n+s}}-(\xi^{q^n+q^{n+s}}+\xi^{q^s+1})} = \alpha,$$

which can be written as

$$(1 - \alpha)(T + T^{q^s}) - \alpha S^{q^s + 1} + (1 + \alpha)(AS - 2BT) = 0.$$

Hence, we have the following result.

Theorem 3.6. Let $\alpha \in \mathbb{F}_{q^n}^*$ with $\alpha \neq 1$ and s a positive integer with gcd(s,n) = 1. If there exist $T, S, A, B \in \mathbb{F}_{q^n}$ such that

- 1. $(1-\alpha)(T+T^{q^s}) \alpha S^{q^s+1} + (1+\alpha)(AS 2BT) = 0;$
- 2. $X^2 SX T \in \mathbb{F}_{q^n}[X]$ is irreducible over \mathbb{F}_{q^n} ;
- 3. $S^{q^s} = 2A + BS;$
- 4. $-T^{q^s} = A^2 + B(AS BT),$

then for every $\delta \in \mathbb{F}_{q^{2n}}$, with $N_{q^{2n}/q^n}(\delta) = \alpha$, there exists $a \in \mathbb{F}_{q^{2n}}^*$ such that $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}(x)) = 2$, where $b = \delta a$.

In the rest of this section $q = p^h$ with p prime. We will show that the existence of the parameters $T, S, A, B \in \mathbb{F}_{q^n}$ satisfying the hypothesis of Theorem 3.6 is equivalent to the existence of a suitable affine \mathbb{F}_{q^n} -rational point of the algebraic plane curve with equation (4) or (5), for q odd or q even respectively.

3.1 Proof of Theorem 1.1 for *q* odd

Denote by $\Delta = S^2 + 4T$. By 3. and 4. of Theorem 3.6, we get

$$B = \epsilon \Delta^{\frac{q^s - 1}{2}}, \quad A = \frac{1}{2} (S^{q^s} - \epsilon S \Delta^{\frac{q^s - 1}{2}})$$

where $\epsilon \in \{1, -1\}$. Hence we get $AS - 2BT = \frac{1}{2}\epsilon \Delta^{\frac{q^s-1}{2}} - \frac{1}{2}S^{q^s+1}$. Replacing such values in 1. of Theorem 3.6, we get

$$2(T+T^{q^s})(1-\alpha) + (1-\alpha)S^{q^s+1} = \epsilon(\alpha+1)\Delta^{\frac{q^s+1}{2}}.$$
(3)

Also, the irreducibility of $X^2 - SX - T$ over \mathbb{F}_{q^n} is equivalent to the existence of a nonsquare element η of \mathbb{F}_{q^n} and a nonzero element Z of \mathbb{F}_{q^n} such that $\Delta = \eta Z^2$. Therefore, (3) becomes

$$2(T + T^{q^s}) + S^{q^s+1} = \beta \eta^{\frac{q^s+1}{2}} Z^{q^s+1},$$

where $\beta = \epsilon \frac{\alpha + 1}{1 - \alpha}$. Using that $T = \frac{\eta Z^2 - S^2}{4}$, we get the following equation:

$$-(S^{q^s}-S)^2 + \eta Z^2 + \eta^{q^s} Z^{2q^s} - 2\beta \eta^{\frac{q^s+1}{2}} Z^{q^s+1} = 0.$$
(4)

Theorem 3.7. Let $\beta \in \mathbb{F}_{q^n} \setminus \{1, -1\}$ and η a non-square in \mathbb{F}_{q^n} . The plane curve \mathcal{C} with affine equation (4) is absolutely irreducible and has genus $g(\mathcal{C}) = q^{2s} - q^s - 1$.

Proof. Let $G(Z) = \eta Z^2 + \eta^{q^s} Z^{2q^s} - 2\beta \eta^{\frac{q^s+1}{2}} Z^{q^s+1} \in \mathbb{F}_{q^n}[Z]$, and let \mathcal{C}_1 be the plane curve with affine equation $F_1(U, Z) = 0$, where $F_1(U, Z) = U^2 - G(Z)$. By direct computation using the assumption $\beta \neq \pm 1$ follows that 0 is the unique multiple root of G(Z), with multiplicity 2; the other $2q^s - 2$ roots $\lambda_1, \ldots, \lambda_{2q^s-2}$ of G(Z) are simple. Then G(Z) is not a square in $\mathbb{K}[Z]$, whence $F_1(U, Z)$ is irreducible over $\mathbb{K} = \overline{\mathbb{F}}_{q^n}$, i.e. \mathcal{C}_1 is absolutely irreducible.

The genus of the quadratic Kummer cover C_1 of the projective line is computed as follows. Let z, u be the coordinate functions of C_1 , so that the function field of C_1 is $\mathbb{K}(C_1) = \mathbb{K}(z, u)$. The valuation of G(z) at the zero of $z - \lambda_i$ in $\mathbb{K}(\mathbb{P}^1_z) = \mathbb{K}(z)$ is 1, for every $i = 1, \ldots, 2q^s - 2$. The valuation of G(z) at any other place of \mathbb{P}^1_z is even; namely, it is 2 at the zero of $z, -2q^s$ at the pole of z, and 0 at the zero of $z - \mu$ whenever $G(\mu) \neq 0$. By Theorem 2.2, the only ramified places in $C_1 \to \mathbb{P}^1_z$ are the zeros of $z - \lambda_1, \ldots, z - \lambda_{2q^s-2}$; hence,

$$g(\mathcal{C}_1) = 1 + 2(g(\mathbb{P}_z^1) - 1) + \frac{1}{2}(2q^s - 2)(2 - 1) = q^s - 2.$$

Since \mathcal{C} has equation $(S^{q^s} - S)^2 = G(Z)$, it is enough to show that \mathcal{C} is an Artin-Schreier cover of \mathcal{C}_1 , with covering $\varphi : \mathcal{C} \to \mathcal{C}_1$, $(Z, S) \mapsto (Z, U = S^{q^s} - S)$, of degree q^s . To this aim, consider the two poles P_{∞} and Q_{∞} of u on \mathcal{C}_1 ; the rational function 1/z is a local parameter at each of them, i.e. $v_{P_{\infty}}(1/z) = v_{Q_{\infty}}(1/z) = 1$. By direct computation, the Laurent series of uat P_{∞} with respect to 1/z is

$$u = \sqrt{\eta^{q^s}} (1/z)^{-q^s} - \beta \sqrt{\eta} (1/z)^{-1} + \frac{\eta - \beta^2 \eta}{2\sqrt{\eta^{q^s}}} (1/z)^{q^s - 2} + w,$$

for some $w \in \mathbb{K}(\mathcal{C}_1)$ with $v_{P_{\infty}}(w) > q^s - 2$. By choosing $\omega_{P_{\infty}} = \sqrt{\eta}z$ one has that $u - (\omega_{P_{\infty}}^{q^s} - \omega_{P_{\infty}})$ has valuation -1 at P_{∞} , because $\beta \neq 1$. Analogously, there exists $\omega_{Q_{\infty}}$ such that $v_{Q_{\infty}}(u - (\omega_{Q_{\infty}}^{q^s} - \omega_{Q_{\infty}})) = -1$. Hence, by Theorem 2.3, \mathcal{C} is an absolutely irreducible Artin-Schreier extension \mathcal{C}_1 of degree q^s .

The ramified places in $\mathcal{C} \to \mathcal{C}_1$ are exactly P_{∞} and Q_{∞} , which are totally ramified; any other place of \mathcal{C}_1 is unramified under \mathcal{C} . Therefore,

$$g(\mathcal{C}) = q^s \cdot g(\mathcal{C}_1) + \frac{q^s - 1}{2} (-2 + 2 \cdot 2) = q^{2s} - q^s - 1.$$

		-	

Proposition 3.8. Let C be the plane curve with affine equation (4). If

$$n \ge \begin{cases} 4s+1 & \text{if } q > 3, \\ 4s+2 & \text{if } q = 3, s > 1, \\ 5 & \text{if } q = 3, s = 1; \end{cases}$$

then there exists an \mathbb{F}_{q^n} -rational affine point (\bar{z}, \bar{s}) of \mathcal{C} such that $\bar{t} = \frac{\eta \bar{z}^2 - \bar{s}^2}{4}$ is different from zero.

Proof. By Theorem 3.7, C is absolutely irreducible with genus $g(C) = q^{2s} - q^s - 1$. By Theorem 2.4, the number N_{q^n} of \mathbb{F}_{q^n} -rational places of C satisfies

$$N_{q^n} \ge q^n + 1 - 2(q^{2s} - q^s - 1)\sqrt{q^n}.$$

From the proof of Theorem 3.7 the following facts follow.

- z has exactly 2 poles on \mathcal{C} , which coincide with the poles of s, namely the places lying over P_{∞} and Q_{∞} .
- Using the equation of C, the zeros of $t = \frac{\eta z^2 s^2}{4} = \frac{(\sqrt{\eta}z s)(\sqrt{\eta}z + s)}{4}$ on C are also zeros of $(\beta 1)z^{q^s+1}$ and hence of z as $\beta \neq 1$; thus, they are the common zeros of z and s on C, and there are exactly 2 of them.

Altogether, there are 4 places of C which are either poles of s or z or t, or zeros of t. The assumption on n implies that

$$q^{n} + 1 - 2(q^{2s} - q^{s} - 1)\sqrt{q^{n}} > 4,$$

whence $N_{q^n} > 4$. Then there exists an \mathbb{F}_{q^n} -rational place P which is not a pole of z, s, or t, and is not a zero of t. Then the point $(\bar{z}, \bar{s}) = (z(P), s(P))$ yields the claim.

From Theorem 3.6 and Proposition 3.8 follows Corollary 3.9, which is our main result Theorem 1.1 when q is odd.

Corollary 3.9. Let q be an odd prime power, $s \ge 1$ be such that gcd(s, n) = 1. Suppose that

$$n \ge \begin{cases} 4s+2 & \text{if } q = 3 \text{ and } s > 1;\\ 4s+1 & \text{otherwise.} \end{cases}$$

Then for every $\delta \in \mathbb{F}_{q^{2n}}$ satisfying $N_{q^{2n}/q^n}(\delta) \notin \{0,1\}$ there exists $a \in \mathbb{F}_{q^{2n}}^*$ such that $\dim_{\mathbb{F}_q} \ker(f_{a,b}) = 2$, where $b = \delta a$.

3.2 Proof of Theorem 1.1 for *q* even

Let q be a power of 2. The conditions of Theorem 3.6 read:

- 1. $T + T^{q^s} + \beta S^{q^s+1} + AS = 0$, with $\beta = \frac{\alpha}{1+\alpha} \notin \{0, 1\};$
- 2. $S \neq 0$ and $\operatorname{Tr}_{q^n/2}(T/S^2) = 1;$
- 3. $B = S^{q^s 1};$
- 4. $A^2 + AS^{q^s} + S^{2q^s 2}T + T^{q^s} = 0.$

By 1. we get

$$A = \beta S^{q^s} + \frac{T + T^{q^s}}{S},$$

which can be replaced in 4. obtaining

$$(\beta^2 + \beta)S^{2(q^s+1)} + S^{q^s+1}(T^{q^s} + T) + S^{2q^s}T + S^2T^{q^s} + T^{2q^s} + T^2 = 0.$$
(5)

Set $T = S^2 Y$. Then (5) reads H(S, Y) = 0, where

$$H(S,Y) = Y^{2} + S^{4(q^{s}-1)}Y^{2q^{s}} + \beta^{2}S^{2(q^{s}-1)} + S^{q^{s}-1}Y +$$

$$S^{3(q^{s}-1)}Y^{q^{s}} + \beta S^{2(q^{s}-1)} + S^{2(q^{s}-1)}Y + S^{2(q^{s}-1)}Y^{q^{s}}.$$
(6)

Straightforward computation using $\operatorname{Tr}_{q^s/2}(Y) + \operatorname{Tr}_{q^s/2}(Y)^2 = Y^{q^s} + Y$ shows that the polynomial H(S, Y) in (6) splits as follows.

Lemma 3.10. We have $H(S, Y) = G(S, Y) \cdot G'(S, Y)$, where

$$G(S,Y) = S^{2(q^s-1)}Y^{q^s} + S^{q^s-1}(1+\beta + \operatorname{Tr}_{q^s/2}(Y)) + Y,$$
$$G'(S,Y) = S^{2(q^s-1)}Y^{q^s} + S^{q^s-1}(\beta + \operatorname{Tr}_{q^s/2}(Y)) + Y.$$

The condition 2. is equivalent to the existence of an element $Z \in \mathbb{F}_{q^n}$ such that

$$T = S^2 (Z^2 + Z + \epsilon), \tag{7}$$

for some fixed $\epsilon \in \mathbb{F}_{q^n}$ such that $\operatorname{Tr}_{q^n/2}(\epsilon) = 1$. Let \mathcal{C} be the plane curve with affine equation $F(S, Z) = G'(S, Z^2 + Z + \epsilon)$.

In order to prove Theorem 1.1 when q is even, by Theorem 3.6 and the arguments at the beginning of Section 3.2, it is enough to prove the existence

of an \mathbb{F}_{q^n} -rational affine point (\bar{s}, \bar{z}) of \mathcal{C} such that $\bar{s} \neq 0$ and $\bar{z}^2 + \bar{z} + \epsilon \neq 0$. This is done by showing that \mathcal{C} is absolutely irreducible, computing its genus, and applying the Hasse-Weil lower bound. To this aim, we consider the following subcovers:

$$\varphi_2 : \mathcal{C} \to \mathcal{C}_2, \quad (S, Z) \mapsto (X = S^{q^s - 1}, Z),$$

 $\varphi_1 : \mathcal{C}_2 \to \mathcal{C}_1, \quad (X, Z) \mapsto (X, Y = Z^2 + Z + \epsilon).$

The curves C_2, C_1 have equation $C_2: F_2(X, Z) = 0$ and $C_1: F_1(X, Y) = 0$, where

$$F_2(X,Z) = X^2 (Z^2 + Z + \epsilon)^{q^s} + X(\beta + \operatorname{Tr}_{q^s/2}(Z^2 + Z + \epsilon)) + Z^2 + Z + \epsilon,$$

$$F_1(X,Y) = X^2 Y^{q^s} + X(\beta + \operatorname{Tr}_{q^s/2}(Y)) + Y.$$

We first prove that C_2 is absolutely irreducible by direct inspection, and that C is absolutely irreducible being a Kummer cover of C_2 . To compute the genus of C, we start by the genus of the absolutely irreducible subcover C_1 of C_2 , which is computed with the Hurwitz genus formula. Then we compute the genus of C_2 as an Artin-Schreier cover of C_1 . Finally, the genus of the Kummer cover C of C_2 is computed.

Let $\gamma, \gamma + 1$ be the roots of $Z^2 + Z + \epsilon \in \mathbb{F}_{q^n}[Z]$. Hence, $\operatorname{Tr}_{q^{2n}/q^n}(\gamma) = 1$ and $\operatorname{N}_{q^{2n}/q^n}(\gamma) = \epsilon$; also, $\operatorname{Tr}_{q^n/2}(\epsilon) = 1$ implies $\gamma \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^n}$.

Lemma 3.11. The curve C_2 is absolutely irreducible.

Proof. By contradiction, suppose $F_2(X, Z) = \hat{F}(X, Z) \cdot \tilde{F}(X, Z)$ for some non-constant polynomials $\hat{F}, \tilde{F} \in \mathbb{K}[X, Z]$. Then $\hat{F}(X, Z) = X \cdot A(Z) + B(Z)$ and $\tilde{F}(X, Z) = X \cdot C(Z) + D(Z)$; up to scalar multiplication,

$$A(Z) = (Z + \gamma)^{a} (Z + \gamma + 1)^{b}, \qquad C(Z) = (Z + \gamma)^{c} (Z + \gamma + 1)^{d},$$

where $a, b, c, d \ge 0$ satisfy $a + c = b + d = q^s$. Also,

$$(Z+\gamma)^{a}(Z+\gamma+1)^{b}D(Z) + (Z+\gamma)^{c}(Z+\gamma+1)^{d}B(Z) = \beta + \operatorname{Tr}_{q^{s}/2}(Z^{2}+Z+\epsilon).$$

Clearly, $a, b \in \{0, q^s\}$ as $\beta \neq 0$. If a = 0, then $b = c = q^s$ and d = 0, since $B(Z)D(Z) = Z^2 + Z + \epsilon$; hence

$$(Z + \gamma + 1)^{q^s} D(Z) + (Z + \gamma)^{q^s} B(Z) = \beta + \operatorname{Tr}_{q^s/2}(Z^2 + Z + \epsilon).$$
(8)

This implies $D(\gamma) \neq 0$ and $B(\gamma + 1) \neq 0$, whence $D(Z) = \lambda(Z + \gamma + 1)$ and $B(Z) = \lambda^{-1}(Z + \gamma)$ for some $\lambda \in \mathbb{K}^*$. With $Z = \gamma$ in (8), we get $\beta = 1$, a contradiction. If $a = q^s$, the same arguments yield a contradiction.

Proposition 3.12. The curve C_1 is absolutely irreducible with genus $q^s/2$.

Proof. As C_1 is a subcover of C_2 , the absolute irreducibility of C_1 follows from Lemma 3.11. Let (X : Y : V) be the homogeneous coordinates of the affine point (X, Y). By direct computation, the affine points of C_1 are simple, and hence we identify each of them with the unique place of C_1 centered at it; the points at infinity $E_0 = (1: 0: 0)$ and $E_1 = (0: 1: 0)$ of C_1 are singular. The point E_0 is q^s -fold with unique tangent line $\ell_Y : Y = 0$ having intersection multiplicity $q^s + 1$ with C_1 at E_0 ; the point E_1 is double with unique tangent line $\ell_X : X = 0$ having intersection multiplicity $q^s + 1$ with C_1 at E_1 .

We compute the genus of \mathcal{C}_1 by applying Theorem 2.1 to the covering $\varphi_0: \mathcal{C}_1 \to \mathbb{P}^1_y, (X:Y:V) \mapsto (Y:V)$, where y is the coordinate function of Y; φ_0 has degree 2. We describe the places of \mathbb{P}^1_y which ramify in φ_0 ; we denote by $P_{\mu}, \mu \in \mathbb{K}$, the zero of $y - \mu$ on \mathbb{P}^1_y , and by P_{∞} the pole of y on \mathbb{P}^1_y .

- If $\bar{y} \in \mathbb{K}$ satisfies $\bar{y} \neq 0$ and $\beta + \operatorname{Tr}_{q^s/2}(\bar{y}) \neq 0$, then $F_1(X, \bar{y})$ has two distinct roots $\bar{x}_1, \bar{x}_2 \in \mathbb{K}$, and hence $P_{\bar{y}}$ does not ramify in φ_0 .
- If $\bar{y} \in \mathbb{K}$ is one of the $q^s/2$ distinct roots of $\beta + \operatorname{Tr}_{q^s/2}(Y)$, then $(\bar{x}, \bar{y}) \in \varphi_0^{-1}(P_{\bar{y}})$, with $\bar{x} = \sqrt{\bar{y}^{1-q^s}}$. The tangent line to \mathcal{C}_1 at (\bar{x}, \bar{y}) is $\ell_{\bar{y}} : Y \bar{y} = 0$, having multiplicity intersection 2 with \mathcal{C}_1 at (\bar{x}, \bar{y}) . Hence,

$$e((\bar{x},\bar{y}) \mid P_{\bar{y}}) = e((\bar{x},\bar{y}) \mid P_{\bar{y}}) \cdot v_{P_{\bar{y}}}(y-\bar{y}) = v_{(\bar{x},\bar{y})}(y-\bar{y}) = 2,$$

so that $P_{\bar{y}}$ totally ramifies in φ_0 .

• The change of coordinates $(X: Y: V) \mapsto (X: V: Y)$ maps E_1 to the origin $E_2 = (0: 0: 1) = (0, 0)$ and C_1 to the curve \overline{C}_1 with affine equation $\overline{F}_1(X, Y) = 0$ where

$$\overline{F}_1(X,Y) = X^2 + X(\beta Y^{q^s+1} + \sum_{i=0}^{sh-1} Y^{q^s+1-2^i}) + Y^{q^s+1}.$$

The point E_2 is double for $\overline{\mathcal{C}}_2$ with tangent line $\ell_X \colon X = 0$; while this holds, we apply iteratively the quadratic transformation $(X \colon Y \colon V) \mapsto$ $(XV \colon Y^2 \colon YV)$ which maps $\overline{\mathcal{C}}_1$ to the curve with equation $\overline{F}_1(XY,Y)/Y^2 =$ 0. After k times, the curve has equation

$$X^{2} + X(\beta Y^{q^{s}+1-k} + \sum_{i=0}^{sh-1} Y^{q^{s}+1-2^{i}-k}) + Y^{q^{s}+1-2k} = 0.$$

Hence, after $q^s/2$ times, the curve has only one affine point on the line $\ell_Y: Y = 0$, which is a simple point. This means that \mathcal{C}_1 has exactly one place centered at E_1 , which we identify with E_1 . Since the intersection multiplicity of \mathcal{C}_1 at E_1 with $\ell_{\infty}: V = 0$ and ℓ_Y is 2 and 0 respectively, we have that $v_{E_1}(y) = -2$; see [19, Theorem 4.36]. Thus, the pole of y in \mathbb{P}^1_y is totally ramified in φ_0 .

• The unique place centered at $E_2 = (0,0)$ is clearly a zero of y. The only place of \mathbb{P}^1_y which can be covered by E_0 is the zero of y. Therefore, the zero of y in \mathbb{P}^1_y is not ramified in φ_0 , and E_0 and E_2 are the simple zeros of y on \mathcal{C}_1 .

The $q^s/2+1$ ramification places P', namely E_1 and (\bar{x}, \bar{y}) with $\beta + \operatorname{Tr}_{q^2/2}(\bar{y}) = 0$, are wildly ramified. For each of them, we choose a local parameter t' at P', a local parameter t at the place P lying under P' in φ_0 , and compute $v_{P'}(d\varphi_0^*(t)/dt')$, where the pull-back φ_0^* of φ_0 is the identity on $\mathbb{K}(\mathbb{P}^1_y) = \mathbb{K}(y)$.

- Let $P' = E_1$, lying over $P = P_{\infty}$. We choose t = 1/y. From $v_{E_1}(y) = -2$ we get $v_{E_1}(x) = q^s 1$; hence we can choose $t' = 1/(xy^{q^s/2})$. By direct computation, the Laurent series of t at E_1 with respect to t' is $t = (t')^2 + (t')^3 + w$, with $v_{E_1}(w) \ge 4$. Thus, $\frac{dt}{dt'} = (t')^2 + \frac{dw}{dt'}$ has valuation 2 at E_1 .
- Let $P' = (\bar{x}, \bar{y})$, lying over $P = P_{\bar{y}}$ with $\beta + \operatorname{Tr}_{q^s/2}(\bar{y}) = 0$. We choose $t = y \bar{y}$ and $t' = x \bar{x}$. By direct computation, the Laurent series of t at P' with respect to t' is $t = \frac{\bar{y}^{q^s}}{\bar{x}+1}(t')^2 + \frac{\bar{y}^{q^s}}{\bar{x}^2+1}(t')^3 + w$, with $v_{P'}(w) \ge 4$. Thus, $\frac{dt}{dt'} = \frac{\bar{y}^{q^s}}{\bar{x}^2+1}(t')^2 + \frac{dw}{dt'}$ has valuation 2 at P'.

Theorem 2.1 now yields

$$2g(\mathcal{C}_1) - 2 = \deg(\varphi_0) \cdot (2g(\mathbb{P}_y^1) - 2) + \left(\frac{q^s}{2} + 1\right) \cdot 2,$$

whence $g(\mathcal{C}_1) = \frac{q^s}{2}$.

Proposition 3.13. The curve C_2 has genus $q^s - 1$.

Proof. By Lemma 3.11, C_2 is absolutely irreducible; hence, the covering φ_1 : $C_2 \to C_1$, $(X, Z) \mapsto (X, Y = Z^2 + Z + \epsilon)$, is an Artin-Schreier covering of degree 2. Every place of C_1 which is not a pole of $y - \epsilon$ is unramified in φ_1 . We consider the unique pole of $y - \epsilon$ on C_1 , namely E_1 . By direct computation, the Laurent series of y at E_1 with respect to the local parameter $t' = 1/(xy^{q^{s/2}})$ is $y - \epsilon = (t')^{-2} + (t')^{-1} + w$, with $v_{E_1}(w) \ge 0$. Choosing $\omega = (t')^{-1}$, we have $v_{E_1}((y - \epsilon) - (\omega^2 + \omega)) = v_{E_1}(w) \ge 0$. Thus, by Theorem 2.3, E_1 is unramified in φ_1 . Altogether, the covering $\varphi_1 : C_2 \to C_1$ is unramified. This implies

$$2g(\mathcal{C}_2) - 2 = \deg(\varphi_1) \cdot (2g(\mathcal{C}_1) - 2),$$

whence $g(\mathcal{C}_2) = q^s - 1$.

Theorem 3.14. For the curve C the following holds.

- (a) C is absolutely irreducible with genus $q^{2s} q^s 1$.
- (b) Let s and z be the coordinate functions of C, and let $t = s^2(z^2 + z + \epsilon)$. The number of \mathbb{F}_{q^n} -rational places of C which are zeros of t, or poles of either s or z or t, is at most $2q^s + 2$.

Proof. We compute the valuation of x at the places of C_2 . From the proofs of Propositions 3.12 and 3.13 follows that x has exactly 2 zeros on C_1 , namely E_2 with $v_{E_2}(x) = 1$ and E_1 with $v_{E_1}(x) = q^s - 1$; also, x has exactly 4 zeros on C_2 , namely the two places Q_1, Q_2 lying over E_2 and the two places Q_3, Q_4 over E_1 . Using the ramification indices, this implies $v_{Q_1}(x) = v_{Q_2}(x) = 1$ and $v_{Q_3}(x) = v_{Q_4}(x) = q^s - 1$. The unique pole of x on C_1 is E_0 ; since $v_{E_0}(y) = 1$, this implies $v_{E_0}(x) = -q^s$. The poles of x on C_2 are the two places R_1, R_2 lying over E_0 , with $v_{R_1}(x) = v_{R_2}(x) = -q^s$.

Therefore, by Theorem 2.2, C is absolutely irreducible and $\varphi_2 : C \to C_2$ is a Kummer covering of degree $q^s - 1$. The places of C_2 which ramify in φ_2 are exactly Q_1, Q_2, R_1, R_2 , and they are totally ramified; any other place is unramified in φ_2 . The genus of C is

$$g(\mathcal{C}) = 1 + \deg(\varphi_2) \cdot (g(\mathcal{C}_2) - 1) + \frac{1}{2} \cdot 4 \cdot (\deg(\varphi_2) - 1) = q^{2s} - q^s - 1.$$

Using the proofs of Propositions 3.12 and 3.13, we obtain that:

- s has exactly 2 poles on C, namely the places over R_1 or R_2 ;
- z has exactly $2(q^s 1)$ poles on \mathcal{C} , namely the places over Q_3 or Q_4 ;
- s^2 has exactly $2(q^s 1) + 2$ zeros on C; namely, two of them lie over Q_1 or Q_2 , while $2(q^s 1)$ of them lie over Q_3 or Q_4 (and have been been already considered above);

• $z^2 + z + \epsilon = y$ has exactly 4 zeros on C, namely the places over Q_1, Q_2, R_1, R_2 (which have been already considered above).

Altogether, the number of \mathbb{F}_{q^n} -rational places of \mathcal{C} which are poles of s, z, t, or are zeros of t, is smaller than or equal to $2q^s + 2$.

From Theorems 3.6 and 3.14 follows Corollary 3.15, which is our main result Theorem 1.1 when q is even.

Corollary 3.15. Let q be an even prime power, $s \ge 1$ be such that gcd(s, n) = 1. Suppose that

$$n \ge \begin{cases} 4s+2 & if q = 2 \ and \ s > 2; \\ 4s+1 & otherwise. \end{cases}$$

Then for every $\delta \in \mathbb{F}_{q^{2n}}$ satisfying $N_{q^{2n}/q^n}(\delta) \notin \{0,1\}$ there exists $a \in \mathbb{F}_{q^{2n}}^*$ such that $\dim_{\mathbb{F}_q} \ker(f_{a,b,s}) = 2$, where $b = \delta a$.

Proof. By Theorems 2.4 and 3.14(a), the number N_{q^n} of \mathbb{F}_{q^n} -rational places of \mathcal{C} satisfies

$$N_{q^n} \ge q^n + 1 - 2(q^{2s} - q^s - 1)\sqrt{q^n} > 2q^s + 2.$$

By Theorem 3.14(b), there exists an \mathbb{F}_{q^n} -rational affine poin (\bar{s}, \bar{z}) of \mathcal{C} such that $\bar{t} = \bar{s}^2(\bar{z}^2 + \bar{z} + \epsilon)$ is different from zero. The claim follows.

4 Applications to linear sets and rank metric codes

4.1 Linear sets

Let $\Lambda = \operatorname{PG}(V, \mathbb{F}_{q^m}) = \operatorname{PG}(1, q^m)$, where V is a vector space of dimension 2 over \mathbb{F}_{q^m} . A point set L of Λ is said to be an \mathbb{F}_q -linear set of Λ of rank k if it is defined by the non-zero vectors of a k-dimensional \mathbb{F}_q -vector subspace U of W, i.e.

$$L = L_U = \{ \langle \mathbf{u} \rangle_{\mathbb{F}_{q^m}} \colon \mathbf{u} \in U \setminus \{\mathbf{0}\} \}.$$

We say that two linear sets L_U and L_W of $\Lambda = PG(1, q^m)$ are PFL-equivalent if there exists $\varphi \in PFL(2, q^m)$ such that $\varphi(L_U) = L_W$. We start by pointing out that if the point $\langle (0,1) \rangle_{\mathbb{F}_{q^m}}$ is not contained in a linear set L_U of rank m of $\mathrm{PG}(1,q^m)$ (which we can always assume after a suitable projectivity), then $U = U_f = \{(x, f(x)) : x \in \mathbb{F}_{q^m}\}$ for some q-polynomial $f(x) = \sum_{i=0}^{m-1} a_i x^{q^i} \in \tilde{\mathcal{L}}_{m,q}$. In this case we will denote the associated linear set by L_f . Also, recall that the weight of a point $P = \langle \mathbf{u} \rangle_{\mathbb{F}_{q^m}}$ is $w_{L_U}(P) = \dim_{\mathbb{F}_q}(U \cap \langle \mathbf{u} \rangle_{\mathbb{F}_{q^m}}).$

One of the most studied classes of linear sets of the projective line, especially because of its applications (see e.g. [27, 29]), is the family of maximum scattered linear sets. A maximum scattered \mathbb{F}_q -linear set of $\mathrm{PG}(1, q^m)$ is an \mathbb{F}_q -linear set of rank m of $\mathrm{PG}(1, q^m)$ of size $(q^m - 1)/(q - 1)$, or equivalently a linear set of rank m in $\mathrm{PG}(1, q^m)$ all of whose points have weight one. If L_f is a maximum scattered linear set in $\mathrm{PG}(1, q^m)$, we also say that f is a scattered polynomial. The known scattered polynomials of \mathbb{F}_{q^m} are

- 1. $f_1(x) = x^{q^s} \in \hat{\mathcal{L}}_{m,q}$, with gcd(s,m) = 1, see [5];
- 2. $f_2(x) = x^{q^s} + \alpha x^{q^{m-s}} \in \tilde{\mathcal{L}}_{m,q}$, with $m \ge 4$, gcd(s,m) = 1, $N_{q^m/q}(\alpha) \notin \{0,1\}$, see [20, 23, 29];
- 3. $f_3(x) = x^{q^s} + \alpha x^{q^{s+\frac{m}{2}}} \in \tilde{\mathcal{L}}_{m,q}, m \in \{6,8\}, \text{gcd}(s,\frac{m}{2}) = 1$ and some conditions on α , see [8] and below;
- 4. $f_4(x) = x^q + x^{q^3} + \alpha x^{q^5} \in \tilde{\mathcal{L}}_{6,q}, q \text{ odd and } \alpha^2 + \alpha = 1, \text{ see } [13, 24];$
- 5. $f_5(x) = h^{q-1}x^q h^{q^2-1}x^{q^2} + x^{q^4} + x^{q^5} \in \tilde{\mathcal{L}}_{6,q}, q \text{ odd}, h^{q^3+1} = -1$, see [3, 35].

In [8], the authors introduced the family of linear sets $L_{\delta,s}$ of rank 2n in $PG(1, q^{2n})$ mentioned in 3., i.e. those linear sets defined by the \mathbb{F}_q -subspace

$$U_{\delta,s} = \{ (x, f_{\delta,s}(x)) \colon x \in \mathbb{F}_{q^{2n}} \} \subset \mathbb{F}_{q^{2n}} \times \mathbb{F}_{q^{2n}}, \tag{9}$$

where

$$f_{\delta,s}(x) = x^{q^s} + \delta x^{q^{n+s}} \in \tilde{\mathcal{L}}_{2n,q},$$

with $N_{q^{2n/q^n}}(\delta) \notin \{0,1\}, 1 \leq s \leq 2n-1 \text{ and } \gcd(s,n) = 1$. The relevance of this family relies on the property that each point of $L_{\delta,s}$ has weight at most two; see [8, Proposition 4.1]. In [8, Section 7] the authors proved that for n = 3 and q > 4 there exists $\delta \in \mathbb{F}_{q^2}$ such that $L_{s,\delta}$ is scattered; for n = 4,

q odd and $\delta^2 = -1$ the linear set $L_{\delta,s}$ is scattered. In [28, Theorem 7.3] the authors completely determined for n = 3 necessary and sufficient conditions on δ ensuring $L_{\delta,s}$ to be scattered. Note that for n = 3 we may restrict to the case s = 2, since every linear set $L_{\delta,s}$ is equivalent to $L_{\delta',2}$ for some $\delta' \in \mathbb{F}_{q^{2n}}^*$. More precisely, if $N_{q^6/q^3}(\delta) \notin \{0,1\}$ and we denote $A = -\frac{1}{\delta^{q^3+1}-1}$, one has that $L_{\delta,2}$ is scattered if and only if the equation

$$Y^{2} - (\operatorname{Tr}_{q^{3}/q}(A) - 1)Y + \operatorname{N}_{q^{3}/q}(A) = 0$$
(10)

admits two distinct roots in \mathbb{F}_q .

Theorem 4.1. Let q be a prime power and n, s be two relatively prime positive integers. Suppose that

$$n \ge \begin{cases} 4s+2 & \text{if } q=3 \text{ and } s>1, \text{ or } q=2 \text{ and } s>2;\\ 4s+1 & \text{otherwise.} \end{cases}$$

Then, for every $\delta \in \mathbb{F}_{q^{2n}}^*$, the \mathbb{F}_q -linear set $L_{\delta,s}$ in $\mathrm{PG}(1,q^{2n})$ is not scattered.

Proof. For every $m \in \mathbb{F}_{q^{2n}}$, the weight of the point $\langle (1,m) \rangle_{\mathbb{F}_{q^{2n}}}$ in $L_{\delta,s}$ coincides with the dimension over \mathbb{F}_q of the kernel of $f_{\delta,s}(x) - mx$.

If $N_{q^{2n}/q^n}(\delta) = 1$, then the point $\langle (1,0) \rangle_{\mathbb{F}_{q^{2n}}}$ has weight n in $L_{\delta,s}$. Let $N_{q^{2n}/q^n}(\delta) \neq 1$. By Theorem 1.1, there exists $a \in \mathbb{F}_{q^{2n}}^*$ such that $\dim_{\mathbb{F}_q} \ker(f_{a,\delta a,s}(x)) = 2$, whence

$$\dim_{\mathbb{F}_q} \ker \left(a \left(f_{\delta,s}(x) + \frac{1}{a} x \right) \right) = 2.$$

This implies that the point $\langle (1, -\frac{1}{a}) \rangle_{\mathbb{F}_{q^{2n}}}$ has weight 2 in $L_{\delta,s}$. The claim is proved.

Hence, we have the following description for the linear set $L_{\delta,s}$.

Corollary 4.2. Let q be a prime power and n, s be two relatively prime positive integers.

- If n = 3, then L_{δ,s} is a scattered linear set if and only if Equation 10 admits two distinct roots in F_q.
- If n = 4, q is odd and $\delta^2 = -1$ then $L_{\delta,s}$ is scattered.

• *If*

$$n \ge \begin{cases} 4s+2 & if q=3 \ and \ s>1, \ or \ q=2 \ and \ s>2, \\ 4s+1 & otherwise, \end{cases}$$

then, for every $\delta \in \mathbb{F}_{a^{2n}}^*$, $L_{\delta,s}$ is not scattered.

Proof. The claim follows from [28, Theorem 7.3], [8, Theorem 7.2], and Theorem 4.1. \Box

Among the known scattered polynomials listed above, the families in 3., 4. and 5. provide scattered polynomials for infinitely many q's, but only over a specific extension of \mathbb{F}_q , namely either \mathbb{F}_{q^6} or \mathbb{F}_{q^8} . Unlike this situation, the families in 1. and 2. provide scattered polynomials over infinitely many extensions \mathbb{F}_{q^m} of \mathbb{F}_q ; they are named respectively as scattered polynomials of pseudoregulus type, and as scattered polynomials of LP type (after Lunardon and Polverino).

The scattered polynomials of pseudoregulus or LP type have raised the following question: which polynomials over \mathbb{F}_{q^m} are scattered over infinitely many extensions of \mathbb{F}_{q^m} ?

Definition 4.3. [4, Section 1] Let $f(x) \in \tilde{\mathcal{L}}_{m,q}$, $0 \leq t \leq m-1$, $\ell \geq 1$, and $U_{\ell} = \{(x^{q^t}, f(x)) : x \in \mathbb{F}_{q^{m\ell}}\}$. We say that f(x) is an exceptional scattered polynomial of index t if $L_{U_{\ell}}$ is a scattered \mathbb{F}_q -linear set in $\mathrm{PG}(1, q^{m\ell})$ for infinitely many ℓ 's.

Clearly, the scattered polynomials of pseudoregulus type are exceptional scattered of index 0. Also, for the scattered polynomial $f_2(x)$ of LP type,

$$U_{f_2} = \{ (x^{q^s}, x^{q^{2s}} + \alpha x) \colon x \in \mathbb{F}_{q^m} \};$$

thus, the polynomial $x^{q^{2s}} + \alpha x$ is exceptional scattered of index s.

For a scattered polynomial $f(x) \in \tilde{\mathcal{L}}_{m,q}$ of index t, we say that f(x) is t-normalized if the following properties hold: f(x) is monic; the coefficient of x^{q^t} in f(x) is zero; if t > 0, the coefficient of x in f(x) is nonzero. Up to PGL-equivalence of the corresponding scattered linear set, we may always assume that f(x) is t-normalized.

Theorem 4.4. Let $f(x) \in \tilde{\mathcal{L}}_{m,q}$ be a t-normalized exceptional scattered polynomial of index t. Then the following holds.

- If t = 0, then f(x) is of pseudoregulus type; see [4, Corollary 3.4] for q > 5, [2, Section 4] for $q \le 5$.
- If t = 1 or t = 2, then f(x) is either of pseudoregulus type or of LP type; see [4, Corollary 3.7] for t = 1, [2, Corollary 1.4] for t = 2.
- If $t \ge 3$, q is odd, and $\max\{\deg_q f(x), t\}$ is an odd prime, then f(x) = x; see [15, Theorem 1.2].

Recall that the polynomials $f_3(x)$ of family 3. in the list above are scattered under certain assumptions for $m \in \{6, 8\}$; even when $f_3(x)$ is not scattered, still all the points of L_{f_3} have weight at most 2. Thus, one may conjecture that family 3. contains scattered polynomials over \mathbb{F}_{q^m} for every even m. Note that, even if this is the case, the arising scattered polynomials are not exceptional: not only the coefficients but also the degree depend heavily on the underlying field \mathbb{F}_{q^m} .

Our asymptotic result Theorem 1.1 shows that the family of scattered polynomial in 3. cannot be extended to any higher extension \mathbb{F}_{q^m} when m is large enough with respect to s.

4.2 Rank metric codes

Rank metric codes were introduced by Delsarte [14] in 1978 and they have been intensively investigated in recent years because of their applications; we refer to [30] for a recent survey on this topic. The set of $m \times n$ matrices $\mathbb{F}_q^{m \times n}$ over \mathbb{F}_q may be endowed with a metric, called *rank metric*, defined by

$$d(A, B) = \operatorname{rk}(A - B).$$

A subset $\mathcal{C} \subseteq \mathbb{F}_q^{m \times n}$ equipped with the rank metric is called a *rank metric code* (shortly, a *RM*-code). The minimum distance of \mathcal{C} is defined as

$$d = \min\{d(A, B) \colon A, B \in \mathcal{C}, \ A \neq B\}.$$

Denote the parameters of a RM-code $\mathcal{C} \subseteq \mathbb{F}_q^{m,n}$ with minimum distance d by (m, n, q; d). We are interested in \mathbb{F}_q -linear RM-codes, i.e. \mathbb{F}_q -subspaces of $\mathbb{F}_q^{m \times n}$. Delsarte showed in [14] that the parameters of these codes must obey a Singleton-like bound, i.e.

$$|\mathcal{C}| < q^{\max\{m,n\}(\min\{m,n\}-d+1)}$$

When equality holds, we call \mathcal{C} a maximum rank distance (MRD for short) code. Examples of \mathbb{F}_q -linear MRD-codes were first found in [14, 16]. We say that two \mathbb{F}_q -linear RM-codes \mathcal{C} and \mathcal{C}' are equivalent if there exist $X \in$ $\mathrm{GL}(m,q), Y \in \mathrm{GL}(n,q)$, and $\sigma \in \mathrm{Aut}(\mathbb{F}_q)$ such that

$$\mathcal{C}' = \{ X C^{\sigma} Y \colon C \in \mathcal{C} \}.$$

The *left* and *right* idealisers of C are defined in [21] as $L(C) = \{A \in GL(m,q) : AC \subseteq C\}$ and $R(C) = \{B \in GL(n,q) : CB \subseteq C\}$. They are invariant under the equivalence of rank metric codes, and have been investigated in [22]; further invariants have been introduced in [17, 26].

Much of the focus on MRD-codes of $\mathbb{F}_q^{m \times m}$ to date has been on codes which are \mathbb{F}_{q^m} -linear, i.e. codes in which the left (or right) idealiser contains a field isomorphic to \mathbb{F}_{q^m} , since for such codes a fast decoding algorithm has been developed in [16]. Very few examples of such codes are known, see [3, 8, 9, 13, 14, 16, 23, 24, 29, 35].

In [29, Section 5] Sheekey showed that scattered \mathbb{F}_q -linear sets of PG(1, q^m) of rank m yield \mathbb{F}_q -linear MRD-codes with parameters (m, m, q; m - 1) with left idealiser isomorphic to \mathbb{F}_{q^m} ; see [10, 12, 31] for further details on such kind of connections. We briefly recall here the construction from [29]. Let $U_f = \{(x, f(x)): x \in \mathbb{F}_{q^m}\}$, where f(x) is a scattered q-polynomial. The choice of an \mathbb{F}_q -basis for \mathbb{F}_{q^m} defines a canonical ring isomorphism between End($\mathbb{F}_{q^m}, \mathbb{F}_q$) and $\mathbb{F}_q^{m \times m}$. Thus, the set

$$\mathcal{C}_f = \{x \mapsto af(x) + bx \colon a, b \in \mathbb{F}_{q^m}\} \subset \operatorname{End}(\mathbb{F}_{q^m}, \mathbb{F}_q)$$

corresponds to a set of $m \times m$ matrices over \mathbb{F}_q forming an \mathbb{F}_q -linear MRDcode with parameters (m, m, q; m - 1). Also, as \mathcal{C}_f is an \mathbb{F}_{q^m} -subspace of End $(\mathbb{F}_{q^m}, \mathbb{F}_q)$, its left idealiser $L(\mathcal{C}_f)$ is isomorphic to \mathbb{F}_{q^m} ; see also [8, Section 6].

Now consider the set

$$\mathcal{C}_{f_{\delta,s}} = \{ x \mapsto a(x^{q^s} + \delta x^{q^{s+n}}) + bx \colon a, b \in \mathbb{F}_{q^{2n}} \},\$$

which corresponds to a set of $2n \times 2n$ matrices over \mathbb{F}_q forming an \mathbb{F}_q -linear rank metric code with parameters (2n, 2n, q; 2n - i), where

$$i = \max\{w_{L_{\delta,s}}(P) \colon P \in \mathrm{PG}(1,q^{2n})\}$$

The following theorem is a consequence of Corollary 4.2 and states that, when n is large enough, $C_{f_{\delta,s}}$ is not an MRD-code.

Theorem 4.5. Let q be a prime power and n, s be two relatively prime positive integers.

- If n = 3, then C_{f_{δ,s}} is an MRD-code if and only if Equation 10 admits two distinct roots in F_q; see [8] and [28].
- If n = 4, q odd and $\delta^2 = -1$ then $C_{f_{\delta,s}}$ is an MRD-code; see [8].
- *If*

$$n \ge \begin{cases} 4s+2 & \text{if } q=3 \text{ and } s>1, \text{ or } q=2 \text{ and } s>2\\ 4s+1 & \text{otherwise;} \end{cases}$$

then, for every $\delta \in \mathbb{F}_{q^{2n}}^*$, $\mathcal{C}_{f_{\delta,s}}$ is not an MRD-code and its minimum distance is n-2.

References

- D. BARTOLI, M. GIULIETTI, G. MARINO AND O. POLVERINO: Maximum scattered linear sets and complete caps in Galois spaces, *Combinatorica* 38(2) (2018), 255–278.
- [2] D. BARTOLI AND M. MONTANUCCI: Towards the full classification of exceptional scattered polynomials, arXiv:1905.11390.
- [3] D. BARTOLI, C. ZANELLA AND F. ZULLO: A new family of maximum scattered linear sets in $PG(1, q^6)$, Ars Math. Contemp. (forthcoming).
- [4] D. BARTOLI AND Y. ZHOU: Exceptional scattered polynomials, J. Algebra 509 (2018), 507–534.
- [5] A. BLOKHUIS AND M. LAVRAUW: Scattered spaces with respect to a spread in PG(n, q), Geom. Dedicata **81** (2000), 231–243.
- [6] B. CSAJBÓK: Scalar q-subresultants and Dickson matrices, J. Algebra 547 (2020), 116–128.
- [7] B. CSAJBÓK, G. MARINO AND O. POLVERINO: Classes and equivalence of linear sets in $PG(1, q^n)$, J. Combin. Theory Ser. A **157** (2018), 402–426.

- [8] B. CSAJBÓK, G. MARINO, O. POLVERINO AND C. ZANELLA: A new family of MRD-codes, *Linear Algebra Appl.* 548 (2018), 203–220.
- [9] B. CSAJBÓK, G. MARINO, O. POLVERINO AND Y. ZHOU: Maximum rank-distance codes with maximum left and right idealisers, arXiv:1807.08774.
- [10] B. CSAJBÓK, G. MARINO, O. POLVERINO AND F. ZULLO: A special class of scattered subspaces, arXiv:1906.10590.
- [11] B. CSAJBÓK, G. MARINO, O. POLVERINO AND F. ZULLO: A characterization of linearized polynomials with maximum kernel, *Finite Fields Appl.* 56 (2019), 109–130.
- [12] B. CSAJBÓK, G. MARINO, O. POLVERINO AND F. ZULLO: Maximum scattered linear sets and MRD-codes, J. Algebraic Combin. 46 (2017), 1–15.
- [13] B. CSAJBÓK, G. MARINO AND F. ZULLO: New maximum scattered linear sets of the projective line, *Finite Fields Appl.* 54 (2018), 133–150.
- [14] P. DELSARTE: Bilinear forms over a finite field, with applications to coding theory, J. Combin. Theory Ser. A 25 (1978), 226–241.
- [15] A. FERRAGUTI AND G. MICHELI: Exceptional scatteredness in prime degree, arXiv:2002.00500.
- [16] E. GABIDULIN: Theory of codes with maximum rank distance, Problems Inform. Transmission, 21(3) (1985), 3–16.
- [17] L. GIUZZI AND F. ZULLO: Identifiers for MRD-codes, *Linear Algebra Appl.* 575 (2019), 66–86.
- [18] R. GOW AND R. QUINLAN: Galois extensions and subspaces of alterning bilinear forms with special rank properties, *Linear Algebra Appl.* 430 (2009), 2212–2224.
- [19] J.W.P. HIRSCHFELD, G. KORCHMÁROS, AND F. TORRES: Algebraic Curves over a Finite Field, *Princeton Series in Applied Mathematics*, Princeton (2008).

- [20] M. LAVRAUW, G. MARINO, O. POLVERINO AND TROMBETTI: Solution to an isotopism question concerning rank 2 semifields, J. Combin. Des. 23(2) (2015), 60–77.
- [21] D. LIEBHOLD AND G. NEBE: Automorphism groups of Gabidulin-like codes, Arch. Math. 107(4) (2016), 355–366.
- [22] G. LUNARDON, R. TROMBETTI AND Y. ZHOU: On kernels and nuclei of rank metric codes, J. Algebraic Combin. 46 (2017), 313–340.
- [23] G. LUNARDON AND O. POLVERINO: Blocking Sets and Derivable Partial Spreads, J. Algebraic Combin. 14 (2001), 49–56.
- [24] G. MARINO, M. MONTANUCCI AND F. ZULLO: MRD-codes arising from the trinomial $x^q + x^{q^3} + cx^{q^5} \in \mathbb{F}_{q^6}[x]$, Linear Algebra Appl. **591** (2020), 99–114.
- [25] G. MCGUIRE AND J. SHEEKEY: A Characterization of the Number of Roots of Linearized and Projective Polynomials in the Field of Coefficients, *Finite Fields Appl.* 57 (2019), 68–91.
- [26] A. NERI, S. PUCHINGER, A. HORLEMANN-TRAUTMANN: Equivalence and Characterizations of Linear Rank-Metric Codes Based on Invariants, arXiv:1911.13059.
- [27] O. POLVERINO: Linear sets in finite projective spaces, *Discrete Math.* 310(22) (2010), 3096–3107.
- [28] O. POLVERINO AND F. ZULLO: On the number of roots of some linearized polynomials, arXiv:1909.00802.
- [29] J. SHEEKEY: A new family of linear maximum rank distance codes, Adv. Math. Commun. 10(3) (2016), 475–488.
- [30] J. SHEEKEY: MRD codes: constructions and connections, Combinatorics and finite fields: Difference sets, polynomials, pseudorandomness and applications, Radon Series on Computational and Applied Mathematics, K.-U. Schmidt and A. Winterhof (eds.).
- [31] J. SHEEKEY AND G. VAN DE VOORDE: Rank-metric codes, linear sets and their duality, *Des. Codes Cryptogr.* 88 (2020), 655–675.

- [32] H. STICHTENOTH: Algebraic Function Fields and Codes, Volume 254 of Graduate Texts in Mathematics, 2nd edition, Springer, Berlin (2009).
- [33] B. WU AND Z. LIU:, Linearized polynomials over finite fields revisited, *Finite Fields Appl.* 22 (2013), 79–100.
- [34] C. ZANELLA: A condition for scattered linearized polynomials involving Dickson matrices, J. Geom. (2019) **110**:50. https://doi.org/10.1007/s00022-019-0505-z.
- [35] C. ZANELLA AND F. ZULLO: Vertex properties of maximum scattered linear sets of $PG(1, q^n)$, Discrete Math. **343(5)** (2020).

Olga Polverino, Giovanni Zini and Ferdinando Zullo Dipartimento di Matematica e Fisica, Università degli Studi della Campania "Luigi Vanvitelli", Viale Lincoln 5, I-81100 Caserta, Italy {olga.polverino,giovanni.zini,ferdinando.zullo}@unicampania.it