

RESOLUTION ANALYSIS OF INVERTING THE GENERALIZED RADON TRANSFORM FROM DISCRETE DATA IN \mathbb{R}^3

ALEXANDER KATSEVICH¹

ABSTRACT. A number of practically important imaging problems involve inverting the generalized Radon transform (GRT) \mathcal{R} of a function f in \mathbb{R}^3 . On the other hand, not much is known about the spatial resolution of the reconstruction from discretized data. In this paper we study how accurately and with what resolution the *singularities* of f are reconstructed. The GRT integrates over a fairly general family of surfaces \mathcal{S}_y in \mathbb{R}^3 . Here y is the parameter in the data space, which runs over an open set $\mathcal{V} \subset \mathbb{R}^3$. Assume that the data $g(y) = (\mathcal{R}f)(y)$ are known on a regular grid y_j with step-sizes $O(\epsilon)$ along each axis, and suppose $\mathcal{S} = \text{singsupp}(f)$ is a piecewise smooth surface. Let f_ϵ denote the result of reconstruction from the discrete data. We obtain explicitly the leading singular behavior of f_ϵ in an $O(\epsilon)$ -neighborhood of a generic point $x_0 \in \mathcal{S}$, where f has a jump discontinuity. We also prove that under some generic conditions on \mathcal{S} (which include, e.g. a restriction on the order of tangency of \mathcal{S}_y and \mathcal{S}), the singularities of f do not lead to non-local artifacts. For both computations, a connection with the uniform distribution theory turns out to be important. Finally, we present a numerical experiment, which demonstrates a good match between the theoretically predicted behavior and actual reconstruction.

1. INTRODUCTION

A large number of practically important imaging problems involve inversion of the generalized Radon transform (GRT), i.e. recovering an unknown function f from its integrals over a family of surfaces. The reconstruction may involve finding f itself, or finding f modulo smoother terms. Most of the times, the surfaces are not planes. Below is a list of some of the most common integral transforms with some of the most prominent examples of their use.

- (1) Integration over spheres. Applications include ultrasound imaging (or, SONAR) (see [15] and references therein) as well as thermoacoustic and photoacoustic tomography [12, 21].
- (2) Integration over ellipses. This transform arises in linearized seismic imaging with a common offset between the sources and receivers [7].
- (3) Integration over cones arises in Compton camera imaging. Applications are single-scattering optical tomography, Compton camera medical imaging, and homeland security (see [19] for a recent review).

In all of the above cases, one collects a discrete data set and reconstructs f using a numerical algorithm. Frequently, reconstruction is achieved by applying a linear inversion formula (as opposed to a non-linear reconstruction algorithm based on fidelity functional minimization). In all of the above examples it is of fundamental importance to know the resolution of the method as a function of (1) the data sampling rate, and (2) specific implementation of the inversion formula that is used. Despite the significance of this problem, not much is known about the resolution of reconstruction from discrete data. The main reason for this is

¹Department of Mathematics, University of Central Florida, Orlando, FL 32816. This work was supported in part by NSF grant DMS-1615124.

that the classical sampling theory, which addresses such problems, can be applied only in a few simplest settings of the classical Radon transform (CRT) [14]. The known results are quite scarce, and they are of semi-qualitative nature (see e.g. pp. 784–786 in [6]). Very recently, a more flexible approach to sampling based on semiclassical analysis was proposed in [18]. Let A be a Fourier Integral Operator (FIO). The idea of [18] is to determine how the data Af should be sampled to allow for accurate interpolation of its values on a lattice provided that f is semiclassically bandlimited. If the sampling condition is violated, then reconstruction from the discrete values of Af (i.e., applying a parametrix A^{-1} to the interpolated Af) leads to aliasing artifacts, which are also analyzed in [18].

An alternative approach to the analysis of resolution was proposed recently in [10, 11]. The idea is to investigate how accurately and with what resolution the *singularities* of f are reconstructed. For some of the above problems there is no exact inversion formula, and inversion modulo smoother terms is the most one can hope for. In such cases, spatial resolution of the recovery of singularities is all one needs. Note that in this paper both f and $g = \mathcal{R}f$ are assumed to have singularities in the sense of a conventional, classical wavefront set (see e.g. [8]). In contrast, the main assumption in [18] is that f and, consequently, the data Af have only semiclassical singularities (see e.g. [23]). It is possible to apply the approach of [18] to the analysis of classical singularities, but this would require summing a series over “folded” frequencies in the Fourier domain, which is complicated.

In [10, 11] the author considers the inversion of the CRT of f in \mathbb{R}^2 and \mathbb{R}^3 . The parametrization of the data is standard, i.e. in terms of the affine and angular variables. Suppose the step-sizes along the angular and affine variables are $O(\epsilon)$. Let f_ϵ denote the result of reconstruction from the discrete data. The author picks a point x_0 , where f has a jump singularity, and obtains explicitly the leading singular behavior of f_ϵ in an $O(\epsilon)$ -neighborhood of x_0 as $\epsilon \rightarrow 0$. The obtained behavior, which we call *edge response*, provides the desired resolution of the reconstruction algorithm. It is shown also that convex parts of the singular support of f do not create non-local artifacts. The case when f changes during the scan (so-called, dynamic CT) is considered in the 2D setting as well [10].

In this paper we generalize the approach of [10, 11]. The reconstruction problem is now formulated in terms of the GRT \mathcal{R} , which integrates f over a fairly general family of surfaces \mathcal{S}_y in \mathbb{R}^3 . Here $\text{supp}(f) \subset \mathcal{U}$, where $\mathcal{U} \subset \mathbb{R}^3$ is an open set, and y is the parameter in the data space. For the problem to be well-determined, we assume that y runs over an open set $\mathcal{V} \subset \mathbb{R}^3$. As is seen, our setting is fairly general and covers all the problems mentioned above. The GRT in this paper is very close to that considered by Beylkin in [2], only the parametrization of the surfaces \mathcal{S}_y is slightly different. This gives us more flexibility to connect our results with practical applications, where GRTs arise.

Assume that the data $g = \mathcal{R}f$ are known on a regular grid y_j with step-sizes $O(\epsilon)$ along each axis. Suppose $\mathcal{S} = \text{singsupp}(f)$ is a piecewise smooth surface. Similarly to [10, 11], we obtain explicitly the leading singular behavior of f_ϵ in an $O(\epsilon)$ -neighborhood of a generic point $x_0 \in \mathcal{S}$, where f has a jump discontinuity. We also prove that under some generic conditions on \mathcal{S} (which include, e.g. a restriction on the order of tangency of \mathcal{S}_y and \mathcal{S}), the singularities of f do not lead to non-local artifacts. For both computations, a connection with the uniform distribution theory [13] turns out to be important. It is possible that violation of the imposed conditions leads to artifacts. Analysis of such artifacts and analysis of more general surfaces \mathcal{S} will be the subject of future research.

The reconstruction formula $g \rightarrow \check{f}$, which contains a suitably adapted adjoint \mathcal{R}^* , is one specific example of an FIO. Here \check{f} is such that $\check{f} - f$ is smoother than

f. Thus, the reconstruction algorithm can be viewed as an application of an FIO to discrete data $(\mathcal{R}f)(y_j)$. A number of methods for computing the action of FIOs on discrete data have been proposed, see e.g. [4, 5, 1, 22] and references therein. To the best of the author's knowledge, proposed is the first method to compute the *resolution* of the reconstruction obtained by applying an FIO to discrete data that comes from an image with classical singularities. Extension of the method to more general FIOs will also be the subject of future work.

The paper is organized as follows. In Section 2 we define the GRT \mathcal{R} via an incidence relation $\mathcal{C} \subset \mathcal{U} \times \mathcal{V}$, list the properties of the function $\Phi(x, y)$ that defines the incidence relation, define generic points, and specify the continuous and discrete inversion formulas that are used in the analysis. The main result is formulated in Section 3, which also contains the beginning of the proof. The entire proof spans Sections 3–7. In Section 3 we obtain the behavior of g near its singular support, which generalizes one of the results of [16, 17] from the CRT to the GRT. The behavior of the interpolated data g_ϵ near $\text{singsupp}(g)$ is obtained in Section 4. The contribution of the leading singular term to the edge response at a generic point $x_0 \in \mathcal{S}$ is computed in Section 5. In Section 6 we show that lower order terms do not contribute to the edge response. In Section 7 we prove that, under some assumptions, remote singularities do not contribute to the edge response as well. Results of a numerical experiment, which show a good match between the theoretically predicted behavior and actual reconstruction, are in Section 8.

2. PRELIMINARY CONSTRUCTION

Let $\mathcal{U}, \mathcal{V} \subset \mathbb{R}^3$ be two open connected sets, where \mathcal{U} is the image domain, and \mathcal{V} is the data domain. Each $y \in \mathcal{V}$ determines a smooth surface $\mathcal{S}_y \subset \mathcal{U}$. Let \mathcal{C} be the corresponding incidence relation $\mathcal{C} \subset \mathcal{U} \times \mathcal{V}$, which is defined in terms of a smooth function $\Phi(x, y) \in C^\infty(\mathcal{U} \times \mathcal{V})$:

$$(2.1) \quad \mathcal{C} := \{(x, y) \in \mathcal{U} \times \mathcal{V} : \Phi(x, y) = 0\}.$$

Another way to state (2.1) is that $x \in \mathcal{S}_y$ if and only if $\Phi(x, y) = 0$. Define the submanifold:

$$(2.2) \quad \mathcal{T}_x := \{y \in \mathcal{V} : \Phi(x, y) = 0\}, \quad x \in \mathcal{U}.$$

Thus, \mathcal{T}_x is the collection of all $y \in \mathcal{V}$ such that \mathcal{S}_y contains x . The main assumptions about Φ are as follows (“DF” stands for Defining Function):

DF1. Φ is real-valued and non-degenerate, i.e.

$$(2.3) \quad \Phi'_x(x, y) \neq 0, \quad \Phi'_y(x, y) \neq 0, \quad (x, y) \in \mathcal{C};$$

DF2. For each $x \in \mathcal{U}$, the map $\mathcal{T}_x \rightarrow S^2$ defined by $y \rightarrow \pm \Phi'_x(x, y) / |\Phi'_x(x, y)|$, $y \in \mathcal{T}_x$, is surjective;

DF3. For each $y \in \mathcal{V}$, the vectors $\Phi'_y(x, y)$ and $\Phi'_y(z, y)$ are not parallel whenever $x, z \in \mathcal{S}_y$, $x \neq z$; and

DF4. The mixed Hessian of Φ is non-degenerate,

$$(2.4) \quad \det \left(\frac{\partial^2 \Phi(x, y)}{\partial x_i \partial y_j} \right) \neq 0, \quad (x, y) \in \mathcal{C},$$

where $\partial/\partial x_i$, $i = 1, 2$, and $\partial/\partial y_j$, $j = 1, 2$, are basis vectors in the tangent spaces to the submanifolds \mathcal{S}_{y_0} and \mathcal{T}_{x_0} at x_0 and y_0 , respectively.

Condition DF2 means that the tomographic data are complete, i.e. any singularity is visible. Condition DF3 says that there are no conjugate points. Condition DF4 is a local version of the Bolker condition. Conditions DF3 and DF4 imply the (global) Bolker condition. Conditions DF1–DF4 are analogous to Conditions (I)–(IV) in [2].

Conditions DF1 and DF4 combined are equivalent to the condition (cf. eq. (4.23), [20], p. 335) that at every point $(x, y) \in \mathcal{C}$:

$$(2.5) \quad \det \begin{pmatrix} \Phi''_{xy} & (\Phi'_x)^T \\ \Phi'_y & 0 \end{pmatrix} \neq 0.$$

In the paper we consider functions, which can be represented as a finite sum

$$(2.6) \quad f(x) = \sum_j \chi_{D_j} f_j(x),$$

where χ_{D_j} is the characteristic function of the domain $D_j \subset \mathcal{U}$. For each j :

- (1) D_j is bounded,
- (2) The boundary of D_j is piecewise C^∞ ,
- (3) f_j is C^∞ in a domain containing the closure of D_j .

Denote $\mathcal{S} := \cup_j \partial D_j$. By construction, $\text{singsupp}(f) \subset \mathcal{S}$.

The GRT of f is given by:

$$(2.7) \quad g(y) = (\mathcal{R}f)(y) := \int_{\mathcal{S}_y} b(x, y) f(x) dx, \quad y \in \mathcal{V},$$

where the weight b is smooth (i.e., C^∞) and non-vanishing, and dx is the area element on \mathcal{S}_y . The discrete data are given by

$$(2.8) \quad g(\epsilon j), \quad j \in r + \mathbb{Z}^3,$$

for some $r \in \mathbb{R}^3$.

Even though (2.8) assumes that the stepsize along each data axis equals ϵ , this is a non-restrictive assumption. Indeed, consider a smooth diffeomorphism $\psi: \mathcal{V} \rightarrow \tilde{\mathcal{V}}$ for some open $\tilde{\mathcal{V}} \subset \mathbb{R}^2$, so that ψ maps an irregular grid covering \mathcal{V} into a regular, square grid covering $\tilde{\mathcal{V}}$. Introducing a new defining function $\tilde{\Phi}(x, \tilde{y}) := \Phi(x, \psi^{-1}(\tilde{y}))$, we can transform any smoothly sampled data set into the one with a square grid. Clearly, if Φ satisfies DF1–DF4, then $\tilde{\Phi}$ satisfies DF1–DF4 as well.

Conditions DF1–DF4 imply that (cf. [2] and [20], Sections VIII.5 and VIII.6):

- (1) The GRT \mathcal{R} is a Fourier Integral Operator (FIO) with phase function $\lambda\Phi(x, y)$;
- (2) The corresponding canonical relation is

$$(2.9) \quad C := \{((x, \lambda\Phi'_x(x, y)), (y, -\lambda\Phi'_y(x, y))) : \Phi(x, y) = 0, \lambda \in \mathbb{R} \setminus 0, x \in \mathcal{U}, y \in \mathcal{V}\},$$

which is a local canonical graph;

- (3) Any suitably modified adjoint of \mathcal{R} , denoted \mathcal{R}^* , is also an FIO, whose canonical relation C^* is obtained from (2.9) by switching the $(x, \xi) \in T^*\mathcal{U}$ and $(y, \eta) \in T^*\mathcal{V}$ variables; and
- (4) The composition $\mathcal{R}^*(\dots)\mathcal{R}$, where the dots denote a cut-off combined with a suitable differential operator, is a pseudo-differential operator (Ψ DO), i.e. $C^* \circ C$ is a subset of the diagonal in $T^*\mathcal{U}$.

Given a point $x \in \mathcal{S}$, find $y = y(x)$ (which is smooth locally) such that \mathcal{S}_y is tangent to \mathcal{S} at x . Denote

$$(2.10) \quad N(x) := \Pi_{\mathcal{S}_y}(x) - \Pi_{\mathcal{S}}(x),$$

where $\Pi_{\mathcal{S}}(x)$ is the matrix of the second fundamental form of \mathcal{S} at $x \in \mathcal{S}$ written in an orthonormal basis of $T_x\mathcal{S}$.

For any $x \in \mathcal{U}$, introduce the set

$$(2.11) \quad \Gamma_x := \{y \in \mathcal{V} : x \in \mathcal{S}_y, \mathcal{S}_y \text{ is tangent to } \mathcal{S} \text{ at some } z, z \neq x\}.$$

Definition 1. A pair $(x_0, y_0) \in \mathcal{C}$ is **globally generic** if whenever \mathcal{S}_{y_0} is tangent to \mathcal{S} at some $z \neq x_0$ the following conditions hold:

- GG1. \mathcal{S} is smooth at z , and $N(z)$ is either positive or negative definite; and
 GG2. Let $\dot{\Gamma}_{x_0}$ be a non-vanishing at any point tangent vector field along Γ_{x_0} .
 There exists an open set \mathcal{V}_1 , $y_0 \in \mathcal{V}_1 \subset \mathcal{V}$, such that for each $m \in \mathbb{Z}^3$,
 $|m| > 0$, and all $\delta > 0$ sufficiently small,
 (1) The set $\{y \in \Gamma_{x_0} \cap \mathcal{V}_1 : |m \cdot \dot{\Gamma}_{x_0}(y)| \leq \delta\}$ is contained in a finite number
 of segments of Γ_{x_0} (this number may depend on m and δ), and
 (2) The sum of the lengths of these segments goes to zero as $\delta \rightarrow 0$.

As is shown in Section 7, Conditions GG1 and DF3 imply that Γ_{x_0} is a smooth curve, so Condition GG2 makes sense.

An example when Condition GG2 is violated is when Γ_{x_0} contains a straight line segment and $m \cdot \dot{\Gamma}_{x_0}(y) \equiv 0$ on this segment for some $m \in \mathbb{Z}^3$, $|m| > 0$.

Definition 2. A pair $(x_0, y_0) \in \mathcal{C}$, $x_0 \in \mathcal{S}$, is **locally generic** if whenever \mathcal{S}_{y_0} is tangent to \mathcal{S} at x_0 the following conditions hold:

- LG1. \mathcal{S} is smooth at x_0 , and $N(z)$ is either positive or negative definite; and
 LG2. There is no $\lambda \neq 0$ such that $\lambda \Phi'_y(x_0, y_0) \in \mathbb{Z}^3$.

Definition 3. A pair $(x_0, y_0) \in \mathcal{C}$ is **generic** if it is both locally and globally generic.

Let φ be an interpolating kernel (IK), i.e. $\varphi(0) = 1$ and $\varphi(j) = 0$ for all $j \in \mathbb{Z}$, $j \neq 0$. Suppose also that φ satisfies the following assumptions:

IK1. φ is exact up to the order 2, i.e.

$$(2.12) \quad \sum_{j \in \mathbb{Z}^3} j^m \varphi(u - j) = u^m, \quad m \in (0 \cup \mathbb{N})^3, |m| \leq 2, u \in \mathbb{R}^3;$$

IK2. φ is compactly supported;

IK3. All partial derivatives of φ up to the order 2 are continuous;

IK4. All partial derivatives of φ of order 3 are piecewise continuous and bounded;
 and

IK5. φ is normalized, i.e. $\int \varphi(y) dy = 1$.

The interpolated version of g can be written in the form

$$(2.13) \quad g_\epsilon(y) := \sum_{j \in r + \mathbb{Z}^3} g(\epsilon j) \varphi\left(\frac{y - \epsilon j}{\epsilon}\right).$$

First, we derive a microlocal inversion formula for the GRT \mathcal{R} , which reconstructs exactly the leading singularities of f . Pick any $(x_0, y_0) \in \mathcal{C}$. Let α_0 be a unit vector normal to \mathcal{S}_{y_0} at x_0 . For $(x, \alpha) \in \mathcal{U} \times S^2$ close to (x_0, α_0) and for t , $|t| \ll 1$, find the local solution $y = Y(\alpha, t; x)$ such that $x + t\alpha \in \mathcal{S}_y$ and α is normal to \mathcal{S}_y at $x + t\alpha$. By construction, $y_0 = Y(\alpha_0, t = 0; x_0)$. Here we use the assumption that the data are complete, i.e. such a solution exists. It is shown below (see (4.2)) that the map $(\alpha, t) \rightarrow y = Y(\alpha, t; x)$ is a local diffeomorphism that depends smoothly on x .

Let \mathcal{V}_1 be a small neighborhood of y_0 . Pick any $\chi \in C_0^\infty(\mathcal{V}_1)$ such that $\chi \equiv 1$ near y_0 . The inversion formula with continuous data is given by

$$(2.14) \quad f_\chi(x) = -\frac{1}{4\pi^2} \int_{S^2_+} \frac{\chi(Y(\alpha, 0; x))}{b(x, Y(\alpha, 0; x))} \left(\frac{\partial}{\partial t} \right)^2 g(Y(\alpha, t; x)) \Big|_{t=0} d\alpha.$$

This inversion formula emulates the CRT inversion formula by backprojecting a second order derivative of the GRT. The affine variable t is computed relative to x as opposed to the origin, as is the case with the CRT. Hence the GRT analogue of the usual term $\alpha \cdot x$ is missing from (2.14), because it is absorbed by the function

Y . Due to the symmetry $g(Y(\alpha, t; x)) = g(Y(-\alpha, -t; x))$, in (2.14) we integrate over half of the unit sphere S_+^2 .

Using the argument following (2.8), it is easy to show that the map $f \rightarrow f_\chi$ is a Ψ DO of degree zero with principal symbol 1 microlocally near (x_0, α_0) (see e.g. [2, 9]). Thus, the singularities of f and f_χ are the same to leading order (e.g., in the scale of Sobolev spaces) microlocally near (x_0, α_0) . An inversion formula that recovers all the singularities of f can be obtained by combining (2.14) with a microlocal partition of unity. In the case of discrete data, we use the same inversion formula (2.14), but replace g with g_ϵ . The corresponding reconstruction is denoted $f_{\chi\epsilon}$.

3. STATEMENT OF MAIN RESULT. BEGINNING OF PROOF

3.1. Statement of main result. Pick $(x_0, y_0) \in \mathcal{C}$ such that \mathcal{S} is smooth at $x_0 \in \mathcal{S}$, \mathcal{S}_{y_0} is tangent to \mathcal{S} at x_0 , and $N(x_0)$ is either positive definite or negative definite. Fix some orthonormal basis in the common plane tangent to \mathcal{S} and \mathcal{S}_y at x_0 . Let α_0 be the unit vector normal to \mathcal{S} at x_0 . For convenience of calculations, we sometime use the Cartesian coordinates (x_1, x^\perp) determined by

$$(3.1) \quad x = x_1 \alpha_0 + x^\perp, \quad x^\perp \in \alpha_0^\perp,$$

where α_0^\perp is the plane through x_0 and normal to α_0 . This plane is tangent to both \mathcal{S} and \mathcal{S}_{y_0} at x_0 . The direction of α_0 is chosen so that $N(x_0)$ is *negative definite*. The side of \mathcal{S} where α_0 points is called *interior*. The other side of \mathcal{S} is called *exterior*. If necessary, multiply Φ by (-1) so that $\Phi'_x(x_0, y_0)/|\Phi'_x(x_0, y_0)| = -\alpha_0$.

Consider the point

$$(3.2) \quad x_\epsilon := x_0 + \epsilon \tilde{x}, \quad \tilde{x} \in \tilde{\mathcal{U}},$$

where $\tilde{\mathcal{U}}$ is a bounded set. Denote:

$$(3.3) \quad \begin{aligned} f_\chi(x_{0^\pm}) &:= \lim_{\epsilon \rightarrow 0^\pm} f_\chi(x_0 + \epsilon \alpha_0), \quad f_0 := \lim_{\epsilon \rightarrow 0^+} (f(x_0 + \epsilon \alpha_0) - f(x_0 - \epsilon \alpha_0)), \\ \nu &:= \frac{|\Phi'_x|}{|\Phi'_y|}, \quad \beta_0 = \frac{\Phi'_y}{|\Phi'_y|}. \end{aligned}$$

Here and in what follows the convention is that if the arguments of Φ and its derivatives are omitted, then they are evaluated at x_0, y_0 .

We also introduce local y -coordinates with the origin at y_0 :

$$(3.4) \quad y = (y_1, y^\perp) = y_1 \beta_0 + y^\perp.$$

Thus, equation $y_1 = 0$ determines the plane tangent to the submanifold \mathcal{T}_{x_0} at y_0 . We frequently denote this plane β_0^\perp .

Finally, we use the extension of the CRT to all of $\mathbb{R}^3 \setminus 0$ according to:

$$(3.5) \quad \hat{f}(u, s) := \int f(x) \delta(u \cdot x - s) dx, \quad u \in \mathbb{R}^3 \setminus 0,$$

for a sufficiently smooth f .

The main result of the paper is the following

Theorem 1. *Pick a generic pair $(x_0, y_0) \in \mathcal{C}$ such that \mathcal{S}_{y_0} is tangent to \mathcal{S} at $x_0 \in \mathcal{S}$. Then*

$$(3.6) \quad \lim_{\epsilon \rightarrow 0} f_{\chi\epsilon}(x_\epsilon) = f_\chi(x_{0^+}) - f_0 \int_{\nu h}^{\infty} \hat{\varphi}(\beta_0, s) ds,$$

where $h = \tilde{x} \cdot \alpha_0$, and $\hat{\varphi}$ is the CRT of φ .

By IK5, $\int_{\mathbb{R}} \hat{\varphi}(\beta_0, s) ds = 1$. The inversion formula (2.14) reconstructs jumps of f accurately, so $f_{\chi}(x_{0+}) - f_{\chi}(x_{0-}) = f_0$ and (3.8) can be written as follows

$$(3.7) \quad \lim_{\epsilon \rightarrow 0} f_{\chi\epsilon}(x_{\epsilon}) = f_{\chi}(x_{0-}) + f_0 \int_{-\infty}^{\nu h} \hat{\varphi}(\beta_0, s) ds.$$

By linearity, the proof of Theorem 1 can be split into two parts: local and global. The local part is formulated as follows.

Theorem 2. *Pick a locally generic pair $(x_0, y_0) \in \mathcal{C}$ such that \mathcal{S}_{y_0} is tangent to \mathcal{S} at $x_0 \in \mathcal{S}$. Suppose $\text{supp}(f)$ is contained in a sufficiently small neighborhood of x_0 . Then*

$$(3.8) \quad \lim_{\epsilon \rightarrow 0} f_{\chi\epsilon}(x_{\epsilon}) = f_{\chi}(x_{0+}) - f_0 \int_{\nu h}^{\infty} \hat{\varphi}(\beta_0, s) ds,$$

where $h = \tilde{x} \cdot \alpha_0$, and $\hat{\varphi}$ is the CRT of φ .

Since \mathcal{R} is an FIO with canonical relation (2.9), $g = \mathcal{R}f$ is singular only when \mathcal{S}_y is tangent to \mathcal{S} . Therefore, we are interested in the behavior of g in a neighborhood of y_0 .

Remark. *Strictly speaking, one has to distinguish between the original coordinates that describe points $x \in \mathcal{U}$, $y \in \mathcal{V}$ and those in (3.1), (3.4), respectively. For example, one should write $x = \hat{x}_1 \alpha_0 + \hat{x}^{\perp}$ instead of $x = x_1 \alpha_0 + x^{\perp}$. Such notation would emphasize that \hat{x}_1 is not the first component of x in the original coordinates, i.e. $\hat{x}_1 \neq x_1$. Similarly, a derivative like $\partial\Phi(x, y)/\partial x_1$, if written in full, becomes $\partial\Phi(x(\hat{x}_1, \hat{x}^{\perp}), y(\hat{y}_1, \hat{y}^{\perp}))/\partial \hat{x}_1$. To avoid burdensome notations, whenever the coordinates $(\hat{x}_1, \hat{x}^{\perp})$ and $(\hat{y}_1, \hat{y}^{\perp})$ are used, we will stick with the simplified notation and assume that the above convention holds.*

3.2. Behavior of g near its singular support. Because \mathcal{S} is smooth in a neighborhood of $x_0 \in \mathcal{S}$, there is a smooth local diffeomorphism $x \rightarrow (z, p)$ so that

$$(3.9) \quad x = z + pn(z), \quad z \in \mathcal{S}, \quad n(z) \text{ is normal to } \mathcal{S} \text{ at } z, \quad |n(z)| \equiv 1.$$

The normal $n(z)$ is chosen so that $N(z)$ is negative definite. Thus $n(x_0) = \alpha_0$. Clearly, we can extend the function $n(z)$, $z \in \mathcal{S}$, to $n(x)$ defined in a neighborhood of \mathcal{S} by the formula $n(z + pn(z)) := n(z)$. With a slight abuse of notation, the extended function will also be denoted $n(\cdot)$.

Using (3.9), define $\Psi(z + pn(z)) := p$. Then $\Psi(x) = 0$ is the equation of \mathcal{S} near x_0 , and Ψ is smooth. By construction, $\Psi(x) > 0$ on the interior side of \mathcal{S} .

Consider the system of equations

$$(3.10) \quad \Phi'_x(z + pn(z), y) - \lambda n(z) = 0, \quad \Phi(z + pn(z), y) = 0, \quad \Psi(z) = 0.$$

If we set $p = 0$ and solve (3.10) for y , we find submanifolds \mathcal{S}_y tangent to \mathcal{S} . We also need to solve these equations for z, p , and λ in terms of y .

Lemma 1. *Pick $(x_0, y_0) \in \mathcal{C}$ such that (1) \mathcal{S}_{y_0} is tangent to \mathcal{S} at $x_0 \in \mathcal{S}$, (2) \mathcal{S} is smooth at x_0 , and (3) $N(x_0)$ is negative definite. There exists an open set \mathcal{V}_1 , $y_0 \in \mathcal{V}_1 \subset \mathcal{V}$, such that*

- (1) *The set of $y \in \mathcal{V}_1$ such that \mathcal{S}_y is tangent to \mathcal{S} in a neighborhood of x_0 is a smooth submanifold of \mathcal{V} through y_0 . The vector $\Phi'_y(x_0, y_0)$ is normal to this submanifold at y_0 .*
- (2) *The solutions $z = Z(y)$, $p = P(y)$, and $\lambda = \Lambda(y)$ to (3.10) depend smoothly on $y \in \mathcal{V}_1$, and*

$$(3.11) \quad P'_y(y_0) = \frac{1}{|\Phi'_x|} \Phi'_y \neq 0;$$

- (3) Equations (3.10) determine a smooth function $y = \bar{Y}(z, p)$, $(z, p) \in \mathcal{S} \times \mathbb{R}$, in a neighborhood of $(x_0, 0)$.

Proof. Differentiate (3.10) with respect to z , p , λ , and y , and set $z = x_0$, $p = 0$, $y = y_0$ to obtain the 5×8 matrix

$$(3.12) \quad \begin{bmatrix} \Phi''_{xx} - \lambda n'_x & \Phi''_{xx} \alpha_0 & -\alpha_0 & \Phi''_{xy} \\ \Phi'_x & \Phi'_x \cdot \alpha_0 & 0 & \Phi'_y \\ \Psi'_x & 0 & 0 & 0 \end{bmatrix}.$$

Here n'_x is the derivative of the function $n(x)$ extended to a neighborhood of \mathcal{S} as described following (3.9). Since \mathcal{S}_{y_0} is tangent to \mathcal{S} at $x_0 \in \mathcal{S}$, we have $\Phi'_x \parallel \alpha_0$, so $\Phi'_x \cdot \alpha_0 = \lambda = -|\Phi'_x| \neq 0$. This also gives the value of λ to be used in (3.12).

To prove the first part of the first assertion we need to show that z , λ , and y_1 are smooth functions of y^\perp . Remove the columns corresponding to the derivatives with respect to p (because $p = 0$ is fixed) and y^\perp to obtain a 5×5 submatrix. Calculation in coordinates shows that $\Psi'_x(z + pn(z)) \equiv n(z)$ (cf. (3.9)). By applying elementary row and column operations, it is clear that this submatrix is full-rank if and only if the following matrix has rank two:

$$(3.13) \quad \Phi''_{x^\perp x^\perp} - \lambda(n^\perp)'_{x^\perp}.$$

Here $\Phi''_{x^\perp x^\perp} : \alpha_0^\perp \rightarrow \alpha_0^\perp$ is the appropriate submatrix of Φ''_{xx} in the coordinates (3.1), and n^\perp is the projection of n onto α_0^\perp . It is easy to see that

$$(3.14) \quad \Phi''_{x^\perp x^\perp} - \lambda(n^\perp)'_{x^\perp} = -\lambda N(x_0).$$

The desired assertion follows from Condition LG1 (see Definition 2).

Next, set $p = 0$ in (3.10) and assume that z, y_1 are functions of y^\perp . Differentiating the last two equations in (3.10) with respect to y^\perp and using that $\Phi'_y \cdot \beta_0 = |\Phi'_y| \neq 0$ gives $\Phi'_x z'_{y^\perp} = 0$ and $\partial y_1 / \partial y^\perp = 0$. This proves the second part of the first assertion.

The first part of the second assertion follows by retaining the columns corresponding to the derivatives with respect to z, p , and λ . As before, the resulting 5×5 submatrix is full-rank because the matrix in (3.13) has rank two. The second part of the second assertion follows by considering z, p , and λ as functions of y , differentiating the last two equations in (3.10) with respect to y , and using that $\Phi'_x, \Phi'_y \neq 0$.

The third assertion follows immediately from data completeness and the Bolker condition (Conditions DF2 and DF4, respectively). \square

We need the following lemma, which generalizes one of the results of Ramm and Zaslavsky [16, 17] from the CRT to the GRT.

Lemma 2. *Pick $(x_0, y_0) \in \mathcal{C}$ such that (1) \mathcal{S}_{y_0} is tangent to \mathcal{S} at $x_0 \in \mathcal{S}$, (2) \mathcal{S} is smooth at x_0 , and (3) $N(x_0)$ is negative definite. Suppose $\text{supp}(f)$ is contained in a small neighborhood of x_0 . For any $z \in \mathcal{S}$ and p in small neighborhoods of x_0 and 0, respectively, one has:*

$$(3.15) \quad g(\bar{Y}(z, p)) = p_+ G(z, p) + G_1(z, p) \text{ and } G(z, 0) = f_0(z) b(z, \bar{Y}(z, 0)) \frac{2\pi}{\sqrt{\det N(z)}},$$

for some smooth $G(z, p)$, $G_1(z, p)$.

Proof. Recall that $\bar{Y}(z, p)$ is the smooth function of $(z, p) \in \mathcal{S} \times \mathbb{R}$ defined by the conditions that $z + pm(z) \in \mathcal{S}_y$ and $n(z)$ be normal to \mathcal{S}_y at the point $z + pm(z)$, see Assertion (3) of Lemma 1. Recall also that in coordinates (3.1), $N(x_0)$ is negative definite. By linearity, we may assume that $f \equiv 0$ on the exterior side of \mathcal{S} . In

particular, $f(x_0 - \epsilon\alpha_0) \equiv 0$, $\epsilon > 0$, in (3.3). In this case we have to prove (3.15) with $G_1 \equiv 0$. By construction,

$$(3.16) \quad g(y) = \int_{\mathcal{S}_y} f(x)b(x, y)\theta(\Psi(x)) dx,$$

where θ is the unit step function (Heaviside function). Consider the system

$$(3.17) \quad \Psi'_x(x) - \mu\Phi'_x(x, y) = 0, \quad \Phi(x, y) = 0,$$

which we solve for x and μ in terms of y . Equations (3.17) determine the stationary point $x_*(y)$ of $\Psi(x)$ on the surface \mathcal{S}_y (the parameter μ , which corresponds to $1/\lambda$ in (3.10), is the Lagrange multiplier). From (3.10) and the property of Ψ'_x , the solution $x_*(y)$, $\mu(y)$ to (3.17) can be obtained from the solution $z = Z(y)$, $p = P(y)$, $\lambda = \Lambda(y)$ to (3.10): $x_*(y) = Z(y) + P(y)n(Z(y))$, $\mu(y) = 1/\Lambda(y)$. By Assertion (2) of Lemma 1, $x_*(y)$ is smooth near y_0 .

As is easily checked, when the matrix $\Psi''_{xx}(x_0) - \mu(y_0)\Phi''_{xx}(x_0, y_0)$ is viewed as a quadratic form on α_0^\perp , it coincides with $N(x_0)$. The latter is negative definite, hence $x_*(y)$ is the local maximum of $\Psi(x)$ on \mathcal{S}_y . By the Morse lemma, find local coordinates ω on \mathcal{S}_y , which depend smoothly on y , such that $\omega(x_*(y)) = 0$ and $\Psi(x) = \Psi(x_*(y)) - |\omega|^2$, $x = x(\omega; y) \in \mathcal{S}_y$. Since f, b , and $x(\omega; y)$ are all smooth, we get from (3.16)

$$(3.18) \quad g(y) = \int \theta(\Psi(x_*(y)) - |\omega|^2) F(\omega, y) d\omega$$

for some smooth F . Expand F in the Taylor series around $\omega = 0$ and integrate in spherical coordinates $\omega = r\Theta$. Integration with respect to Θ removes all the odd powers of r , i.e. only the even powers of r remain. By construction,

$$(3.19) \quad \Psi(x_*(y)) = \Psi(z + pn(z)) = p, \quad y = \bar{Y}(z, p),$$

and the first statement in (3.15) follows.

To prove the second statement, we use the local coordinates (3.1). In these coordinates, the local equation of \mathcal{S}_y becomes

$$(3.20) \quad x_1 = x_1(x^\perp, y) = \phi(y) + a(y) \cdot x^\perp + \frac{A(y)x^\perp \cdot x^\perp}{2} + O(|x^\perp|^3)$$

for some smooth ϕ , a , and A . Suppose y is such that $x_*(y) = x_0 + p\alpha_0$. Substitute such a pair $(x_*(y), y)$ into (3.17) and use (3.20) to conclude that $\phi(y) \equiv p$ and $a(y) \equiv 0$. Let $x_1 = Q(x^\perp)$ be the equation of \mathcal{S} in the coordinates (3.1). By construction, $Q'(0) = 0$. Clearly, $A(y_0)$ and $Q''(0)$ are the matrices of the second fundamental form of \mathcal{S}_{y_0} and \mathcal{S} , respectively, at x_0 in the coordinates (3.1). From (3.16),

$$(3.21) \quad g(y) = \int f(x)b(x, y)\theta(\psi(x^\perp, y)) \sqrt{1 + |Q'(x^\perp)|^2} dx^\perp,$$

$$\psi(x^\perp, y) := x_1(x^\perp, y) - Q(x^\perp), \quad x = (x_1(x^\perp, y), x^\perp).$$

Integrating in (3.21) by diagonalizing $N = \psi''_{x^\perp x^\perp}(0, y_0)$ and changing variables, we find

$$(3.22) \quad G(x_0, 0) = f(x_0)b(x_0, y_0) \frac{2\pi}{\sqrt{\det N}},$$

which finishes the proof. \square

4. LOCAL BEHAVIOR OF INTERPOLATED DATA.

Similarly to (3.10), consider the equations

$$(4.1) \quad \Phi'_x(x + t\alpha, y) - \lambda\alpha = 0, \quad \Phi(x + t\alpha, y) = 0,$$

which we solve to find $y = Y(\alpha, t; x)$ assuming (α, t, x) is in a neighborhood of $(\alpha_0, 0, x_0)$. Differentiating (4.1) with respect to λ and y and setting $x = x_0, t = 0, \alpha = \alpha_0, \lambda = -|\Phi'_x|$, we obtain similarly to (3.12) a matrix, which is non-degenerate. As opposed to (3.12), the key reason why it is non-degenerate is the Bolker condition. Hence $Y(\alpha, t; x)$ is a smooth function of α, t and x . Next we substitute $y = Y(\alpha, t; x)$ into (4.1) and obtain a few useful properties of Y . The first one is that $\partial Y_1 / \partial \alpha^\perp = 0$. Indeed, differentiate the second equation in (4.1) with respect to α^\perp and set $x = x_0, t = 0, \alpha = \alpha_0$ to obtain $\Phi'_y \partial Y / \partial \alpha^\perp = 0$. The desired assertion follows from (3.4). In a similar fashion, we have

$$(4.2) \quad \det(\partial Y^\perp / \partial \alpha^\perp) \neq 0, \quad \partial Y_1 / \partial t = |\Phi'_x| / |\Phi'_y| \neq 0, \quad \det(\partial Y / \partial (\alpha^\perp, t)) \neq 0.$$

The first result is obtained by differentiating the first equation in (4.1) with respect to α^\perp and using the Bolker condition and that $\partial Y_1 / \partial \alpha^\perp = 0$. The second result is obtained by differentiating the second equation in (4.1) with respect to t and using the properties of the selected coordinates (3.1), (3.4). The last result is an obvious consequence of the first two and that $\partial Y_1 / \partial \alpha^\perp = 0$.

In view of (4.2), given any small $\omega > 0$, we can find a sufficiently small open set $\mathcal{V}_1, y_0 \in \mathcal{V}_1 \subset \mathcal{V}$, such that $Y(\alpha, t; x_\epsilon) \in \mathcal{V}_1$ implies $|\alpha^\perp| < \omega$ for all $t = O(\epsilon)$ and x_ϵ provided that ϵ is sufficiently small. This relationship between \mathcal{V}_1 and ω is assumed in what follows.

Since $\text{supp}(\chi) \subset \mathcal{V}_1$, the integral with respect to α in (2.14) can be split into two sets:

$$(4.3) \quad \Omega_1 := \{\alpha \in S_+^2 : |\alpha^\perp| < A\epsilon^{1/2}\}, \quad \Omega_2 := \{\alpha \in S_+^2 : A\epsilon^{1/2} < |\alpha^\perp| < \omega\},$$

for some small (but fixed) $\omega > 0$. Here $A > 0$ is a large parameter. Let $f_{\chi_\epsilon}^{(j)}$ denote the result of integrating in (2.14) (with g replaced by g_ϵ) over $\Omega_j, j = 1, 2$. The behavior of $f_{\chi_\epsilon}^{(1)}$ is investigated first. This is done in Section 5. In the remainder of this section we lay the groundwork for that investigation by deriving the behavior of $Y(\alpha, t; x_\epsilon)$ and $g_\epsilon(y)$ in a neighborhood of $(\alpha_0, 0, x_0)$ and y_0 , respectively.

4.1. Local behavior of $Y(\alpha, t; x_\epsilon)$. The first step is to obtain the leading term behavior of the function $Y(\alpha, t; x_\epsilon)$ for $t = O(\epsilon)$ and $\alpha \in \Omega_1$, i.e. for $|\alpha^\perp| = O(\epsilon^{1/2})$. In this section we continue using the coordinates (3.1).

Expanding $y = Y(\alpha, t; x_\epsilon)$ in the Taylor series around $x = x_0, t = 0, \alpha = \alpha_0$ and using that $|x_\epsilon - x_0| = O(\epsilon), t = O(\epsilon), |\alpha^\perp| = O(\epsilon^{1/2})$, and $\partial Y_1 / \partial \alpha^\perp = 0$ gives

$$(4.4) \quad y_1 = O(\epsilon), \quad |y^\perp| = O(\epsilon^{1/2}).$$

For x in an $O(\epsilon)$ neighborhood of the origin (i.e., x_0) and for y in an $O(\epsilon^{1/2})$ neighborhood of the origin (i.e., y_0) we have

$$(4.5) \quad \begin{aligned} \Phi(x, y) &= \Phi'_x \cdot x + \Phi'_y \cdot y + \frac{\Phi''_{yy} y \cdot y}{2} + O(\epsilon^{3/2}), \\ \Phi'_x(x, y) &= \Phi'_x + \Phi''_{xy} y + O(\epsilon). \end{aligned}$$

To find $y = Y(\alpha, t; x_\epsilon)$, substitute $x = x_\epsilon$ into (4.5) and solve

$$(4.6) \quad \begin{aligned} \Phi'_x \cdot (\epsilon \tilde{x} + t\alpha) + \Phi'_y \cdot y + \frac{\Phi''_{yy} y \cdot y}{2} &= O(\epsilon^{3/2}), \\ \Phi'_x + \Phi''_{xy} y &= \lambda(1, \alpha^\perp) \pmod{O(\epsilon)}. \end{aligned}$$

Recall that $h = \tilde{x} \cdot \alpha_0$. Switching to the coordinates (3.1), (3.4), using (4.4), and keeping only the terms of order $O(\epsilon)$ in the first equation in (4.6) gives

$$(4.7) \quad -(\epsilon h + t)|\Phi'_x| + |\Phi'_y|y_1 + \frac{\Phi''_{y^\perp y^\perp} y^\perp \cdot y^\perp}{2} = O(\epsilon^{3/2}).$$

Projecting the second equation in (4.6) onto α_0 and onto α_0^\perp implies

$$(4.8) \quad \lambda = -|\Phi'_x| + O(\epsilon^{1/2}), \quad \Phi''_{x^\perp y^\perp} y^\perp = \lambda \alpha^\perp + O(\epsilon),$$

leading to

$$(4.9) \quad Y^\perp(\alpha, t; x_\epsilon) = -|\Phi'_x|(\Phi''_{x^\perp y^\perp})^{-1} \alpha^\perp + O(\epsilon).$$

By the Bolker condition (2.4), $\Phi''_{x^\perp y^\perp}$ is nondegenerate. Substitution into (4.7) now yields:

$$(4.10) \quad Y_1(\alpha, t; x_\epsilon) = \nu \left(\epsilon h + t - \frac{M \alpha^\perp \cdot \alpha^\perp}{2} \right) + O(\epsilon^{3/2}),$$

$$M := |\Phi'_x|(\Phi''_{x^\perp y^\perp})^{-T} \Phi''_{y^\perp y^\perp} (\Phi''_{x^\perp y^\perp})^{-1}, \quad M : \alpha_0^\perp \rightarrow \alpha_0^\perp.$$

Recall that ν is defined in (3.3).

4.2. The leading local behavior of $g(Y(\alpha, t; x_\epsilon))$. We plan to substitute $y = Y(\alpha, t = 0; x_\epsilon)$ into (2.13). Hence the second step is to find the leading behavior of $g(y)$ when $|y - Y(\alpha, 0; x_\epsilon)| = O(\epsilon)$ and $|\alpha^\perp| = O(\epsilon^{1/2})$. This is done by finding the asymptotics of $z = Z(y)$ and $p = P(y)$, which are determined by solving (3.10).

Recall that the local equation of \mathcal{S} in the coordinates (3.1) and the interior unit normal are given by

$$(4.11) \quad z_1 = \frac{Q z^\perp \cdot z^\perp}{2} + O(|z^\perp|^3), \quad n(z) = (1 + O(|z^\perp|^2), -Q z^\perp + O(|z^\perp|^2)).$$

By (4.9), (4.10), $|y_1| = O(\epsilon)$, $|y^\perp| = O(\epsilon^{1/2})$. From (3.11), $\partial p / \partial y^\perp = 0$ at $y = y_0$, so this implies $|p| = O(\epsilon)$, $x_1 = O(\epsilon)$, and $|z^\perp|, |x^\perp| = O(\epsilon^{1/2})$, where $x = z + pn(z)$. The equation (4.11) leads to:

$$(4.12) \quad x = \left(\frac{Q z^\perp \cdot z^\perp}{2} + O(\epsilon^{3/2}), z^\perp \right) + p(1 + O(\epsilon^{1/2}), -Q z^\perp + O(\epsilon))$$

$$= \left(p + \frac{Q z^\perp \cdot z^\perp}{2} + O(\epsilon^{3/2}), z^\perp + O(\epsilon^{3/2}) \right).$$

From (4.12), $x^\perp = z^\perp + O(\epsilon^{3/2})$.

Given that now $|x^\perp| = O(\epsilon^{1/2})$, the expansion in the first line in (4.5) should include additional terms. The second equation in (3.10) becomes

$$(4.13) \quad -|\Phi'_x|x_1 + |\Phi'_y|y_1 + \frac{\Phi''_{x^\perp x^\perp} x^\perp \cdot x^\perp}{2} + \Phi''_{x^\perp y^\perp} y^\perp \cdot x^\perp + \frac{\Phi''_{y^\perp y^\perp} y^\perp \cdot y^\perp}{2} = O(\epsilon^{3/2}).$$

Solving for x_1 we find:

$$(4.14) \quad x_1 = \frac{1}{\nu} y_1 + \frac{1}{|\Phi'_x|} \left\{ \frac{\Phi''_{x^\perp x^\perp} x^\perp \cdot x^\perp}{2} + \Phi''_{x^\perp y^\perp} y^\perp \cdot x^\perp + \frac{\Phi''_{y^\perp y^\perp} y^\perp \cdot y^\perp}{2} \right\} + O(\epsilon^{3/2}).$$

Using (4.9) and that $|y - Y(\alpha, 0; x_\epsilon)| = O(\epsilon)$, we have

$$(4.15) \quad y^\perp = -|\Phi'_x|(\Phi''_{x^\perp y^\perp})^{-1} \alpha^\perp + O(\epsilon),$$

and the unit normal vector to \mathcal{S}_Y is thus:

$$(4.16) \quad \left(1 + O(\epsilon^{1/2}), \alpha^\perp - \frac{\Phi''_{x^\perp x^\perp} x^\perp}{|\Phi'_x|} + O(\epsilon) \right).$$

The big- O terms in (4.16) follow by noticing that differentiation with respect to x_1 in (4.13), (4.14) converts $O(\epsilon^{3/2})$ into $O(\epsilon^{1/2})$, and differentiation with respect to x^\perp converts $O(\epsilon^{3/2})$ into $O(\epsilon)$. This follows by looking at the terms absorbed by $O(\epsilon^{3/2})$.

From (4.11) and (4.16), the two normal vectors are parallel (the first equation in (3.10)) if

$$(4.17) \quad \alpha^\perp - \frac{\Phi''_{x^\perp x^\perp} z^\perp}{|\Phi'_x|} + O(\epsilon) = -Qz^\perp + O(\epsilon),$$

which implies

$$(4.18) \quad z^\perp = N^{-1}\alpha^\perp + O(\epsilon), \quad x^\perp = N^{-1}\alpha^\perp + O(\epsilon).$$

Matching the first components in (4.12) and using (4.14), (4.15), (4.18) gives after simple transformations:

$$(4.19) \quad p = \frac{1}{\nu} \left(y_1 + \frac{M_1 \alpha^\perp \cdot \alpha^\perp}{2} \right) + O(\epsilon^{3/2}), \quad M_1 := \nu(M - N^{-1}),$$

where M is defined in (4.10). Summarizing (4.18) and (4.19) we have:

$$(4.20) \quad \begin{aligned} Z^\perp(y) &= N^{-1}\alpha^\perp + O(\epsilon), \quad P(y) = \frac{1}{\nu} \left(y_1 + \frac{M_1 \alpha^\perp \cdot \alpha^\perp}{2} \right) + O(\epsilon^{3/2}), \\ |y - Y(\alpha, 0; x_\epsilon)| &= O(\epsilon), \quad |\alpha^\perp| = O(\epsilon^{1/2}). \end{aligned}$$

Substituting into (3.15) we find

$$(4.21) \quad g(y) = f(Z(y))b(Z(y), y^*) \frac{2\pi}{\sqrt{\det N}} P_+(y) + O(P_+^2(y)),$$

where $y^* = \bar{Y}(z, p = 0)$ whenever $y = \bar{Y}(z, p)$.

From (4.11), (4.18) and (4.9), (4.10) it follows that $|Z(y) - x_0| = O(\epsilon^{1/2})$ and $|y - y_0| = O(\epsilon^{1/2})$ whenever $|y - Y(\alpha, 0; x_\epsilon)| = O(\epsilon)$, hence

$$(4.22) \quad f(Z(y))b(Z(y), y^*) = f(x_0)b(x_0, y_0) + O(\epsilon^{1/2}).$$

4.3. Local behavior of the interpolated data. In this subsection we find the behavior of the interpolated data near $y_0 = 0$. Combining (4.20)–(4.22) we get

$$(4.23) \quad \begin{aligned} g(\epsilon j) &= \frac{(f(x_0)b(x_0, y_0) + O(\epsilon^{1/2}))2\pi/\nu}{\sqrt{\det N}} \left(\epsilon j_1 + \frac{M_1 \alpha^\perp \cdot \alpha^\perp}{2} + O(\epsilon^{3/2}) \right)_+ \\ &\quad + O(P_+^2(\epsilon j)), \end{aligned}$$

where M_1 is defined in (4.19). From (4.10) and (4.20), $|P(\epsilon j)| = O(\epsilon)$. Using (4.9), (4.10) and (4.23) gives:

$$(4.24) \quad \begin{aligned} &g_\epsilon(Y(\alpha, t; x_\epsilon)) \\ &= \frac{(f(x_0)b(x_0, y_0) + O(\epsilon^{1/2}))2\pi\epsilon/\nu}{\sqrt{\det N}} \sum_{j \in r + \mathbb{Z}^3} \left(j_1 + \frac{M_1 \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} + O(\epsilon^{1/2}) \right)_+ \\ &\quad \times \varphi \left(\nu \left(h + \tilde{t} - \frac{M \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right) - j_1 + O(\epsilon^{1/2}), -\frac{|\Phi'_x|}{\epsilon^{1/2}} (\Phi''_{x^\perp y^\perp})^{-1} \tilde{\alpha}^\perp - j^\perp + O(1) \right) \\ &\quad + O(\epsilon^2), \quad \tilde{t} := t/\epsilon, \quad \tilde{\alpha}^\perp := \alpha^\perp/\epsilon^{1/2}. \end{aligned}$$

Since φ is compactly supported, the number of terms in the sum in (4.24) is bounded. Using additionally that $|\tilde{\alpha}^\perp| \leq A < \infty$, the sum itself is bounded as well.

Finally, combining with the fact that $|(a + O(\epsilon^{1/2}))_+ - a_+| = O(\epsilon^{1/2})$ uniformly in $a \in \mathbb{R}$ and using (4.9) and (4.10) gives

$$(4.25) \quad \begin{aligned} & g_\epsilon(Y(\alpha, t; x_\epsilon)) \\ &= \frac{f(x_0)b(x_0, y_0)2\pi\epsilon/\nu}{\sqrt{\det N}} \sum_{j \in r + \mathbb{Z}^3} \left(j_1 + \frac{M_1 \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right)_+ \\ & \times \varphi \left(\nu \left(h + \tilde{t} - \frac{M \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right) - j_1 + O(\epsilon^{1/2}), -\frac{|\Phi'_x|}{\epsilon^{1/2}} (\Phi''_{x^\perp y^\perp})^{-1} \tilde{\alpha}^\perp - j^\perp + O(1) \right) \\ & + O(\epsilon^{3/2}). \end{aligned}$$

In view of (4.25), denote

$$(4.26) \quad \psi(q, u) := \sum_{j \in r + \mathbb{Z}^3} (j \cdot \beta_0 + q)_+ \varphi(u - j), \quad q \in \mathbb{R}, \quad u \in \mathbb{R}^3.$$

Here we will need a higher order approximation of $Y(\alpha, t; x_\epsilon)$ than the one in (4.9), (4.10):

$$(4.27) \quad \begin{aligned} Y^\perp/\epsilon &= -|\Phi'_x| (\Phi''_{x^\perp y^\perp})^{-1} \tilde{\alpha}^\perp + A_1(\tilde{\alpha}^\perp, \tilde{\alpha}^\perp) + A_2 \tilde{t} + A_3 h + O(\epsilon^{1/2}), \\ Y_1/\epsilon &= \nu \left(h + \tilde{t} - \frac{M \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right) + O(\epsilon^{1/2}), \end{aligned}$$

where A_1 is a bilinear map $\mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, and $A_2, A_3 \in \mathbb{R}^2$. Moreover, the two $O(\epsilon^{1/2})$ terms in (4.27) depend smoothly on \tilde{t} and $\tilde{\alpha}^\perp$. In particular, differentiation with respect to \tilde{t} does not change the order of these terms as $\epsilon \rightarrow 0$. Hence we rewrite (4.25) as follows

$$(4.28) \quad \begin{aligned} g_\epsilon(Y(\alpha, t; x_\epsilon)) &= \frac{f_0 b_0 2\pi\epsilon/\nu}{\sqrt{\det N}} \psi(q(\tilde{\alpha}^\perp), u(\tilde{\alpha}^\perp, \tilde{t})) + O(\epsilon^{3/2}), \quad q(\tilde{\alpha}^\perp) := \frac{M_1 \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2}, \\ u(\tilde{\alpha}^\perp, \tilde{t}) &:= \left(\nu \left(h + \tilde{t} - \frac{M \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right) + O(\epsilon^{1/2}), \right. \\ & \quad \left. -\frac{|\Phi'_x|}{\epsilon^{1/2}} (\Phi''_{x^\perp y^\perp})^{-1} \tilde{\alpha}^\perp + A_1(\tilde{\alpha}^\perp, \tilde{\alpha}^\perp) + A_2 \tilde{t} + A_3 h + O(\epsilon^{1/2}) \right). \end{aligned}$$

We also have

$$(4.29) \quad \begin{aligned} \partial u(\tilde{\alpha}^\perp, \tilde{t}) / \partial \tilde{t} \Big|_{\tilde{t}=0} &= \nu \left(1 + O(\epsilon^{1/2}), A_2/\nu + O(\epsilon^{1/2}) \right), \\ \partial^2 u(\tilde{\alpha}^\perp, \tilde{t}) / \partial \tilde{t}^2 \Big|_{\tilde{t}=0} &= O(\epsilon^{1/2}). \end{aligned}$$

5. ESTIMATING THE TERM $f_{\chi\epsilon}^{(1)}$.

To study $f_{\chi\epsilon}^{(1)}$ we need the following lemma, which follows immediately from (4.26) and the properties IK1–IK3 of φ .

Lemma 3. *Partial derivatives of $\psi(q, u)$ with respect to u up to the order two are continuous. Also, one has*

$$(5.1) \quad \psi(q, u) = \psi(q + m \cdot \beta_0, u - m), \quad \forall m \in \mathbb{Z}^3,$$

and, for some $c > 0$,

$$(5.2) \quad \psi(q, u) = 0 \text{ if } u \cdot \beta_0 + q < -c; \quad \psi(q, u) = u \cdot \beta_0 + q \text{ if } u \cdot \beta_0 + q > c.$$

Denote

$$(5.3) \quad U(q, u) := - \left. \frac{\partial^2}{\partial \tau^2} \psi(q, u + \tau(1, A_2/\nu)) \right|_{s=0}.$$

The following lemma is a direct consequence of Lemma 3 (see also properties IK3, IK4 of φ).

Lemma 4. *The function $U(q, u)$ has piecewise continuous bounded first order partial derivatives with respect to q and u . Also,*

$$(5.4) \quad U(q, u) = U(q + m \cdot \beta_0, u - m), \quad \forall m \in \mathbb{Z}^3,$$

and, for some $c > 0$,

$$(5.5) \quad U(q, u) \equiv 0 \text{ if } |u \cdot \beta_0 + q| > c.$$

Note that the derivative in the inversion formula (2.14) is with respect to t . Using that $t = \epsilon \tilde{t}$ (cf. (4.24)) and taking into account (4.29), (5.3), in the formula below we will acquire the factor $(\epsilon/\nu)^2$. Thus, in terms of U , the expression for $f_{\chi\epsilon}^{(1)}$ becomes after changing variables $\alpha^\perp \rightarrow \tilde{\alpha}^\perp$ (this brings the factor ϵ), setting $\tilde{t} = 0$, and using (4.28), (4.29), (5.3):

$$(5.6) \quad \begin{aligned} f_{\chi\epsilon}^{(1)}(x_\epsilon) &= \frac{1}{4\pi^2} \frac{f_0(2\pi\epsilon/\nu)}{\sqrt{\det N}} \frac{\epsilon}{(\epsilon/\nu)^2} \int_{|\tilde{\alpha}^\perp| < A} U(q(\tilde{\alpha}^\perp), u_0(\tilde{\alpha}^\perp)) d\tilde{\alpha}^\perp + O(\epsilon^{1/2}), \\ u_0(\tilde{\alpha}^\perp) &:= \left(\nu \left(h - \frac{M\tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} \right), -\frac{|\Phi'_x|}{\epsilon^{1/2}} (\Phi''_{x^\perp y^\perp})^{-1} \tilde{\alpha}^\perp + A_1(\tilde{\alpha}^\perp, \tilde{\alpha}^\perp) + A_3 h \right). \end{aligned}$$

Two simplifications have been made in deriving (5.6). First, using that first and second order derivatives of ψ are bounded, it follows from (4.29) that

$$(5.7) \quad \left. \frac{\partial^2}{\partial \tilde{t}^2} \psi(\cdot, u(\tilde{\alpha}^\perp, \tilde{t})) \right|_{\tilde{t}=0} = \nu^2 \left. \frac{\partial^2}{\partial \tau^2} \psi(\cdot, u(\tilde{\alpha}^\perp, 0) + \tau(1, A_2/\nu)) \right|_{\tau=0} + O(\epsilon^{1/2}).$$

Second, since the derivatives of U are bounded, the integral with respect to $\tilde{\alpha}^\perp$ is over a bounded set, and the coefficient in front of the integral is bounded, using (5.7) and then replacing $u(\tilde{\alpha}^\perp, 0)$ with $u_0(\tilde{\alpha}^\perp)$ in the arguments of U leads to the term $O(\epsilon^{1/2})$ outside the integral.

Approximate the domain $|\tilde{\alpha}^\perp| < A$ by a union of non-overlapping small squares of size δ . Let these squares be denoted B_k , $k = 1, 2, \dots, O(\delta^{-2})$. By (5.6),

$$(5.8) \quad \begin{aligned} u_0(\tilde{\alpha}^\perp) &= u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp) + O(\delta), \quad \tilde{\alpha}^\perp \in B_k, \\ \Delta u_k(\tilde{\alpha}^\perp) &:= \left(0, -|\Phi'_x| (\Phi''_{x^\perp y^\perp})^{-1} ((\tilde{\alpha}^\perp - \tilde{\alpha}_k^\perp)/\epsilon^{1/2}) \right) \in \beta_0^\perp, \end{aligned}$$

where $\tilde{\alpha}_k^\perp$ is the center of B_k . By Lemma 4,

$$(5.9) \quad U(q(\tilde{\alpha}^\perp), u_0(\tilde{\alpha}^\perp)) = U(q(\tilde{\alpha}_k^\perp), u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp)) + O(\delta), \quad \tilde{\alpha}^\perp \in B_k.$$

Therefore

$$\begin{aligned}
(5.10) \quad & \int_{B_k} U(q(\tilde{\alpha}^\perp), u_0(\tilde{\alpha}^\perp)) d\tilde{\alpha}^\perp \\
&= \int_{B_k} [U(q(\tilde{\alpha}_k^\perp), u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp)) + O(\delta)] d\tilde{\alpha}^\perp \\
&= \int_{B_k} [U(q(\tilde{\alpha}_k^\perp) + \lfloor u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp) \rfloor \cdot \beta_0, \{u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp)\}) + O(\delta)] d\tilde{\alpha}^\perp \\
&= \int_{B_k} [U(q(\tilde{\alpha}_k^\perp) + u_0(\tilde{\alpha}_k^\perp) \cdot \beta_0 - \{u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp)\} \cdot \beta_0, \\
&\quad \{u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp)\}) + O(\delta)] d\tilde{\alpha}^\perp,
\end{aligned}$$

where we have used that $\Delta u_k(\tilde{\alpha}^\perp) \cdot \beta_0 = 0$. In (5.10) and below the fractional part of a vector is computed component wise: $\{u\} = (\{u_1\}, \{u_2\}, \{u_3\})$, where $\{u_i\} = u_i - \lfloor u_i \rfloor$ and $\lfloor u_i \rfloor$ is the largest integer not exceeding u_i .

Pick any $m \in \mathbb{Z}^3$, $m \neq 0$. Let m^\perp be the projection of m onto the plane β_0^\perp . Condition LG2 in Definition 2 implies that $m^\perp \neq 0$. By the local Bolker condition DF4, $\Phi''_{x^\perp y^\perp}$ is non-degenerate, so the vector $(\Phi''_{x^\perp y^\perp})^{-T} m^\perp \in \mathbb{R}^2$ is not zero. Using (5.8), the standard Weyl-type argument (cf. [13]) implies that

$$\begin{aligned}
(5.11) \quad & \lim_{\epsilon \rightarrow 0} \int_{B_k} U(q(\tilde{\alpha}^\perp), u_0(\tilde{\alpha}^\perp)) d\tilde{\alpha}^\perp \\
&= \left(\int_{[0,1]^3} U(q(\tilde{\alpha}_k^\perp) + u_0(\tilde{\alpha}_k^\perp) \cdot \beta_0 - \omega \cdot \beta_0, \omega) d\omega + O(\delta) \right) \text{Vol}(B_k).
\end{aligned}$$

Indeed, consider the function $U_1(q, \omega) := U(q - \omega \cdot \beta_0, \omega)$. Clearly, $U_1(q, \omega)$ is periodic: $U_1(q, \omega) = U_1(q, \omega + m)$, $m \in \mathbb{Z}^3$. Expand $U_1(q, \omega)$ in a Fourier series:

$$(5.12) \quad U_1(q, \omega) = \sum_{m \in \mathbb{Z}^3} A_m(q) \exp(2\pi i m \cdot \omega).$$

By the argument preceding (5.11),

$$\begin{aligned}
(5.13) \quad & m \cdot \left(0, -|\Phi'_x|(\Phi''_{x^\perp y^\perp})^{-1}(\tilde{\alpha}^\perp - \tilde{\alpha}_k^\perp) \right) = -\frac{1}{|\Phi'_x|} \left[(\Phi''_{x^\perp y^\perp})^{-T} m^\perp \right] \cdot (\tilde{\alpha}^\perp - \tilde{\alpha}_k^\perp), \\
& (\Phi''_{x^\perp y^\perp})^{-T} m^\perp \neq 0, |m| > 0.
\end{aligned}$$

Therefore,

$$\begin{aligned}
(5.14) \quad & \lim_{\epsilon \rightarrow 0} \left| \int_{B_k} \exp(2\pi i m \cdot (u_0(\tilde{\alpha}_k^\perp) + \Delta u_k(\tilde{\alpha}^\perp))) d\tilde{\alpha}^\perp \right| \\
&= \lim_{\epsilon \rightarrow 0} \left| \int_{B_k} \exp\left(-\frac{2\pi i}{|\Phi'_x| \epsilon} \left[(\Phi''_{x^\perp y^\perp})^{-T} m^\perp \right] \cdot \tilde{\alpha}^\perp \right) d\tilde{\alpha}^\perp \right| = 0, |m| > 0.
\end{aligned}$$

In other words, only the term corresponding to $m = 0$ survives, and the desired assertion follows from the standard approximation argument.

By (4.23), (4.28) and (5.6),

$$(5.15) \quad q(\tilde{\alpha}_k^\perp) + u_0(\tilde{\alpha}_k^\perp) \cdot \beta_0 = \nu h - \frac{\nu N^{-1} \tilde{\alpha}_k^\perp \cdot \tilde{\alpha}_k^\perp}{2}.$$

Add the integrals over all the squares B_k and use (5.6), (5.11), and (5.15):

$$\begin{aligned}
(5.16) \quad & \lim_{\epsilon \rightarrow 0} f_{\chi^\epsilon}^{(1)}(x_\epsilon) \\
&= \kappa \sum_k \int_{B_k} U(q(\tilde{\alpha}^\perp), u_0(\tilde{\alpha}^\perp)) d\tilde{\alpha}^\perp + O(\delta) \\
&= \kappa \sum_k \left(\int_{[0,1]^3} U\left(\nu h - \frac{\nu N^{-1} \tilde{\alpha}_k^\perp \cdot \tilde{\alpha}_k^\perp}{2} - \omega \cdot \beta_0, \omega\right) d\omega + O(\delta) \right) \text{Vol}(B_k) + O(\delta), \\
\kappa &:= \frac{\nu}{2\pi} \frac{f_0}{\sqrt{\det N}}.
\end{aligned}$$

Since $\delta > 0$ can be as small as we like, (5.16) implies

$$(5.17) \quad \lim_{\epsilon \rightarrow 0} f_{\chi^\epsilon}^{(1)}(x_\epsilon) = \kappa \int_{|\tilde{\alpha}^\perp| < A} \int_{[0,1]^3} U\left(\nu h - \frac{\nu N^{-1} \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} - \omega \cdot \beta_0, \omega\right) d\omega d\tilde{\alpha}^\perp.$$

The second argument of U is bounded and N is negative definite, so by (5.5) the integral with respect to $\tilde{\alpha}^\perp$ over the set $|\tilde{\alpha}^\perp| < A$, when $A > 0$ is large enough (but fixed), can be replaced by the integral over all \mathbb{R}^2 . Changing variables and integrating in spherical coordinates gives

$$\begin{aligned}
(5.18) \quad \lim_{\epsilon \rightarrow 0} f_{\chi^\epsilon}^{(1)}(x_\epsilon) &= \kappa \int_{\mathbb{R}^2} \int_{[0,1]^3} U\left(\nu h - \frac{\nu N^{-1} \tilde{\alpha}^\perp \cdot \tilde{\alpha}^\perp}{2} - \omega \cdot \beta_0, \omega\right) d\omega d\tilde{\alpha}^\perp \\
&= \kappa \frac{2\sqrt{\det N}}{\nu} \int_{\mathbb{R}^2} \int_{[0,1]^3} U(\nu h + v \cdot v - \omega \cdot \beta_0, \omega) d\omega dv \\
&= f_0 \int_0^\infty \int_{[0,1]^3} U(\nu h + \tau - \omega \cdot \beta_0, \omega) d\omega d\tau.
\end{aligned}$$

The integral with respect to ω can be evaluated explicitly. Recall that in our coordinates, $\beta_0 = (1, 0, 0)$ (cf. (3.4)). From (4.26) and (5.3),

$$\begin{aligned}
(5.19) \quad & - \int_{[0,1]^3} U(q - \omega \cdot \beta_0, \omega) d\omega \\
&= \int_{[0,1]^3} \frac{\partial^2}{\partial \tau^2} \psi(q - \omega \cdot \beta_0, \tau(1, A_2/\nu) + \omega) \Big|_{\tau=0} d\omega \\
&= \frac{\partial^2}{\partial \tau^2} \int_{[0,1]^3} \sum_{j \in \tau + \mathbb{Z}^3} (q - \beta_0 \cdot (\omega - j))_+ \varphi(\tau(1, A_2/\nu) + \omega - j) d\omega \Big|_{\tau=0} \\
&= \frac{\partial^2}{\partial \tau^2} \int_{\mathbb{R}^3} (q - \beta_0 \cdot \omega)_+ \varphi(\tau(1, A_2/\nu) + \omega) d\omega \Big|_{\tau=0} \\
&= \frac{\partial^2}{\partial \tau^2} \int_{\mathbb{R}^3} (q - \beta_0 \cdot \omega)_+ \varphi(\tau(1, 0) + \omega) d\omega \Big|_{\tau=0} = \hat{\varphi}(\beta_0, q).
\end{aligned}$$

Substitute (5.19) into (5.18)

$$(5.20) \quad \lim_{\epsilon \rightarrow 0} f_{\chi^\epsilon}^{(1)}(x_\epsilon) = -f_0 \int_0^\infty \hat{\varphi}(\beta_0, \nu h + \tau) d\tau = f_0 \left(- \int_{\nu h}^\infty \hat{\varphi}(\beta_0, \tau) d\tau \right).$$

Since φ is normalized and compactly supported,

$$(5.21) \quad - \int_{\nu h}^\infty \hat{\varphi}(\beta_0, \tau) d\tau \rightarrow \begin{cases} 0, & h > c, \\ -1, & h < -c. \end{cases}$$

for some $c > 0$.

Using the definition (3.5) and some simple transformations, we can rewrite the integral in (5.20) in two different forms

$$(5.22) \quad \int_{\nu h}^{\infty} \hat{\varphi}(\beta_0, \tau) d\tau = \int_{|\Phi'_x|/h}^{\infty} \hat{\varphi}(\Phi'_y, \tau) d\tau = \int_{\Phi'_x \tilde{x} + \Phi'_y \tilde{y} > 0} \varphi(\tilde{y}) d\tilde{y}.$$

6. ANALYSIS OF THE TERM $f_{\chi^\epsilon}^{(2)}$.

Lemma 5. *One can find $\omega > 0$, $\epsilon_0 > 0$ small enough and $A > 0$ large enough so that $g(y)$ is smooth in a neighborhood of all y such that $(y - Y(\alpha, 0; x_\epsilon))/\epsilon \in \text{supp}(\varphi)$ for any $\alpha \in \Omega_2$, $\tilde{x} \in \tilde{U}$ (cf. (3.2)), and $0 < \epsilon < \epsilon_0$.*

Proof. Fix some $c > 0$ sufficiently large. Using that $N(x_0)$ is negative definite, equation (4.20) implies that we can find $A > 0$ large enough and $\omega > 0$ small enough so that $P(Y(\alpha, 0; x_\epsilon)) > c\epsilon$ for all $\alpha \in \Omega_2$ and all $\tilde{x} \in \tilde{U}$ provided that ϵ is small enough. Since φ is compactly supported and $P(y)$ is smooth, $P(y) > 0$ for all y such that $(y - Y(\alpha, 0; x_\epsilon))/\epsilon \in \text{supp}(\varphi)$ (this is where we use that $c > 0$ is sufficiently large). Therefore, g is a smooth function in a neighborhood of all such y because in this case we can drop the subscript '+' from $P_+(y)$ in (3.15). \square

Lemma 5 implies that the inversion formula (2.14) and its discrete analogue do not see singularities in the data when $\alpha \in \Omega_2$ provided that ω, A are selected as in the proof of Lemma 5. Since $|\Omega_1| = O(\epsilon)$, clearly the limit $\lim_{\epsilon \rightarrow 0^+} f_{\chi^\epsilon}^{(2)}(x_\epsilon)$ exists and is independent of $\tilde{x} \in \tilde{U}$.

Using again that $N(x_0)$ is negative definite, there exist sufficiently small neighborhoods \mathcal{U}_1 of x_0 and \mathcal{V}_1 of y_0 such that $\mathcal{S}_y \cap \mathcal{U}_1$ is on the exterior side of \mathcal{S} whenever \mathcal{S}_y is tangent to \mathcal{S} and $y \in \mathcal{V}_1$. This implies that if $x \in \mathcal{U}_1$ and x is on the interior side of \mathcal{S} , then there is no $y \in \mathcal{V}_1$ such that \mathcal{S}_y contains x and is tangent to \mathcal{S} . In turn, this implies that the data $g(y)$, $y = Y(\alpha, t = 0; x) \in \mathcal{V}_1$, which is used to compute $f_\chi(x)$ is also smooth. Then, clearly, $\lim_{\epsilon \rightarrow 0^+} (f_{\chi^\epsilon}^{(2)}(x_0 + \epsilon h \alpha_0) - f_\chi(x_0 + \epsilon h \alpha_0)) = 0$ for any $h > 0$, and

$$(6.1) \quad \lim_{\epsilon \rightarrow 0^+} f_{\chi^\epsilon}^{(2)}(x_\epsilon) = f_\chi(x_{0+}).$$

This concludes the proof of the first part of the theorem.

7. CONTRIBUTION OF REMOTE SINGULARITIES

Suppose \mathcal{S}_{y_0} is tangent to \mathcal{S} at some $z_0 \in \mathcal{S}_{y_0}$, $z_0 \neq x_0$, \mathcal{S} is smooth at z_0 , and $N(z_0)$ is either positive definite or negative definite. Set $\alpha_0 = \Phi'_x(z_0, y_0)/|\Phi'_x(z_0, y_0)|$, so that $y_0 = Y(\alpha_0, 0; x_0)$. As before, \mathcal{V}_1 is a small neighborhood of y_0 , and $\text{supp}(\chi) \subset \mathcal{V}_1$.

As follows from assertion (1) of Lemma 1 (with x_0 replaced by z_0 as the point of tangency), the set of $y \in \mathcal{V}_1$ such that \mathcal{S}_y is tangent to \mathcal{S} near z_0 is a smooth submanifold of \mathcal{V}_1 through y_0 , and the vector $\Phi'_y(z_0, y_0)$ is normal to it at y_0 . The local equation of the manifold is $P(y) = 0$, where the function P is the same as in Lemma 1. By assertion (2) of the lemma, $P'_y(y_0) \neq 0$.

Additionally, \mathcal{T}_{x_0} is another smooth submanifold through y_0 , and $\Phi'_y(x_0, y_0)$ is normal to it at y_0 . By the assumption DF3 of no conjugate points, $\Phi'_y(x_0, y_0)$ and $\Phi'_y(z_0, y_0)$ are not parallel, so the intersection of the two submanifolds is a smooth curve Γ_{x_0} through y_0 . This is the same curve Γ_{x_0} , which was introduced in (2.11). From this argument it is easy to see that Γ_x depends smoothly on x near $x = x_0$.

Theorem 3. *Pick a globally generic pair $(x_0, y_0) \in \mathcal{C}$ such that $x_0 \notin \mathcal{S}$ and \mathcal{S}_{y_0} is tangent to \mathcal{S} at $z_0 \in \mathcal{S}$. Suppose $\text{supp}(f)$ is contained in a sufficiently small neighborhood of z_0 . One has*

$$(7.1) \quad \lim_{\epsilon \rightarrow 0} f_{\chi\epsilon}(x_\epsilon) = f_\chi(x_0).$$

Proof. Suppose first that the reconstruction point is x_0 . Since the reconstruction point is fixed, the dependence of various quantities on x_0 is omitted from notations in most places when there is no risk of confusion. Then

$$(7.2) \quad \begin{aligned} f_{\chi\epsilon}(x_0) &= \int_{S_+^2} B(\alpha) \sum_j g(\epsilon j) \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha, \\ B(\alpha) &:= - \frac{\chi(Y(\alpha, 0))}{4\pi^2} \frac{1}{b(x_0, Y(\alpha, 0))}. \end{aligned}$$

By Lemma 2,

$$(7.3) \quad g(y) := P_+(y)G(y), \quad G(y) := G(Z(y), P(y)),$$

Since φ is compactly supported, we can expand the factor $P(y)$ in (7.3) in the Taylor series centered at $Y(\alpha, 0)$. Let $L(y)$ be its linear term:

$$(7.4) \quad L(y) := P(Y(\alpha, 0)) + P'_y(Y(\alpha, 0)) \cdot (y - Y(\alpha, 0)).$$

We begin by looking at the expression, which is obtained by ignoring the second and higher order terms in the expansion of P :

$$(7.5) \quad J_\epsilon^{(1)} := \int_{S_+^2} B(\alpha) \sum_j G(\epsilon j) L_+(\epsilon j) \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha.$$

Clearly,

$$(7.6) \quad \begin{aligned} &\left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha \\ &= \frac{1}{\epsilon^2} \nabla_{u(\alpha)}^2 \varphi \left(\frac{Y(\alpha, 0)}{\epsilon} - j \right) + \frac{1}{\epsilon} \varphi'_y \left(\frac{Y(\alpha, 0)}{\epsilon} - j \right) \cdot Y''_{tt}(\alpha, 0), \\ u(\alpha) &:= Y'_t(\alpha, 0), \quad \nabla_u^2 \varphi(y) = \left(\frac{\partial}{\partial t} \right)^2 \varphi(y + tu) \Big|_{t=0}. \end{aligned}$$

Consider the most singular part of $J_\epsilon^{(1)}$, which is obtained by using the first term on the right in (7.6) and replacing $G(\epsilon j)$ with $G(Y(\alpha, 0))$:

$$(7.7) \quad \begin{aligned} J_\epsilon^{(1a)} &:= \int_{S_+^2} B_1(\alpha) \sum_j L_+(\epsilon j) \frac{1}{\epsilon^2} \nabla_{u(\alpha)}^2 \varphi \left(\frac{Y(\alpha, 0)}{\epsilon} - j \right) d\alpha, \\ B_1(\alpha) &:= B(\alpha) G(Y(\alpha, 0)). \end{aligned}$$

In view of (7.4) and (7.7), similarly to (4.26) and (5.3), introduce the function

$$(7.8) \quad \psi(q, v; \alpha) := \sum_j (e(\alpha) \cdot (j - v) + q)_+ \nabla_{u(\alpha)}^2 \varphi(v - j), \quad e(\alpha) := P'_y(Y(\alpha, 0)).$$

Clearly,

- (1) ψ is compactly supported in q (by (2.12)),
- (2) ψ has bounded first order partial derivatives, and
- (3) $\psi(q, v; \alpha) = \psi(q, v - m; \alpha)$ for any $m \in \mathbb{Z}^3$.

Using (7.8) in (7.7) yields:

$$(7.9) \quad J_\epsilon^{(1a)} = \frac{1}{\epsilon} \int_{S_+^2} B_1(\alpha) \psi \left(\frac{P(Y(\alpha, 0))}{\epsilon}, \frac{Y(\alpha, 0)}{\epsilon}; \alpha \right) d\alpha.$$

Introduce local coordinates $s = (s_1, s_2)$ on S_+^2 so that $s_1 \equiv P(Y(\alpha, 0))$ in a neighborhood of α_0 . As is shown at the beginning of this section, Γ is the transverse intersection of the submanifolds \mathcal{T}_{x_0} and $\{y \in \text{supp}(\chi) : P(y) = 0\}$. By the first equation in (4.2), $\alpha \rightarrow Y(\alpha, 0) \in \mathcal{T}_{x_0}$ is a regular parametrization near α_0 . In (4.2), $y_0 = Y(\alpha_0, 0; x_0)$, and Y^\perp is determined by the projection onto the plane $(\Phi'_y(x_0, y_0))^\perp$ (as opposed to $(\Phi'_y(z_0, y_0))^\perp$). Hence $P'_y(y_0) \neq 0$ (cf. (3.11)) implies $\partial P(Y(\alpha, 0))/\partial \alpha^\perp \neq 0$ near α_0 . Therefore the preimage of $\Gamma \cap \mathcal{V}_1$, given by $\{\alpha \in S_+^2 : P(Y(\alpha, 0)) = 0, Y(\alpha, 0) \in \mathcal{V}_1\}$ is also a smooth curve, and local coordinates (s_1, s_2) with the required property do exist. Then

$$(7.10) \quad \begin{aligned} J_\epsilon^{(1a)} &= \frac{1}{\epsilon} \int_{\mathbb{R}^2} B_1(\alpha(s)) \psi \left(\frac{P(Y(\alpha(s), 0))}{\epsilon}, \frac{Y(\alpha(s), 0)}{\epsilon}; \alpha(s) \right) \left| \frac{\partial \alpha}{\partial s} \right| ds + O(\epsilon) \\ &= \int_{\mathbb{R}^2} B_2(s_2) \psi \left(\tilde{s}_1, \frac{Y(\alpha(0, s_2), 0)}{\epsilon} + \frac{\partial Y(\alpha(s_1, s_2), 0)}{\partial s_1} \Big|_{s_1=0} \tilde{s}_1; \alpha(0, s_2) \right) d\tilde{s}_1 ds_2 \\ &\quad + O(\epsilon), \\ B_2(s_2) &:= B_1(\alpha(0, s_2)) \left| \frac{\partial \alpha(s_1 = 0, s_2)}{\partial s} \right|, \quad \tilde{s}_1 = s_1/\epsilon. \end{aligned}$$

In the first line, the integral is over the bounded set $\{s \in \mathbb{R}^2 : Y(\alpha(s), 0) \in \text{supp}(\chi)\}$. In the second line, the integral can be confined to a bounded set $\{(\tilde{s}_1, s_2) \in \mathbb{R}^2 : |\tilde{s}_1| < \tilde{A}, Y(\alpha(0, s_2)) \in \text{supp}(\chi)\}$ for some $\tilde{A} > 0$ large enough.

Lemma 6. *Let D be a rectangle $D := [a_1, b_1] \times [a_2, b_2]$. Consider a function $\psi \in C(D \times \mathbb{R}^3)$. Suppose ψ is periodic: $\psi(s, y) = \psi(s, y + m)$ for any $m \in \mathbb{Z}^3$ and $(s, y) \in D \times \mathbb{R}^3$. Let $Y : [a_2, b_2] \rightarrow \mathbb{R}^3$ be a C^1 function with the following property. For any $m \in \mathbb{Z}^3$, $|m| > 0$,*

- (1) *The set $\{s_2 \in [a_2, b_2] : |m \cdot Y'(s_2)| \leq \delta\}$ is contained in a finite number of intervals for all $\delta > 0$ sufficiently small (this number may depend on m and δ), and*
- (2) *The sum of the lengths of these intervals goes to zero as $\delta \rightarrow 0$.*

Then one has

$$(7.11) \quad \lim_{\epsilon \rightarrow 0^+} \int_D \psi \left(s, \frac{Y(s_2)}{\epsilon} \right) ds = \int_D \int_{[0,1]^3} \psi(s, y) dy ds.$$

Proof. Pick any $\delta_1 > 0$. Let $\Psi_m(s)$ denote the coefficients of the Fourier expansion of $\psi(s, y)$ with respect to y . We can find $M > 0$ large enough and a partition of D into sufficiently small rectangles such that

$$(7.12) \quad \sup_{s, y} \left| \psi(s, y) - \sum_{|m| \leq M} \tilde{\Psi}_m(s) \exp(2\pi i m \cdot y) \right| \leq \delta_1.$$

Here $\tilde{\Psi}_m$ is an approximation of Ψ_m , which is constant on each rectangle of the partition. Thus, the lemma will be proven if we show that

$$(7.13) \quad \int_a^b \exp(2\pi i m \cdot Y(s_2)/\epsilon) ds_2 \rightarrow 0, \quad \epsilon \rightarrow 0, \quad \text{for any } [a, b] \subset [a_2, b_2], \quad m \in \mathbb{Z}^3, |m| > 0.$$

Using assumptions (1) and (2) of the lemma, partition $[a, b]$ into a finite collection of non-overlapping intervals so that (i) their union is as close to $[a, b]$ as we like,

and (ii) in each of these intervals $m \cdot Y'(s_2)$ is bounded away from zero. The result now follows immediately. \square

Clearly, Condition GG2 in Definition 1 is independent of the choice of the vector field $\dot{\Gamma}$ as long as it does not vanish at any point of Γ . In the s -coordinates, $Y(\alpha(0, s_2), 0)$ is a regular parametrization of $\Gamma \cap \mathcal{V}_1$ because $|\partial Y^\perp / \partial \alpha^\perp|_{\alpha=\alpha_0} \neq 0$ and $|\partial \alpha^\perp / \partial s| \neq 0$ (see also the argument following (7.9)). The latter determinant is computed at s such that $\alpha(s) = \alpha_0$. Therefore $\partial Y / \partial s_2$ never vanishes on $\Gamma \cap \mathcal{V}_1$. Condition GG2 implies that $Y(\alpha(0, s_2))$ satisfies Conditions (1), (2) in Lemma 6. Set

$$(7.14) \quad \psi_1(s, y) := B_2(s_2) \psi \left(\tilde{s}_1, y + \frac{\partial Y(\alpha(s_1, s_2), 0)}{\partial s_1} \Big|_{s_1=0} \tilde{s}_1; \alpha(0, s_2) \right), \quad s = (\tilde{s}_1, s_2).$$

Using the properties (1)–(3) of ψ , we see that Lemma 6 applies to ψ_1 . Also, ψ_1 is compactly supported. Compact support along \tilde{s}_1 is due to the property (1) of ψ , and along s_2 – due to the cut-off χ . Substituting ψ_1 into (7.10), using Lemma 6, and then expressing ψ_1 in term of ψ yields

$$(7.15) \quad \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1a)} = \int_{\mathbb{R}^2} B_2(s_2) \left[\int_{[0,1]^3} \psi(\tilde{s}_1, v; \alpha(0, s_2)) dv \right] d\tilde{s}_1 ds_2.$$

By (7.8), similarly to (5.19),

$$(7.16) \quad \begin{aligned} & \int_{[0,1]^3} \sum_j (e \cdot (j - v) + q)_+ \nabla_u^2 \varphi(v - j) dv \\ &= \int_{\mathbb{R}^3} (-e \cdot v + q)_+ \nabla_u^2 \varphi(v) dv = (e \cdot u)^2 \hat{\varphi}(e, q). \end{aligned}$$

As $|e|$ not necessarily equals one, (7.16) assumes the extended definition of the CRT, cf. (3.5). By (7.6) and (7.8),

$$(7.17) \quad e(\alpha) \cdot u(\alpha) = \partial P(Y(\alpha, t)) / \partial t|_{t=0} =: P'_t(\alpha), \quad \alpha = \alpha(0, s_2).$$

With φ normalized, using (7.16) with $q = \tilde{s}_1$ and (7.17) in (7.15) gives

$$(7.18) \quad \begin{aligned} \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1a)} &= \int_{\mathbb{R}} B_2(s_2) (P'_t(\alpha(0, s_2)))^2 \int \hat{\varphi}(e(\alpha(0, s_2)), \tilde{s}_1) d\tilde{s}_1 ds_2 \\ &= \int_{\mathbb{R}} B_2(s_2) (P'_t(\alpha(0, s_2)))^2 ds_2. \end{aligned}$$

Consequently, from (7.10) we get

$$(7.19) \quad \begin{aligned} \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1a)} &= \int_{\mathbb{R}} B_1(\alpha(0, s_2)) \left| \frac{\partial \alpha(0, s_2)}{\partial s} \right| (P'_t(\alpha(0, s_2)))^2 ds_2 \\ &= \int_{\mathbb{R}^2} B_1(\alpha(s)) (P'_t(\alpha(s)))^2 \delta(P(Y(\alpha(s), 0))) \left| \frac{\partial \alpha(s)}{\partial s} \right| ds \\ &= \int_{S_+^2} B_1(\alpha) \left(\frac{\partial P(Y(\alpha, t))}{\partial t} \Big|_{t=0} \right)^2 \delta(P(Y(\alpha, 0))) d\alpha. \end{aligned}$$

In the second line we used that $s = (s_1, s_2)$ and $s_1 \equiv P(Y(\alpha(s), 0))$.

Next, consider the second part of $J_\epsilon^{(1)}$, which is obtained by using the second term on the right in (7.6) and replacing $G(\epsilon j)$ with $G(Y(\alpha, 0))$:

$$(7.20) \quad J_\epsilon^{(1b)} := \int_{S_+^2} B_1(\alpha) \sum_j L_+(\epsilon j) \frac{1}{\epsilon} \varphi'_y \left(\frac{Y(\alpha, 0)}{\epsilon} - j \right) \cdot Y''_{tt}(\alpha, 0) d\alpha.$$

The function $L_+(y)$ has bounded first derivatives, hence the limit of $J_\epsilon^{(1b)}$ can be easily found:

$$\begin{aligned}
(7.21) \quad \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1b)} &= \lim_{\epsilon \rightarrow 0} \int_{S_+^2} B_1(\alpha) \sum_j L_+(\epsilon j) \frac{1}{\epsilon} \varphi'_y \left(\frac{Y(\alpha, 0) - \epsilon j}{\epsilon} \right) \cdot Y''_{tt}(\alpha, 0) d\alpha \\
&= \lim_{\epsilon \rightarrow 0} \int_{S_+^2} B_1(\alpha) \partial_y \sum_j L_+(\epsilon j) \varphi \left(\frac{Y(\alpha, 0) + y - \epsilon j}{\epsilon} \right) \Big|_{y=0} \cdot Y''_{tt}(\alpha, 0) d\alpha \\
&= \int_{S_+^2} B_1(\alpha) \partial_y L_+(Y(\alpha, 0) + y) \Big|_{y=0} \cdot Y''_{tt}(\alpha, 0) d\alpha \\
&= \int_{S_+^2} B_1(\alpha) P'_y(Y(\alpha, 0)) \cdot Y''_{tt}(\alpha, 0) \theta(P(Y(\alpha, 0))) d\alpha.
\end{aligned}$$

Recall that $P > 0$ on the interior side of \mathcal{S} .

The final piece of $J_\epsilon^{(1)}$ is

$$\begin{aligned}
(7.22) \quad J_\epsilon^{(1c)} &:= \int_{S_+^2} B(\alpha) \sum_j (G(\epsilon j) - G(Y(\alpha, 0))) L_+(\epsilon j) \\
&\quad \times \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha.
\end{aligned}$$

In an $O(\epsilon)$ neighborhood of Γ , we have $|L(y)| = O(\epsilon)$. For any fixed $y \in \mathcal{V}_1$, $P(y) > 0$, we compute by dropping the subscript '+' from L :

$$\begin{aligned}
(7.23) \quad &\sum_j (G(\epsilon j) - G(Y(\alpha, 0))) L(\epsilon j) \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} \\
&= \left(\frac{\partial}{\partial t} \right)^2 [(G(Y(\alpha, t)) - G(Y(\alpha, 0))) L(Y(\alpha, t))] \Big|_{t=0} + O(\epsilon).
\end{aligned}$$

Substitution into (7.22) gives

$$\begin{aligned}
(7.24) \quad &\lim_{\epsilon \rightarrow 0} J_\epsilon^{(1c)} \\
&= \int_{S_+^2} B(\alpha) \left(\frac{\partial}{\partial t} \right)^2 [(G(Y(\alpha, t)) - G(Y(\alpha, 0))) L(Y(\alpha, t))] \Big|_{t=0} \theta(P(Y(\alpha, 0))) d\alpha.
\end{aligned}$$

We can apply the limit as $\epsilon \rightarrow 0$ inside the integral in (7.22) to obtain (7.24) because the integrand is uniformly bounded. This follows because the function $(G(y) - G(Y(\alpha, 0)))L(y)$ is smooth away from an $O(\epsilon)$ neighborhood of Γ , and the integrand is $O(1)$ within that neighborhood.

The final term to be considered arises because of the difference between $P(y)$ (cf. (7.3)) and its linear approximation $L(y)$ (cf. (7.4)):

$$(7.25) \quad J_\epsilon^{(2)} := \int_{S_+^2} B(\alpha) \sum_j G(\epsilon j) [P_+(\epsilon j) - L_+(\epsilon j)] \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha.$$

In the domain where $P(\epsilon j)$ and $L(\epsilon j)$ are both positive, we have

$$(7.26) \quad P_+(\epsilon j) - L_+(\epsilon j) = \frac{1}{2} P''_{yy}(Y(\alpha, 0)) (\epsilon j - Y(\alpha, 0)) \cdot (\epsilon j - Y(\alpha, 0)) + O(\epsilon^3).$$

This difference is zero if $P(\epsilon j)$ and $L(\epsilon j)$ are both negative. Thus,

$$(7.27) \quad \lim_{\epsilon \rightarrow 0} \sum_j G(\epsilon j) [P(\epsilon j) - L(\epsilon j)] \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t) - \epsilon j}{\epsilon} \right) \Big|_{t=0} \\ = G(Y(\alpha, 0)) P''_{yy}(Y(\alpha, 0)) Y'_t(\alpha, 0) \cdot Y'_t(\alpha, 0), \quad P(Y(\alpha, 0)) > 0.$$

In the region where $P(Y(\alpha, 0)) < 0$, the limit is obviously zero. Hence

$$(7.28) \quad \lim_{\epsilon \rightarrow 0} J_\epsilon^{(2)} = \int_{S^2_+} B_1(\alpha) P''_{yy}(Y(\alpha, 0)) Y'_t(\alpha, 0) \cdot Y'_t(\alpha, 0) \theta(P(Y(\alpha, 0))) d\alpha.$$

As before, we can apply the limit as $\epsilon \rightarrow 0$ inside the integral in (7.25) to obtain (7.28) because the integrand is uniformly bounded. Indeed,

$$(7.29) \quad P_+(y) - L_+(y) = (L(y) + O(\epsilon^2))_+ - L_+(y) = O(\epsilon^2),$$

so the integrand in (7.25) remains bounded as $\epsilon \rightarrow 0$. The domain where $P(y)$ and $L(y)$ are of different signs is a shrinking $O(\epsilon)$ neighborhood of Γ , and the desired result follows.

Combining (7.19), (7.21), (7.24), and (7.28) gives the result, which, in compact form, can be written as follows

$$(7.30) \quad \lim_{\epsilon \rightarrow 0} f_{\chi\epsilon}(x_0) \\ = \int_{S^2_+} B(\alpha) [G_0(\partial_t P)^2 \delta(P_0) + (G_0 P'_y Y''_{tt} + \partial_t^2((G - G_0)L) + G_0 P''_{yy} Y'_t \cdot Y'_t) \theta(P_0)] d\alpha \\ = \int_{S^2_+} B(\alpha) [G_0(\partial_t P)^2 \delta(P_0) + (G_0 \partial_t^2 P + \partial_t^2((G - G_0)L)) \theta(P_0)] d\alpha.$$

Here $G_0 := G(Y(\alpha, 0))$, $G := G(Y(\alpha, t))$, $P := P(Y(\alpha, t))$, $P_0 := P(Y(\alpha, 0))$, and the derivatives with respect to t are evaluated at $t = 0$. This coincides with what we get by substituting $g = P_+ G$ into the continuous inversion formula (2.14). Indeed, representing $P_+ G = (GP)\theta(P)$, we have

$$(7.31) \quad \partial_t^2((GP)\theta(P)) = \partial_t[\partial_t(GP)\theta(P) + (GP)\delta(P)\partial_t P] = \partial_t[\partial_t(GP)\theta(P)] \\ = \partial_t(GP)\partial_t P \delta(P_0) + \partial_t^2(GP)\theta(P_0) \\ = G_0(\partial_t P)^2 \delta(P_0) + \partial_t^2(GP)\theta(P_0).$$

The coefficients in front of the delta-function in (7.30) and (7.31) match. Subtracting the coefficients in front of the Heaviside function gives:

$$(7.32) \quad [G_0 \partial_t^2 P + \partial_t^2((G - G_0)L)] - \partial_t^2(GP) = -\partial_t^2[(G - G_0)(P - L)] = 0.$$

Here we have used that G_0 is independent of t , and the expression under the derivative has a zero of third order at $t = 0$. Thus the theorem is proven in the case $x = x_0$.

Next, consider the case of a general $x_\epsilon := x_0 + \epsilon \tilde{x}$ (cf. (7.1)). We begin by repeating the steps (7.3)–(7.9), where all the auxiliary functions, such as Y , are computed using x_ϵ instead of x_0 . It is clear that in any place where an auxiliary function is not divided by ϵ , e.g. $B(\alpha)$ in (7.5) and $e(\alpha)$, $u(\alpha)$ in (7.8), replacing x_ϵ with x_0 introduces an error of magnitude $O(\epsilon)$. Here we also used the property (2) of ψ . Consequently, the analogue of (7.9) for x_ϵ becomes:

$$(7.33) \quad J_\epsilon^{(1a)}(x_\epsilon) = \frac{1}{\epsilon} \int_{S^2_+} B_1(\alpha) \psi \left(\frac{P(Y(\alpha, 0; x_\epsilon))}{\epsilon}, \frac{Y(\alpha, 0; x_\epsilon)}{\epsilon}; \alpha \right) d\alpha + O(\epsilon),$$

where only Y is different from the analogous function in (7.9). Note that $P(y)$ depends only on the shape of \mathcal{S} in a neighborhood of x_0 and, therefore, is independent of x_ϵ . We have

$$(7.34) \quad \begin{aligned} Y(\alpha, 0; x_\epsilon) &= Y(\alpha, 0) + \epsilon W(\alpha, \tilde{x}) + O(\epsilon^2), \\ P(Y(\alpha, 0; x_\epsilon)) &= P(Y(\alpha, 0)) + \epsilon P'_y(Y(\alpha, 0))W(\alpha, \tilde{x}) + O(\epsilon^2), \end{aligned}$$

for some smooth and bounded W . Here $Y(\alpha, 0)$ is the same as in (7.9). Substituting into (7.33) gives

$$(7.35) \quad \begin{aligned} J_\epsilon^{(1a)}(x_\epsilon) &= \frac{1}{\epsilon} \int_{S_+^2} B_1(\alpha) \psi \left(\frac{P(Y(\alpha, 0))}{\epsilon} + P'_y(Y(\alpha, 0))W(\alpha, \tilde{x}), \right. \\ &\quad \left. \frac{Y(\alpha, 0)}{\epsilon} + W(\alpha, \tilde{x}); \alpha \right) d\alpha + O(\epsilon). \end{aligned}$$

Similarly to (7.14), introduce

$$(7.36) \quad \begin{aligned} \psi_2(s, y) &:= B_1(\alpha) \psi(\tilde{s}_1 + P'_y(Y(\alpha, 0))W(\alpha, \tilde{x}), y + W(\alpha, \tilde{x}); \alpha), \\ \alpha &= \alpha(0, s_2), s = (\tilde{s}_1, s_2). \end{aligned}$$

The point \tilde{x} is fixed, so we do not need to list it in the arguments of ψ_2 . Clearly, ψ_2 satisfies the same properties (1)–(3) as ψ . Hence Lemma (6) applies to ψ_2 as well, and we get similarly to (7.10), (7.14), and (7.15):

$$(7.37) \quad \begin{aligned} \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1a)}(x_\epsilon) &= \int_{\mathbb{R}^2} \left[\int_{[0,1]^3} \psi_2(\tilde{s}_1, v; \alpha(0, s_2)) dv \right] d\tilde{s}_1 ds_2 \\ &= \int_{\mathbb{R}^2} B_2(s_2) \left[\int_{[0,1]^3} \psi(\tilde{s}_1, v; \alpha(0, s_2)) dv \right] d\tilde{s}_1 ds_2. \end{aligned}$$

Here we have used that the integrals with respect to \tilde{s}_1 and v are unaffected by the constant (with respect to \tilde{s}_1 and v) shifts in (7.36). Therefore, (7.19) holds with $J_\epsilon^{(1a)}(x_\epsilon)$ on the left.

To find the limit of $J_\epsilon^{(1b)}(x_\epsilon)$, consider the key step in (7.21):

$$(7.38) \quad \sum_j L_+(\epsilon j; x_\epsilon) \varphi \left(\frac{Y(\alpha, 0; x_\epsilon) + y - \epsilon j}{\epsilon} \right) = L_+(Y(\alpha, 0; x_\epsilon) + y; x_\epsilon),$$

which is rewritten with x_0 replaced by x_ϵ . This equality holds everywhere except in an $O(\epsilon)$ neighborhood of Γ ($= \Gamma_{x_0}$). Here we use that the curve Γ_{x_ϵ} , which is obtained by solving $P(Y(\alpha, 0; x_\epsilon)) = 0$, depends smoothly on x_ϵ , and $\text{dist}(\Gamma_{x_\epsilon}, \Gamma) = O(\epsilon)$ (see the argument preceding the statement of Theorem 3). Similarly to (7.21), the integrand is uniformly bounded, and we get

$$(7.39) \quad \lim_{\epsilon \rightarrow 0} J_\epsilon^{(1b)}(x_\epsilon) = \int_{S_+^2} B_1(\alpha) P'_y(Y(\alpha, 0)) \cdot Y''_{tt}(\alpha, 0) \theta(P(Y(\alpha, 0))) d\alpha.$$

The fact that the limits of $J_\epsilon^{(1c)}(x_\epsilon)$ and $J_\epsilon^{(2)}(x_\epsilon)$ as $\epsilon \rightarrow 0$ are independent of $\tilde{x} \in \tilde{\mathcal{U}}$ can be established in a similar way, and the theorem is proven. \square

8. NUMERICAL EXPERIMENT

We start by constructing an interpolation kernel with the required properties. To obtain φ , we first obtain an interpolation kernel φ_{1D} that has properties IK1–IK5 in \mathbb{R} , and then extend it to \mathbb{R}^3 in a separable fashion. To obtain φ_{1D} we use the result of [11], where such a kernel is obtained following the method in [3]:

$$(8.1) \quad \varphi_{1D}(t) = 0.5(B_3(t) + B_3(t-2)) + 4B_3(t-1) - 2(B_4(t) + B_4(t-1)).$$

Here B_n is the cardinal B-spline of degree n supported on $[0, n+1]$. Then the kernel φ becomes

$$(8.2) \quad \varphi(y) = \prod_{k=1}^3 \varphi_{1D} \left(\frac{y_k}{\Delta_k} + 3 \right), \quad y = (y_1, y_2, y_3),$$

where Δ_k is the data stepsize along the k -th axis. For simplicity, in this paper all the Δ_k are equal, i.e. $\Delta_k = \epsilon$, $k = 1, 2, 3$.

The GRT we consider here integrates a function supported in the half-space $x_3 > 0$ over spheres that are tangent to the plane $x_3 = 0$. The family of such spheres is three-dimensional. We parametrize the spheres (and, consequently, the GRT) by the coordinates of their center y . Thus, the surfaces \mathcal{S}_y are spheres, and the defining function Φ in (2.7) becomes:

$$(8.3) \quad \Phi(x, y) := y_3^2 - (x_1 - y_1)^2 - (x_2 - y_2)^2 - (x_3 - y_3)^2 = 0.$$

Clearly,

$$(8.4) \quad \Phi'_x(x, y) = 2(y - x), \quad \Phi'_y(x, y) = 2(x_1 - y_1, x_2 - y_2, x_3).$$

The test object is the ball with center $x_c = (0, 0, 11)$, radius $R = 5$, and uniform density 1. The point on the boundary x_0 , in a neighborhood of which we compute resolution, is given by

$$(8.5) \quad x_0 = x_c - R\alpha_0, \quad \alpha_0 = (\sin(0.2\pi) \cos(0.7\pi), \sin(0.2\pi) \sin(0.7\pi), \cos(0.2\pi)).$$

In agreement with our convention, α_0 points into the interior of the ball.

There can be two spheres that are tangent to the ball at x_0 . As an example, we consider the sphere whose center y_0 satisfies $(x_0 - y_0) \cdot \alpha_0 > 0$. Thus, for reconstruction near x_0 we use the data in a neighborhood of y_0 . With this choice of y_0 , the condition $\Phi'_x(x_0, y_0)/|\Phi'_x(x_0, y_0)| = -\alpha_0$ (see the text following (3.1)) is satisfied with Φ given by (8.3). For the selected x_0 , α_0 , and y_0 , we compute using (8.4):

$$(8.6) \quad \nu = |\Phi'_x|/|\Phi'_y| = 0.526.$$

To compute the GRT, we use the formula for the area of the spherical cap:

$$(8.7) \quad A = 2\pi R h,$$

where R is the radius of the sphere, and h is the height of the cap. The values of R and h can be computed once the center of the sphere \mathcal{S}_y is chosen (e.g., $R = y_3$). To simulate discrete data, the GRT is computed at the points $y = r + \epsilon j$. The interpolated data g_ϵ is computed using (2.13), where the kernel is given by (8.2) with $\Delta_k = \epsilon$, $k = 1, 2, 3$.

To apply the inversion formula (2.14), we numerically integrate g_ϵ over a neighborhood of α_0 on the unit sphere. To compute $(\partial/\partial t)^2 g_\epsilon(Y(\alpha, t; x))$ at $t = 0$ we use (2.13) and the chain rule as in (7.6). Given x , α , and t , the center of the sphere containing the point $x + t\alpha$ and normal to α at that point (cf. the paragraph following (2.13)) is easily found to be:

$$(8.8) \quad Y(\alpha, t; x) = (x + t\alpha) - \frac{x_3 + t\alpha_3}{1 + \alpha_3} \alpha.$$

Consequently,

$$(8.9) \quad \left. \frac{\partial}{\partial t} Y(\alpha, t; x) \right|_{t=0} = \frac{1}{1 + \alpha_3} \alpha, \quad \left. \frac{\partial^2}{\partial t^2} Y(\alpha, t; x) \right|_{t=0} = 0,$$

and

$$(8.10) \quad \left(\frac{\partial}{\partial t} \right)^2 \varphi \left(\frac{Y(\alpha, t; x) - \epsilon j}{\epsilon} \right) \Big|_{t=0} d\alpha = \frac{1}{\epsilon^2} \frac{1}{(1 + \alpha_3)^2} \sum_{i,k=1}^3 \varphi''_{ik} \left(\frac{Y(\alpha, 0; x)}{\epsilon} - j \right) \alpha_i \alpha_k.$$

The cut-off function χ in (2.14) is constructed as follows. Let α^\perp run through the unit sphere in the plane α_0^\perp . Then any $\alpha \in S_+^2$ (α_0 is the North pole of S_+^2) can be represented in the form $\alpha = (\cos \omega)\alpha_0 + (\sin \omega)\alpha^\perp$, $0 \leq \omega \leq \pi/2$. In the code we use

$$(8.11) \quad \chi(\alpha) = \begin{cases} 1, & 0 \leq \omega < 0.8\omega_{\text{mx}}, \\ \frac{1 + \cos((\omega - 0.8\omega_{\text{mx}})/(0.2\omega_{\text{mx}}))}{2}, & 0.8\omega_{\text{mx}} \leq \omega < \omega_{\text{mx}}, \\ 0, & \omega \geq \omega_{\text{mx}}. \end{cases}$$

Finally, the predicted response is computed using (3.8). The results corresponding to $\epsilon = 0.01$ are shown in Figure 1. We see a good match between the predicted and actual transition curves.




FIGURE 1. Comparison of the predicted and actual transition curves for $\epsilon = 0.01$.

REFERENCES

- [1] F. Andersson, M. V. De Hoop, and H. Wendt. Multiscale Discrete Approximation of Fourier Integral Operators. *Multiscale Modeling and Simulation*, 10:111–145, 2012.
- [2] G. Beylkin. The inversion problem and applications of the generalized Radon transform. *Comm. Pure and Appl. Math.*, 37:579–599, 1984.
- [3] T. Blu, P. Thévenaz, and M. Unser. Complete Parameterization of Piecewise-Polynomial Interpolation Kernels. *IEEE Transactions on Image Processing*, 12:1297–1309, 2003.
- [4] E. Candes, L. Demanet, and L. Ying. Fast computation of Fourier integral operators. *SIAM Journal on Scientific Computing*, 29:2464–2493, 2007.
- [5] E. Candes, L. Demanet, and L. Ying. A fast butterfly algorithm for the computation of Fourier integral operators. *SIAM Multiscale Modeling and Simulation*, 7:1727–1750, 2009.
- [6] M. Cheney and B. Borden. Synthetic Aperture Radar Imaging. In O. Scherzer, editor, *Handbook of Mathematical Methods in Imaging*, pages 763–799. Springer, New York, NY, 2015.
- [7] C. Grathwohl, P. Kunstmann, E. T. Quinto, and A. Rieder. Approximate inverse for the common offset acquisition geometry in 2D seismic imaging. *Inverse Problems*, 34, 2018. article id 014002.
- [8] L. Hormander. *The Analysis of Linear Partial Differential Operators, Vol I*. Springer Verlag, New York, 1983.

- [9] A. Katsevich. An accurate approximate algorithm for motion compensation in two-dimensional tomography. *Inverse Problems*, 26, 2010. article ID 065007 (16 pp).
- [10] A. Katsevich. A local approach to resolution analysis of image reconstruction in tomography. *SIAM Journal on Applied Mathematics*, 77:1706–1732, 2017.
- [11] A. Katsevich. Analysis of reconstruction from discrete radon transform data in \setminus^3 when the function has jump discontinuities. *SIAM Journal on Applied Mathematics*, 2019. to appear.
- [12] P. Kuchment and L. Kunyansky. Mathematics of Photoacoustic and Thermoacoustic Tomography. In O. Scherzer, editor, *Handbook of Mathematical Methods in Imaging*, pages 1117–1167. Springer, New York, NY, 2015.
- [13] L. Kuipers and H. Niederreiter. *Uniform Distribution of Sequences*. Dover Publications, Inc., Mineola, NY, 2006.
- [14] F. Natterer. *The Mathematics of Computerized Tomography*. SIAM, Philadelphia, 2001.
- [15] E. T. Quinto, A. Rieder, and Th. Schuster. Local inversion of the sonar transform regularized by the approximate inverse. *Inverse Problems*, 27, 2011. article id 035006.
- [16] A.G. Ramm and A.I. Zaslavsky. Reconstructing singularities of a function given its Radon transform. *Math. and Comput. Modelling*, 18(1):109–138, 1993.
- [17] A.G. Ramm and A.I. Zaslavsky. Singularities of the Radon transform. *Bull. Amer. Math. Soc.*, 25:109–115, 1993.
- [18] P. Stefanov. Semiclassical sampling and discretization of certain linear inverse problems. arXiv:1811.01240, 2018.
- [19] F. Terzioglu, P. Kuchment, and L. Kunyansky. Compton camera imaging and the cone transform: a brief overview. *Inverse Problems*, 34, 2018. article id 054002.
- [20] F. Trèves. *Introduction to Pseudodifferential and Fourier Integral Operators. Volume 2: Fourier Integral Operators*. The University Series in Mathematics. Plenum, New York, 1980.
- [21] K. Wang and M. A. Anastasio. Photoacoustic and Thermoacoustic Tomography: Image Formation Principles. In O. Scherzer, editor, *Handbook of Mathematical Methods in Imaging*, pages 1081–1116. Springer, New York, NY, 2015.
- [22] H. Yang. *Oscillatory Data Analysis and Fast Algorithms for Integral Operators*. PhD thesis, Stanford University, 2015.
- [23] M. Zworski. *Semiclassical analysis*, volume 138 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.