ON THE COMPLEXITY OF PROPERTIES OF TRANSFORMATION SEMIGROUPS

LUKAS FLEISCHER

University of Stuttgart, FMI Universitätsstraße 38, 70569 Stuttgart, Germany

School of Computer Science, University of Waterloo 200 University Avenue West, Waterloo, ON N2L 3G1, Canada

TREVOR JACK

University of Colorado, Department of Mathematics 2300 Colorado Avenue, Boulder, USA

ABSTRACT. We investigate the computational complexity for determining various properties of a finite transformation semigroup given by generators. We introduce a simple framework to describe transformation semigroup properties that are decidable in AC^0 . This framework is then used to show that the problems of deciding whether a transformation semigroup is a group, commutative or a semilattice are in AC^0 . Deciding whether a semigroup has a left (resp. right) zero is shown to be NL-complete, as are the problems of testing whether a transformation semigroup is nilpotent, \mathcal{R} -trivial or has central idempotents. We also give NL algorithms for testing whether a transformation semigroup is idempotent, orthodox, completely regular, Clifford or has commuting idempotents. Some of these algorithms are direct consequences of the more general result that arbitrary fixed semigroup equations can be tested in NL. Moreover, we show how to compute left and right identities of a transformation semigroup in polynomial time. Finally, we show that checking whether an element is regular is PSPACE-complete.

1. INTRODUCTION

Given a permutation group by generators, many of its properties, like size and membership, can be determined in polynomial time using Sims' stabilizer chains. In contrast, the known algorithms for the corresponding problems for transformation semigroups given by generators often rely on an enumeration of the \mathcal{R} -classes of the

E-mail addresses: fleischer@fmi.uni-stuttgart.de, trevor.jack@colorado.edu. *Date*: August 29, 2019.

²⁰¹⁰ Mathematics Subject Classification. Primary: 20M20; Secondary 68Q25.

Key words and phrases. transformation semigroups, algorithms, computational complexity, PSPACE-completeness, NL-completeness, regular elements, semigroup identities.

This material is based upon work supported by the National Science Foundation under Grant No. DMS 1500254.

semigroup, which already requires exponential time [4, 9]. Moreover, the membership problem for transformation semigroups is known to be PSPACE-complete [8], as is checking whether two elements are \mathcal{R} -related [2] and whether a transformation semigroup is aperiodic [3]. In this work, we will show that many similar problems, such as checking whether a specific element is regular and variants thereof, are also PSPACE-complete (Theorem 7.3).

While the hardness results above indicate that problems concerning Green's relations, inverses and idempotent elements are generally hard in the transformation semigroup setting, this is certainly not true in general. Clearly, the problem of checking whether a given transformation is idempotent is in AC^0 , the class of all sets decidable by unbounded fan-in Boolean circuits of constant depth: it suffices to verify that the transformation is the identity on its image. Similarly, we show that the problems of testing whether a transformation semigroup is a group, commutative or a semilattice are in AC^0 . Our results even hold in the *uniform* setting where so-called *direct connection languages* of the circuits are required to be decidable in logarithmic time using a deterministic random-access Turing machine.

Aside from that, we prove that many decision problems for transformation semigroups are complete for NL, the class of all sets decidable in non-deterministic logarithmic space. This includes testing whether a transformation semigroup

- contains a left zero, a right zero or a zero (Theorem 4.6);
- is nilpotent (Theorem 4.10);
- is \mathcal{R} -trivial (Theorem 4.13);
- has central idempotents (Theorem 5.4).

For some problems, we establish membership in NL, but we leave hardness results as open problems. These include testing whether a transformation semigroup

- is a band (Corollary 5.2);
- has commuting idempotents (Corollary 5.2);
- is orthodox (Corollary 5.2);
- is completely regular (Theorem 5.6);
- is a Clifford semigroup (Corollary 5.8).

We also describe an NL algorithm to check whether a transformation semigroup models a fixed identity or a fixed quasi-identity involving idempotents (Theorem 5.1) and prove that this problem is NL-complete for very simple identities already (Theorem 5.3).

Additionally, we show that the left and right identities of a transformation semigroup can be enumerated in polynomial time (Theorems 6.2 and 6.4).

2. Preliminaries

Let S be a semigroup. An element ℓ of a semigroup S is a *left identity* if $\ell s = s$ for all $s \in S$. An element r of a semigroup S is a *right identity* if sr = s for all $s \in S$. An element ℓ of a semigroup S is a *left zero* if $\ell s = \ell$ for all $s \in S$. An element r of a semigroup S is a *right zero* if sr = r for all $s \in S$. An element that is both a left and a right zero is called zero. If a semigroup contains a zero element, this element is unique and usually denoted by 0. If a semigroup S that has a zero, ℓ and a right zero r, then $\ell = \ell r = r$ is a zero. A semigroup S that has a zero, $0 \in S$, is called *d*-nilpotent if $S^d = \{0\}$ for a fixed $d \in \mathbb{N}$. We call S nilpotent if it is *d*-nilpotent for some $d \in \mathbb{N}$. The smallest such d is called its nilpotency degree.

3

An element $e \in S$ is *idempotent* if $e^2 = e$. A semigroup is *idempotent* if all elements are idempotent. Following classical terminology, idempotent semigroups are often also referred to as *bands*. A semigroup is *orthodox* if the product of any two idempotents is again idempotent. It is well-known that in every finite semigroup S, there is a natural number ω_S such that s^{ω_S} is idempotent for every $s \in S$. If the reference to S is clear from the context, we usually write ω instead of ω_S .

Two elements $s, t \in S$ commute if st = ts. An element $s \in S$ is central if it commutes with every other element of S. A semigroup S is commutative if all pairs of elements of S commute. A semigroup that is both idempotent and commutative is also called a *semilattice*.

An element $s \in S$ is regular if there exists $t \in S$ such that sts = s. An element $t \in S$ such that tst = t is called a *weak inverse* of s. If sts = s and tst = t, then t is an *inverse* of s. A semigroup is called *regular* if all of its elements are regular. A semigroup is *inverse* if every element has a unique inverse. A semigroup S is *completely regular* if every element of S belongs to some subgroup of S. A semigroup is a *Clifford semigroup* if it is completely regular and its idempotents commute.

A semigroup is called *aperiodic* if it does not contain any nontrivial subsemigroup that is a group. A semigroup S is \mathcal{R} -trivial if it does not contain two distinct elements $s, t \in S$ such that $sS \cup \{s\} = tS \cup \{t\}$.

A variety of finite semigroups is a class of finite semigroups that is closed under finite direct products and under taking divisors. In the literature, such classes of semigroups are often also referred to as pseudovarieties.

The full transformation semigroup over some set Q is the set of all mappings $f: Q \to Q$, the so-called transformations over Q, together with function composition. Subsemigroups of the full transformation semigroup are often also referred to as transformation semigroups. The elements of Q are sometimes referred to as points.

For $n \in \mathbb{N}$, we define $[n] := \{1, \ldots, n\}$ and we use T_n to denote the full transformation semigroup over the set [n]. For elements $a_1, \ldots, a_k \in T_n$, let $\langle a_1, \ldots, a_k \rangle$ be the subsemigroup of T_n generated by a_1, \ldots, a_k .

For $s \in T_n$ and $A \subseteq T_n$, define the kernel of s (resp. A) as

$$\ker(s) := \left\{ (p,q) \in [n]^2 \mid ps = qs \right\}$$
$$\ker(A) := \bigcap_{a \in A} \ker(a).$$

For $S = \langle a_1, \ldots, a_k \rangle$, note that ker $(S) = \text{ker}(\{a_1, \ldots, a_k\})$ and that ker(S) forms an equivalence relation on [n]. Then [n]/ker(S) is a partition into equivalence classes, $[\![q]\!] := \{p \in [n] \mid ps = qs \text{ for every } s \in S\}$. We can define a natural action of S on these classes by the following homomorphism:

$$S \to T_{[n]/\ker(S)}, s \mapsto \overline{s}$$

with

$$\llbracket q \rrbracket \overline{s} = \llbracket qs \rrbracket$$
 for $q \in [n]$.

Let $\overline{S} := \{\overline{s} : s \in S\}.$

Denote the image of a semigroup element $s \in S$ as

 $[n]s := \{q \in [n] \mid q = ps \text{ for some } p \in [n] \text{ and some } s \in S\}$

Define $[n]S := \bigcup_{s \in S} [n]s$. We can define a natural action of S on its image by the following homomorphism:

$$S \to T_{[n]S}, s \mapsto \widetilde{s}$$

with

$$q\tilde{s} = qs$$
 for $q \in [n]S$

and

$$\widetilde{S} := \{ \widetilde{s} : s \in S \}$$

Let S be a transformation semigroup, i.e., $S := \langle a_1, \ldots, a_k \rangle \leq T_n$ for some $n, k \in \mathbb{N}$. For $d \in \mathbb{N}$, we define S to act upon d-tuples $[n]^d$ component-wise: for $(q_1, \ldots, q_d) \in [n]^d$ and $s \in S$,

$$(q_1,\ldots,q_d)s = (q_1s,\ldots,q_ds)$$

For a set of transformations $A := \{a_1, \ldots, a_k\}$ acting on a set Q, define the transformation graph $\Gamma(A, Q)$ as having vertices Q and directed edges

$$E := \{ (p,q) \in Q^2 \mid \exists i \in [k] (pa_i = q) \}.$$

Denote the *pre-image* of an element $q \in Q$ as

$$S^{-1}(q) := \{ q \in Q \mid ps = q \text{ for some } s \in S \}.$$

We use the notation

$$Fix(A,Q) := \{ q \in Q \mid qa = q \text{ for all } a \in A \}$$

to denote the set of fixed points of A.

3. FIRST-ORDER DEFINABLE PROPERTIES

In this section, we will investigate classes of transformation semigroups with very efficient membership tests. To this end, we introduce a variant of first-order logic that is used to define classes of semigroups. We allow quantification over the set of generators A and over points of the underlying set Q. The only allowed predicates are of the form $p \cdot a_1 \cdots a_k = q \cdot b_1 \cdots b_\ell$ where p, q are variables corresponding to points and $a_1, \ldots, a_k, b_1, \ldots, b_\ell$ are variables corresponding to generators. For example, the formula

$$\forall a \in A \,\forall p, q \in Q \colon p \cdot a = q \cdot a$$

can be used to express that the image of every element of S is a singleton. It follows immediately from the well-known result $FO = AC^0$ [6, Theorem 5.22] that the membership problem for classes of semigroups defined by such formulas is in AC^0 .

Remark 3.1. It is important to note that the logical formalism described above does not allow quantification over elements of the generated transformation semigroup S. In fact, allowing such quantifiers yields a much more expressive formalism. For example, the following formula can be used to define the class of all aperiodic transformation semigroups:

$$\forall s, t \in S \colon (s \ \mathcal{H} \ t \to s = t).$$

5

Here, $s \mathcal{H} t$ is a shorthand for the formula

$$(s = t) \lor ((\exists r \in S : sr = t) \land (\exists r \in S : rs = t) \land (\exists r \in S : tr = s) \land (\exists r \in S : rt = s)),$$

s = t is a short form for $\forall q \in Q: q \cdot s = q \cdot t$, and formulas of the form rs = t are short forms for $\forall q \in Q: q \cdot rs = q \cdot t$. Testing whether a transformation semigroup is aperiodic has been shown to be PSPACE-complete in [3].

The class of commutative transformation semigroups is easily described using this formalism. Therefore, we can design an efficient algorithm for the following decision problem:

Commutative

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Is $\langle a_1, \ldots, a_k \rangle$ commutative?

Theorem 3.2. Commutative is in AC^0 .

Proof. Clearly, a semigroup is commutative if and only if all generators commute. Therefore, commutativity can be expressed by the formula

$$\forall a, b \in A \, \forall q \in Q \colon q \cdot ab = q \cdot ba.$$

Thus, testing whether a transformation semigroup is commutative is in AC^0 .

Theorem 3.3. The problem of determining if a transformation semigroup is a semilattice is in AC^0 .

Proof. We can use the formula

$$\forall a \in A \,\forall q \in Q \colon q \cdot a^2 = q \cdot a$$

to express that every generator is idempotent. Together with the previous theorem, this yields the desired statement. $\hfill \Box$

Our formula for permutation groups will be based on the following lemma.

Lemma 3.4. Let $a_1, \ldots, a_k \in T_n$. Then, $S := \langle a_1, \ldots, a_k \rangle$ is a group if and only if the following three properties hold:

- (1) all generators have the same image $X \subseteq Q$,
- (2) all generators are permutations on X, and
- (3) all generators have the same kernel $\ker(a_i)$.

Proof. Suppose that S is a group. If one of the generators a_i is not a permuation on its image, then the image of a_i^2 is strictly smaller than the image of a_i , which yields $a_i^{\omega+1} \neq a_i$. Since in a group, the only idempotent element is the identity element, this is a contradiction. We may therefore assume that all generators are permutations on their images. Therefore, if two generators a_i and a_j do not have the same image, then the same holds for a_i^{ω} and a_j^{ω} , contradicting the fact that a group contains a unique idempotent element. If there exist $p, q \in [n]$ and $i, j \in [k]$ such that $pa_i = qa_i$ but $pa_j \neq qa_j$, we have $pa_i^{\omega} = qa_i^{\omega}$ and $pa_j^{\omega} \neq qa_j^{\omega}$ using the observation that a_j is a bijection on its image. This again yields two different idempotent elements.

Conversely, suppose that the three properties stated above hold. By induction, all three properties then also hold for every element of S. Let $s \in S$. We claim

that s^{ω} is the identity element which suffices to conclude the proof. To this end, we show that $s^{\omega}a_i = a_i = a_i s^{\omega}$ for all $i \in [k]$. Let $q \in [n]$. Clearly, since s is a bijection on the image of a_i , the transformation s^{ω} is the identity on qa_i , thus $qa_i s^{\omega} = qa_i$. Let $p = qs^{\omega}$. Clearly, $ps^{\omega} = qs^{\omega}$, i.e., $(p,q) \in \ker(s^{\omega})$. Since all elements have the same kernel, this yields $pa_i = qa_i$ and thus, $qs^{\omega}a_i = qa_i$.

Theorem 3.5. The problem of determining whether a transformation semigroup is a group is in AC^0 .

Proof. We provide formulas for all the conditions stated in Lemma 3.4. The first condition can be expressed by the formula

$$\forall a, b \in A \,\forall q \in Q \colon (\exists p \in Q \colon pa = q) \to (\exists p \in Q \colon pb = q).$$

The second property is expressed by

$$\forall a \in A \,\forall p, q \in Q \colon pa \neq qa \rightarrow pa^2 \neq qa^2.$$

The property of matching kernels is defined by the

$$\forall a, b \in A \,\forall p, q \in Q \colon (pa = qa \leftrightarrow pb = qb)$$

This suffices to show that the problem of deciding whether a transformation semigroup is a group is in AC^0 .

4. Locally Testable Properties

In this section, we investigate properties of a transformation semigroup S with an underlying set [n] that can be checked simply by verifying certain conditions on each vertex of the graph $\Gamma(S, [n])$. We first investigate the problem of testing whether a transformation semigroup has a right zero.

RightZero

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Does $\langle a_1, \ldots, a_k \rangle$ have a right zero?

Lemma 4.1. Let $k, n \in \mathbb{N}$ and let $a_1, \ldots, a_k \in T_n$. An element r is a right zero of $S := \langle a_1, \ldots, a_k \rangle$ iff $[n] = \bigcup_{p \in [n]_r} S^{-1}(p)$ is a disjoint union.

Proof. For the forward direction, assume r is a right zero. Note that $r^2 = r$. Hence, r fixes its images. Pick any $q \in [n]$ and $s, t \in S$ such that $qs, qt \in [n]r$. Then qs = qsr = qr = qtr = qt and thus $[n] = \bigcup_{p \in [n]r} S^{-1}(p)$ is a disjoint union.

Conversely, assume S has an element, r, such that $[n] = \bigcup_{p \in [n]_r} S^{-1}(p)$ is disjoint. Pick any $q \in [n]$ and any $s \in S$. Then $q \in S^{-1}(qr)$ and $q \in S^{-1}(qsr)$. But since the union is disjoint, this means qr = qsr. Thus, r is a right zero.

Lemma 4.2. Let $k, n \in \mathbb{N}$ and let $a_1, \ldots, a_k \in T_n$. Then $S := \langle a_1, \ldots, a_k \rangle$ has a right zero iff for every pair $p, q \in [n]$ that are in the same connected component of $\Gamma(\{a_1, \ldots, a_k\}, [n])$, there is some $s \in S$ such that ps = qs.

Proof. For the forward direction, assume r is a right zero of S. Then, by Lemma 4.1, $[n] = \bigcup_{q' \in [n]_r} S^{-1}(q')$ is a disjoint union. Thus, the $S^{-1}(q')$ sets are the vertex sets of the connected components of $\Gamma(\{a_1, \ldots, a_k\}, [n])$. Moreover, for each $p, q \in S^{-1}(q'), pr = q' = qr$.

For the converse, let $w \in S$ have minimal image size. Pick any $p, q \in [n]$ in the same connected component V. Note that $pw, qw \in V$. By assumption, then, there

is some $s \in S$ such that pws = qws. Since |[n]w| is minimal, then |[n]w| = |[n]ws|and hence pw = qw. That is, |Vw| = 1 for each connected component V. Thus, $[n] = \bigcup_{q' \in [n]w} S^{-1}(q')$ is disjoint. Therefore, by Lemma 4.1, w is a right zero. \Box

Lemma 4.3. RightZero is in NL.

Proof. By Lemma 4.2, we need only show every pair of vertices from the same connected component can be collapsed by some semigroup element to a common image. First, compute the edges of $\Gamma(\{a_1, \ldots, a_k\}, [n])$ with a log-space transducer. Then for each pair $(p,q) \in [n]$, use a log-space algorithm for undirected graph connectivity [10] to determine if p and q are in the same connected component. If not, proceed to the next pair. If so, guess generators for an $s \in S$ such that ps = qs.

Similarly, one can ask whether a semigroup has a left zero. Due to lack of leftright symmetry in transformation semigroups, this requires a different approach.

LeftZero

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Does $\langle a_1, \ldots, a_k \rangle$ have a left zero?

Lemma 4.4. Let $k, n \in \mathbb{N}$. Let $a_1, \ldots, a_k \in T_n$ and $S := \langle a_1, \ldots, a_k \rangle$. Then, S contains a left zero iff for each $q \in [n]$, there exists $s \in S$ such that $qs \in Fix(S, [n])$.

Proof. For the forward direction, assume $\ell \in S$ is a left zero. Then for any $q \in [n]$ and any $s \in S$, $(q\ell)s = q\ell$. Thus, $q\ell \in Fix(S, [n])$.

Conversely, assume that for any $q \in [n]$, there exists $s \in S$ such that $qs \in Fix(S, [n])$. Let $s \in S$ be an element such that [n]s is of minimal size. Note that $Fix(S, [n]) \subseteq [n]s$. Assume for contradiction there is $q \in [n]s \setminus Fix(S, [n])$. We know there is an $t \in S$ such that $qt \in Fix(S, [n])$. Since Fix(S, [n])t = Fix(S, [n]), then |[n]st| < |[n]s|, a contradiction. Therefore, [n]s = Fix(S, [n]). Consequently, (qs)t = qs for every $t \in S$ and thus s is a left zero.

Lemma 4.5. LeftZero is in NL.

Proof. By Lemma 4.4, we need only verify that for each $q \in [n]$, there is an $s \in S$ such that $qs \in \text{Fix}(S, [n])$. Iterate through $q \in [n]$, initializing p := q. Then, repeatedly guess generators $a_i \in \{a_1, \ldots, a_k\}$ and let $p = pa_i$. Non-deterministically stop guessing generators and check that $p \in \text{Fix}(S, [n])$ by iterating through $i \in [k]$ and verifying that $pa_i = p$.

We also consider the problem of deciding whether a semigroup contains a zero. **Zero**

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Does $\langle a_1, \ldots, a_k \rangle$ have a zero?

We now prove that LeftZero, RightZero, and Zero are all NL-hard by a reduction of the following problem known to be NL-complete [7, Theorem 26].

DFAEmptiness

- Input: A deterministic finite automaton (DFA) \mathcal{A} over an alphabet Σ .
- Problem: Is there $w \in \Sigma^*$ that is accepted by \mathcal{A} ?

Theorem 4.6. LeftZero, RightZero, and Zero are NL-complete.

ON THE COMPLEXITY OF PROPERTIES OF TRANSFORMATION SEMIGROUPS 8

Proof. A semigroup contains a zero element iff it contains both a left zero and a right zero. Then Zero is in NL as a corollary to Lemmas 4.5 and 4.3. It remains to prove that all three problems are NL-hard.

Given a DFA with state set [n], initial state $q_o \in [n]$, accepting states $F \subseteq [n]$, and transformations $a_1, \ldots, a_k \in T_n$, we construct a subsemigroup of T_{n+1} as follows. First extend a_1, \ldots, a_k by defining them to fix n+1. Then define the following new transformations:

$$xb := \begin{cases} q_0 & \text{if } q \in [n], \\ n+1 & \text{if } q = n+1. \end{cases} \qquad xc := \begin{cases} q & \text{if } q \in [n+1] \setminus F, \\ n+1 & \text{if } q \in F \text{ or } q = n+1. \end{cases}$$

We claim $S = \langle a_1, \ldots, a_k, b, c \rangle$ has a zero iff the language of the DFA is nonempty; that is, there exists $w \in \{a_1, \ldots, a_k\}^*$ such that $q_0 w \in F$.

Assume there exists $w \in \{a_1, \ldots, a_k\}^*$ such that $q_0 w \in F$. Then [n+1]bwc = $\{n+1\}$ and thus S has a zero element. Conversely, assume S has a zero element $0 \in S$. Since n+1 is the only point fixed by all transformations and $(q_0)s = q_00$ for every $s \in S$, then $q_0 = n + 1$. The only transformation that sends something from [n] to n+1 is c, which only sends elements of F to n+1. So, $0 = w_1 c w_2$ where $w_1, w_2 \in S, q_0 w_1 \in F$, and $w_1 \in a_1, \ldots, a_k, b^*$. Because $[n]b = \{q_0\}$, then either (case 1) $w_1 \in \{a_1, \ldots, a_k\}^*$ or (case 2) $w_1 = w_3 b w_4$ where $w_4 \in \{a_1, \ldots, a_k\}^*$ and $q_0w_4 \in F$. In either case, we have a word from $\{a_1, \ldots, a_k\}^*$ that is accepted by the DFA.

Finally, note that $\Gamma(\{a_1, \ldots, a_k, b, c\}, [n+1])$ has a single connected component. By Lemma 4.1, any right zero $r \in S$ must have an image size of a single point. Since $Fix(S, [n+1]) = \{n+1\}$, the only right zero of S would be the zero element, if it exists. Similarly, since the images of left zeroes are fixed points, the only left zero of S would be the zero element, if it exists. \square

We define the following problem and show that it is in NL:

NilpotentSemigroup

- Input: a₁,..., a_k ∈ T_n
 Problem: Is ⟨a₁,..., a_k⟩ nilpotent?

Lemma 4.7. Let $S := \langle a_1, \ldots, a_k \rangle \leq T_n$. The following properties are equivalent:

- (1) S is nilpotent;
- (2) S has a zero element, 0, and $\Gamma(\{a_1, \ldots, a_k\}, [n] \setminus [n]0)$ is acyclic;
- (3) S is *n*-nilpotent.

Proof. We prove $(1) \Rightarrow (2)$ by contraposition. Certainly, if S does not have a zero element, it is not nilpotent by definition. Now assume there is a zero element, 0, but also a cycle in $\Gamma(\{a_1,\ldots,a_k\},[n]\setminus[n])$. Let the cycle be (q_1,\ldots,q_i,q_1) . Then there are generators $b_1, \ldots, b_i \in \{a_1, \ldots, a_k\}$ such that $q_j b_j = q_{j+1}$ for $1 \leq j < i$ and $q_i b_i = q_1$. Let $s = b_1 \cdots b_i$ and note that $q_1 s = q_1$ and that $q_1 \notin [n]0$. So, $q_1 s^d = q_1 \neq q_1 0$ for any $d \in \mathbb{N}$.

For (2) \Rightarrow (3), assume S has a zero element, 0, and assume that the graph $\Gamma(\{a_1,\ldots,a_k\},[n]\setminus[n]0)$ is acyclic. By Lemma 4.1, the connected components of $G := \Gamma(\{a_1, \ldots, a_k\}, [n])$ are formed by the vertex sets $S^{-1}(q)$ for $q \in [n]0$. Hence the only cycles in G are the loops around each vertex in [n]0. Let d be the diameter of G; that is, the length of the longest path from any $p \in [n]$ to p0. Clearly $d \leq n$. We claim that S is nilpotent of degree d, and hence n-nilpotent. Pick any $b_1, \ldots, b_d \in \{a_1, \ldots, a_k\}$. For each $1 \leq i \leq d$, let $t_i := b_1 \ldots b_i$. For each $p \in [n]$ the directed path (p, pt_1, \ldots, pt_d) has d vertices. Since d is the diameter, the final vertex must be p0. Thus $t_d = 0$ and $S^d = \{0\}$.

Certainly, $(3) \Rightarrow (1)$.

9

We are now able to describe an NL algorithm for testing nilpotency.

Lemma 4.8. NilpotentSemigroup is in NL.

Proof. By Lemma 4.7, S is nilpotent if and only if $x_1 \cdots x_n = x_{n+1} \cdots x_{2n+1}$ for all $x_1, \ldots, x_{2n+1} \in S$. Since NL is closed under complementation, it suffices to show that we can decide in non-deterministic logarithmic space whether there exist $x_1, \ldots, x_{2n+1} \in S$ with $x_1 \cdots x_n \neq x_{n+1} \cdots x_{2n+1}$.

To this end, we first guess an integer $q \in [n]$ and let p := q. Then, in a loop, we guess an integer $i \in [k]$ and let $q := qa_i$. This loop is repeated n times in total. In another loop that is iterated n + 1 times, we repeatedly guess an integer $i \in [k]$ and let $p := qa_i$. Finally, we verify that $p \neq q$.

To show that NilpotentSemigroup is NL-complete, we will use the following lemma. The statement is slightly more general than needed. This allows us to reuse the lemma in the next section.

Lemma 4.9. There exists a log-space transducer which, given a DFA A, produces a transformation semigroup S given by generators with the following properties:

- (1) If $L(\mathcal{A}) \neq \emptyset$, then there exists an idempotent in S that is not a left zero.
- (2) If $L(\mathcal{A}) = \emptyset$, then every square in S is zero.

Proof. Suppose the input consists of a DFA with state set [n], initial state $q_0 \in [n]$, accepting states $F \subseteq [n]$ and transformations $a_1, \ldots, a_k \in T_n$. We assume without loss of generality that $q_0 \notin F$. We define n^2 transformations $a_{i,j}$ with $i, j \in [n]$ on $[n] \times [n] \cup \{0\}$ by setting $0a_{i,j} = 0$ and

$$(q,k)a_{i,j} := \begin{cases} (qa_i, k+1) & \text{if } j = k < n \text{ and } q \notin F, \\ 0 & \text{otherwise} \end{cases}$$

for all $q, k, i, j \in [n]$. We define an additional transformation b by 0b = 0 and

$$(q,k)b := \begin{cases} (q_0,1) & \text{if } q \in F, \\ 0 & \text{otherwise} \end{cases}$$

for all $q, k \in [n]$. We now show that $S = \langle a_{i,j}, b \mid i, j \in [n] \rangle$ satisfies the two properties stated in the lemma.

Suppose that the language accepted by the DFA is non-empty, i.e., there exists a word $u \in \{a_1, \ldots, a_k\}^*$ with $q_0 u \in F$. Without loss of generality, we may assume that |u| < n, i.e., $u = a_{i_1} \ldots a_{i_\ell}$ for some $\ell \in [n-1]$ and $i_1, \ldots, i_\ell \in [k]$. Let $v = a_{i_1,1} \ldots a_{i_\ell,\ell}b$. Note that b^2 is the transformation that maps every element to 0. It is easy to verify that $v^2 = v$. By construction, we have $(q_0, 1)v = (q_0, 1)$ and $(q_0, 1)vb^2 = 0$. Thus, $vb^2 \neq v$ and v is not a left zero.

Conversely, it is easy to see that if the language accepted by the DFA is empty, then $s^2 = b^2$ for every $s \in S$. As mentioned before, b^2 is a zero element.

We conclude by proving that testing nilpotency is NL-complete.

Theorem 4.10. NilpotentSemigroup is NL-complete.

Proof. Lemma 4.8 shows that NilpotentSemigroup belongs to NL. Lemma 4.9 immediately yields a reduction from DFAEmptiness to NilpotentSemigroup. If case (1) from Lemma 4.9 holds, then let e be the idempotent that is not a left zero and note that $e^d \neq 0$ for any $d \in \mathbb{N}$. Thus, the semigroup is not nilpotent. If case (2) holds, then the only idempotent element is the zero element. Since every element has an idempotent power and S has finitely many elements, then S is nilpotent. \Box

We will now show that the following problem is NL-complete.

\mathcal{R} -trivial

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Is $\langle a_1, \ldots, a_k \rangle$ *R*-trivial?

Lemma 4.11. Let $a_1, \ldots, a_k \in T_n$. Then, $S := \langle a_1, \ldots, a_k \rangle$ is \mathcal{R} -trivial if and only if each cycle in $\Gamma(\{a_1, \ldots, a_k\}, [n])$ is of length 1.

Proof. For the implication from left to right, suppose that $\Gamma(\{a_1, \ldots, a_k\}, [n])$ contains a cycle longer than length 1. Then, there exist $q \in Q$ and $i_1, \ldots, i_{\ell} \in [k]$ such that $q \cdot a_{i_1} \neq q$ and $q \cdot a_{i_1} \cdots a_{i_{\ell}} = q$. Let $e := (a_{i_1} \cdots a_{i_{\ell}})^{\omega}$. Since $q \cdot e = q$ and $q \cdot ea_{i_1} \neq q$, the elements e and ea_{i_1} are distinct. But $ea_{i_1}a_{i_2}\cdots a_{i_{\ell}}(a_{i_1}\cdots a_{i_{\ell}})^{\omega-1} = e^2 = e$, thus $eS^1 = ea_{i_1}S^1$.

For the converse direction, suppose $sS^1 = tS^1$ and $s \neq t$. Then there exists $u, v \in S^1$ and $q \in [n]$ such that su = t, tv = s, and $qs \neq qt$. Then qs = qsuv and $qs \neq qt = qsu$, yielding a cycle longer than length 1.

Lemma 4.12. There exists a log-space transducer which, given a directed graph G = (V, E), produces a transformation semigroup given by generators with the following properties:

- (1) If the graph is acyclic, then the semigroup is nilpotent.
- (2) If the graph is not acyclic, then the semigroup is not \mathcal{R} -trivial and there exists an idempotent that is not central.

Proof. The underlying set is $Q = V \cup \{0\}$ and we add one generator $a_{(v,w)}$ for each edge $(v,w) \in E$ which we define by $va_{(v,w)} = w$ and $ua_{(v,w)} = 0$ for all $u \neq v$.

Clearly, if G is acyclic, any product of more than |V| generators is the zero element, i.e., the transformation that maps each point to 0. In this case, the semigroup is nilpotent.

Conversely, if G contains a cycle v_0, \ldots, v_ℓ with $(v_{i-1}, v_i) \in E$ for all $i \in [\ell]$ and with $v_0 = v_\ell$, then the element $s = a_{(v_0, v_1)} \cdots a_{(v_{\ell-1}, v_\ell)}$ is the identity on v_0 and maps all other points to 0. Therefore, s is idempotent. Note that $a_{(v_0, v_1)}s$ is the zero element but $sa_{(v_0, v_1)}$ is not which means that s is not central. Moreover, the sequence of generators defining s forms a cycle at v_0 that is longer than length 1. Thus, the semigroup is not \mathcal{R} -trivial by Lemma 4.11.

Theorem 4.13. *R*-trivial is NL-complete.

Proof. The graph $\Gamma(\{a_1, \ldots, a_k\}, [n])$ can be computed by a log-space transducer. In another log-space transduction, we remove all cycles of length 1. We then use the NL algorithm for graph acyclicity. So, \mathcal{R} -trivial is in NL by Lemma 4.11. For NL-hardness, note that nilpotent semigroups are \mathcal{R} -trivial and thus Lemma 4.12 reduces the NL-complete problem of directed graph acyclicity to \mathcal{R} -trivial. \Box

5. Semigroup Equations

Let $X = \{x_1, \ldots, x_m\}$ be a nonempty set of variables. Let X^+ denote the set of all nonempty words over X, and let u and v be two fixed nonempty words over X. We say that a semigroup S models u = v if h(u) = h(v) holds for all homomorphisms $h: X^+ \to S$. For a fixed identity u = v, define the following problem:

Model(u = v)

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Does $\langle a_1, \ldots, a_k \rangle$ model u = v?

This problem is the dual of the well-known identity checking problem for a fixed semigroup S: given words u, v as input, decide whether S models u = v. See [1] for background and complexity results on identity checking; in particular, examples of semigroups for which it is coNP-complete. In contrast, we will prove that Model(u = v) is in NL for any fixed identity u = v and NL-complete for specific identities. Before describing the algorithm, we will consider two generalizations of this problem.

So called ω -identities are often used to define varieties of finite semigroups. In addition to concatenating variables, they also allow for using the ω -operator to take idempotent powers. However, testing whether a semigroup models a fixed ω -identity is PSPACE-complete for very simple identities already: the ω -identity $x^{\omega}x = x^{\omega}$ defines the class of finite aperiodic semigroups and testing aperiodicity of transformation semigroups is known to be PSPACE-complete [3]. Part of the hardness of this problem results from the fact that the same variable can appear both inside and outside the scope of an ω -operator. We will prove that the complexity drops to NL if we disallow such occurrences of variables as well as nesting of ω -operators. This variant of ω -identities will be called *quasi-identities* and is formalized using slightly different notation below.

For containment in NL, we will actually prove that the following generalization is in NL. A semigroup S models $x_1 = x_1^2, \ldots, x_s = x_s^2 \Rightarrow u = v$ if for all homomorphisms $h: X^+ \to S$ with $h(x_1), \ldots, h(x_s)$ idempotent, we have h(u) = h(v).

 $\mathbf{Model}(x_1 = x_1^2, \dots, x_s = x_s^2 \Rightarrow u = v)$

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Does $\langle a_1, \ldots, a_k \rangle$ model $x_1 = x_1^2, \ldots, x_s = x_s^2 \Rightarrow u = v$?

Theorem 5.1. Let $X = \{x_1, \ldots, x_m\}$ be a nonempty finite set of variables. Let $e \in \{0, \ldots, m\}$ and let $u, v \in X^+$. Then, $Model(x_1 = x_1^2, \ldots, x_e = x_e^2 \Rightarrow u = v)$ belongs to NL.

Proof. We describe an NL algorithm to test whether a semigroup $S = \langle a_1, \ldots, a_k \rangle$ does not satisfy a fixed quasi-identity

$$x_1 = x_1^2, \dots, x_e = x_e^2 \Rightarrow x_{i_1} \cdots x_{i_\ell} = x_{j_1} \cdots x_{j_r}$$

where $X = \{x_1, \ldots, x_m\}$ is a fixed set of variables, $i_1, \ldots, i_\ell, j_1, \ldots, j_r \in [m]$, and $0 \le e \le m$. Since NL is closed under complementation, this implies that the decision problem $\operatorname{Model}(x_1 = x_1^2, \ldots, x_e = x_e^2 \Rightarrow u = v)$ belongs to NL. For each $i \in [m]$, we let $P_1(i) = \{p \in [\ell] \mid i_p = i\}$ and $P_2(i) = \{p \in [r] \mid j_p = i\}$. The algorithm is depicted in Algorithm 1. Since $\ell + r$ is a constant, the algorithm only requires logarithmic space.

Algorithm 1 coNL algorithm for $Model(x_1 = x_1^2, \dots, x_e = x_e^2 \Rightarrow u = v)$

Input: $a_1, \ldots, a_k \in T_n$ **Output:** Does $\langle a_1, \ldots, a_k \rangle$ not model $x_1 = x_1^2, \ldots, x_e = x_e^2 \Rightarrow u = v$? 1: guess integers $p_1, ..., p_{\ell+1}, q_1, ..., q_{r+1} \in [n]$ 2: if $p_1 \neq q_1$ or $p_{\ell+1} = q_{r+1}$ then reject end if 3: for all $i \in [m]$ do for all $j \in [\ell]$ do $p'_j := p_j, p''_j := p_{j+1}$ end for for all $j \in [r]$ do $q'_j := q_j, q''_j := q_{j+1}$ end for 4: 5:6: guess $c \in [k]$ 7: for all $j \in P_1(i)$ do $p'_j := p'_j a_c$, $p''_j := p''_j a_c$ end for for all $j \in P_2(i)$ do $q'_j := q'_j a_c$, $q''_j := q''_j a_c$ end for until $\forall j \in P_1(i) : p'_j = p_{j+1}$ and $\forall j \in P_2(i) : q'_j = q_{j+1}$ and $i \in [e] \Rightarrow (\forall j \in P_1(i) : p''_j = p_{j+1}$ and $\forall j \in P_2(i) : q''_j = q_{j+1})$ 8: 9: 10: 11: 12: end for 13: accept

The process corresponds to nondeterministically replacing each variable x_i by an element of S such that the left-hand side and the right-hand side of the equation map the point $p_1 = q_1 \in [n]$ to distinct points $p_{\ell+1}, q_{r+1} \in [n]$. The variables p''_i and q''_i ensure that the substitutions can be performed in a way such that all variables x_i with $i \leq e$ are substituted by idempotent elements. A formal correctness proof follows.

First, suppose that the input $S = \langle a_1, \ldots, a_k \rangle$ does not model the quasi-identity $x_1 = x_1^2, \ldots, x_e = x_e^2 \to x_{i_1} \cdots x_{i_\ell} = x_{j_1} \cdots x_{j_r}$. This means that there are elements $s_1, \ldots, s_m \in S$ such that $s_1 = s_1^2, \ldots, s_e = s_e^2$ and $s_{i_1} \cdots s_{i_\ell} \neq s_{j_1} \cdots s_{j_r}$. Pick a $p_1 \in [n]$ such that $p_1 s_{i_1} \cdots s_{i_\ell} \neq p_1 s_{j_1} \cdots s_{j_r}$. Let $q_1 := p_1$. For each $f \in [\ell]$, let $p_f = p_1 s_{i_1} \cdots s_{i_{f-1}}$.

To verify that the algorithm will accept the input, consider any $s_i \in \{s_1, \ldots, s_m\}$. Let $s_i = a_{c_1} \cdots a_{c_g}$ with $c_1, \ldots, c_g \in [k]$. Lines 6–10 will successively guess the generators and transform p'_j for each $j \in P_1(i)$; likewise for q'_i . When this loop completes, the algorithm will have transformed each p'_j and q'_j to $p_j s_i$ and $q_j s_i$, respectively. Consequently, $p'_j = p_j s_i = p_{j+1}$ for each $j \in P_1(i)$ and $q'_j = q_j s_i = q_{j+1}$ for each $j \in P_2(i)$. Furthermore, this loop transforms p''_j and q''_j , which tracks how the generators act on p_{j+1} and q_{j+1} . By the end, $p''_j = p_{j+1}s_i$ and $q''_j = q_{j+1}s_i$. For each $j \in P_1(i) \cap [e]$, s_i will be idempotent so that $p''_j = p_{j+1}s_i = (p_j s_i)s_i = p_j s_i = p_{j+1}$. Likewise, $q''_j = q_{j+1}$ for each $j \in P_2(i) \cap [e]$.

We now prove that if the algorithm accepts, then $S = \langle a_1, \ldots, a_k \rangle$ does not model $x_1 = x_1^2, \ldots, x_e = x_e^2 \Rightarrow u = v$. Let $p_1, \ldots, p_{\ell+1}, q_1, \ldots, q_{r+1}$ be the guessed integers in Line 1. For each $i \in [m]$, let $s_i = a_{c_1} \cdots a_{c_g}$ be the sequence of guessed generators in Line 7. Then for each $j \in P_1(i)$, $p_j s_i = p_{j+1}$ and for each $j \in P_2(i)$, $q_j s_i = q_{j+1}$. Let s_i^{ω} be the idempotent power of s_i . Then for each $j \in P_1(i)$ with $j \leq e$, we have $p_{j+1} s_i^{\omega} = p_{j+1} s_i = p_{j+1}$ and for each $j \in P_2(i)$ with $j \leq e$, we have $q_{j+1} s_i^{\omega} = q_{j+1} s_i = q_{j+1}$.

By the definitions of $P_1(i)$ and $P_2(i)$, this demonstrates that $p_1h(u) = p_{\ell+1}$ and $q_1h(v) = q_{r+1}$ where $h: X^+ \to S$ is the homomorphism defined by $h(x_i) = s_i^{\omega}$ for

all $i \in [e]$ and $h(x_i) = s_i$ for all $i \in \{e + 1, \dots, m\}$. By Line 2 of the algorithm, we obtain $h(u) \neq h(v)$, thereby concluding the proof.

Theorem 5.1 immediately yields the following.

Corollary 5.2. Given generators $a_1, \ldots, a_k \in T_n$, there are NL algorithms to check the following properties of $S = \langle a_1, \ldots, a_k \rangle$:

- (1) all elements in S are idempotent;
- (2) all idempotents are central in S;
- (3) all idempotents commute;
- (4) the product of any two idempotents is idempotent.

We now give lower bounds for $Model(x_1 = x_1^2, \ldots, x_s = x_s^2 \Rightarrow u = v)$. We certainly cannot state such a result without restricting the class of considered equations. For example, Model(x = x) is certainly not complete for any meaningful complexity class. A transformation semigroup is commutative if and only if all generators commute, which implies $Model(xy = yx) \in AC^0$. Another interesting setting is the class of finite groups which is defined by the ω -identities $x^{\omega}y = y = yx^{\omega}$. While the identities look similar to the identity for aperiodicity, no variables appear both inside and outside the scope of an ω -operator, so the problem is in NL by Theorem 5.1. And using a different approach mentioned in Section 1, it is easy to see that deciding whether a transformation semigroup is a group is actually in AC^0 .

At the same time, Model(u = v) is NL-complete for some equations. The following two theorems are direct consequences of Lemma 4.9 and Lemma 4.12.

Theorem 5.3. Model $(x^2y = x^2)$ is NL-complete.

Proof. Given a DFA \mathcal{A} , Lemma 4.9 yields a transformation semigroup S that reduces DFAEmptiness to Model($x^2y = x^2$). If $L(\mathcal{A}) = \emptyset$, then S satisfies $x^2y = x^2$. If $L(\mathcal{A}) \neq \emptyset$, then there is an idempotent $e \in S$ and some $s \in S$ such that $es \neq e$. Then, $e^2s = es \neq e = e^2$.

Theorem 5.4. Testing whether all idempotents are central is NL-complete.

Proof. Given a directed graph G, Lemma 4.12 yields a transformation semigroup Sthat reduces the directed acyclic graph problem to testing whether all idempotents are central. If G is not acyclic, S has a non-central idempotent. If G is acyclic, S is nilpotent and thus has only one idempotent; its zero, which is central.

This suggests that a classification of the exact complexity of testing whether a given semigroup satisfies a fixed ω -identity (in terms of the structure of the considered identity) is hard.

We return to a problem mentioned at the end of Section 7: determining if a commutative semigroup is regular. We show this is in NL by proving a harder result; determining whether a semigroup is completely regular.

CompletelyRegular

- Input: a₁,..., a_k ∈ T_n
 Problem: Is ⟨a₁,..., a_k⟩ completely regular?

Lemma 5.5. A transformation $s \in T_n$ generates a subgroup iff $s|_{[n]s}$ is a permutation.

14 ON THE COMPLEXITY OF PROPERTIES OF TRANSFORMATION SEMIGROUPS

Proof. If s permutes [n]s, then $\langle s \rangle$ is isomorphic to a subgroup of the symmetric group over |[n]s| elements.

Conversely, let $s \in T_n$ generate a subgroup. Then $s = s^k$ for some natural number k > 1. Pick any $p, q \in [n]$ such that $ps \neq qs$. Because $ps^k \neq qs^k$, then $ps^2 \neq qs^2$. Thus, $ps \neq qs$ implies $ps^2 \neq qs^2$. That is, s permutes [n]s. \Box

Theorem 5.6. CompletelyRegular is in NL.

Proof. Let $S := \langle a_1, \ldots, a_k \rangle \leq T_n$. By Lemma 5.5, this semigroup is completely regular if and only if the restriction of each $s \in S$ to its image is a permutation. Since NL is closed under complementation, we only need to give an NL algorithm for the complement of CompletelyRegular. Thus, it suffices to verify that there exists some element of S that is not a permutation on its image.

Corollary 5.7. Deciding whether a commutative semigroup given by generators is regular is in NL.

Proof. A commutative semigroup is regular iff it is completely regular by [5, p.107]. So, this follows directly from Theorem 5.6.

Thus, Theorems 5.6 and 5.2 yield:

Corollary 5.8. Given generators $a_1, \ldots, a_k \in T_n$, there is an NL algorithm to check whether $\langle a_1, \ldots, a_k \rangle$ is a Clifford semigroup.

6. Polynomial Time Problems

We will prove that the following problem can be executed in polynomial time.

LeftIdentities

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Enumerate the left identities of $\langle a_1, \ldots, a_k \rangle$.

Lemma 6.1. Let $k, n \in \mathbb{N}$, let $a_1, \ldots, a_k \in T_n$, and $S := \langle a_1, \ldots, a_k \rangle$. Then an element $\ell \in S$ is a left identity of S iff there is an $i \in [k]$ such that $\overline{a_i}$ permutes $[n]/\ker(S)$ and $\ell = a_i^{\omega}$.

Proof. For the backward direction, let $\overline{a_i}$ permute $[n]/\ker(S)$. Then $\overline{a_i}^{\omega}$ is the identity map on $[n]/\ker(S)$. That is, for any $q \in [n]$, we have that $[\![qa_i^{\omega}]\!] = [\![q]\!]\overline{a_i}^{\omega} = [\![q]\!]$. Thus, $qa_i^{\omega}s = qs$ for every $s \in S$, making a_i^{ω} a left identity of S.

For the forward direction, let ℓ be a left identity of S. Then $q\ell s = qs$ for every $q \in [n]$ and every $s \in S$, meaning $[\![q\ell]\!] = [\![q]\!]$. Thus, $\overline{\ell}$ is the identity map on $[n]/\ker(S)$. Consequently, $\ell = sa_i$ for some generator a_i where $\overline{a_i}$ is a permutation of $[n]/\ker(S)$. As proved in the paragraph above, $\overline{a_i}^{\omega}$ is also the identity map on $[n]/\ker(S)$ and thus $\overline{sa_i} = \overline{a_i}^{\omega}$. Then the following holds for every $q \in [n]$:

$$\llbracket qs \rrbracket \overline{a_i} = \llbracket q \rrbracket \overline{sa_i} = \llbracket q \rrbracket \overline{a_i}^{\omega} = \llbracket qa_i^{\omega-1} \rrbracket \overline{a_i}.$$

Since $\overline{a_i}$ is a permutation of $[n]/\ker(S)$, then $[\![qs]\!] = [\![qa_i^{\omega-1}]\!]$ and thus $qst = qa_i^{\omega-1}t$ for every $t \in S$. In particular, $qsa_i = qa_i^{\omega}$, proving $\ell = a_i^{\omega}$.

Theorem 6.2. LeftIdentities can be computed in polynomial time.

Proof. By Lemma 6.1, we need only produce the idempotent powers of generators that induce permutations of $[n]/\ker(S)$. Certainly, we can produce $[n]/\ker(S)$ in polynomial time. Let a_i be a generator such that $\overline{a_i}$ is a permutation of $[n]/\ker(S)$. Represent $\overline{a_i}$ in cycle notation and let m be the least common multiple of the lengths of the cycles. Then $\overline{a_i}^m$ is idempotent and we claim that a_i^m is idempotent as well. Pick any $q \in [n]$. Because $\overline{a_i}^m$ is idempotent:

$$\llbracket qa_i^{m-1} \rrbracket \overline{a_i} = \llbracket q \rrbracket \overline{a_i}^m = \llbracket q \rrbracket \overline{a_i}^{2m} = \llbracket qa_i^{2m-1} \rrbracket \overline{a_i}.$$

Because $\overline{a_i}$ is a permutation, then $[\![qa_i^{m-1}]\!] = [\![qa_i^{2m-1}]\!]$. That is, $qa_i^{m-1}s = qa_i^{2m-1}s$ for every $s \in S$. In particular, $qa_i^m = qa_i^{2m}$ and thus a_i^m is idempotent. Because we can find the idempotent power of a_i in polynomial time, we can compute all left identities in polynomial time.

We will prove that the following problem can be executed in polynomial time.

RightIdentities

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Enumerate the right identities of $\langle a_1, \ldots, a_k \rangle$.

Note that $r \in S$ is a right identity of S iff \tilde{r} is the identity of \tilde{S} , since for any $q \in [n]$ and any $s \in S$, qsr = qs iff $qs\tilde{r} = qs$.

Lemma 6.3. Let $k, n \in \mathbb{N}$, let $a_1, \ldots, a_k \in T_n$, and $S := \langle a_1, \ldots, a_k \rangle$. Then an element $r \in S$ is a right identity of S iff there is an $i \in [k]$ such that \tilde{a}_i permutes [n]S and r equals the idempotent power of a_i .

Proof. For the backward direction, let \tilde{a}_i permute [n]S. Then \tilde{a}_i^{ω} is the identity map of \tilde{S} . Then, $qsa_i^{\omega} = qs$ for every $q \in [n]$ and every $s \in S$, making a_i^{ω} a right identity of S.

For the forward direction, let r be a right identity of S. Then qsr = qs for every $q \in [n]$ and every $s \in S$, making \tilde{r} the identity map on [n]S. Then, $r = a_i b$ for some generator a_i where \tilde{a}_i is a permutation of [n]S. As proved in the paragraph above, \tilde{a}_i^{ω} is also the identity map of [n]S and thus $\tilde{r} = \tilde{a}_i^{\omega}$. Then the following holds for every $q \in [n]$:

$$qa_ib = qa_i\widetilde{a_i}^{\omega}b = qa_i^{\omega}a_ib = qa_i^{\omega}$$

Therefore, r equals the idempotent power of a_i .

Theorem 6.4. RightIdentities can be computed in polynomial time.

Proof. By Lemma 6.3, we need only produce the idempotent powers of generators that induce permutations of [n]S. Note that $[n]S = \bigcup_{i \in [k]} [n]a_i$ can be produced in polynomial time. Let a_i be a generator such that \tilde{a}_i is a permutation of [n]S. Represent \tilde{a}_i in cycle notation and let m be the least common multiple of the lengths of the cycles. Then \tilde{a}_i^m is idempotent and we claim that a_i^m is idempotent as well. Because \tilde{a}_i^m is the identity map of [n]S, then for any $q \in [n]$, we have that $qa_i^m = qa_i^m \tilde{a}_i^m = qa_i^{2m}$. Thus, a_i^m is idempotent. Since we found a_i and m in polynomial time, then we can compute all right identities in polynomial time. \Box

7. Global Properties

In this section, we consider properties of a transformation semigroup S that cannot be checked locally in the sense of Section 4. We now define the following four problems:

RegularElement

- Input: $a_1, \ldots, a_k \in T_n$
- Problem: Is there $s \in \langle a_1, \ldots, a_k \rangle$ such that $a_k s a_k = a_k$?

GeneralizedRegular

- Input: $a_1, \ldots, a_k, s \in T_n$
- Problem: Is there $t \in \langle a_1, \ldots, a_k \rangle$ such that sts = s?

GeneralizedWeakInverse

- Input: $a_1, \ldots, a_k, s \in T_n$
- Problem: Is there $t \in \langle a_1, \ldots, a_k \rangle$ such that tst = t?

GeneralizedInverse

- Input: $a_1, \ldots, a_k, s \in T_n$
- Problem: Is there $t \in \langle a_1, \ldots, a_k \rangle$ that is an inverse of s?

We now prove that all four of these problems are PSPACE-complete. We begin by showing that RegularElement and GeneralizedWeakInverse are PSPACE-hard by adapting a proof by Christian Brandl and Hans Ulrich Simon [2]. We reduce the following problem known to be PSPACE-complete [8].

DFAIntersection

- Input: Deterministic finite automata (DFA) $\mathcal{A}_1, \ldots, \mathcal{A}_k$ over a shared alphabet Σ , each with a unique final state.
- Problem: Is there $w \in \Sigma^*$ that sends each initial state to its corresponding final state?

Lemma 7.1. RegularElement and GeneralizedWeakInverse are PSPACE-hard.

Proof. Let $\mathcal{A}_1, \ldots, \mathcal{A}_k$ be DFAs with corresponding sets of states Q_1, \ldots, Q_k , initial states p_1, \ldots, p_k , final states q_1, \ldots, q_k , and a common alphabet $\Sigma = \{a_1, \ldots, a_m\}$. We assume without loss of generality that the sets of states are pairwise disjoint, i.e., $Q_i \cap Q_j = \emptyset$ for $1 \leq i < j \leq k$. Let

$$Q := \{0\} \cup \bigcup_{i=1}^k Q_i$$

be the disjoint union of the DFA states along with a new state, denoted by 0. Extend the action of Σ to transformations on Q by defining $0a_i = 0$ for each $i \in \{1, \ldots, m\}$. Define the following additional transformation:

$$qb := \begin{cases} p_j & \text{if } q = q_j, \\ 0 & \text{otherwise.} \end{cases}$$

We claim there is a word w in the intersection of the DFA languages iff b is regular in $S := \langle a_1, \ldots, a_m, b \rangle$. Assume there is a word w accepted by each $\mathcal{A}_1, \ldots, \mathcal{A}_k$ so that $p_j w = q_j$ for each $j \in [k]$. Then we can verify that qb = qbwb for all $q \in Q$. If q is not a final state, then qbwb = 0wb = 0b = 0 = qb. If $q = q_j$ for some $j \in [k]$, then $q_jbwb = p_jwb = q_jb$.

Now let b = bsb for some transformation $s \in S$. If $q = q_j$, then $q_jb = p_j$. So, $p_j = p_j sb$. Because b sends everything to 0 except for q_j and since 0t = 0 for any $t \in S$, then s must contain a word w that sends p_j to q_j . This is true for each $j \in [k]$ so that w must be in each of the DFA languages. Thus, the reduction is complete and RegularElement is PSPACE-hard.

We now reduce DFAIntersection to GeneralizedWeakInverse. Define:

$$qc := \begin{cases} s_j & \text{if } q \in Q_j, \\ 0 & \text{if } q = 0. \end{cases}$$

We claim there is a word w accepted by each $\mathcal{A}_1, \ldots, \mathcal{A}_k$ iff b has a weak inverse in $S := \langle a_1, \ldots, a_k, c \rangle$. Assume w is accepted by each $\mathcal{A}_1, \ldots, \mathcal{A}_k$. We claim cwis a weak inverse for b; that is, qcwbcw = qcw for each $q \in Q$. If $q \in Q_j$, then $qcwbcw = p_jwbcw = q_jbcw = p_jcw = p_jw = qcw$. And clearly 0cwbcw = 0 = 0cw. Conversely, assume there is a $t \in S$ satisfying tbt = t. Note that $p_jt \in Q_j$ for all $j \in [k]$. If $p_jt \neq q_j$, then $p_jtbt = 0t = 0 \notin Q_j$. Thus, $p_jt = q_j$. Because ccan only reset states in Q_j back to p_j , then t must end with a word w that sends p_j to q_j . Then w is accepted by $\mathcal{A}_1, \ldots, \mathcal{A}_k$. Thus, GeneralizedWeakInverse is PSPACE-hard. \Box

Lemma 7.2. RegularElement, GeneralizedRegular, GeneralizedWeakInverse, and GeneralizedInverse are in PSPACE.

Proof. The non-deterministic Algorithm 2 correctly decides GeneralizedInverse. Since it only requires space to store a single transformation c in T_n and compute products ca_i, bcb, cbc , Algorithm 2 requires working space $O(n \log(n))$. Hence, GeneralizedInverse is in NPSPACE, which is PSPACE by Savitch's Theorem.

Algorithm 2

Function GeneralizedInverse (a_1, \ldots, a_k, b) Input: $a_1, \ldots, a_k, s \in T_n$ Output: Is there $t \in \langle a_1, \ldots, a_k \rangle$ such that sts = s and tst = t? 1: guess $i \in [k]$ and let $t := a_i$ 2: while $sts \neq s$ or $tst \neq t$ do 3: guess $i \in [k]$ and let $t := ta_i$ 4: end while 5: accept

Straighforward adaptations of line 2 in Algorithm 2 yield non-deterministic polynomial space algorithms for the other three decision problems. Hence these problems are in PSPACE as well. $\hfill \Box$

Theorem 7.3. RegularElement, GeneralizedRegular, GeneralizedWeakInverse, and GeneralizedInverse are all PSPACE-complete.

Proof. Note that every regular element in a semigroup has an inverse in that semigroup [5, p.51]. So, RegularElement can be reduced to GeneralizedRegular and GeneralizedInverse by simply letting $s = a_k$. Because RegularElement is PSPACEhard by Lemma 7.1, then the other two are as well. And because GeneralizedRegular and GeneralizedInverse are in PSPACE by Lemma 7.2, then all three problems are PSPACE-complete. GeneralizedWeakInverse is PSPACE-complete since it is PSPACE-hard by Lemma 7.1 and it is in PSPACE by Lemma 7.2.

Note that we did not define the problem of determining if an element $s \in S$ has a weak inverse in S, because this is always true for any element of a finite semigroup: we have $s^{\omega-1}ss^{\omega-1} = s^{2\omega-1} = s^{\omega-1}$, so $s^{\omega-1}$ is a weak inverse of s.

We can show that the complexity of determining if a semigroup given by generators is regular is in PSPACE. Note that we can nondeterministically guess an element of S and, by Theorem 7.3, verify in polynomial time that it is not regular. Thus, this problem is in coNPSPACE, which, by Savitch's Theorem, is PSPACE. It is unknown whether checking semigroup regularity is PSPACE-hard.

8. Open Problems

The problems of testing whether a transformation semigroup is a band, has commuting idempotents, is orthodox, is completely regular or a Clifford semigroup were shown to be decidable in non-deterministic logarithmic space. The problems of testing whether a transformation semigroup is regular or inverse were shown to be decidable in polynomial space. Lower bounds for each of these problems are still open.

Acknowledgments

We would like to thank Peter Mayr and Nik Ruškuc for discussions on the material in this paper.

References

- Jorge Almeida, Mikhail Volkov, and S. V. Goldberg. Complexity of the identity checking problem for finite semigroups. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 358(Issledovaniya po Konstruktivnooĭ Matematike i Matematicheskoĭ Logike. XI):5– 22, 301, 2008.
- [2] Christian Brandl and Hans Ulrich Simon. Complexity analysis: transformation monoids of finite automata. In *Developments in language theory*, volume 9168 of *Lecture Notes in Comput. Sci.*, pages 143–154. Springer, Cham, 2015.
- [3] Sang Cho and Dũng T. Hunh. Finite-automaton aperiodicity is PSPACE-complete. Theoret. Comput. Sci., 88(1):99–116, 1991.
- [4] James East, Attila Egri-Nagy, James D. Mitchell, and Yann Presse. Computing finite semigroups. Journal of Symbolic Computation, 2018.
- [5] John M. Howie. Fundamentals of semigroup theory, volume 12 of London Mathematical Society Monographs. New Series. The Clarendon Press, Oxford University Press, New York, 1995. Oxford Science Publications.
- [6] Neil Immerman. Descriptive complexity. Graduate Texts in Computer Science. Springer-Verlag, New York, 1999.
- [7] Neil D. Jones. Space-bounded reducibility among combinatorial problems. J. Comput. System Sci., 11(1):68-85, 1975.
- [8] Dexter Kozen. Lower bounds for natural proof systems. IEEE Comput. Sci., Long Beach, Calif., 1977.
- [9] James D. Mitchell et al. Semigroups GAP package, Version 3.0.20, Oct 2018.
- [10] Omer Reingold. Undirected connectivity in log-space. J. ACM, 55(4):17:1–17:24, September 2008.