

# High-resolution optical spectroscopy of the post-AGB supergiant V340 Ser (=IRAS 17279–1119)

V.G. Klochkova, V.E. Panchuk, N.S. Tavalzhanskaya, and M.V. Yushkin  
e-mail: valenta@sao.ru

Special Astrophysical Observatory RAS, Nizhnij Arkhyz, 369167 Russia

October 16, 2020

**Abstract** Some evidences of wind variability and velocity stratification in the extended atmosphere has been found in the spectra of the supergiant V340 Ser (=IRAS 17279–1119) taken at the 6-m BTA telescope with a spectral resolution  $R \geq 60000$ . The  $H\alpha$  line has a P Cyg profile whose absorption component ( $V = +34$  km/s) is formed in the upper layers of the expanding atmosphere close to the circumstellar environment. For four dates the mean velocity has been derived from the positions of 300–550 symmetric metal absorptions with an accuracy better than  $\pm 0.1$  km/s:  $V_{\odot} = 59.30, 60.09, 58.46, \text{ and } 55.78$  km/s. A lot of low-excitation metal lines have an inverse P Cyg profile. The mean positions of their emission components,  $V_{\odot} = 46.3 \pm 0.4$  km/s, differ systematically from the velocity inferred from symmetric absorptions, suggesting the presence of a velocity gradient in the supergiant extended atmosphere. The multicomponent profile of the NaID lines contains the interstellar,  $V_{\odot} = -11.2$  km/s, and circumstellar,  $V_{\odot} = +10$  km/s, components and the component forming in the upper atmospheric layers,  $V_{\odot} = +34.0$  km/s. The mean velocity from 20–30 diffuse interstellar bands (DIBs) identified in the spectra,  $V_{\odot}(\text{DIBs}) = -11.6 \pm 0.2$  km/s, agrees with the velocity from interstellar NaI and KI components. The equivalent width of the oxygen triplet  $W(\text{OI } 7774) = 1.25 \text{ \AA}$  corresponds to an absolute magnitude of the star  $M_V \approx -4.6^m$ , which, taking into account the total (interstellar + circumstellar) extinction, leads to a distance to the star  $d \approx 2.3$  kpc.

**Key words.** stars, evolution, post-AGB stars, atmospheres, nucleosynthesis, envelopes, optical spectra.

## 1. INTRODUCTION

V340 Ser, the central star of the infrared (IR) source IRAS 17279–1119, was entered into a list of stars with IR excesses by Oudmaijer et al. (1992). Later, having studied the Hubble Space Telescope data for a large sample of IR sources, Siódmiak et al. (2008) showed IRAS 17279–1119 to have a point image and classified V340 Ser as a postasymptotic giant branch (hereafter post-AGB) star. These authors pointed out the presence of a near-IR excess, suggesting the presence

of warm circumstellar dust. Highly evolved stars with initial masses in the range  $2 \div 8 M_{\odot}$  are observed at the post-AGB stage.

According to Blöcker (1995a), at the preceding AGB evolutionary stage these stars are observed as red supergiants with effective temperatures  $T_{\text{eff}} \approx 3000 \div 4500$  K. The AGB stage for stars with the above masses is the final evolution phase with nucleosynthesis in the stellar cores. The interest in AGB stars and their closest descendants stems primarily from the fact that in the interiors of these stars, which are at a short evolutionary stage, there are physical conditions for the synthesis of heavy-metal nuclei and the dredge-up of nuclear reaction products into the stellar atmosphere and then into the circumstellar and interstellar medium. Due to these processes, AGB stars with initial masses below  $3 \div 4 M_{\odot}$  are the main suppliers of all the elements heavier than iron synthesized through the s-process, which basically consists in slow (compared to  $\beta$  decay) neutronization of nuclei. Fe nuclei serve as initial nuclei for the chain of s-process reactions. In stars with initial mass below  $3 \div 4 M_{\odot}$  the required neutron flux is provided by the  $^{13}\text{C}(\alpha, n)^{16}\text{O}$  reaction, while in the case of more massive stars, with initial mass above  $4 \div 5 M_{\odot}$ , the analogous reaction proceeds on  $^{22}\text{Ne}$  nuclei. It should be emphasized that these more massive AGB stars can also be the sources of lithium. The details of the evolution of stars near the AGB and the results of present-day calculations of the synthesis and dredge-up of elements are presented in Herwig (2005), di Criscienzo et al. (2016), and Liu et al. (2018).

The spectral type F2–3 II is given for the supergiant V340 Ser in the SIMBAD database. This supergiant lies outside the Galactic plane, which already suggests that it probably belongs to low mass evolved stars. By now the star has been studied by both photometric and spectroscopic methods. A number of authors (in particular, Kazarovets et al. 2000; de Smedt et al. 2016) classify V340 Ser as an RV Tau variable. The post-AGB stars of this category are located on the Hertzsprung–Russell diagram in (or near) the instability strip and have near-IR excesses, which serves as evidence for the presence of warm circumstellar dust that has not yet detached from the photospheric layers of the star.

The abundances of chemical elements in the atmosphere of V340 Ser have been studied repeatedly and quite reliably. Based on high-resolution spectra, but with a moderate S/N ratio, Arellano Ferro et al. (2001) derived the model atmosphere parameters for this star:  $T_{\text{eff}} = 7300$  K, surface gravity  $\log g = 1.25$ , and microturbulence  $\xi_t = 4.6$  km/s. They found a lowered metallicity,  $[\text{Fe}/\text{H}]_{\odot} = -0.60$ , an overabundance of heavy s-process elements, and a carbon-to-oxygen abundance ratio  $\text{C}/\text{O} \approx 1$ . As a result, these authors supported the previous conclusion (van Winckel, 1997) that the star belongs to the post-AGB stars that have passed the third dredge-up.

The paper by Arellano Ferro et al. (2001) is also important in that its authors pointed out the absence of selective depletion of chemical elements in the atmosphere of this star, which usually distorts the elemental abundances in the atmospheres of related post-AGB stars (for examples, see Klochkova (1995), Giridhar et al. (2000), Rao et al. (2012), Klochkova and Tavalzhanskaya (2019)).

Based on high-resolution spectra, Rao et al. (2012) determined the model parameters and detailed atmospheric chemical composition for a sample of post-AGB stars, including V340 Ser. The model parameters adopted by them for this star are, i.e.,  $T_{\text{eff}} = 7300 \pm 150$  K, surface gravity  $\log g = 2.25 \pm 0.25$ , and microturbulence  $\xi_t = 4.7 \pm 0.25$  km/s, agree, within the error limits, with the previously derived parameters from Arellano Ferro et al. (2001). At a lowered metallicity,  $[\text{Fe}/\text{H}]_{\odot} = -0.60$ , a nontrivial atmospheric chemical composition was found for V340 Ser: moderate, but significant enhancements of  $\alpha$ -process elements,  $[\alpha/\text{Fe}] = +0.44$ , and heavy s-process metals,  $[\text{s}/\text{Fe}] = +0.69$ .

Later, using VLT+UVES spectra, de Smedt et al. (2016) again confirmed the parameters and chemical abundances of V340 Ser. The main goal of their study, the search for evidence of lead in the atmospheres of the program stars, was not achieved, because for 14 post-AGB stars with heavy-metal-enriched atmospheres the authors obtained only upper limits for the abundance of this element. The paper of these authors is important in our study of the optical spectra for V340 Ser, because, having performed a spectroscopic monitoring with the HERMES spectrograph of the 1.2-m telescope, de Smedt et al. (2016) concluded that this star is a binary by determining its orbital parameters. The period found by them is  $P = 365$  days, the systemic velocity is  $\gamma = 56.8 \pm 0.5$  km/s, and the velocity amplitude is  $\Delta V_r = 7.92 \pm 0.44$  km/s.

The behavior of the photometric parameters for V340 Ser has also been studied reliably. Having performed a 7-year-long UBV monitoring of the star, Arkhipova et al. (2011) concluded that its brightness is variable with an amplitude  $\Delta V \approx 0.2^m$  typical of post-AGB stars and a fundamental period  $P = 89.6$  days. Appending the ASAS data distributed more uniformly in time to their analysis, Arkhipova et al. (2011) determined the amplitude,  $\Delta V \approx 0.3^m$ , and the fundamental period,  $P = 89.8$  days.

In this paper we present the results of our analysis of the optical spectra for V340 Ser taken at the 6 meter BTA telescope in 2018–2020. The main goal of our work is the search for spectral peculiarities and their variability with time. In Section 2 we briefly describe the methods of observations and data analysis. In Section 3 we present the results by comparing them with the previously published ones and in Section 3 we provide our conclusions.

## 2. OBSERVATIONS, DATA REDUCTION, AND ANALYSIS OF SPECTRA

**Table 1.** Results of our measurements of the heliocentric radial velocity  $V_\odot$  in the spectra of V340 Ser based on sets of lines of various types. The number of features of various types used is given in parentheses. The positions of the absorption components forming outside the stellar atmosphere are given for the NaI D lines  $V_\odot$ , km/s

JD −2458000	$V_\odot$ , km/s						
	Metal absorptions	Metal emissions	H $\alpha$ (abs) H $\alpha$ (em)	H $\beta$	NaI	KI	DIBs
213.5	59.30 $\pm 0.05$ (557)	45.0 $\pm 0.2$ (59)	34.2 101.7	53.6	−11.3, 10.0, 33.7	−9.2	−11.9 $\pm 0.2$ (23)
220.4	60.09 $\pm 0.05$ (459)	46.0 $\pm 0.3$ (43)	34.9 100.1	54.8	−11.2, 10.2, 33.4	−10.0	−11.3 $\pm 0.2$ (32)
574.4	58.46 $\pm 0.06$ (283)	48.0 $\pm 0.4$ (49)	34.5 101.2	53.9	−11.4, 10.3, 35.0	−10.7	−12.1 $\pm 0.3$ (18)
924.2	55.78 $\pm 0.06$ (408)	46.0 $\pm 0.3$ (28)	33.6 97.5	49.3	−11.1, 9.1, 32.8	−9.3	−11.3 $\pm 0.3$ (15)

The spectra of V340 Ser were taken with the Nasmyth echelle spectrograph (NES) (Panchuk et al. 2017) at the 6-m BTA telescope. The dates of observations of the star are listed in Table 1. At these dates the NES was equipped with a large format 4608×2048 CCD with a pixel size of 0.0135×0.0135 mm, the readout noise is 1.8 e<sup>−</sup>. The recorded spectral range is  $\Delta\lambda = 470$ –778 nm. To reduce the light losses without any loss of the spectral resolution, the NES is


equipped with an image slicer forming three slices. Each spectral order in the 2-dimensional image of the spectrum is repeated three times with a shift along the cross dispersion. The spectral resolution is  $\approx 60000$ , the signal-to-noise ratio is  $S/N > 100$ , it changes along the echelle order from 100 to 150.

The one-dimensional data were extracted from the 2-dimensional echelle spectra with the modified (taking into account the peculiarities of the NES echelle frames) ECHELLE context of the MIDAS software package. The details of the procedure were described by Yushkin and Klochkova (2005). The cosmic-ray particle hits were removed by a median averaging of two spectra taken successively one by one. The wavelength calibration was carried out using the spectra of a hollow-cathode Th-Ar lamp. The entire subsequent reduction, including the photometric and positional measurements, was performed with the up-to-date version of the DECH20t code developed by Galazutdinov (1992). Note that this code traditionally used by us to reduce the spectra allows the radial velocities for individual features in the line profiles to be measured. The systematic measurement errors of the heliocentric velocities  $V_{\odot}$  estimated from sharp interstellar Na I components do not exceed 0.25 km/s (from one line); the random errors for shallow absorptions are  $\approx 0.5$  km/s, the mean value per line. Thus, for the average values of  $V_{\odot}$  in Table 1 the random errors are  $0.05 \div 0.3$  km/s, depending on the number of measured lines. We identified the features in the spectrum of V340 Ser using the atlas of the optical spectrum for the canonical post-AGB star HD 56126 (=IRAS 07134+1005, Sp = F5 Iab) published by Klochkova et al. (2007a), which is based on the observational data of the 6-m BTA telescope in combination with the NES.

### 3. RESULTS

#### 3.1. Peculiarities of the optical spectrum for V340 Ser and radial velocity stratification

The spectral anomalies inherent in RV Tau stars are the emission components of the complex and variable (with phase)  $H\alpha$  profiles, the He I 5876 Å emission, the splitting of selected absorptions, the multicomponent profiles of the Na I D lines, and the weak metal emissions appearing at certain times of the cycle. The spectra of the prototype stars RV Tau, U Mon, and ACHer studied by Baird (1982, 1984), Pollard et al. (1997), Klochkova and Panchuk (1998), and Kipper and Klochkova (2013) serve as examples. Both the absorption splitting and the emission in the He I 5876 Å line are absent in our spectra of V340 Ser. The optical spectrum of this star, which largely corresponds to the expected spectrum of an F2–3 supergiant, contains some of the peculiarities listed above. First, the  $H\alpha$  profile is complex, which is typical of post-AGB stars. The profile of this line presented in Fig. 1 in relative intensity – radial velocity coordinates includes the broad absorption wings on which the absorption and emission components are superimposed. As follows from Fig. 1, the positions of both components do not change with time, but a weak variability of their intensity can be noted. The change in the shape of the blue wing of the absorption component is more pronounced, reflecting a change in the outflow velocity. A large difference between the velocities corresponding to the cores of the emission and absorption components in the  $H\alpha$  profile should be emphasized. As can be seen from the data in Table 1 and Fig. 1, this difference is about 65 km/s and exceeds the same parameter for the RV Tau stars studied by Pollard et al. (1997), including the star AD Aql, in the spectrum of which the  $H\alpha$  profile is close to the one recorded by us in the spectrum of V340 Ser, by many times. As can be seen from Fig. 2, the  $H\beta$  profile is a purely absorption one,




**Figure 1.**  $H\alpha$  profile in relative intensity – radial velocity coordinates in the spectra of V340 Ser taken at different dates: April 5, 2018 (black line), April 11, 2018 (green line), March 31, 2019 (red line), and April 2, 2020 (indigo line). The two solid vertical bars indicate the positions of the TiI 6554.23 Å and CaI 6572.80 Å emissions. The vertical line marks the systemic velocity  $\gamma = 56.8 \pm 0.5$  km/s from de Smedt et al. (2016).

it contains no visible emission features. As follows from Table 1, the position of the  $H\beta$  core does not coincide quite closely with the positions of the atmospheric metal absorptions, it is blueshifted by 5–6 km/s, suggesting a stratification of the velocity pattern.

The circumstellar envelopes of evolved stars are often the sources of molecular and maser emission, which allows their systemic velocity to be reliably recorded. However, for V340 Ser there is no information about any envelope features in the radio band. The optical spectrum of this star does not contain any evidence of molecules either and, therefore, we use  $\gamma = 56.8 \pm 0.5$  km/s from de Smedt et al. (2016) as the systemic velocity for this star.

Second, the  $H\alpha$  profile in the spectrum of V340 Ser in Fig. 1 is a P Cyg type one. This profile type is typical of high-luminosity stars with an extended and expanding atmosphere.  $H\alpha$  is formed in the upper layers of the extended stellar atmosphere. The intense absorption component results from the absorption of radiation in the region in front of the star. The position of the core of the absorption component (+34 km/s) points to its formation in the high layers of the expanding atmosphere close to the circumstellar medium. The velocity of about 34 km/s differs from the adopted systemic velocity, suggesting an expansion of the region in which this absorption is formed. Such a profile type is rarely encountered in the spectra of post-AGB stars. Numerous examples of the varieties of the  $H\alpha$  profile in the spectra of post-AGB stars are given by Klochkova (1997), Sanchez Contreras et al. (2008), and Molina et al. (2014): purely absorption profiles, complex P Cyg profiles, inverse P Cyg profiles, etc.

However, in this extensive collection of recorded profiles there is no profile similar to the  $H\alpha$  one in the spectra of V340 Ser. A variety of profile types is observed in the papers by Gonzalez




**Figure 2.**  $H\beta$  profile in the spectra of V340 Ser taken at different dates: April 5, 2018 (black line), April 11, 2018 (green line), and March 31, 2019 (red line). The vertical line marks the systemic velocity  $\gamma = 56.8 \pm 0.5$  km/s from de Smedt et al. (2016).


et al. (1997) and Pollard et al. (1997) devoted to the spectroscopy of RV Tau stars, as in the community of post-AGB stars as a whole. An  $H\alpha$  profile with a strong redshifted emission similar to the profile in the spectrum of V340 Ser is encountered rarely, but, as has been noted above, it was recorded by Pollard et al. (1997) in the spectrum of AD Aql. A significant phase variability of the  $H\alpha$  profile, including the variability of the positions of both its components, was recorded in the spectrum of this star. So far we have recorded no such significant variability of the positions and intensities of the  $H\alpha$  profile components for V340 Ser. A change only in the extent of the blue absorption wing is clearly seen in Fig. 1, tracing the variability of the outflow velocity. A close  $H\alpha$  profile type with a large velocity difference,  $>100$  km/s, for the absorption and emission components is also observed in the spectrum of the post-AGB star 89 Her (=IRAS 17534+2603). As an illustration, Fig. 3 shows a fragment of the spectrum for 89 Her taken on June 3, 2010, with the NES at the BTA telescope.

We see the influence of the wind distorting the blue wings of the strong SiII(2) 6347 and 6371 Å absorptions in the spectrum of V340 Ser in the right fragment of Fig. 4, where the FeI 6359 Å emission and the  $\lambda = 6376$  and 6379 Å DIBs are also marked. The distortion of the SiII(2) 6347 and 6371 Å absorption wings in the profiles in relative intensity – radial velocity coordinates is also illustrated by Fig. 5. Here the absorption cores of both profiles are shifted relative to the adopted systemic velocity, but their positions agrees well with the velocity from different metal absorptions in the spectrum: in Table 1 for this date  $V_{\odot}(\text{abs}) = 60.09$  km/s.

Having obtained the set of radial velocities for V340 Ser by the cross-correlation method, de Smedt et al. (2016) concluded that the star is binary and classified V340 Ser as a spectroscopic binary with a period of 365 days. Later, using the same technique, Oomen et al. (2018) improved the orbital elements for a sample of post-AGB stars in binary systems. For V340 Ser they provide




**Figure 3.** Fragment of the spectrum for 89 Her containing H $\alpha$  and TiI, VI, FeI, and NiI emissions. The identification of the main absorptions is indicated. The dots mark the telluric features.



**Figure 4.** Fragments of the spectrum for V340 Ser containing peculiar features: FeI and NiI emissions as well as strong SiII(2) 6347 and 6371 Å absorptions with an extended blue wing and DIBs in the right fragment. The identification of the main features is indicated.

the following orbital elements: a period of 363 days, a  $\gamma$  velocity of 56.1 km/s, and a velocity amplitude  $K = 8.4 \pm 1.0$  km/s.

Note two points. First, having high-quality spectra of this star, we found no evidence of spectroscopic binarity. Second, as follows from the data in Table 1, having taken the spectra of V340 Ser at arbitrary dates in 2018–2020, we recorded no radial velocity variability in the star expected from the results by de Smedt et al. (2016) and Oomen et al. (2018). Given the period about 1 year, we can assume that all our spectra were taken at close orbital phases.



**Figure 5.** SiII(2) 6347 (solid line) and 6371 Å absorption profiles in the April 11, 2018 spectrum of V340 Ser. The vertical line marks the systemic velocity  $\gamma = 56.8 \pm 0.5$  km/s (de Smedt et al., 2016).

The necessity of continuing the spectral observations of V340 Ser and a detailed study of the variability of the velocity pattern is obvious.

### 3.2. Metal emissions

The spectral fragments in Fig. 4 illustrate the following peculiarity of the optical spectrum for V340 Ser that has not been noted previously in publications: the presence of weak emissions of neutral metals with a low excitation potential of the lower level. Two such emissions in the H $\alpha$  wings are also clearly seen in Fig. 1. Table 2 lists all of such emissions identified by us in the recorded wavelength range. The last column in this table gives the velocities corresponding to the position of the emission feature. As follows from the data in Table 1, the mean velocity from the emissions changes insignificantly from date to date, from  $V_{\odot}(\text{em}) = 45.0 \pm 0.2$  to  $48.0 \pm 0.4$  km/s. For each date  $V_{\odot}(\text{em})$  differs systematically by  $\geq 10$  km/s from the mean velocity inferred from the absorptions, suggesting a velocity stratification in the extended stellar atmosphere. The full widths at half maximum (FWHMs) of these emissions forming in the outer layers of the extended atmosphere of V340 Ser are about  $0.25 \text{ \AA}$  or  $\delta V_{\odot} \approx 10 \div 12$  km/s, which is twice the FWHM of the forbidden ionospheric [OI] emissions in the spectrum. It should be emphasized that emissions are also present in several ion lines, for example, NdII 5102, YII 5509 and EuII 6645 Å. This peculiarity can affect the abundances of chemical elements and, therefore, it should be confirmed in subsequent observations.



Such low-excitation emissions of neutral metals were previously recorded in the spectra of the post-AGB candidates LN Hya (=IRAS 12538–2611) (Klochkova and Panchuk, 2012) and V1648 Aql (Klochkova and Tavalzhanskaya, 2019), whose spectral types are close to that of V340 Ser. For LN Hya metal emissions appeared in the spectra taken at its active phases in 2010, when the inverse P Cyg profile for the metal lines differed significantly from the one observed at quiescent phases. The position of the H $\alpha$  absorption component also differed significantly from its position at other dates of observations. Furthermore, the H $\alpha$  core was significantly (approximately by 15 km/s) redshifted relative to the symmetric metal absorptions.

The spectrum of the above mentioned post-AGB star 89 Her also contains a lot of low excitation emissions, as was already noted previously by Waters et al. (1993). As an illustration, Fig. 3 presents a fragment with narrow emissions of the spectrum for this star taken with the NES. Emissions of metal atoms were also recorded in the 1980-s in the spectra of RV Tau stars. We will also add the paper by Bopp (1984), whose author emphasized the appearance of such emissions during deep minima of the star R Sct, which also belongs to RV Tau stars, to the already mentioned papers by Baird (1982, 1984) and Klochkova and Panchuk (1998).

Some of the Fe, Co, and Ni emissions from Table 2 are also presented in the spectrum of the yellow hypergiant  $\rho$  Cas with an extended envelope (for details and references, see Klochkova et al. 2018). Besides, in the spectrum of  $\rho$  Cas the mean velocity from these emissions changes insignificantly with time and differs little from the systemic velocity of the hypergiant. The low width of these emissions in the spectrum of  $\rho$  Cas and the coincidence of the velocities suggest that these weak emissions are formed in the outer extended gaseous envelope whose sizes exceed considerably the photometric radius of the star. Emissions are observed predominantly in the periods of brightness decline in  $\rho$  Cas, which may point to a relative stability of the emission measure observed against the background of a weakened photospheric spectrum.

Table 2: Weak emissions identified in the spectrum of V340 Ser


$\lambda$	Element	$V_{\odot}$ , km/s
4953.200	NiI	46.20
5079.682	CeII	45.86
5092.800	NdII	42.86
5102.390	NdII	42.27
5107.446	FeI	45.85
5150.840	FeI	43.08
5151.910	FeI	45.20
5198.710	FeI	43.00
5216.270	FeI	43.39
5409.784	CrI	42.29
5434.523	FeI	40.45
5462.485	NiI	41.89
5490.150	TiI	44.67
5644.140	TiI	48.95
5711.883	NiI	46.81
5847.000	NiI	47.13
5999.040	TiI	42.84
6007.310	NiI	43.97
<i>see next page</i>		

Table 2, cont		
6065.482	FeI	44.89
6108.107	NiI	48.32
6108.530	Cl	40.30
6128.990	NiI	42.83
6137.691	FeI	43.60
6191.190	NiI	46.14
6191.558	FeI	45.09
6200.313	FeI	43.19
6213.429	FeI	43.55
6219.270	FeI	44.89
6230.722	FeI	42.47
6261.098	TiI	43.08
6265.131	FeI	45.17
6312.240	TiI	43.95
6330.130	CrI	45.75
6358.693	FeI	48.43
6393.600	FeI	43.43
6400.000	FeI	43.00
6430.844	FeI	44.75
6494.980	FeI	42.93
6498.950	FeI	48.96
6531.410	VI	47.03
6532.890	FeI	41.82
6546.238	FeI	45.25
6554.230	TiI	46.92
6572.800	CaI	45.91
6574.240	FeI	46.57
6592.913	FeI	43.03
6643.629	NiI	46.40
6677.985	FeI	43.14
6767.768	NiI	43.86
6770.960	CoI	46.38
6814.950	CoI	47.39
7052.870	CoI	46.81
7138.910	TiI	46.60
7291.449	NiI	41.72
7357.740	TiI	47.17
7714.308	NiI	48.73

### 3.3. DIBs and the multicomponent profile of the NaI D lines

Despite its significant distance from the Galactic plane (the Galactic latitude of the star  $|b| > 12^\circ$ ), numerous interstellar features were detected in the optical spectrum of V340 Ser. In particular, two such features with wavelengths  $\lambda = 6376$  and  $6379 \text{ \AA}$  are clearly seen in Fig. 4. Table 3 lists all of the DIBs from the well-known list by Jenniskens and Desert (1994) identified

by us and reliably distinguished among the blends in the spectrum of V340 Ser. For these features the table gives the radial velocity corresponding to the DIBs positions and their equivalent widths  $W_\lambda$ .



**Figure 6.** NaI 5889 and 5895 Å line profiles (dots) in the spectrum of V340 Ser. The positions of the profile absorption components forming in the interstellar medium (1), the circumstellar medium (2 and 3), and the stellar atmosphere (4) are marked. The vertical line marks the systemic velocity  $\gamma = 56.8 \pm 0.5$  km/s (de Smedt et al., 2016).

A high quality of our spectra for V340 Ser allowed us to resolve the NaI 5889 and 5895 Å D lines into components and to measure the position of the interstellar KI 7696 Å absorption. Individual absorptions, whose average positions are given in Table 1 and Fig. 6, are confidently distinguished in the NaI 5889 and 5895 Å lines profile in our spectra. The position of the long-wavelength absorption component (designated as 4 in this figure) for each date of observations agrees, within the above error limits, with the mean velocity  $V_\odot(\text{abs})$  measured from a large set of metal absorptions. Such agreement suggests the formation of this NaI D line component in the stellar atmosphere.

According to the data from Table 1, the position of the shortest-wavelength component 1 of the NaI doublet lines,  $V_\odot = -11.3$  km/s, is stable. Given the measurement errors, its position coincides with the average velocity from the set of DIBs identified in the spectrum, suggesting that this NaI doublet component is formed in the interstellar medium. In Fig. 6 we can see a differing steepness of the component wings, which also confirms the proposed interpretation of their formation regions.

Components 2 and 3 of the NaI D lines with velocities  $V_\odot = +10.1$  and  $+33.7$  km/s are redshifted relative to the mean radial velocity  $V_\odot(\text{abs})$  from the photospheric absorptions. It is natural to assume that both these components are formed in the circumstellar medium and the outer layers of the expanding stellar atmosphere. The component with a velocity of 10.1 km/s is

**Table 3.** Parameters of DIBs in the spectrum of V340 Ser

$\lambda$ , Å	$V_{\odot}$ , km/s	$W_{\lambda}$ , mÅ
4984.79	−11.86	23
4987.42	−12.70	6
5780.48	−12.55	263
5797.06	−12.11	49
5818.75	−9.61	1
5849.81	−10.69	5
6195.98	−11.35	36
6203.05	−10.18	52
6211.60	−11.39	29
6269.85	−11.04	12
6283.84	−10.27	85:
6329.97	−12.75	7
6376.08	−12.41	18
6379.32	−10.40	41
6439.48	−11.48	4
6445.28	−13.03	7
6532.10	−12.14	11
6613.62	−12.85	114
6632.86	−11.52	9
6770.05	−10.20	7
6827.30	−11.26	7
7224.03	−10.76	9
7357.20	−10.48	17
7367.13	−13.50	13
7581.30	−10.22	3
7721.85	−9.80	4

formed in an envelope expanding with a velocity typical of the envelopes of post-AGB stars (for comparison, see the results by Bakker et al. (1995, 1997) and Klochkova (2014)). The component with a velocity of 33.7 km/s is formed in the uppermost layers of the stellar atmosphere that pass into the circumstellar medium undetached from the star. The absorption of the complex H $\alpha$  profile is also formed here. These layers move with a velocity  $\approx 23$  km/s relative to the deep atmospheric layers, where symmetric absorptions are formed. The multicomponent profile of the NaI doublet lines also contains the emission component 5 exceeding the local continuum in Fig. 6. The position of this emission,  $V_{\odot} \approx 104$  km/s, is close to the position of the H $\alpha$  emission component.

Note the intensity ratio of the components in the NaI D1 (5895 Å) and D2 (5889 Å) lines. If the corresponding NaI atomic levels are populated proportionally to the statistical weights, then the D2/D1 ratio in the emission spectrum must be 2. In the spectra of V340 Ser the emission feature 5 is clearly seen in the D2 line and is at the detection limit in the D1 line. As the number density of NaI atoms rises, the role of the resonant scattering and self-absorption processes increases and, as a result, the intensity ratio of the NaI emission doublet lines will tend to 1.

In the spectra of stellar atmospheres we observe the D1 and D2 absorptions of equal intensities (for example, component 4 in Fig. 6). As a rule, a similar picture is also observed in the

spectra of the interstellar medium (for example, component 1 in Fig. 6), suggesting a saturation of the NaI doublet absorption profiles 4 and 1 due to self-absorption. In the spectra of the circumstellar envelopes of V340 Ser the intensity ratios of the doublet absorption components differs significantly:  $\approx 1.3$  and  $1.5$  for components 2 and 3, respectively. We conclude that no saturation in the line cores has been achieved for both emission feature 5 and absorption features 2 and 3.

Such a multicomponent profile of the NaI D lines with a fragment of the long-wavelength emission was recorded previously by Klochkova and Chentsov (2004) in the spectrum of the post-AGB star V510 Pup (=IRAS 08005–2356). This star has evolved farther from the AGB than V340 Ser and, therefore, has a detached envelope, which manifests itself, in particular, in the presence of circumstellar Swan C<sub>2</sub> absorption bands. Furthermore, emission from the OH bands simultaneously at 1612 and 1667 MHz was recorded for the source IRAS 08005–2356, which, according to Lewis (1989), in the absence of SiO and H<sub>2</sub>O masers suggests that the object approaches the planetary nebula stage. Note that in the spectrum of V510 Pup Klochkova and Chentsov (2004) identified the forbidden [CaII] 7291 and 7324 Å doublet emissions rarely observed in the spectra of post-AGB stars. These low excitation forbidden emissions are typical for the spectra of stars of selected types with a mass outflow and serve as evidence for the presence of a circumstellar disk. In particular, these emissions were identified in the spectra of the yellow supergiants V1302 Aql (=IRC+10420) (Klochkova et al. 1999) and V509 Cas (=IRC+60379) (Aret et al. 2017; Klochkova et al. 2019; Klochkova 2019) and the spectra of B supergiants with the B[e] phenomenon (Aret et al. 2012). In the spectra of V340 Ser we also see an emission that could be identified with the forbidden 7291 Å line, which could point to the probable presence of a circumstellar disk in the system. Unfortunately, the second line of this doublet, [CaII] 7324 Å, which could confirm the presence of a forbidden emission, falls between adjacent orders in our echelle frame. Additional observations are needed to confirm the presence of this important spectral feature. At present, we have to identify the 7291 Å emission in the spectrum with the low excitation NiI line with  $\lambda=7291.449$  Å.

Emission components in the NaI resonance lines with a FWHM  $\Delta \approx 120$  km/s were recorded by Klochkova et al. (2006) in the spectrum of the cool variable AGB star associated with the IR source IRAS 20508+2011. The multicomponent profile of the NaI D lines in the spectrum of the semiregular variable QY Sge is even more interesting. A broad emission with a FWHM  $\Delta \approx 120$  km/s was recorded by Menzies and Whitelock (1988), who assumed that the formation region of the emission in the resonance lines is a fairly hot and inhomogeneous circumstellar envelope, while the large emission width is due to the scattering of photons by moving dust grains in the envelope. Having recorded both broad and narrow metal emissions in the spectrum of QY Sge, Rao et al. (2002) developed the assumption of Menzies and Whitelock (1988) by proposing a model with a circumstellar torus and bipolar jets. Subsequently, based on their spectroscopic monitoring of QY Sge, Klochkova et al. (2007b) confirmed the conclusion about the emission formation in the circumstellar medium by showing the stability of its position in time.

### 3.4. Distance to the star and its luminosity

We will estimate the luminosity of the star using the luminosity – OI 7774 Å triplet intensity criterion known for F-G supergiants. In the spectra of V340 Ser the total equivalent width of the oxygen triplet is  $W_\lambda(7774) = 1.25$  Å, which is a typical value for post-AGB stars (for comparison, see the data from Molina et al. (2018)). The luminosity estimate for V340 Ser from

$W_\lambda(7774)$  is fairly accurate, because, according to van Winckel (1997), there is no oxygen abundance anomaly in the atmosphere of this star. Applying the calibration  $M_V \longleftrightarrow W_\lambda(7774)$  from Takeda et al. (2018), we get an absolute magnitude  $M_V \approx -4.6^m$  for V340 Ser. The luminosity of V340 Ser derived in this way,  $\log L/L_\odot = 3.75$ , agrees well with its spectral classification and the luminosity expected from the theoretical views of Blöcker (1995b) about the evolutionary status of post-AGB stars.

To estimate the color excess  $E(B - V)$  due to the interstellar extinction, we used the equivalent widths  $W_\lambda(\text{DIBs})$  measured in the spectra of V340 Ser for a sample of DIBs and the calibration relations  $E(B - V) \longleftrightarrow W_\lambda(\text{DIBs})$  from Kos and Zwitter (2013). Invoking seven confidently measured bands from Table 3 (4984, 5780, 5797, 6196, 6203, 6379, 6613 Å), for which the calibration relations by Kos and Zwitter (2013) are available, we found the mean color excess to be  $E(B - V) = 0.59^m$ . This color excess is definitely below the total reddening  $E(B - V)$ , because some fraction of the stellar reddening is also attributable to the extinction in the circumstellar medium. As follows from Fig. 6, the intensity of the circumstellar components 2 and 3 is only slightly lower than that of the interstellar one, suggesting a significant contribution of the circumstellar extinction to the total reddening  $E(B - V)$ .

By modeling the spectral energy distribution (SED) for V340 Ser, de Smedt et al. (2016) estimated the total reddening:  $E(B - V) = 0.76^m$ . With this estimate of the color excess we find the total extinction to be  $A_V \approx 2.43^m$  (for the standard ratio  $R = 3.2$ ). Taking into account the mean apparent magnitude  $V = 9.63^m$  from Arkhipova et al. (2011), we estimate the distance to the star to be  $d \approx 2.3$  kpc. The distance estimate  $d \approx 3.43$  kpc for V340 Ser from the extensive catalogue by Vickers et al. (2015) should be mentioned here. While modeling the SED for post-AGB stars, these authors adopted the luminosity range  $L/L_\odot = 6000 \pm 1500$  for them. Our luminosity estimate for V340 Ser from the IR oxygen triplet lies within this range. While modeling the SED, Vickers et al. (2015) took into account the anomalous SED pattern for IRAS 17279–1119 due to the presence of warm ( $\approx 855$  K) and cold ( $\approx 171$  K) dust. At the same time, Vickers et al. (2015) found the reddening to be  $E(B - V) = 0.45^m$ , which is considerably lower than the reddening from de Smedt et al. (2016) and even lower than the interstellar reddening fraction derived by us from DIBs.

The Gaia DR2 catalogue (2018) gives a reliable parallax for V340 Ser,  $\pi = 0.1577 \pm 0.0427$  mas, which leads to a significant distance of the star:  $d \geq 6.3$  kpc. However, in the 2016 version the data for the star differ significantly:  $\pi = 0.340 \pm 0.260$  mas. This parallax gives a distance to the star close to our estimate, but the parallax accuracy is too low. Here we should allude to the paper by Xu et al. (2019), in which they compared the Gaia DR2 parallaxes with the VLBI data and noted a large difference in these data for AGB stars with extended dust envelopes. Furthermore, the large parallax offset for Gaia DR2 relative to VLBI,  $-0.075 \pm 0.029$  mas (Xu et al. 2019), should be taken into account for distant objects.

On the whole, it can be asserted that the set of parameters for V340 Ser, i.e., its luminosity, distance from the Galactic plane, lowered metallicity, and chemical peculiarities, is consistent with its belonging to the post-AGB stage in the Galactic thick disk. Note that the set of main parameters and chemical composition of V340 Ser allows this star to be included in the homogeneous group of post-AGB stars with large enhancements of carbon and heavy s-process metals identified by Klochkova (1997) and van Winckel and Reyniers (2000). All stars of this group have structured, often bipolar, envelopes, where a lot of molecular and maser features are formed. Furthermore, an as yet unidentified emission at  $21\mu$  is contained in the IR spectra of all these stars (for more details, see Hrivnak et al. 2009). The circumstellar medium of V340 Ser has no these features, i.e., its envelope has not yet detached from the stars and has not cooled

to the requires extent. Hrivnak et al. (2009) suggest that the feature at  $21\mu$  is formed in the IR spectra of post-AGB stars with cold dust envelopes, with a temperature  $\approx 120\text{--}150$  K.

It is necessary to check whether V340 Ser belongs to RV Tau stars, because a star with the main parameters  $T_{\text{eff}} = 7200$  K and  $\log L/L_{\odot} = 3.75$  lies outside the instability strip, according to the data by Kiss et al. (2007). Recall that, previously, Arkhipova et al. (2011) also questioned this status of V340 Ser, while Siódmiak et al. (2008) classified V340 Ser as a post-AGB star, but not as an RV Tau one.

## 4. CONCLUSIONS

We studied the optical spectra of V340 Ser (=IRAS 17279–1119) taken at the 6-m BTA telescope with a spectral resolution  $R \geq 60000$ . For four dates we measured the heliocentric radial velocity from numerous metal absorptions (from 300 to 550 lines):  $V_{\odot} = 59.30 \pm 0.05$ ,  $60.09 \pm 0.05$ ,  $58.46 \pm 0.06$ , and  $55.78 \pm 0.06$  km/s. Given the high accuracy of the velocity, these values point to a weak variability of the velocity with time.

The interstellar,  $V_{\odot} = -11.20$  km/s, and circumstellar,  $V_{\odot} = +10$  km/s, components and the component forming in the uppermost atmospheric layers,  $V_{\odot} = +34.0$  km/s, were identified in the NaI D line profile. The values averaged over four spectra have an accuracy of  $\pm 0.2$  km/s. The mean velocity from 20–30 DIBs identified in the spectra,  $V_{\odot}(\text{DIBs}) = -11.6 \pm 0.2$  km/s, coincides with the velocity from the interstellar NaI component.

The  $H\alpha$  line has a PCyg type profile, with the position of its absorption component ( $V_{\odot} = +34$  km/s) pointing to its formation in the upper layers of the expanding atmosphere.

The weak emissions with an intensity  $\approx 10\%$  of the local continuum level were identified with low excitation lines of metal atoms. Their stable positions,  $V_{\odot} = 46.3 \pm 0.4$  km/s, systematically differ from the velocity inferred from atmospheric absorptions, suggesting the presence of a velocity gradient in the upper layers of the stellar atmosphere.

Based on the equivalent width triplet  $W_{\lambda}(7774) = 1.25 \text{ \AA}$  of the IR oxygen, we estimated the absolute magnitude to be  $M_v \approx -4.6^m$ , which, when taking into account the total (in the interstellar and circumstellar media) extinction, leads to a distance estimate for the star  $d \approx 2.3$  kpc. This value is half the distance corresponding to the parallax of V340 Ser from DR2 Gaia (2018). Such a difference can be a consequence of the correction for the extinction in the interstellar medium and the object's circumstellar environment.

It is necessary to check whether V340 Ser belongs to RV Tau type stars, because a star with the main parameters  $T_{\text{eff}} = 7200$  K and  $\log L/L_{\odot} = 3.75$  lies outside the instability strip.

*Acknowledgements.* We thank the Russian Foundation for Basic Research for its partial financial support (project no. 18-02-00029 a). We used the SIMBAD, SAO/NASA ADS, ASAS, and Gaia DR2 astronomical databases.

## References

- A. Arellano Ferro, S. Giridhar, and P. Mathias, *Astron. Astrophys.* 368, 250 (2001).
- F. Arenou, X. Luri, C. Babusiaux, et al., *Astron. Astrophys.* 616, A17 (2018);
- A. Aret, M. Kraus, M. F. Muratore, and M. Borges Fernandes, *MNRAS*, 423, 284 (2012).
- A. Aret, M. Kraus, I. Kolka, and G. Maravelias, *ASP Conf. Ser.* 508, 357 (2017).
- V.P. Arkhipova, N. P. Ikonnikova, and G. V. Komissarova, *Astron. Lett.* 37, 635 (2011).
- S.R. Baird, *PASP*, 94, 850 (1982).
- S.R. Baird, *PASP* 96, 72 (1984).
- 8.E. Bakker, H. J. G. L. M. Lamers, L. B. F. M. Waters, and T. Schoenmaker, *Astrophys. Space Sci.* 224, 335 (1995).
- E. Bakker, E. F. van Dishoeck, L. B. F. M. Waters, and T. Schoenmaker, *Astron. Astrophys.* 323, 469 (1997).
- T. Blöcker, *Astron. Astrophys.* 297, 727 (1995a).
- T. Blöcker, *Astron. Astrophys.* 299, 755 (1995b).
- B.W. Bopp, *PASP*, 96, 432 (1984).
- M. di Criscienzo, P. Ventura, D. A. Garcia Hernandez, et al. *MNRAS*, 462, 395 (2016).
- G.A. Galazutdinov, *SAO Preprint No.* 92 (1992).
- S. Giridhar, D. L. Lambert, and G. Gonzalez, *Astrophys. J.* 531, 521 (2000).
- G. Gonzalez, D. L. Lambert, and S. Giridhar, *Astrophys. J.* 479, 427 (1997).
- F. Herwig, *Ann. Rev. Astron. Astrophys.* 43, 435 (2005).
- B.J. Hrivnak, K. Volk, and S. Kwok, *Astrophys. J.* 694, 1147 (2009).
- P. Jenniskens and F.-X. Desert, *Astron. Astrophys. Suppl. Ser.* 106, 39 (1994).
- N. Rao Kameswara, A. Goswami, and D. L. Lambert, *MNRAS*, 334, 129 (2002)
- E.V. Kazarovets, N. N. Samus, and O. V. Durlevich, *Inform. Bull. Var. Stars*, No. 4870, 1 (2000).
- T. Kipper and V. G. Klochkova, *Baltic Astron.* 22, 77 (2013).
- L.L. Kiss, A. Derekas, Gy. M. Szabo, T. R. Bedding, and L. Szabados, *MNRAS*. 375, 1338 (2007).
- V.G. Klochkova, *MNRAS*, 272, 710 (1995).
- V.G. Klochkova, *Bull. Spec. Astrophys. Observ.* 44, 5 (1997).
- V.G. Klochkova, *Astrophys. Bull.* 69, 279 (2014).
- V.G. Klochkova, *Astrophys. Bull.* 74, 475 (2019).
- V.G. Klochkova and V. E. Panchuk, *Astron. Lett.* 24, 650 (1998).
- V.G. Klochkova and V. E. Panchuk, *Astron. Rep.* 56, 104 (2012).



- V.G. Klochkova and N. S. Tavalzhanskaya, *Astrophys. Bull.* 74, 277 (2019).
- V.G. Klochkova and E. L. Chentsov, *Astron. Rep.* 48, 301 (2004).
- V.G. Klochkova, E. L. Chentsov, and N. S. Tavganskaya, *Astrophys. Bull.* 48, 25 (1999).
- V.G. Klochkova, V. E. Panchuk, N. S. Tavalzhanskaya, and G. Zhao, *Astron. Rep.* 50, 232 (2006).
- V.G. Klochkova, E. L. Chentsov, N. S. Tavganskaya, and M. V. Shapovalov, *Astrophys. Bull.* 62, 162 (2007a).
- V.G. Klochkova, V. E. Panchuk, E. L. Chentsov, and M. V. Yushkin, *Astrophys. Bull.* 62, 162 (2007b).
- V.G. Klochkova, V. E. Panchuk, and N. S. Tavalzhanskaya, *Astron. Rep.* 62, 623 (2018).
- V.G. Klochkova, E. L. Chentsov, and V. E. Panchuk, *Astrophys. Bull.* 74, 41 (2019).
- J. Kos and T. Zwitter, *Astrophys. J.* 774, 72 (2013).
- B.M. Lewis, *Astrophys. J.* 338, 234 (1989).
- N. Liu, R. Gallino, S. Bisterzo, A. M. Davis, R. Trappitsch, and L. R. Nittler, *Astrophys. J.* 865, 112 (2018).
- J.W. Menzies and P. A. Whitelock, *MNRAS*, 233, 697 (1988).
- R.E. Molina, S. Giridhar, C. B. Pereira, A. Arellano Ferro, and S. Muneer, *Rev. Mex. Astron. Astrofis.* 50, 293 (2014).
- G.-M. Oomen, H. van Winckel, O. Pols, et al. , *Astron. Astrophys.* 620, A85 (2018).
- R.D. Oudmaijer, W. E. C. J. van der Veen, L. B. F. M. Waters, et al. *Astron. Astrophys. Suppl. Ser.* 96, 625 (1992).
- V.E. Panchuk, V. G. Klochkova, and M. V. Yushkin, *Astron. Rep.* 61, 820 (2017).
- K.R. Pollard, P. L. Cottrell, W. A. Lawson, M. D. Albow, and W. Tobin, *MNRAS*, 286, 1 (1997).
- S. Sumangala Rao, S. Giridhar, and D. L. Lambert, *MNRAS*, 419, 1254 (2012).
- C. Sanchez Contreras, R. Sahai, R. Goodrich, and A. Gil de Paz, *Astrophys. J. Suppl. Ser.* 179, 166 (2008).
- N. Siódmiak, M. Meixner, T. Ueta, et al. *Astrophys. J.* 677, 382 (2008).
- K. de Smedt, H. van Winckel, D. Kamath, et al. *Astron. Astrophys.* 587, A6 (2016).
- Y. Takeda, G. Jeong, and I. Han, *Publ. Astron. Soc. Jpn.* 70, 8 (2018).
- S.B. Vickers, D. J. Frew, O. A. Parker, and I. S. Bojicic, *MNRAS*, 447, 1673 (2015).
- L.B.F.M. Waters, C. Waelkens, M. Mayor, and N. R. Trams, *Astron. Astrophys.* 269, 242 (1993).
- H. van Winckel, *Astron. Astrophys.* 319, 561 (1997).
- H. van Winckel and M. Reyniers, *Astron. Astrophys.* 354, 135 (2000).
- S. Xu, B. Zhang, M. J. Reid, Z. Xingwu, and W. Guangli, *Astrophys. J.* 875, 114 (2019).
- M.V. Yushkin and V. G. Klochkova, *SAO Preprint No.206 (Spec. Astrophys. Observ., 2005).*