Unitary evolution to a state with a fixed mean number of particles

Bogusław Broda

Department of Theoretical Physics, Faculty of Physics and Applied Informatics, University of Łódź, 90-236 Łódź, Pomorska 149/153, Poland

Abstract

In the framework of finite-dimensional Fock space models, for a predefined fixed mean number of particles \bar{n}_k , it is shown that there is a "large" multi-dimensional subspace $s_{\bar{n}_k}$ of initial pure states, in the space S of all pure states, unitarily evolving to a subspace $S_{\bar{n}_k}$ of final pure states which yield \bar{n}_k . As an example, in particular it follows that the blackbody form of the mean number of particles \bar{n}_k does not by itself contradict unitarity of black hole evaporation.

Keywords: states with a fixed mean number of particles, unitary evolution in finite-dimensional Hilbert spaces, unitarity of black hole evaporation PACS: 03.65.Aa (Quantum systems with finite Hilbert space), 04.70.Dy (Quantum aspects of black holes, evaporation, thermodynamics) $2010\ MSC:\ 81Q99$ (None of the above, but in this section), 83C57 (Black holes)

1. Introduction

The aim of the work is to show that there are many (initial) pure states unitarily evolving to a (pure) state with a predefined fixed mean number of particles \bar{n}_k . Although primary motivation to pose such a problem has resulted from analysis of the black hole information paradox [1, 2, 3, 4], actually the problem directly refers to quantum mechanics and is entirely independent from black hole context.

In short, the black hole information (loss) paradox (problem/puzzle) consists in difficulty in explaining the status of unitarity of the process of black hole evaporation. According to standard picture of semiclassical gravity, a black hole (quantumly) evaporates due to the Hawking effect, and finally transmutes into blackbody radiation. In consequence, in general distinct initial pure states forming a black hole could possibly be transformed into the same final "structureless" blackbody (thermal) radiation. Due to this seemingly "many to one" process,

Email addresses: boguslaw.broda@uni.lodz.pl (Bogusław Broda), http://merlin.phys.uni.lodz.pl/BBroda (Bogusław Broda)

unitarity (of evolution) could be lost in contradistinction with fundamentals of quantum mechanics.

In the present paper we analyze the issue of unitarity of the process of evolution of a pure state to a (pure) state with a predefined fixed mean number of particles \bar{n}_k . We present our arguments in three steps, in the form of the following three models defined in finite-dimensional Hilbert spaces: (1) Toy Model (a model defined on the Bloch sphere [5, 6, 7]); (2) a more realistic, Fermion Fock space model; and (3) Boson Fock space model with a cutoff [8]. More precisely, we will show that there is a "large" (in the sense of dimension) subspace $S_{\bar{n}_k}$ of distinct pure states in the whole space S of pure states (or in the corresponding Hilbert space \mathcal{H}) yielding the same (almost arbitrary) mean number of particles \bar{n}_k . Therefore, a process which could seem, at first glance, to be "many to one" can actually be "one to one" (and unitary), because there is "enough room" in the space S to "accommodate" this process.

One should stress that we exclusively operate on pure states, i.e., nowhere do density matrices, nor mixed states or purifications of thereof appear in our considerations, explicitly or implicitly. In our analysis, \bar{n}_k , blackbody or another, is a mean of the particle number operator \hat{n}_k in a pure state (see (2)), and possible thermality of the spectrum is, in a sense, simulated by an appropriately chosen pure state $|\bar{n}_k\rangle$ (see (9), (16), (27)).

In the context of black holes, we do not prove that the actual process of black hole evaporation is "one to one" and unitary, but as an illustration of our analysis we argue that the blackbody(-like) shape of the Hawking spectrum does not by itself imply non-unitarity of black hole evaporation. Obviously, our analysis is quite general, because no explicit particular form of the mean number of particles \bar{n}_k (blackbody or another) enters our analysis.

2. Primary motivation and the Toy Model

In his famous work on black hole radiation spectrum [9], Hawking derived an explicit formula for the mean number of particles \bar{n}_k understood as a quantum average

$$\bar{n}_k \equiv \langle \hat{n}_k \rangle \,, \tag{1}$$

where \hat{n}_k is the particle number operator for the mode k. With a good approximation, the mean number of particles of black hole radiation, \bar{n}_k , appears to be blackbody. Actually, what counts from our perspective is the *total mean* number of particles [10], rather than usually discussed temporary quantity, but as it is mentioned in Introduction our analysis is insensitive to any particular form of \bar{n}_k .

2.1. General idea

Our main claim is that we have a "huge multitude" of pure states yielding a predefined fixed mean number of particles \bar{n}_k . More precisely, we have a "large", in the sense of low codimension, subspace denoted by $S_{\bar{n}_k}$ in the space S of all pure states (or in the Hilbert space \mathcal{H}) corresponding to almost any arbitrarily

2.2 Toy Model

chosen \bar{n}_k (the only restriction on \bar{n}_k is imposed, depending on the case, by the mild condition (10), (23) or (32)). Therefore, having given the mean \bar{n}_k , and provided we are able to determine the corresponding ("large") subspace $S_{\bar{n}_k}$, we choose any state $|\bar{n}_k\rangle \in S_{\bar{n}_k}$ yielding, by virtue of the definition of $S_{\bar{n}_k}$, the average with the expected predefined value(s), i.e.,

$$\bar{n}_k \left| \hat{n}_k \right| \bar{n}_k \rangle = \bar{n}_k. \tag{2}$$

In the next step we can perform a (thought) unitary transformation U(-t) on $S_{\bar{n}_k}$, now interpreted as a subspace of final states, obtaining another subspace $s_{\bar{n}_k}$ (incidentally, because of unitarity of U(-t), $s_{\bar{n}_k}$ is isometric to $S_{\bar{n}_k}$ in the sense of the complex metric on \mathcal{H}) interpreted as a subspace of possible initial states. The unitary transformation U(-t) corresponds to evolution backward in time (the minus sign). Thus, we can conclude that the "huge multitude" of distinct initial pure states belonging to $s_{\bar{n}_k}$ unitarily (according to U(t)) evolves towards the "huge multitude" of distinct pure states belonging to the subspace $S_{\bar{n}_k}$ of states yielding, by virtue of the construction, the fixed predefined mean number of particles \bar{n}_k .

2.2. Toy Model

Now, let us introduce the Toy Model (a model on the Bloch sphere). Its sole role is to explicitly elucidate and visualize (because of low dimension) our main idea. As a chief postulate of the model we assume that the whole Universe consists of only one fermion mode (2-level system). Its Hilbert space $\mathcal{H} = \mathbb{C}^2$ is 4-dimensional in real sense (in this paper we only operate real dimensions), and in the Fock space base $\{|0\rangle, |1\rangle\}$ any state $|\psi\rangle \in \mathcal{H}$ can be expressed as

$$|\psi\rangle = \alpha_0 |0\rangle + \alpha_1 |1\rangle, \qquad \alpha_0, \alpha_1 \in \mathbb{C}.$$
(3)

Because of normalization ($\langle \psi | \psi \rangle = 1$) and of arbitrariness of phase, pure states for this system are parameterized by points on the 2-dimensional Bloch sphere $S^2 (= \mathbb{C}P^1)$ [5, 6, 7]. Then, the general state can be specified as

$$|\psi\rangle = \cos\frac{\theta}{2}|0\rangle + e^{i\varphi}\sin\frac{\theta}{2}|1\rangle, \qquad 0 \le \theta \le \pi, \quad 0 \le \varphi < 2\pi,$$
 (4)

where θ and φ are the polar and azimuthal angle on S^2 , respectively. From the point of view of quantum mechanics, any (pure) state is uniquely given by a point on S^2 , and arbitrary continuous unitary (e.g. time) evolution U(t)corresponds to rotation of S^2 , i.e., $U(t) \in SO(3)$.

Now, we would like to determine the entire (sub)space $S_{\bar{n}}$ of the states $|\bar{n}\rangle$ yielding the fixed mean number of particles \bar{n} , where obviously

$$0 \le \bar{n} \le 1. \tag{5}$$

Since k = 1, the mode number k has been skipped in this subsection, and the mean number of particles \bar{n} is now a single number belonging to the interval (5). In general, the state we are looking for, expressed as

$$\left|\bar{n}\right\rangle = \bar{\alpha}_0 \left|0\right\rangle + \bar{\alpha}_1 \left|1\right\rangle \tag{6}$$

(in our paper the "bar" symbol over coefficients denotes their specific values, instead, complex conjugation is denoted by the "asterisk"), should satisfy the two conditions:

$$\langle \bar{n} | \bar{n} \rangle = 1, \qquad \langle \bar{n} | \hat{n} | \bar{n} \rangle = \bar{n}.$$
 (7)

Utilizing the angular parametrization (4) on the Bloch sphere, we easily find

$$\bar{n} = \left|\bar{\alpha}_1\right|^2 = \sin^2\frac{\theta}{2}.\tag{8}$$

Thus, finally the solution of the problem (full set of solutions of Eqs.(7) modulo phase) assumes the following explicit form

$$\left|\bar{n}\right\rangle = \sqrt{1-\bar{n}}\left|0\right\rangle + e^{i\varphi}\sqrt{\bar{n}}\left|1\right\rangle.$$
(9)

Eq.(9) says that all the points (interpreted by us as final pure states) on the circle ("parallel") $S_{\bar{n}}$, parameterized by the azimuthal angle φ and determined by the "latitude" (on S^2) given by the polar angle (see (8)) $\theta = 2 \arcsin \sqrt{\bar{n}}$, yield the same predefined \bar{n} . In the next step we can (thought) rotate the circle (parallel) $S_{\bar{n}}$ ("backward in time evolution U(-t)") obtaining another circle (not a parallel, in general) $s_{\bar{n}}$ parametrizing all the states (interpreted as initial states) which can be transformed back (in the course of the "proper time evolution U(t)") onto states on $S_{\bar{n}}$ with the predefined \bar{n} . Thus, in general, we have a "one to one" unitary relation between points on isometric circles on S^2 , where parallels play a distinguished role of "thermality imitating" states. Since for $\bar{n} = 0, 1$ the circle $S_{\bar{n}}$ degenerates to a point (poles), we can impose a mild restriction on admissible values of \bar{n} , removing the boundary values of the interval (5), putting

$$0 < \bar{n} < 1. \tag{10}$$

An example situation is explicitly illustrated in Fig.1.

3. Fock space models

In this section we introduce two more realistic models based on fermion Fock space and boson Fock space, respectively. To technically simplify our discussion (algebraization of the problem), as well as to make it more quantitative and rigorous, we impose some cutoffs on the Fock spaces, which implies finite dimensionality of corresponding Hilbert spaces.

3.1. Fermion Fock space model

First, we consider a fermion model defined on the antisymmetric Fock space \mathcal{F}_A^m , where $m \ (\geq 1)$ is a number of fermion modes. Here, the cutoff simply means that the number of modes m is finite. Generalizing the linear expansion (3), we can express any state $|\psi\rangle \in \mathcal{F}_A^m$ as a linear combination

$$|\psi\rangle = \sum_{\boldsymbol{n}} \alpha_{\boldsymbol{n}} |\boldsymbol{n}\rangle, \qquad \alpha_{\boldsymbol{n}} \in \mathbb{C},$$
 (11)



Figure 1: Toy Model (a model on the Bloch sphere S^2). Geometric statement: Rotation $U(\alpha)$ through the angle $\alpha = \pi/2$ in the plane of Figure transforms the circles s_1 and s_2 onto the circles S_1 and S_2 , respectively. Polar angle coordinates of the circles S_1 and S_2 are θ_1 and θ_2 , respectively, and the points on the circles are parametrized by the azimuthal angle φ . In particular, the distinct points $x_1, x_2 \in s_2$ are transformed (rotated) onto the points $X_1, X_2 \in S_2$, respectively. Quantum-mechanical statement: Two distinct initial pure states x_1 and x_2 unitarily (after "time $\pi/2$ ") evolve onto two distinct pure states X_1, X_2 , respectively, with the same mean number of particles ("spectrum") $\bar{n} = \sin^2 \frac{\theta_2}{2}$.

where for convenience we have introduced a multi-index $\mathbf{n} \equiv n_1 \cdots n_m$ with $n_k = 0, 1 \ (k = 1, \dots, m)$.

Analogously to the presentation (6), the points/states $|\bar{n}\rangle \in \hat{S}_{\bar{n}}$, i.e., those normalized and satisfying the condition (2), can be expressed by the sum

$$\left| \bar{\boldsymbol{n}} \right\rangle = \sum_{\boldsymbol{n}} \bar{\alpha}_{\boldsymbol{n}} \left| \boldsymbol{n} \right\rangle, \tag{12}$$

where for the multi-parameter $\bar{\boldsymbol{n}} \equiv \bar{n}_1, \ldots, \bar{n}_m$ we assume $0 \leq \bar{n}_k \leq 1$, and the "tilde" over $S_{\bar{\boldsymbol{n}}}$ denotes the space of states before identification (symbolized by "/~") of states differing by phase, i.e., $S_{\bar{\boldsymbol{n}}} \equiv \tilde{S}_{\bar{\boldsymbol{n}}}$ /~. Henceforth, we use the symbol $S_{\bar{\boldsymbol{n}}}$ instead of $S_{\bar{n}_k}$. Normalization condition for (12) reads

$$\langle \bar{\boldsymbol{n}} | \bar{\boldsymbol{n}} \rangle = \sum_{\boldsymbol{n}} | \bar{\alpha}_{\boldsymbol{n}} |^2 = 1,$$
 (13)

whereas the condition (2) yields the following system of m quadratic equations

$$\langle \bar{\boldsymbol{n}} | \hat{n}_k | \, \bar{\boldsymbol{n}} \rangle = \sum_{\boldsymbol{n}_k} | \bar{\alpha}_{\boldsymbol{n}_k} |^2 = \bar{n}_k, \qquad (14)$$

where we have introduced another multi-index $\mathbf{n}_k \equiv n_1 \cdots 1_k \cdots n_m$, i.e., the kth index assumes the constant value $n_k = 1$, and consequently there is no summation with respect to n_k in (14). Furthermore, a bit extending the domain of the index k introducing a new auxiliary index $p = 0, 1, \ldots, m$ instead of k $(=1,\ldots,m)$, and additionally defining $\mathbf{n}_0 \equiv \mathbf{n}$ as well as $\bar{n}_0 \equiv 1$, we can rewrite

3.1 Fermion Fock space model

the quadratic equation (13) and the system (14) in the following compact unified form:

$$\sum_{\boldsymbol{n}_p} \left| \bar{\alpha}_{\boldsymbol{n}_p} \right|^2 = \bar{n}_p, \qquad p = 0, 1, \dots m.$$
(15)

Thus, the (sub)space $\tilde{S}_{\bar{n}}$ is defined as a (sub)space of solutions of the system of m+1 quadratic equations (15). Fortunately, to proceed further we do not need an explicit form of $S_{\bar{n}}$, as in the case of the Toy Model of Section 2, where global analysis has been performed. Since we only aim to determine the (co)dimension of $\tilde{S}_{\bar{n}}$ (and of $S_{\bar{n}}$), we can confine ourselves to purely local analysis.

Our strategy is first to find only a single non-degenerate (in the sense explained latter) solution of the quadratic system (15), and next to show that it can be infinitesimally extended in sufficiently many dimensions/directions. It is straightforward to check that the following "(tensor product) Bloch-type" state (its symmetrized version is known as the Dicke state [11, 12])

$$\left|\bar{\boldsymbol{n}};\boldsymbol{\phi}\right\rangle \equiv \sum_{\boldsymbol{n}} \bar{\alpha}_{\boldsymbol{n}}\left(\boldsymbol{\phi}\right) \left|\boldsymbol{n}\right\rangle, \qquad \boldsymbol{\phi} = \varphi_{1}, \dots, \varphi_{m}, \quad 0 \leq \varphi_{k} < 2\pi \quad \left(k = 1, \dots, m\right),$$
(16)

where (cf. (4))

$$\bar{\alpha}_{\boldsymbol{n}}\left(\boldsymbol{\phi}\right) \equiv \bar{\alpha}_{n_{1}\cdots n_{m}}\left(\varphi_{1},\dots,\varphi_{m}\right) = \prod_{k=1}^{m} \left(\delta_{n_{k}}^{0}\cos\frac{\theta_{k}}{2} + \delta_{n_{k}}^{1}e^{i\varphi_{k}}\sin\frac{\theta_{k}}{2}\right), \quad (17)$$

with (cf. (8))

$$\sin^2 \frac{\theta_k}{2} = \bar{n}_k,\tag{18}$$

solves the system (15). Actually, the formulas (16–18) define the whole *m*-dimensional torus $\mathcal{T}^m \equiv \underbrace{\mathcal{S}^1 \times \cdots \times \mathcal{S}^1}_m$ of solutions of the system (15), parameterized by ϕ . Since our analysis is supposed to be local, we only need a single

terized by ϕ . Since our analysis is supposed to be local, we only need a single point/solution of the system (15), and therefore, to simplify our further considerations we put $\phi = 0$ henceforth.

To find a solution of the system (15) in an infinitesimal vicinity of the Blochtype solution (16–18) at the point $\phi = 0$ on the torus \mathcal{T}^m , we insert (into (15)) the expansion

$$\bar{\alpha}_{\boldsymbol{n}} = \bar{\alpha}_{\boldsymbol{n}} \left(0 \right) + z_{\boldsymbol{n}},\tag{19}$$

where z_n is a (complex) infinitesimal variation around the solution $\bar{\alpha}_n(0)$. Thus, we get a system of m + 1 linear equations

Re
$$\sum_{\boldsymbol{n}_{p}} \bar{\alpha}_{\boldsymbol{n}_{p}}(0) z_{\boldsymbol{n}_{p}}^{*} = 0, \qquad p = 0, 1, \dots, m,$$
 (20)

which define m + 1 hyperplanes tangent to the respective m + 1 quadrics determined by the system (15). The maximal possible rank of the matrix of the coefficients entering the system (20) is obviously m + 1, and such a situation

(the most desirable one) geometrically corresponds to a non-degenerate intersection of the hyperplanes tangent to the quadrics. Since to determine the rank of a matrix, one usually invokes determinants, let us calculate the determinant of a matrix constructed from the columns containing the following coefficients: $\bar{\alpha}_{00...0}(0), \bar{\alpha}_{10...0}(0), \bar{\alpha}_{01...0}(0), ..., \bar{\alpha}_{00...1}(0)$. The matrix reads

$$M_{A} = \begin{bmatrix} \bar{\alpha}_{00\dots0} (0) & * & \cdots & * \\ & \bar{\alpha}_{10\dots0} (0) & & \\ & & \ddots & \\ & & & \bar{\alpha}_{00\dots1} (0) \end{bmatrix},$$
(21)

where M_A appears to be an upper triangular matrix (more precisely, all entries of M_A , possibly except the 1st row and the main diagonal, are zero). Then, by virtue of (17)

$$\det M_A = \bar{\alpha}_{00\dots0} (0) \prod_{j=1}^m \bar{\alpha}_{00\dots1_j\dots0} (0) = \prod_{j=1}^m \sin \frac{\theta_j}{2} \cos^m \frac{\theta_j}{2}.$$
 (22)

From (22) it immediately follows that the rank of the system (20) is really maximal (= m + 1), and hence there is no degeneracy, provided we impose the condition $0 < \theta_k < \pi$, which corresponds (by virtue of the relationship (18)) to a very mild restriction on \bar{n}_k (cf. (10)),

$$0 < \bar{n}_k < 1, \qquad k = 1, \dots, m,$$
 (23)

in comparison with all theoretically admissible values: $0 \leq \bar{n}_k \leq 1$.

3.2. Boson Fock space model

Let us now switch to a boson model defined on the symmetric Fock space $\mathcal{F}_{S}^{m,N}$ with cutoffs $m (\geq 1)$ and $N (\geq 1)$, where m is a finite number of boson modes, and a fixed finite number of possible levels, the same for each boson mode, is equal to N + 1 (see [8]). In principle the cutoff N can be arbitrary, but for our needs it should be sufficiently large, i.e.,

$$\bar{n}_k < N < +\infty, \qquad k = 1, \dots, m. \tag{24}$$

As a single-mode Bloch-type state (with $\varphi = 0$) in the cutoff boson case $(\mathcal{F}_{S}^{m,N})$ we assume

$$|\psi\rangle = \cos\frac{\theta}{2}|0\rangle + \sin\frac{\theta}{2}|N\rangle,$$
 (25)

where $0 \le \theta \le \pi$ (cf. (4)). Executing calculations similar to those in Section 2, we obtain as analog of (8)

$$\bar{n} = N \sin^2 \frac{\theta}{2}.$$
(26)

In turn, the tensor product Bloch-type state (with $\phi = 0$) is now (cf. (16))

$$|\bar{\boldsymbol{n}};0\rangle = \sum_{\boldsymbol{n}} \bar{\alpha}_{\boldsymbol{n}}(0) |\boldsymbol{n}\rangle,$$
 (27)

where (cf. (17))

$$\bar{\alpha}_{\boldsymbol{n}}\left(0\right) = \prod_{k=1}^{m} \left(\delta_{n_{k}}^{0} \cos\frac{\theta_{k}}{2} + \delta_{n_{k}}^{N} \sin\frac{\theta_{k}}{2}\right),\tag{28}$$

with (cf. (18) and (29))

$$\sin^2 \frac{\theta_k}{2} = \frac{\bar{n}_k}{N}.$$
(29)

In the boson case, for the multi-index \boldsymbol{n} we assume $n_k = 0, 1, \ldots N$, and for the multi-parameter $\bar{\boldsymbol{n}}, 0 \leq \bar{n}_k < +\infty$, respectively.

The matrix analogous to (21) is now the (upper triangular) matrix

$$M_{S} = \begin{bmatrix} \bar{\alpha}_{00\dots0}(0) & * & \cdots & * \\ & \bar{\alpha}_{N0\dots0}(0) & & & \\ & & \ddots & & \\ & & & \bar{\alpha}_{00\dots N}(0) \end{bmatrix},$$
(30)

and its determinant,

$$\det M_S = \prod_{j=1}^m \sin \frac{\theta_j}{2} \cos^m \frac{\theta_j}{2}, \qquad (31)$$

is exactly of the same form as for M_A (see (22)). The relations (26) and (31) impose a very mild restriction on \bar{n}_k (cf. (23)),

$$0 < \bar{n}_k < +\infty, \qquad k = 1, \dots, m, \tag{32}$$

in comparison with all theoretically admissible values: $0 \leq \bar{n}_k < +\infty$.

3.3. Summary of the Fock space models

In the case of the fermion Fock space model as well as in the case of the boson one, for arbitrary fixed mean number of particles \bar{n}_k , mildly restricted by (23) and (32), respectively, we have shown that the tensor product Bloch-type state (16–18) and (27–29), respectively, determines a non-degenerate intersection point in the corresponding (finite dimensional) Hilbert space $\mathcal{H} = \mathcal{F}_A^m$ and $\mathcal{H} = \mathcal{F}_{S}^{m,N}$, respectively. More precisely, intersecting hyperplanes tangent to the intersecting quadrics defined by the system (15) (with indices $n_k = 0, 1$ and $n_k = 0, 1, \ldots, N$, respectively) are "in general position" (genuine intersection, no overlappings). Therefore, the intersection of the quadrics is also non-degenerate (genuine intersection, no contact points) and hence each quadric (equation in the system of the m + 1 equations (15)) imposes one condition reducing dimension of the subspace by one (codimension increases by one). Consequently, codimension of the intersection of the whole set of the quadrics (15) equals codimension of the intersection of the set of the hyperplanes tangent to these quadrics, and it is equal m + 1 (number of the equations). Then, $\dim \tilde{S}_{\bar{n}} = \dim \mathcal{H} - (m+1)$ and

$$\dim S_{\bar{\boldsymbol{n}}} = \dim \tilde{S}_{\bar{\boldsymbol{n}}} - 1 = \dim \mathcal{H} - m - 2 \tag{33}$$

 $(S_{\bar{n}} \equiv \tilde{S}_{\bar{n}} / \sim)$. Strictly speaking, the subtraction in (33) is justified provided the action of the group U(1) corresponding to the identification ("~") of states differing by phase proceeds tangentially to $\tilde{S}_{\bar{n}}$ at the point $\bar{\alpha}_n(0)$. Glancing at the system (20) we can immediately observe that imaginary parts $(y_n \equiv \text{Im } z_n)$ of infinitesimal variations z_n are absent from the system (20) (unrestricted), and therefore (infinitesimally) $\tilde{S}_{\bar{n}}$ can freely extend in imaginary directions, and this is exactly the direction of (infinitesimal) action of the (phase) group U(1).

For illustrative purposes, let us apply Eq.(33) to the simplest non-trivial case, namely, to our Toy Model (Subsection 2.2). Since now dim $\mathcal{H} = 4$ and m = 1, we get dim $S_{\bar{n}} = 4 - 1 - 2 = 1$, which obviously agrees with dimension of a circle ("parallel").

To better quantify and justify the term "huge multitude" or "large" introduced in Section 2 in the context of dimension of $S_{\bar{n}}$, we should compare dimensions of all relevant spaces. To begin with, for the Hilbert space of states \mathcal{H} we have dim $\mathcal{H} = 2 (N+1)^m$, where N = 1 or N = cutoff (see (24)), in the fermion case (\mathcal{F}_A^m) or in the boson case $(\mathcal{F}_S^{m,N})$, respectively. Since the normalization condition for states lowers dimension by one, and so does also identification of states differing by phase [5, 6], (dimension of the space S of all states) dim $S = \dim \mathcal{H} - 2 = 2 (N+1)^m - 2$. In turn, according to (33) dim $S_{\bar{n}} = 2 (N+1)^m - m - 2$. Then, $S_{\bar{n}}$ is a $[2 (N+1)^m - m - 2]$ -dimensional subspace in the $[2 (N+1)^m - 2]$ -dimensional space S of all states, and consequently codimension of $S_{\bar{n}}$ in S is equal to the number m of the modes. Then, codim $S_{\bar{n}} = m \sim C \log \dim S$ (where $C = \log^{-1} (N+1)$) and exactly in this ("logarithmic") sense is dimension of $S_{\bar{n}}$ large.

Interpreting $S_{\bar{n}}$ as a subspace of final (pure) states with a fixed spectrum, and performing a (thought) unitary transformation U(-t) (understood as evolution backward in time), we obtain (an isometric to $S_{\bar{n}}$ subspace) $s_{\bar{n}}$, which can be interpreted as a subspace of initial states. By virtue of the construction, unitary (time) evolution determined by U(t) transforms all pure states belonging to $s_{\bar{n}}$ onto (pure) states belonging to $S_{\bar{n}}$ which yield the same predefined fixed mean number of particles \bar{n}_k .

4. Final remarks

In the presented analysis we have shown that there is a huge multitude of distinct pure states which can unitarily be evolved to states with a predefined fixed mean number of particles \bar{n}_k . Comparing in the framework of our finitedimensional Fock space models a subspace $S_{\bar{n}}$ of final pure states yielding \bar{n}_k , e.g. blackbody-like or another, to the whole space S of pure states, we observe that asymptotically the order of growth of codimension of $S_{\bar{n}}$ ($\subset S$) is (actually) only logarithmic function of dimension of S. Then, as $S_{\bar{n}}$ is a (sub)space which parametrizes a really "huge multitude" of all final states with given \bar{n}_k , and $s_{\bar{n}}$ (isometric to $S_{\bar{n}}$) is a (sub)space of all possible initial states, there is "enough room" in the space S to "accommodate" unitarily realized time evolution.

REFERENCES

To illustrate this result in the context of black hole evaporation, we should adopt the following scenario. In the beginning of evolution, as an input state we have matter forming a black hole in a pure state. In turn, the final state only consists of the radiation (in a pure state) which is solely characterized by its blackbody(-like) spectrum (mean number of particles) \bar{n}_k . Since according to our analysis the space of all possible input pure states yielding the given \bar{n}_k is large, the black hole evaporation can be unitarily realized with an ease. Actually, black hole radiation is characterized not only by its blackbody spectrum but also by thermal density matrix. Therefore, for preserving unitarity, not only the blackbody spectrum \bar{n}_k should be reproduced properly (and in principle it can), but so should also the averages of all observables. Obviously, this later property cannot be reproduced by any pure state.

I am grateful for Reviewer's very accurate and valuable remarks.

References

- S. Chakraborty, K. Lochan, Black Holes: Eliminating Information or Illuminating New Physics?, Universe 3 (3) (2017) 55. arXiv:1702.07487, doi:10.3390/universe3030055.
- D. Harlow, Jerusalem Lectures on Black Holes and Quantum Information, Rev. Mod. Phys. 88 (2016) 015002. arXiv:1409.1231, doi:10.1103/ RevModPhys.88.015002.
- [3] D. Marolf, The Black Hole information problem: past, present, and future, Rept. Prog. Phys. 80 (9) (2017) 092001. arXiv:1703.02143, doi:10.1088/ 1361-6633/aa77cc.
- [4] J. Polchinski, The Black Hole Information Problem, in: Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings (TASI 2015): Boulder, CO, USA, June 1– 26, 2015, 2017, pp. 353–397. arXiv:1609.04036, doi:10.1142/10270.
- [5] I. Bengtsson, K. Życzkowski, Geometry of Quantum States: An Introduction to Quantum Entanglement, 2nd Edition, Cambridge University Press, 2017. doi:10.1017/9781139207010.
- [6] D. Chruściński, A. Jamiołkowski, Geometric phases in classical and quantum mechanics, Vol. 36 of Progress in Mathematical Physics, Birkhäuser Boston, Inc., Boston, MA, 2004. doi:10.1007/978-0-8176-8176-0.
- [7] M. A. Nielsen, I. L. Chuang, Quantum Computation and Quantum Information: 10th Anniversary Edition, Cambridge University Press, 2010. doi:10.1017/CB09780511976667.
- [8] M. Trzetrzelewski, J. Wosiek, Quantum systems in a cut Fock space, Acta Phys. Polon. B35 (2004) 1615–1624. arXiv:hep-th/0308007.

- [9] S. W. Hawking, Particle Creation by Black Holes, Commun. Math. Phys. 43 (1975) 199-220. doi:10.1007/BF02345020.
- [10] B. Broda, Total spectral distributions from Hawking radiation, Eur. Phys. J. C77 (11) (2017) 756. arXiv:1706.04557, doi:10.1140/epjc/ s10052-017-5336-1.
- [11] R. H. Dicke, Coherence in spontaneous radiation processes, Physical Review 93 (1) (1954) 99.
- [12] J. Martin, O. Giraud, P. Braun, D. Braun, T. Bastin, Multiqubit symmetric states with high geometric entanglement, Physical Review A 81 (6) (2010) 062347.