ON THE COMPARISON OF THE FRIDMAN INVARIANT AND THE SQUEEZING FUNCTION

FENG RONG, SHICHAO YANG

ABSTRACT. Let D be a bounded domain in \mathbb{C}^n , $n \geq 1$. In this paper, we study two biholomorphic invariants on D, the Fridman invariant $e_D(z)$ and the squeezing function $s_D(z)$. More specifically, we study the following two questions about the quotient invariant $m_D(z) = s_D(z)/e_D(z)$: 1) If $m_D(z_0) = 1$ for some $z_0 \in D$, is D biholomorphic to the unit ball? 2) Is $m_D(z)$ constantly equal to 1? We answer both questions negatively.

1. INTRODUCTION

Let D be a bounded domain in \mathbb{C}^n , $n \ge 1$, and \mathbb{B}^n the unit ball in \mathbb{C}^n . Denote by $B_D^k(z,r)$ the k_D -ball in D centered at $z \in D$ with radius r > 0, where k_D is the Kobayashi distance on D. Denote by $\mathcal{O}_u(D_1, D_2)$ the set of *injective* holomorphic maps from D_1 into D_2 .

In [8], Fridman introduced the following invariant:

$$h_D(z) = \inf\{1/r: B_D^k(z,r) \subset f(\mathbb{B}^n), f \in \mathcal{O}_u(\mathbb{B}^n, D), f(0) = z\}.$$

For comparison purposes, we call the following variant of $h_D(z)$ the Fridman invariant (cf. [6, 15]):

$$e_D(z) = \sup\{ \tanh(r) : B_D^k(z,r) \subset f(\mathbb{B}^n), f \in \mathcal{O}_u(\mathbb{B}^n,D), f(0) = z \}.$$

Obviously, $e_D(z) = \tanh(h_D^{-1}(z))$ and $h_D(z) = (\operatorname{artanh}(e_D(z)))^{-1}$.

In [3], Deng, Guan and Zhang introduced another invariant, called the *squeezing* function, as follows:

$$s_D(z) = \sup\{r: \ r\mathbb{B}^n \subset f(D), \ f \in \mathcal{O}_u(D, \mathbb{B}^n), \ f(z) = 0\}.$$

It is clear from the definitions that both the Fridman invariant and the squeezing function are invariant under biholomorphisms, and both take value in (0, 1]. Both invariants have attracted much attention in recent years, and we refer the readers to the recent survey article [5] and the references therein for various aspects of the current research on this topic.

Since they are similar in spirit to the Kobayashi-Eisenman volume form K_D and the Carathéodory volume form C_D , respectively, it is a natural problem to consider the comparison of these two biholomorphic invariants. Indeed, recently in [15], Nikolov and Verma have shown the following

²⁰¹⁰ Mathematics Subject Classification. 32H02, 32F45.

Key words and phrases. Fridman invariant, squeezing function, quotient invariant.

The authors are partially supported by the National Natural Science Foundation of China (grant no. 11871333).

Theorem 1. [15, Proposition 1] Let D be a bounded domain. Then, for any $z \in D$,

 $s_D(z) \le e_D(z).$

Regarding the quotient $M_D(z) := C_D(z)/K_D(z)$, the following classical result is well-known (cf. [17, Theorem E]).

Theorem 2. [10, Proposition 11.3.3] Let D be a bounded domain in \mathbb{C}^n . If there is a point $z \in D$ such that $M_D(z) = 1$, then D is biholomorphic to the unit ball \mathbb{B}^n .

Obviously, the quotient $m_D(z) := s_D(z)/e_D(z)$ is also a biholomorphic invariant, and Theorem 1 shows that $m_D(z) \leq 1$ for all $z \in D$. It is then natural to ask whether an analogous result to Theorem 2 holds for m_D (cf. [13, Concluding Remarks]). Our first main result answers this question *negatively*. More precisely, we have the following

Theorem 3. Let D be a bounded, balanced and convex domain in \mathbb{C}^n . Then,

 $m_D(0) = 1.$

Recall that a domain D is said to be *balanced* if for any $z \in D$, $\lambda z \in D$ for all $|\lambda| \leq 1$. And D is said to be *homogeneous* if the automorphism group of D is transitive. As an immediate corollary to Theorem 3, we have the following

Corollary 4. Let D be a bounded, balanced, convex and homogeneous domain in \mathbb{C}^n . Then,

$$m_D(z) \equiv 1.$$

On a bounded homogeneous domain D, both the Fridman invariant and the squeezing function are constants, which we call the *Fridman constant* ρ_D^e and the squeezing constant ρ_D^s , respectively. By Corollary 4, on a bounded, balanced, convex and homogeneous domain D, $\rho_D^e = \rho_D^s$, and we simply denote it by ρ_D .

It is well-known that the four types of classical Cartan domains are bounded, balanced, convex and homogeneous domains (see e.g. [14]), and so are their products. Combining Corollary 4 with [11, Theorem 2], we immediately have the following (cf. [2, Theorem 4.5])

Corollary 5. Let D_i , $1 \le i \le m$, be classical Cartan domains and $D = D_1 \times D_2 \times \cdots \times D_m$. Then,

$$\rho_D = (\rho_{D_1}^{-2} + \rho_{D_2}^{-2} + \dots + \rho_{D_m}^{-2})^{-1/2}.$$

Following Theorem 1 and Corollary 4, another natural question one can ask is whether $s_D(z)$ is always equal to $e_D(z)$. To our best knowledge, this question is open, mainly because both invariants are hard to compute explicitly. Our second main result answers this question *negatively*. More precisely, we have the following

Theorem 6. On the punctured unit disk Δ^* , there exists $\delta > 0$ such that

 $m_{\Delta^*}(z) < 1, \quad \forall \ 0 < |z| < \delta.$

Moreover, $\lim_{z\to 0} m_{\Delta^*}(z) = 0.$

Combining this with the stability results from [8] and [4], we also have the following

Corollary 7. Let $\delta > 0$ be as in Theorem 6. Fix $z_0 \in \Delta^*$ with $|z_0| < \delta$. Denote by A_r the annulus $\{z \in \mathbb{C} : r < |z| < 1\}$. Then, there exists $\epsilon > 0$ such that for all $0 < r < \epsilon$, we have

$$m_{A_r}(z_0) < 1.$$

In section 2, we prove Theorem 3. In section 3, we prove Theorem 6 and Corollary 7.

2. Bounded, balanced and convex domains

Let D be a bounded, balanced and convex domain in \mathbb{C}^n . Denote by k_D and c_D the Kobayashi and Carathéodory distance on D, respectively. The following Lempert's theorem is well-known:

Theorem 8. [12, Theorem 1] On a convex domain D, $k_D = c_D$.

The Minkowski function l_D is defined as (see e.g. [9])

$$l_D(z) = \inf\{t > 0 : z/t \in D\}, \quad z \in \mathbb{C}^n.$$

Note that $D = \{z \in \mathbb{C}^n : l_D(z) < 1\}.$

Combining Theorem 8 with [9, Proposition 2.3.1 (c)], we have the following key lemma.

Lemma 9. Let D be a bounded, balanced and convex domain. Then, for all $z \in D$,

$$\tanh(k_D(0,z)) = l_D(z).$$

We also need the following version of the Schwarz Lemma (see e.g. [16, Theorem 8.1.2]).

Lemma 10. Let D_1 and D_2 be bounded, balanced and convex domains, and f a holomorphic map from D_1 to D_2 with f(0) = 0. Then,

$$f(rD_1) \subset rD_2, \quad 0 < r \le 1.$$

We now prove Theorem 3 (cf. [1]).

Proof of Theorem 3. Since $s_D(0) \leq e_D(0)$ by Theorem 1, we only need to show that $e_D(0) \leq s_D(0)$.

For any $\epsilon > 0$, fix $f \in \mathcal{O}_u(\mathbb{B}^n, D)$ with f(0) = 0, such that $B_D^k(0, r) \subset f(\mathbb{B}^n) =: \Omega$ and $\tanh(r) > (1 - \epsilon)e_D(0)$.

By Lemma 9, we have

$$B_D^k(0,r) = \{ z \in D : k_D(0,z) < r \} = \{ z \in D : l_D(z) < \tanh(r) \}.$$

Thus, if we set $L(z) := \tanh(r)z$, then L is a biholomorphism between D and $B_D^k(0,r)$.

Denote by F the holomorphic inverse of f from Ω to \mathbb{B}^n . If $F(B_D^k(0,r)) = \mathbb{B}^n$, then D is biholomorphic to \mathbb{B}^n via $F \circ L$, which implies that $s_D(0) = 1 \ge e_D(0)$.

If $F(B_D^k(0,r)) \neq \mathbb{B}^n$, then let $t\mathbb{B}^n$ be the maximal ball contained in $F(B_D^k(0,r))$. By the definition of the squeezing function, we have $t \leq s_D(0) < 1$. Thus, there exists $p \in \mathbb{B}^n \setminus F(B_D^k(0,r))$ such that $\|p\| < (1+\epsilon)t \leq (1+\epsilon)s_D(0)$.

By Lemma 10, we get

$$(1-\epsilon)e_D(0) < \tanh(r) \le l_D(f(p)) \le ||p|| < (1+\epsilon)s_D(0).$$

Since ϵ is arbitrary, we must have $e_D(0) \leq s_D(0)$.

Remark 1. In the definition of $e_D(z)$, one can also use the Carathéodory distance c_D instead of the Kobayashi distance k_D , and get another related invariant, denoted by $e_D^c(z)$ (cf. [7, 15]). In [15, Proposition 1], it was actually proven that $s_D(z) \leq e_D^c(z) \leq e_D(z)$ for all $z \in D$. When D is a bounded convex domain, by Lempert's theorem, we have $e_D^c(z) = e_D(z)$. Thus, Theorem 3 and Corollary 4 also hold replacing e_D by e_D^c .

Remark 2. In [11], Kubota studied the following invariant:

$$\rho(D) := \sup\{r: \ r\mathbb{B}^n \subset f(D), \ f \in \mathcal{O}_u(D, \mathbb{B}^n)\}.$$

It is clear from the definitions that $\rho(D) = \sup_{z \in D} s_D(z)$. Thus, on a bounded homogeneous domain D, we have $\rho(D) = \rho_D^s$.

Remark 3. In Corollary 5, if each D_i is the unit disk Δ then $D = \Delta^n$ is a unit polydisk, and we get

$$\rho_{\Delta^n} = n^{-1/2}, \quad h_{\Delta^n}(z) \equiv (\operatorname{artanh}(\rho_D))^{-1} = 2\left(\log\frac{\sqrt{n}+1}{\sqrt{n}-1}\right)^{-1},$$

which recovers [13, Lemma 2.1].

3. The punctured disk and the annulus

First, we prove Theorem 6.

Proof of Theorem 6. Let $D = \Delta^*$ and $\Omega = \Delta \setminus (-1, 0]$. For any fixed $a \in (0, 1)$, by the Riemann mapping theorem, there exists a biholomorphic map $f : \Delta \to \Omega$ with f(0) = a. We need to estimate tanh(r) such that $B_D^k(a, r) \subset f(\Delta) = \Omega$.

By [9, Corollary 9.1.10], we have

$$\tanh(k_D(a,z)) = \left[\frac{\theta^2 + (\log|z| - \log a)^2}{\theta^2 + (\log|z| + \log a)^2}\right]^{1/2}, \quad \theta = \operatorname{Arg}(z) \in (-\pi,\pi].$$

In particular, for any $b \in (-1, 0)$, setting $x = -\log(-b)$ and $A = -\log a$, we have

$$\tanh(k_D(a,b)) = \left[\frac{\pi^2 + (x-A)^2}{\pi^2 + (x+A)^2}\right]^{1/2}.$$

Consider the function

$$h(x) = \frac{(x-A)^2 + \pi^2}{(x+A)^2 + \pi^2}.$$

One readily checks that h(x) takes the minimum value at $x = \sqrt{A^2 + \pi^2}$. Thus,

$$\min h(x) = h(\sqrt{A^2 + \pi^2}) = \frac{(\sqrt{A^2 + \pi^2} - A)^2 + \pi^2}{(\sqrt{A^2 + \pi^2} + A)^2 + \pi^2} = 1 - \frac{4A\sqrt{A^2 + \pi^2}}{(\sqrt{A^2 + \pi^2} + A)^2 + \pi^2}$$
$$\ge 1 - \frac{4A\sqrt{A^2 + \pi^2}}{4A\sqrt{A^2 + \pi^2}} = \frac{\pi^2}{4A\sqrt{A^2 + \pi^2} + \pi^2}.$$

Therefore, there exists $\delta > 0$ such that for any $0 < a < \delta$, we have

$$e_D(a)^2 \ge \frac{\pi^2}{4A\sqrt{A^2 + \pi^2} + \pi^2} > e^{-2A}.$$

By [3, Corollary 7.3], we know that $s_D(a) = a = e^{-A}$. Thus, we get $e_D(a) > s_D(a)$ for all $0 < a < \delta$. Moreover, the estimate above clearly implies that $\lim_{a\to 0} m_{\Delta^*}(a) = 0$.

Since D is circular, this completes the proof.

Remark 4. In [13, Lemma 2.2], Mahajan and Verma gave some estimates on $h_{\Delta^*}(z)$ as $z \to 0$. However, we found their estimates hard to use when converted to $e_{\Delta^*}(z)$. Thus, we gave a more straightforward estimate above, using [9, Corollary 9.1.10].

To prove Corollary 7, we need the following two stability theorems from [8] and [4], adapted to our setting.

Theorem 11. [8, Theorem 2.1] Let D be a completely hyperbolic domain and $\{D_k\}_{k=1}^{\infty}$ a sequence of subdomains such that $D_k \subset D_{k+1}$ and $\cup_k D_k = D$. Then, for any $z_0 \in D$,

$$\lim_{k \to \infty} e_{D_k}(z_0) = e_D(z_0).$$

Theorem 12. [4, Theorem 2.1] Let D be a bounded domain and $\{D_k\}_{k=1}^{\infty}$ a sequence of subdomains such that $D_k \subset D_{k+1}$ and $\bigcup_k D_k = D$. Then, for any $z_0 \in D$,

$$\lim_{k \to \infty} s_{D_k}(z_0) = s_D(z_0).$$

Proof of Corollary 7. Suppose that there does not exist $\epsilon > 0$ such that for all $0 < r < \epsilon$ we have $m_{A_r}(z_0) < 1$. Then, there exists a decreasing sequence of $r_k \to 0$ such that $m_{A_{r_k}}(z_0) = 1$ for all k. Since Δ^* is completely hyperbolic, $A_{r_k} \subset A_{r_{k+1}}$ and $\cup_k A_{r_k} = \Delta^*$, Theorems 11 and 12 apply. Thus, $m_{\Delta^*}(z_0) = \lim_{k\to\infty} m_{A_{r_k}}(z_0) = 1$, which contradicts Theorem 6.

References

- H. Alexander; Extremal holomorphic imbeddings between the ball and polydisc, Proc. Amer. Math. Soc. 68 (1978), 200-202.
- [2] C.-H. Chu, K.-T. Kim, S. Kim; Infinite dimensional holomorphic homogeneous regular domains, J. Geom. Anal. 30 (2020), 223-247.
- [3] F. Deng, Q. Guan, L. Zhang; Some properties of squeezing functions on bounded domains, Pacific J. Math. 257 (2012), 319-341.
- [4] F. Deng, Q. Guan, L. Zhang; Properties of squeezing functions and global transformations of bounded domains, Trans. Amer. Math. Soc. 368 (2016), 2679-2696.
- [5] F. Deng, Z. Wang, L. Zhang, X. Zhou; Holomorphic invariants of bounded domains, J. Geom. Anal. 30 (2020), 1204-1217.
- [6] F. Deng, X. Zhang; Fridman's invariant, squeezing functions, and exhausting domains, Acta Math. Sin. (Engl. Ser.) 35 (2019), 1723-1728.

 B.L. Fridman; On the imbedding of a strictly pseudoconvex domain in a polyhedron, Dokl. Akad. Nauk SSSR 249 (1979) = Soviet Math. Dokl. 20 (1979), 1228-1232.

- [8] B.L. Fridman; Biholomorphic invariants of a hyperbolic manifold and some applications, Trans. Amer. Math. Soc. 276 (1983), 685-698.
- [9] M. Jarnicki, P. Pflug; Invariant Distances and Metrics in Complex Analysis, 2nd ext. ed., de Gruyter Expos. in Math. 9, Walter de Gruyter GmbH, Berlin, 2013.
- [10] S.G. Krantz; Function Theory of Several Complex Variables, 2nd ed., Amer. Math. Soc., Providence, RI, 2001.
- [11] Y. Kubota; A note on holomorphic imbeddings of the classical Cartan domains into the unit ball, Proc. Amer. Math. Soc. 85 (1982), 65-68.
- [12] L. Lempert; Holomorphic retracts and intrinsic metrics in convex domains, Anal. Math. 8 (1982), 257-261.

[13] P. Mahajan, K. Verma; A comparison of two biholomorphic invariants, Internat. J. Math. 30 (2019), 1950012, 16 pp.

- [14] N. Mok; Metric Rigidity Theorems on Hermitian Locally Symmetric Manifolds, Series in Pure Math. 6, World Sci. Publ. Co., Inc., Teaneck, NJ, 1989.
- [15] N. Nikolov, K. Verma; On the squeezing function and Fridman invariants, J. Geom. Anal. 30 (2020), 1218-1225.

[16] W. Rudin; Function Theory in The Unit Ball of \mathbb{C}^n , Grund. der Math. Wissen. **241**, Springer-Verlag, New York-Berlin, 1980.

[17] B. Wong; Characterizations of the ball in \mathbb{C}^n by its automorphism group, Invent. Math. 41 (1977), 253-257.

School of Mathematical Sciences, Shanghai Jiao Tong University, 800 Dong Chuan Road, Shanghai, 200240, P.R. China

 $Email \ address: \verb"frong@sjtu.edu.cn" \\$

School of Mathematical Sciences, Shanghai Jiao Tong University, 800 Dong Chuan Road, Shanghai, 200240, P.R. China

Email address: yangshichao@sjtu.edu.cn