

A BIVARIATE GENERATING FUNCTION FOR ZETA VALUES AND RELATED SUPERCONGRUENCES

ROBERTO TAURASO

ABSTRACT. By using the Wilf-Zeilberger method, we prove a novel finite combinatorial identity related to a bivariate generating function for $\zeta(2+r+2s)$ (an extension of a Bailey-Borwein-Bradley Apéry-like formula for even zeta values). Such identity is then applied to show several supercongruences.

1. INTRODUCTION

The bivariate formula

$$\sum_{k=1}^{\infty} \frac{k}{k^4 - a^2 k^2 - b^4} = \frac{1}{2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1} (5k^2 - a^2)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} ((j^2 - a^2)^2 + 4b^4)}{\prod_{j=1}^k (j^4 - a^2 j^2 - b^4)} \quad (1)$$

has been first conjectured by H. Cohen and then proved independently by T. Rivoal [11, Theorem 1.1] and D. M. Bradley [3, Theorem 1] by reducing it to the finite combinatorial identity

$$\sum_{k=1}^n \binom{2k}{k} \frac{(5k^2 - a^2) \prod_{j=1}^{k-1} ((n^2 - j^2)(n^2 + j^2 - a^2))}{\prod_{j=1}^k (n^2 + (n-j)^2 - a^2)(n^2 + (n+j)^2 - a^2)} = \frac{2}{n^2 - a^2} \quad (2)$$

and by Kh. and T. Hessami Pilehrood [6, Theorem 1] by applying the Wilf-Zeilberger theory. Since the left-hand side of (1) can be written as the generating function of $\zeta(3+2r+4s)$,

$$\sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \binom{r+s}{r} \zeta(3+2r+4s) a^{2r} b^{4s},$$

it follows that, by extracting the coefficients for $(r, s) = (0, 0)$ and $(r, s) = (1, 0)$, we obtain the Apéry-like identities

$$\zeta(3) = \frac{5}{2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k^3 \binom{2k}{k}} \quad \text{and} \quad \zeta(5) = \frac{1}{2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{\binom{2k}{k}} \left(\frac{4}{k^5} - \frac{5H_{k-1}(2)}{k^3} \right) \quad (3)$$

where $H_n(s) = \sum_{j=1}^n \frac{1}{j^s}$ is the *harmonic sum* of weight s . For more details about Apéry-like series see also [1, 4, 6, 5].

Date: February 3, 2020.

2010 Mathematics Subject Classification. 11A07, 05A19, 11B65, 11M06.

Key words and phrases. congruences, central binomial coefficients, harmonic numbers, Wilf-Zeilberger method, zeta values, Apéry-like series.

Here we consider a similar bivariate formula

$$\sum_{k=1}^{\infty} \frac{1}{k^2 - ak - b^2} = \sum_{k=1}^{\infty} \frac{(3k - a)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} (j^2 - a^2 - 4b^2)}{\prod_{j=1}^k (j^2 - aj - b^2)} \quad (4)$$

where the left-hand side is the generating function of $\zeta(2 + r + 2s)$,

$$\sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \binom{r+s}{r} \zeta(2 + r + 2s) a^r b^{2s}.$$

For $a = 0$, (4) yields a formula due to D. H. Bailey, J. M. Borwein, and D. M. Bradley,

$$\sum_{s=0}^{\infty} \zeta(2 + 2s) b^{2s} = \sum_{k=1}^{\infty} \frac{1}{k^2 - b^2} = 3 \sum_{k=1}^{\infty} \frac{1}{\binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} (j^2 - 4b^2)}{\prod_{j=1}^k (j^2 - b^2)}$$

which appeared in [2, Theorem 1.1]. Moreover, for $(r, s) = (1, 0)$ and $(r, s) = (0, 1)$, we get the Apéry-like identities

$$\zeta(3) = \sum_{k=1}^{\infty} \frac{1}{\binom{2k}{k}} \left(\frac{2}{k^3} + \frac{3H_{k-1}(1)}{k^2} \right) \quad \text{and} \quad \zeta(4) = 3 \sum_{k=1}^{\infty} \frac{1}{\binom{2k}{k}} \left(\frac{1}{k^4} - \frac{3H_{k-1}(2)}{k^2} \right). \quad (5)$$

Replacing a by $2a$ and then letting $x^2 = a^2 + b^2$ in (4) we find the equivalent identity

$$\sum_{k=1}^{\infty} \frac{1}{(k - a)^2 - x^2} = \sum_{k=1}^{\infty} \frac{(3k - 2a)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} (j^2 - 4x^2)}{\prod_{j=1}^k ((j - a) - x^2)}$$

which has been proved by Kh. and T. Hessami Pilehrood [7, (24)].

Again, in the same spirit of what has been done for (1), our proof of the identity (4) is reduced to show the following novel finite identity

$$\sum_{k=1}^n \binom{2k}{k} \frac{3k - 2n + a}{k^2 - a^2} \cdot \prod_{j=1}^{k-1} \frac{(j - n)(j - n + a)}{j^2 - a^2} = \frac{2}{n - a}. \quad (6)$$

In [15, Theorem 4.2] the author established that for any prime $p > 5$,

$$\sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} \equiv -\frac{8H_{p-1}(1)}{3} \pmod{p^4}, \quad (7)$$

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} \binom{2k}{k} \equiv \frac{4}{5} \left(\frac{H_{p-1}(1)}{p} + 2pH_{p-1}(3) \right) \pmod{p^4}. \quad (8)$$

Thanks to the finite identities (2) and (6), we managed to improve congruence (7) and to show several other congruences. The main results are as follows: for any prime $p > 5$,

$$\sum_{k=1}^{p-1} \frac{1}{k^3} \binom{2k}{k} \equiv -\frac{2H_{p-1}(1)}{p^2} \pmod{p^2}, \quad (9)$$

$$\sum_{k=1}^{p-1} \binom{2k}{k} \frac{H_k(2)}{k} \equiv \frac{2H_{p-1}(1)}{3p^2} \pmod{p^2}. \quad (10)$$

These congruences are known modulo p (see [8, Theorem 2]) and they confirm modulo p^2 the following conjecture by Z.-W. Sun : for each prime $p > 7$,

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{1}{k^3} \binom{2k}{k} &\equiv -\frac{2H_{p-1}(1)}{p^2} - \frac{13H_{p-1}(3)}{27} \pmod{p^4}, \\ \sum_{k=1}^{p-1} \binom{2k}{k} \frac{H_k(2)}{k} &\equiv \frac{2H_{p-1}(1)}{3p^2} - \frac{38H_{p-1}(3)}{81} \pmod{p^3}. \end{aligned}$$

the first one appeared in [13, Conjecture 1.1] and the second one in [14, Conjecture 5.1].

2. PRELIMINARIES CONCERNING MULTIPLE HARMONIC SUMS

We define the *multiple harmonic sum* as

$$H_n(s_1, \dots, s_r) = \sum_{1 \leq k_1 < k_2 < \dots < k_r \leq n} \frac{1}{k_1^{s_1} k_2^{s_2} \dots k_r^{s_r}}$$

where $n \geq r > 0$ and each s_i is a positive integer. The sum $s_1 + s_2 + \dots + s_r$ is the weight of the multiple sum. Furthermore, by $\{s_1, s_2, \dots, s_j\}^m$ we denote the sequence of length mj with m repetitions of (s_1, s_2, \dots, s_j) .

By [12, Theorem 5.1]), for any prime $p > s + 2$ we have

$$H_{p-1}(s) \equiv \begin{cases} -\frac{s(s+1)}{2(s+2)} p^2 B_{p-s-2} \pmod{p^3} & \text{if } s \text{ is odd,} \\ \frac{s}{s+1} p B_{p-s-1} \pmod{p^2} & \text{if } s \text{ is even.} \end{cases}$$

where B_n be the n -th Bernoulli number.

Let $p > 5$ be a prime, then by [15, Theorem 2.1],

$$H_{p-1}(2) \equiv -\frac{2H_{p-1}(1)}{p} - \frac{pH_{p-1}(3)}{3} \pmod{p^4}. \quad (11)$$

Moreover, by [8, Lemma 3],

$$H_{p-1}(1, 2) \equiv -\frac{3H_{p-1}(1)}{p^2} - \frac{5H_{p-1}(3)}{12} \pmod{p^3}$$

and by [16, Proposition 3.7] and [9, Theorem 4.5]

$$H_{p-1}(1, 1, 2) \equiv -\frac{11H_{p-1}(3)}{12p} \pmod{p^2}, \quad H_{p-1}(1, 1, 1, 2) \equiv -\frac{5H_{p-1}(3)}{6p^2} \pmod{p}.$$

Finally, by [16, Theorem 3.2],

$$H_{p-1}(2, 2) \equiv \frac{H_{p-1}(3)}{3p}, \quad H_{p-1}(1, 3) \equiv \frac{3H_{p-1}(3)}{4p} \pmod{p^2}$$

and by [16, Theorem 3.5],

$$H_{p-1}(2, 1, 2) \equiv 0, \quad H_{p-1}(1, 2, 2) \equiv \frac{5H_{p-1}(3)}{4p^2}, \quad H_{p-1}(1, 1, 3) \equiv -\frac{5H_{p-1}(3)}{12p^2} \pmod{p}.$$

3. PROOFS OF THE GENERATING FUNCTION (4) AND THE RELATED COMBINATORIAL IDENTITY (6)

By partial fraction decomposition with respect to b^2 , we get

$$\frac{\prod_{j=1}^{k-1}(j^2 - a^2 - 4b^2)}{\prod_{j=1}^k(j^2 - aj - b^2)} = \sum_{n=1}^k \frac{C_{n,k}(a)}{n^2 - an - b^2}$$

where

$$C_{n,k}(a) = \frac{\prod_{j=1}^{k-1}(j^2 - (a - 2n)^2)}{\prod_{j=1, j \neq n}^k(j - n)(j + n - a)}.$$

Hence, by inverting the summations order, the identity (4) can be written as

$$\sum_{n=1}^{\infty} \frac{1}{n^2 - an - b^2} = \sum_{k=1}^{\infty} \frac{(3k - a)}{k \binom{2k}{k}} \sum_{n=1}^k \frac{C_{n,k}(a)}{n^2 - an - b^2} = \sum_{n=1}^{\infty} \frac{1}{n^2 - an - b^2} \sum_{k=n}^{\infty} \frac{(3k - a)C_{n,k}(a)}{k \binom{2k}{k}}.$$

It follows that (4) holds as soon as

$$1 = \sum_{k=n}^{\infty} \frac{(3k - a)C_{n,k}(a)}{k \binom{2k}{k}} = \sum_{k=n}^{\infty} \frac{(3k - a)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1}(j^2 - (a - 2n)^2)}{\prod_{j=1, j \neq n}^k(j - n)(j + n - a)}. \quad (12)$$

Taking the same approach given in [11] for the proof of (1), the above formula is equivalent to this finite combinatorial identity

$$\sum_{k=1}^n \binom{2k}{k} (3k - a) \frac{\prod_{j=1}^{k-1}(j - n)(j + n - a)}{\prod_{j=1}^k(j^2 - (a - 2n)^2)} = \frac{2}{a - n}. \quad (13)$$

Both identities (12) and (13) are consequences of the next theorem after setting $z = 2n - a$.

Theorem 1. *For any positive integer n ,*

$$\sum_{k=1}^n \binom{2k}{k} (3k - 2n + z) \frac{\prod_{j=1}^{k-1}(j - n)(j - n + z)}{\prod_{j=1}^k(j^2 - z^2)} = \frac{2}{n - z}, \quad (14)$$

and

$$\sum_{k=n}^{\infty} \frac{(3k-2n+z)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} (j^2 - z^2)}{\prod_{j=1, j \neq n}^k (j-n)(j-n+z)} = 1. \quad (15)$$

Proof. Let

$$F(n, k) = \binom{2k}{k} (3k-2n+z) \frac{\prod_{j=0}^{k-1} (j-n)(j-n+z)}{\prod_{j=1}^k (j^2 - z^2)},$$

and

$$G(n, k) = \frac{k(k^2 - z^2)F(n, k)}{(2n-3k-z)(n+1-k)(n+1-k-z)}.$$

Then (F, G) is a Wilf-Zeilberger pair, or WZ pair, which means that they satisfy the relation

$$F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k).$$

In order to prove (14), it suffices to prove that $S_n := \sum_{k=1}^n F(n, k) = 2n$. Now $S_1 = F(1, 1) = 2$. Moreover,

$$S_{n+1} - S_n = \sum_{k=1}^{n+1} F(n+1, k) - \sum_{k=1}^{n+1} F(n, k) = G(n, n+2) - G(n, 1) = 2$$

because $F(n, n+1) = G(n, n+2) = 0$ and $G(n, 1) = -2$.

In a similar way, we show (15) by considering the WZ pair given by

$$F(n, k) = \frac{(3k-2n+z)}{k \binom{2k}{k}} \cdot \frac{\prod_{j=1}^{k-1} (j^2 - z^2)}{\prod_{j=1, j \neq n}^k (j-n)(j-n+z)},$$

and

$$G(n, k) = \frac{2(2k-1)(k-n)F(n, k)}{n(2n-3k-z)(n-z)}.$$

□

4. MORE BINOMIAL IDENTITIES

Here we collect a few identities, apparently new, involving the binomial coefficients $\binom{2k}{k}$ and $\binom{n+k}{k}$ which will play a crucial role in the next sections.

Theorem 2. *For any positive integer n ,*

$$\frac{3}{2} \sum_{k=1}^n \frac{1}{k} \binom{2k}{k} = \sum_{k=1}^n \frac{1}{k} \binom{n+k}{k} + H_n(1) \quad (16)$$

$$\sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(1)}{2k} - \frac{1}{k^2} \right) = \sum_{k=1}^n \binom{n+k}{k} \frac{H_k(1)}{k} - H_n(2) \quad (17)$$

$$\sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(2)}{k} - \frac{1}{2k^3} \right) = \sum_{k=1}^n \binom{n+k}{k} \frac{H_k(2) + H_n(2)}{k} + H_n(2)H_n(1) - H_n(1, 2) \quad (18)$$

Proof. Let us consider the WZ pair

$$F(n, k) = \frac{1}{k} \binom{n+k}{k} \quad \text{and} \quad G(n, k) = \frac{k}{(n+1)^2} \binom{n+k}{k}$$

then

$$\begin{aligned} S_{n+1} - S_n &= F(n+1, n+1) + \sum_{k=1}^n (G(n, k+1) - G(n, k)) \\ &= F(n+1, n+1) + G(n, n+1) - G(n, 1) \\ &= \frac{3/2}{n+1} \binom{2(n+1)}{n+1} - \frac{1}{n+1} \end{aligned}$$

where $S_n := \sum_{k=1}^n F(n, k)$. Thus

$$S_n = \frac{3}{2} \sum_{k=1}^n \frac{1}{k} \binom{2k}{k} - H_n(1)$$

and we may conclude that (16) holds.

Now let $S_n^{(1)} := \sum_{k=1}^n F(n, k)H_k(1)$ then

$$\begin{aligned} S_{n+1}^{(1)} - S_n^{(1)} &= F(n+1, n+1)H_{n+1}(1) \\ &\quad + \sum_{k=1}^n \left(G(n, k+1)H_k(1) - G(n, k) \left(H_{k-1}(1) + \frac{1}{k} \right) \right) \\ &= F(n+1, n+1)H_{n+1}(1) + G(n, n+1)H_n(1) - \sum_{k=1}^n \frac{G(n, k)}{k} \\ &= \binom{2(n+1)}{n+1} \left(\frac{3H_{n+1}(1)}{2(n+1)} - \frac{1}{(n+1)^2} \right) + \frac{1}{(n+1)^2} \end{aligned}$$

where we used $\sum_{k=1}^n \binom{n+k}{k} = \frac{1}{2} \binom{2(n+1)}{n+1} - 1$. Hence we find that

$$S_n^{(1)} = \sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(1)}{2k} - \frac{1}{k^2} \right) + H_n(2)$$

which implies (17).

Let $S_n^{(2)} := \sum_{k=1}^n F(n, k)H_k(2)$ then

$$\begin{aligned}
S_{n+1}^{(2)} - S_n^{(2)} &= F(n+1, n+1)H_{n+1}(2) \\
&\quad + \sum_{k=1}^n \left(G(n, k+1)H_k(2) - G(n, k) \left(H_{k-1}(2) + \frac{1}{k^2} \right) \right) \\
&= F(n+1, n+1)H_{n+1}(2) + G(n, n+1)H_n(2) - \sum_{k=1}^n \frac{G(n, k)}{k^2} \\
&= \binom{2(n+1)}{n+1} \left(\frac{3H_{n+1}(2)}{2(n+1)} - \frac{1}{2(n+1)^3} \right) - \frac{S_n}{(n+1)^2}
\end{aligned}$$

where we applied

$$\sum_{k=1}^n \frac{G(n, k)}{k^2} = \frac{1}{(n+1)^2} \sum_{k=1}^n F(n, k) = \frac{S_n}{(n+1)^2}.$$

Therefore

$$\begin{aligned}
S_n^{(2)} &= \sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(2)}{2k} - \frac{1}{2k^3} \right) - \sum_{k=1}^n \frac{S_{k-1}}{k^2} \\
&= \sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(2)}{2k} - \frac{1}{2k^3} \right) - \frac{3}{2} \sum_{k=1}^n \frac{1}{k^2} \sum_{j=1}^{k-1} \frac{1}{j} \binom{2j}{j} + H_n(1, 2) \\
&= \sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(2)}{2k} - \frac{1}{2k^3} \right) - \frac{3}{2} \sum_{j=1}^n \frac{1}{j} \binom{2j}{j} (H_n(2) - H_j(2)) + H_n(1, 2) \\
&= \sum_{k=1}^n \binom{2k}{k} \left(\frac{3H_k(2)}{k} - \frac{1}{2k^3} \right) - \frac{3H_n(2)}{2} \sum_{k=1}^n \frac{1}{k} \binom{2k}{k} + H_n(1, 2)
\end{aligned}$$

and the proof of (18) is complete. \square

5. PROOFS OF THE MAIN SUPERCONGRUENCES

Theorem 3. *For any prime $p > 3$,*

$$\sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} \equiv -\frac{8H_{p-1}(1)}{3} - \frac{5p^2 H_{p-1}(3)}{3} \pmod{p^5} \quad (19)$$

Moreover, for any prime $p > 5$,

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{1}{k^3} \binom{2k}{k} &\equiv -\frac{2H_{p-1}(1)}{p^2} \pmod{p^2}, \\ \sum_{k=1}^{p-1} \binom{2k}{k} \frac{H_k(2)}{k} &\equiv \frac{2H_{p-1}(1)}{3p^2} \pmod{p^2}. \end{aligned}$$

Proof. We first note that

$$\binom{p-1+k}{k} = \frac{p}{k} \binom{p+k-1}{k-1} = \frac{p}{k} \prod_{j=1}^{k-1} \left(1 + \frac{p}{j}\right) = \frac{1}{k} \sum_{j=0}^{k-1} p^{j+1} H_{k-1}(\{1\}^j). \quad (20)$$

Therefore, by (16) with $n = p-1$ we obtain the desired congruence (19),

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} &= \frac{2}{3} \left(H_{p-1}(1) + \sum_{j=0}^{p-2} p^{j+1} H_{p-1}(\{1\}^j, 2) \right) \\ &\equiv \frac{2}{3} (H_{p-1}(1) + pH_{p-1}(2) + p^2 H_{p-1}(1, 2) \\ &\quad + p^3 H_{p-1}(1, 1, 2) + p^4 H_{p-1}(1, 1, 1, 2)) \\ &\equiv -\frac{8H_{p-1}(1)}{3} - \frac{5p^2 H_{p-1}(3)}{3} \pmod{p^5}. \end{aligned}$$

By letting $z = 2n$ in (14) we have

$$\sum_{k=1}^n \binom{2k}{k} \frac{k}{k^2 - 4n^2} \prod_{j=1}^{k-1} \frac{j^2 - n^2}{j^2 - 4n^2} = -\frac{2}{3n}.$$

Let $n = p > 5$ be a prime and move the p -th term of the sum to the right-hand side,

$$\sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} \frac{1}{1 - \frac{4p^2}{k^2}} \prod_{j=1}^{k-1} \frac{1 - \frac{p^2}{j^2}}{1 - \frac{4p^2}{j^2}} = \frac{2}{3p} \left(\frac{1}{2} \binom{2p}{p} \prod_{j=1}^{p-1} \frac{1 - \frac{p^2}{j^2}}{1 - \frac{4p^2}{j^2}} - 1 \right).$$

The left-hand side modulo p^4 is congruent to

$$\sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} \left(1 + \frac{4p^2}{k^2}\right) \prod_{j=1}^{k-1} \left(1 + \frac{3p^2}{j^2}\right) \equiv \sum_{k=1}^{p-1} \frac{1}{k} \binom{2k}{k} + p^2 \sum_{k=1}^{p-1} \binom{2k}{k} \left(\frac{1}{k^3} + \frac{3H_k(2)}{k}\right).$$

On the other hand, by [15, Theorem 2.4],

$$\frac{1}{2} \binom{2p}{p} \equiv 1 + 2pH_{p-1}(1) + \frac{2p^3 H_{p-1}(3)}{3} \equiv 1 - p^2 H_{p-1}(2) - \frac{p^4 H_{p-1}(4)}{2} \pmod{p^6}, \quad (21)$$

the right-hand side is

$$\begin{aligned}
\frac{1}{2} \binom{2p}{p} \prod_{j=1}^{p-1} \frac{1 - \frac{p^2}{j^2}}{1 - \frac{4p^2}{j^2}} &\equiv \frac{1}{2} \binom{2p}{p} \prod_{j=1}^{p-1} \left(1 + \frac{3p^2}{j^2} + \frac{12p^4}{j^4} \right) \\
&\equiv \left(1 - p^2 H_{p-1}(2) - \frac{p^4 H_{p-1}(4)}{2} \right) \\
&\quad \cdot \left(1 + 3p^2 H_{p-1}(2) + 12p^4 H_{p-1}(4) + 9p^4 H_{p-1}(2, 2) \right) \\
&\equiv 1 + 2p^2 H_{p-1}(2) + p^4 \left(\frac{17 H_{p-1}(4)}{2} + 3 H_{p-1}(2, 2) \right) \\
&\equiv 1 + 2p^2 H_{p-1}(2) \pmod{p^5}.
\end{aligned}$$

where $2H_{p-1}(2, 2) = (H_{p-1}(2))^2 - H_{p-1}(4) \equiv 0 \pmod{p}$. Finally, by (19),

$$\sum_{k=1}^{p-1} \binom{2k}{k} \left(\frac{1}{k^3} + \frac{3H_k(2)}{k} \right) \equiv \frac{8H_{p-1}(1)}{3p^2} + \frac{5H_{p-1}(3)}{3} + \frac{4pH_{p-1}(2)}{3} \equiv 0 \pmod{p^2}. \quad (22)$$

where we used (11).

By (18), with $n = p - 1$, we have that

$$\begin{aligned}
\sum_{k=1}^{p-1} \binom{2k}{k} \left(\frac{3H_k(2)}{k} - \frac{1}{2k^3} \right) &= p \sum_{k=1}^{p-1} \prod_{j=1}^{k-1} \left(1 + \frac{p}{j} \right) \frac{H_k(2) + H_{p-1}(2)}{k^2} \\
&\quad + H_{p-1}(2)H_{p-1}(1) - H_{p-1}(1, 2) \\
&\equiv p \sum_{k=1}^{p-1} \frac{H_k(2)}{k^2} - H_{p-1}(1, 2) \\
&= pH_{p-1}(2, 2) + pH_{p-1}(4) - H_{p-1}(1, 2) \\
&\equiv -H_{p-1}(1, 2) = \frac{3H_{p-1}(1)}{p^2} \pmod{p^2}. \quad (23)
\end{aligned}$$

The proof is completely as soon as we combine properly congruences (22) and (23). \square

6. FINALE - TWO APÉRY-LIKE CONGRUENCES

The following congruences are related to the second series in (3) and to the first series in (5).

Theorem 4. *For any prime $p > 3$,*

$$\sum_{k=1}^{p-1} \binom{2k}{k} \left(\frac{2}{k^2} - \frac{3H_k(1)}{k} \right) \equiv \frac{2H_{p-1}(1)}{p} + 3pH_{p-1}(3) \pmod{p^4}, \quad (24)$$

$$\sum_{k=1}^{p-1} (-1)^k \binom{2k}{k} \left(\frac{4}{k^4} + \frac{5H_k(2)}{k^2} \right) \equiv -H_{p-1}(4) \pmod{p^2}. \quad (25)$$

Proof. By (17) with $n = p - 1$ and by (20)

$$\begin{aligned}
\sum_{k=1}^{p-1} \binom{2k}{k} \left(\frac{3H_k(1)}{2k} - \frac{1}{k^2} \right) &= \sum_{k=1}^{p-1} \frac{H_k(1)}{k^2} \sum_{j=0}^{k-1} p^{j+1} H_{k-1}(\{1\}^j) - H_{p-1}(2) \\
&\equiv p \sum_{k=1}^{p-1} \frac{H_{k-1}(1) + \frac{1}{k}}{k^2} (1 + pH_{k-1}(1) + p^2 H_{k-1}(1, 1)) - H_{p-1}(2) \\
&\equiv -H_{p-1}(2) + pH_{p-1}(1, 2) + pH_{p-1}(3) \\
&\quad + 2p^2 H_{p-1}(1, 1, 2) + p^2 H_{p-1}(2, 2) + p^2 H_{p-1}(1, 3) \\
&\quad + 3p^3 H_{p-1}(1, 1, 1, 2) + p^3 H_{p-1}(2, 1, 2) + p^3 H_{p-1}(1, 2, 2) \\
&\quad + p^3 H_{p-1}(1, 1, 3) \\
&\equiv -\frac{H_{p-1}(1)}{p} - \frac{3pH_{p-1}(3)}{2} \pmod{p^4}
\end{aligned}$$

where at the last step we applied the results mentioned in the preliminaries.

By comparing the coefficient of a^2 in the expansion of both sides of (2) at $a = 0$ we have

$$\sum_{k=1}^n \binom{2k}{k} \frac{5k^2}{4n^4 + k^4} \prod_{j=1}^{k-1} \frac{n^4 - j^4}{4n^4 + j^4} \left(\frac{1}{5k^2} + \sum_{j=1}^{k-1} \frac{1}{n^2 + j^2} - 2 \sum_{j=1}^k \frac{2n^2 + j^2}{4n^4 + j^4} \right) = -\frac{2}{n^4}.$$

Let $n = p > 3$ be a prime then move to the right-hand side the p -th term of the sum on the left. Thus, the left-hand side modulo p^2 is congruent to

$$\sum_{k=1}^{p-1} (-1)^{k-1} \binom{2k}{k} \frac{5}{k^2} \left(\frac{1}{5k^2} + H_{k-1}(2) - 2H_k(2) \right) = \sum_{k=1}^{p-1} (-1)^k \binom{2k}{k} \left(\frac{4}{k^4} + \frac{5H_k(2)}{k^2} \right).$$

The right-hand side multiplied by p^4 is

$$-2 - \binom{2p}{p} \prod_{j=1}^{p-1} \frac{p^4 - j^4}{4p^4 + j^4} \left(\frac{1}{5} + p^2 \sum_{j=1}^{p-1} \frac{1}{p^2 + j^2} - 2p^2 \sum_{j=1}^p \frac{2p^2 + j^2}{4p^4 + j^4} \right) \quad (26)$$

and it remains to verify that it is congruent to $-p^4 H_{p-1}(4)$ modulo p^6 . We note that

$$\begin{aligned}
\prod_{j=1}^{p-1} \frac{p^4 - j^4}{4p^4 + j^4} &\equiv 1 - 5p^4 H_{p-1}(4) \pmod{p^6}, \\
p^2 \sum_{j=1}^{p-1} \frac{1}{p^2 + j^2} &\equiv p^2 H_{p-1}(2) - p^4 H_{p-1}(4) \pmod{p^6}, \\
2p^2 \sum_{j=1}^p \frac{2p^2 + j^2}{4p^4 + j^4} &\equiv \frac{6}{5} + 2p^2 H_{p-1}(2) + 4p^4 H_{p-1}(4) \pmod{p^6},
\end{aligned}$$

Hence, by (21), (26) simplifies to

$$-2 + 2 \left(1 - p^2 H_{p-1}(2) - \frac{p^4 H_{p-1}(4)}{2} \right) (1 + p^2 H_{p-1}(2)) \equiv -p^4 H_{p-1}(4) \pmod{p^6}$$

and the proof is finished. \square

REFERENCES

- [1] G. Almkvist, A. Granville, *Borwein and Bradley's Apéry-like formulae for $\zeta(4n+3)$* , Experiment. Math., **8** (1999), 197–203.
- [2] D. H. Bailey, J. M. Borwein, D. M. Bradley, *Experimental determination of Apéry-like identities for $\zeta(2n+2)$* , Experiment. Math. **15** (2006), 281–289.
- [3] D. M. Bradley, *Hypergeometric functions related to series acceleration formulas*, Contemporary Math. **457** (2008), 113–125.
- [4] J. M. Borwein, D. M. Bradley, *Empirically determined Apéry-like formulae for $\zeta(4n+3)$* , Experiment. Math. **6** (1997), 181–194.
- [5] W. Chu, *Hypergeometric approach to Apéry-like series*, Integral Transforms Spec. Funct. **28** (2017), 505–518.
- [6] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Simultaneous generation for zeta values by the Markov-WZ method*, Discrete Math. Theor. Comput. Sci. **10** (2008), 115–123.
- [7] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Bivariate identities for values of the Hurwitz zeta function and supercongruences* Electron. J. Comb. **18** (2012), P35, 30 p.
- [8] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Congruences arising from Apéry-type series for zeta values*, Adv. Appl. Math. **49** (2012), 218–238.
- [9] Kh. Hessami Pilehrood, T. Hessami Pilehrood, R. Tauraso, *New properties of multiple harmonic sums modulo p and p -analogues of Leshchiner's series*, Trans. Am. Math. Soc. **366** (2014), 3131–3159.
- [10] S. Mattarei, R. Tauraso, *Congruences for central binomial sums and finite polylogarithms*, J. Number Theory **133** (2013), 131–157.
- [11] T. Rivoal, *Simultaneous generation of Koecher and Almkvist-Granville's Apéry-like formulae*, Exp. Math. **13** (2004), 503–508.
- [12] Z.-H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), 193–223.
- [13] Z.-W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54** (2011), 2509–2535.
- [14] Z.-W. Sun, *A new series for π^3 and related congruences*, Internat. J. Math. **26** (2015), 1550055 (23 pages).
- [15] R. Tauraso, *More congruences for central binomial coefficients*, J. Number Theory **130** (2010), 2639–2649.
- [16] J. Zhao, *Wolstenholme type theorem for multiple harmonic sums*, Int. J. Number Theory **4** (2008), 73–106.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY

E-mail address: tauraso@mat.uniroma2.it