FRAMED MOTIVES OF ALGEBRAIC VARIETIES (AFTER V. VOEVODSKY)

GRIGORY GARKUSHA AND IVAN PANIN

In memory of Vladimir Voevodsky

ABSTRACT. Using the theory of framed correspondences developed by Voevodsky [22], we introduce and study framed motives of algebraic varieties. They are the major computational tool for constructing an explicit quasi-fibrant motivic replacement of the suspension \mathbb{P}^1 -spectrum of any smooth scheme $X \in Sm/k$. Moreover, it is shown that the bispectrum

 $(M_{fr}(X), M_{fr}(X)(1), M_{fr}(X)(2), \ldots),$

each term of which is a twisted framed motive of X, has motivic homotopy type of the suspension bispectrum of X. Furthermore, an explicit computation of infinite \mathbb{P}^1 -loop motivic spaces is given in terms of spaces with framed correspondences. We also introduce big framed motives of bispectra and show that they convert the classical Morel-Voevodsky motivic stable homotopy theory into an equivalent local theory of framed bispectra. As a topological application, it is proved that the framed motive $M_{fr}(pt)(pt)$ of the point $pt = \operatorname{Spec} k$ evaluated at pt is a quasifibrant model of the classical sphere spectrum whenever the base field k is algebraically closed of characteristic zero.

CONTENTS

1. Introduction	2
2. Voevodsky's framed correspondences	4
3. The Voevodsky Lemma	8
4. Motivic version of Segal's theorem	11
5. Framed motives	14
6. Additivity Theorem	17
7. Fibrant motivic spaces generated by S^1 -spectra	22
8. Comparing framed motives	24
9. Proof of Theorem 4.1	30
9.1. Proof of Theorem 4.1(2)	30
9.2. Proof of Theorem 4.1(1)	32
10. Computing infinite \mathbb{P}^1 -loop spaces	34
11. Further applications of framed motives	38
12. The big framed motive functor \mathscr{M}_{fr}^b	43
13. Framed \mathbb{P}^1 -spectra	47
14. Concluding remarks	51
References	51

Key words and phrases. Motivic homotopy theory, framed sheaves, triangulated categories.

²⁰¹⁰ Mathematics Subject Classification. 14F42, 14F05, 18G55, 55O10, 55P42.

The second author was supported by the Russian Science Foundation (grant no. 14-21-00035).

1. INTRODUCTION

In [22] Voevodsky develops the theory of (pre-)sheaves with framed correspondences. One of its aims was to suggest another framework for stable motivic homotopy theory more amenable to explicit calculations (see his Nordfjordeid Lectures [4, Remark 2.15] or his unpublished notes [22]). Recall that Voevodsky [22, Section 2] invented a category of framed correspondences $Fr_*(k)$ whose objects are those of Sm/k and morphisms sets $Fr_*(X,Y) = \bigsqcup_{n \ge 0} Fr_n(X,Y)$ are defined by means of certain geometric data (see Section 2 below). Following Voevodsky [22], we put for every $Y \in Sm/k$ (see Section 2 below):

 $Fr(X,Y) := \operatorname{colim}(Fr_0(X,Y) \xrightarrow{\sigma_Y} Fr_1(X,Y) \xrightarrow{\sigma_Y} \dots \xrightarrow{\sigma_Y} Fr_n(X,Y) \xrightarrow{\sigma_Y} \dots)$

and refer to it as the set of stable framed correspondences. Replacing Y by a simplicial object Y^{\bullet} in Sm/k, we get a simplicial set $Fr(X, Y^{\bullet})$. Finally, one can take the diagonal of the pointed bisimplicial set $Fr(\Delta^{\bullet} \times X, Y^{\bullet})$. Voevodsky conjectured that if the motivic space $Fr(\Delta^{\bullet} \times -, Y^{\bullet})$ is locally connected in the Nisnevich topology, then it is isomorphic in $H_{\mathbb{A}^1}(k)$ to the motivic space $\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(Y^{\bullet}_+)$. This shows that the theory of framed correspondences can give a machinery for computing/producing motivic infinite loop spaces.

In this paper we prove the conjecture in affirmative (see Theorem 10.5) providing that the base field is infinite perfect of characteristic different from 2. This paper shows that the theory of framed correspondences indeed gives a machinery for computing/producing motivic infinite loop spaces. To prove the conjecture, we introduce and study the theory of framed motives as well as the theory of big framed motives. The main aim of the machinery of framed motives is to find an explicit \mathbb{A}^1 -local replacement of the functor

$$\Omega^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{S^1} : H_{\mathbb{A}^1}(k) \to SH_{S^1}(k).$$

To this end, we firstly regard $SH_{S^1}(k)$ as *a full subcategory* of the ordinary stable homotopy category $SH_{S^1}^{nis}(k)$ consisting of \mathbb{A}^1 -local spectra. Then in Theorem 11.7(2) we construct an explicit functor $M_{fr}: H_{\mathbb{A}^1}(k) \to SH_{S^1}(k)$ together with a functor isomorphism

$$\alpha: \Omega^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{S^1} \to M_{fr}.$$

To formulate the main goal of the theory of big framed motives, consider a full subcategory $SH_{nis}^{fr}(k)$ of SH(k) consisting of framed bispectra E such that for any $i, j \ge 0$ the simplicial framed sheaf $E_{i,j}$ is \mathbb{A}^1 -local regarded as an ordinary motivic space and $\{E_{i,j}^f\}$ regarded as the ordinary bispectrum is stably motivically fibrant in the stable motivic model structure. Here "f" refers to a *fibrant* replacement in the local model structure on $sShv_{\bullet}(Sm/k)$.

The the main result of the theory of big framed motives are Theorems 12.4 and 12.5. Theorem 12.4 says that an explicitly constructed functor

$$\mathscr{M}_{fr}^{b}: SH(k) \to SH_{nis}^{Jr}(k)$$

converts classical Morel–Voevodsky's stable motivic homotopy theory SH(k) into an equivalent local homotopy theory of \mathbb{A}^1 -local framed bispectra from $SH_{nis}^{fr}(k)$, and thus producing a new approach to stable motivic homotopy theory. The main ingredient of this equivalent local homotopy theory is framed motivic spaces of the form $C_*Fr(-,Y)$ with $Y \in \Delta^{op}Fr_0(k)$ a simplicial scheme as well as their framed motives $M_{fr}(Y)$. Theorem 12.5 states that morphisms in SH(k)between two bispectra E and E' is the set $\pi_0(E^c, \mathcal{M}_{fr}^b(E')^f)$ of ordinary morphisms between bispectra E^c and $\mathcal{M}_{fr}^b(E')_f$ modulo the naive homotopy. Let us indicate some applications of the theory of framed motives and big framed motives. The fact that α is a functor isomorphism yields the following statement: if the base field *k* is perfect infinite with char $k \neq 2$, then for any $X \in Sm/k$ and any simplicial object Y^{\bullet} in Sm/k one has a canonical isomorphism

$$SH(k)(\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^{1}}X_{+},\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^{1}}Y^{\bullet}_{+}[n]) = SH^{\operatorname{nis}}_{S^{1}}(k)(\Sigma^{\infty}_{S^{1}}X_{+},M_{fr}(Y^{\bullet})[n]), \quad n \ge 0,$$
(1)

(see Theorem 11.5). In particular, the isomorphism α yields that for any simplicial object X^{\bullet} in Sm/k the projection $X^{\bullet} \times \mathbb{A}^1 \to X^{\bullet}$ induces an isomorphism $M_{fr}(X^{\bullet} \times \mathbb{A}^1) \cong M_{fr}(X^{\bullet})$. Furthermore, the functor M_{fr} converts any elementary distinguished Nisnevich square of *k*-smooth varieties to the ordinary Mayer–Vietoris exact triangle (see Theorem 8.12). Another important property of framed motives is as follows: given a morphism $\varphi : Y^{\bullet} \to Z^{\bullet}$ of simplicial objects in Sm/k such that the morphism $\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}(\varphi)$ is an isomorphism in SH(k), then the morphism $M_{fr}(\varphi)$ is a local equivalence.

By definition, the framed motive of a smooth scheme $X \in Sm/k$ over a field k is a motivic S^1 spectrum $M_{fr}(X)$ whose terms are certain explicit motivic spaces with framed correspondences (see Definition 5.2). We use framed motives to construct an explicit quasi-fibrant motivic replacement (i.e. an Ω -spectrum in positive degrees) of the suspension \mathbb{P}^1 -spectrum $\Sigma_{\mathbb{P}^1}^{\infty}X_+$ in Theorem 4.1 (here \mathbb{P}^1 is pointed at ∞). Another application is to show in Theorem 11.1 that an explicitly constructed bispectrum

$$M_{fr}^{\mathbb{G}}(X) = (M_{fr}(X), M_{fr}(X)(1), M_{fr}(X)(2), \ldots),$$

each term of which is a twisted framed motive of X, has motivic homotopy type of the suspension bispectrum $\Sigma_{\mathbb{G}}^{\infty} \Sigma_{S^1}^{\infty} X_+$ of X. Moreover, if we take the stable local fibrant replacement $M_{fr}(X)(n)_f$ of each twisted framed motive then the bispectrum

$$M_{fr}^{\mathbb{G}}(X)_f = (M_{fr}(X)_f, M_{fr}(X)(1)_f, M_{fr}(X)(2)_f, \dots)$$

is motivically fibrant by [1, Theorem A]. These definitions and results equally hold for simplicial objects in the category Sm/k. We should point out that for any simplicial object Y^{\bullet} in Sm/k there is a canonical morphism of bispectra

$$M_{fr}^{\mathbb{G}}(Y^{\bullet}) \to \mathscr{M}_{fr}^{b}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}Y_{+}^{\bullet})$$

which is an isomorphism in SH(k). Thus the composite morphism $\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}Y^{\bullet}_+ \to M^{\mathbb{G}}_{fr}(Y^{\bullet}) \to \mathcal{M}^b_{fr}(\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}Y^{\bullet}_+)$ is an isomorphism in SH(k). This yields equalities (see Theorem 12.5)

$$SH(k)(\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}X_+,\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}Y^{\bullet}_+)=\pi_0(\mathscr{M}^b_{fr}(\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}Y^{\bullet}_+)^f_{0,0}(X)).$$

Here "f" refers to a *fibrant* replacement in the local model structure on $sShv_{\bullet}(Sm/k)$.

Let us also give some applications of isomorphism (1). Since $M_{fr}(Y^{\bullet})$ is a sheaf of Segal S^1 -spectra, it follows that for any n < 0 the Nisnevich sheaf $\pi_{n,0}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}Y_+^{\bullet})$ vanishes. By varying Y^{\bullet} one gets a much stronger vanishing property. Namely, for any n < r one has $\pi_{n,r}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}Y_+^{\bullet}) = 0$. We can also compute $\pi_{n,r}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}Y_+^{\bullet})$ for n = r with $r \leq 0$ as

$$\pi^{\mathbb{A}^1}_{-n,-n}(\Sigma^\infty_{S^1}\Sigma^\infty_{\mathbb{G}}Y^\bullet_+)(K) = H_0(\mathbb{Z}F(\Delta^\bullet_K,Y^\bullet\times\mathbb{G}_m^{\wedge n})), \quad n \geqslant 0.$$

Here K/k is any field extension and $\mathbb{Z}F(\Delta_{K}^{\bullet}, Y^{\bullet} \times \mathbb{G}_{m}^{\wedge n}))$ is an explicit chain complex of free abelian groups. If $X = \operatorname{Spec}(k)$, char k = 0, then using Neshitov's computation $H_0(\mathbb{Z}F(\Delta_{K}^{\bullet}, X \times \mathbb{G}_{m}^{\wedge n})) = K_n^{MW}(K)$ [18], we recover the celebrated theorem of Morel [16] for Milnor–Witt *K*-theory for fields of characteristic zero.

We also give an explicit computation of the suspension functor (see Theorem 10.5)

$$\Sigma_{\mathbb{P}^1}^{\infty}: H_{\mathbb{A}^1}(k) \to SH(k).$$

It is isomorphic to an explicitly constructed functor

$$M_{\mathbb{P}^{\wedge 1}}: H_{\mathbb{A}^1}(k) \to SH(k)$$

that takes a motivic space to a spectrum consisting of spaces with framed correspondences. As an application, an explicit computation of the space $\Omega^{\infty}_{\mathbb{P}^{1}} \Sigma^{\infty}_{\mathbb{P}^{1}}(\mathscr{X})$ is given for any motivic space \mathscr{X} (see Theorem 10.5). Thus the machinery of framed motives leads to explicit computations of infinite \mathbb{P}^{1} -loop spaces.

As a topological application, using the machinery of framed motives together with a theorem of Levine [15], one shows in Theorem 11.9 that the framed motive of the point $M_{fr}(pt)(pt)$, $pt = \operatorname{Spec} k$, evaluated at pt is a quasi-fibrant model (i.e. an Ω -spectrum in positive degrees) of the classical sphere spectrum if the base field k is algebraically closed of characteristic zero. In particular, for any $n \ge 1$ the simplicial set $C_*Fr(pt, pt \otimes S^n)$ has the homotopy type of $\Omega_{Sl}^{\infty} \Sigma_{Sl}^{\infty} S^n$.

Finally, we show that every \mathbb{P}^1 -spectrum *E* is actually isomorphic in SH(k) to a framed spectrum in a canonical way, functorially in *E*. Furthermore, there is an equivalence of categories

$$SH(k) \xrightarrow{\cong} SH^{fr}(k),$$

where $SH^{fr}(k)$ is a full subcategory of framed \mathbb{P}^1 -spectra (see Theorem 13.4).

The main result of [7] says that for any \mathbb{A}^1 -invariant quasi-stable radditive framed presheaf of Abelian groups \mathscr{F} , the associated Nisnevich sheaf \mathscr{F}_{nis} is strictly \mathbb{A}^1 -invariant whenever the base field k is infinite perfect of characteristic different from 2. The assumption on the characteristic is needed to prove the "Surjective étale excision theorem" in [7]. It is for this reason that the main results for framed spaces, framed motives or framed spectra/bispectra are formulated and proven in this paper under the assumption that the base field k is infinite perfect of characteristic different from 2 (if chark = 2 we formulate and prove the same results for spaces, framed motives or framed spectra/bispectra "with 1/2-coefficients" only). However, as soon as the mentioned result of [7] about strict \mathbb{A}^1 -invariance of \mathscr{F}_{nis} is extended to (infinite) perfect fields of any characteristic, then the main results of this paper will automatically be extended to such fields as well.

The authors dedicate the paper to the memory of Vladimir Voevodsky who had kindly given his unpublished notes on framed correspondences [22] to us in 2010.

The authors are very grateful to Andrei Suslin for many helpful discussions. This paper was partly written during the visit of the authors in summer 2014 to the University of Duisburg–Essen (Marc Levine's Arbeitsgruppe). The authors also made major revisions of the original text during their visit to the Mittag-Leffler Institute in Stockholm in 2017. They would like to thank the University of Duisburg–Essen and the Mittag-Leffler Institute for the kind hospitality and support.

Throughout the paper we denote by Sm/k the category of smooth separated schemes of finite type over the base field k.

2. VOEVODSKY'S FRAMED CORRESPONDENCES

In this section we collect basic facts for framed correspondences and framed functors in the sense of Voevodsky [22]. We start with preparations.

Let *S* be a scheme and *Z* be a closed subscheme. Recall that an *étale neighborhood of Z in S* is a triple $(W', \pi' : W' \to S, s' : Z \to W')$ satisfying the conditions:

(i) π' is an étale morphism;

(ii) $\pi' \circ s'$ coincides with the inclusion $Z \hookrightarrow S$ (thus s' is a closed embedding);

(iii) $(\pi')^{-1}(Z) = s'(Z)$

A morphism between two étale neighborhoods $(W', \pi', s') \to (W'', \pi'', s'')$ of Z in S is a morphism $\rho: W' \to W''$ such that $\pi'' \circ \rho = \pi'$ and $\rho \circ s' = s''$. Note that such ρ is automatically étale by [10, VI.4.7].

Definition 2.1 (Voevodsky [22]). For *k*-smooth schemes *X*, *Y* and $n \ge 0$ an *explicit framed correspondence* Φ of level *n* consists of the following data:

- (1) a closed subset Z in \mathbb{A}^n_X which is finite over X;
- (2) an etale neighborhood $p: U \to \mathbb{A}^n_X$ of Z in \mathbb{A}^n_X ;
- (3) a collection of regular functions $\varphi = (\varphi_1, \dots, \varphi_n)$ on *U* such that $\bigcap_{i=1}^n {\varphi_i = 0} = Z$;
- (4) a morphism $g: U \to Y$.

The subset *Z* will be referred to as the *support* of the correspondence. We shall also write triples $\Phi = (U, \varphi, g)$ or quadruples $\Phi = (Z, U, \varphi, g)$ to denote explicit framed correspondences.

Two explicit framed correspondences Φ and Φ' of level *n* are said to be *equivalent* if they have the same support and there exists an open neighborhood *V* of *Z* in $U \times_{\mathbb{A}^n_X} U'$ such that on *V*, the morphism $g \circ pr$ agrees with $g' \circ pr'$ and $\varphi \circ pr$ agrees with $\varphi' \circ pr'$. A *framed correspondence* of level *n* is an equivalence class of explicit framed correspondences of level *n*.

Let $Fr_n(X,Y)$ denote the set of framed correspondences from X to Y. We consider it as a pointed set with the basepoint being the class 0_n of the explicit correspondence with $U = \emptyset$.

As an example, the sets $Fr_0(X,Y)$ coincide with the set of pointed morphisms $X_+ \to Y_+$. In particular, for a connected scheme *X* one has

$$Fr_0(X,Y) = \operatorname{Hom}_{Sm/k}(X,Y) \sqcup \{0_0\}.$$

If $f: X' \to X$ is a morphism of schemes and $\Phi = (U, \varphi, g)$ an explicit correspondence from X to Y then

$$f^*(\Phi) := (U' = U \times_X X', \varphi \circ pr, g \circ pr)$$

is an explicit correspondence from X' to Y.

Remark 2.2. Let $\Phi = (Z, \mathbb{A}^n_X \xleftarrow{p} U, \varphi : U \to \mathbb{A}^n_k, g : U \to Y) \in Fr_n(X, Y)$ be an *explicit framed correspondence of level n*. It can more precisely be written in the form

$$((\alpha_1, \alpha_2, \ldots, \alpha_n), f, Z, U, (\varphi_1, \varphi_2, \ldots, \varphi_n), g) \in Fr_n(X, Y)$$

where

 $\diamond Z \subset \mathbb{A}^n_X$ is a closed subset finite over X,

- \diamond an etale neighborhood (*α*₁, *α*₂,..., *α*_n), *f*) = *p* : *U* → $\mathbb{A}_k^n \times X$ of *Z*,
- ♦ a collection of regular functions $\varphi = (\varphi_1, ..., \varphi_n)$ on *U* such that $\bigcap_{i=1}^n {\varphi_i = 0} = Z$; ♦ a morphism $g: U \to Y$.

We shall usually drop $((\alpha_1, \alpha_2, \dots, \alpha_n), f)$ from notation and just write

 $(Z, U, (\varphi_1, \varphi_2, \dots, \varphi_n), g) = ((\alpha_1, \alpha_2, \dots, \alpha_n), f, Z, U, (\varphi_1, \varphi_2, \dots, \varphi_n), g).$

The following definition is to describe compositions of framed correspondences.

Definition 2.3. Let *X*, *Y* and *S* be *k*-smooth schemes and let

$$a = ((\alpha_1, \alpha_2, \dots, \alpha_n), f, Z, U, (\varphi_1, \varphi_2, \dots, \varphi_n), g)$$

be an explicit correspondence of level *n* from *X* to *Y* and let

$$b = ((\beta_1, \beta_2, \dots, \beta_m), f', Z', U', (\psi_1, \psi_2, \dots, \psi_m), g') \in Fr_m(Y, S)$$

be an explicit correspondence of level *m* from *Y* to *S*. We define their composition as an explicit correspondence of level n + m from *X* to *S* by

$$((\alpha_1, \alpha_2, \ldots, \alpha_n, \beta_1, \beta_2, \ldots, \beta_m), f, Z \times_Y Z', U \times_Y U', (\varphi_1, \varphi_2, \ldots, \varphi_n, \psi_1, \psi_2, \ldots, \psi_m), g').$$

Clearly, the composition of explicit correspondences respects the equivalence relation on them and defines associative maps

$$Fr_n(X,Y) \times Fr_m(Y,S) \rightarrow Fr_{n+m}(X,S).$$

Given $X, Y \in Sm/k$, denote by $Fr_+(X, Y)$ the set $\bigvee_n Fr_n(X, Y)$. The composition of framed correspondences defined above gives a category $Fr_+(k)$. Its objects are those of Sm/k and the morphisms are given by the sets $Fr_+(X, Y), X, Y \in Sm/k$. Since the naive morphisms of schemes can be identified with certain framed correspondences of level zero, we get a canonical functor

$$Sm/k \rightarrow Fr_+(k)$$

The category $Fr_+(k)$ has the zero object. It is the empty scheme. One can easily see that for a framed correspondence $\Phi: X \to Y$ and a morphism $f: X' \to X$, one has $f^*(\Phi) = \Phi \circ f$.

Definition 2.4. Let X, Y, S and T be smooth schemes. There is an *external product*

$$Fr_n(X,Y) \times Fr_m(S,T) \xrightarrow{-\boxtimes -} Fr_{n+m}(X \times S, Y \times T)$$

given by

$$((\alpha_1, \alpha_2, \dots, \alpha_n), f, Z, U, (\varphi_1, \varphi_2, \dots, \varphi_n), g) \boxtimes ((\beta_1, \beta_2, \dots, \beta_m), f', Z', U', (\psi_1, \psi_2, \dots, \psi_m), g') = ((\alpha_1, \alpha_2, \dots, \alpha_n, \beta_1, \beta_2, \dots, \beta_m), f \times f', Z \times Z', U \times U', (\varphi_1, \varphi_2, \dots, \varphi_n, \psi_1, \psi_2, \dots, \psi_m), g \times g').$$

For the constant morphism $c: \mathbb{A}^1 \to pt$, we set (following Voevodsky [22])

$$\Sigma = -\boxtimes (t, c, \{0\}, \mathbb{A}^1, t, c) \colon Fr_n(X, Y) \to Fr_{n+1}(X, Y)$$

and refer to it as the suspension.

Also, following Voevodsky [22], one puts

$$Fr(X,Y) = \operatorname{colim}(Fr_0(X,Y) \xrightarrow{\Sigma} Fr_1(X,Y) \xrightarrow{\Sigma} \dots \xrightarrow{\Sigma} Fr_n(X,Y) \xrightarrow{\Sigma} \dots)$$

and refer to it as the *set stable framed correspondences*. The above external product induces external products

$$\begin{aligned} & Fr_n(X,Y) \times Fr(S,T) \xrightarrow{-\boxtimes -} Fr(X \times S,Y \times T), \\ & Fr(X,Y) \times Fr_0(S,T) \xrightarrow{-\boxtimes -} Fr(X \times S,Y \times T). \end{aligned}$$

Definition 2.5. (I) Let *Y* be a *k*-smooth scheme and $S \subset Y$ be a closed subset and let $U \in Sm/k$. An *explicit framed correspondence of level* $m \ge 0$ *from U* to Y/(Y - S) consists of the tuples:

$$(Z, W, \varphi_1, \ldots, \varphi_m; g: W \to Y),$$

where Z is a closed subset of $U \times \mathbb{A}^m$, finite over U, W is an étale neighborhood of Z in $U \times \mathbb{A}^m$, $\varphi_1, \ldots, \varphi_m$ are regular functions on W, g is a regular map such that $Z = Z(\varphi_1, \ldots, \varphi_m) \cap g^{-1}(S)$.

The set Z is called the *support* of the explicit framed correspondence. We shall also write quadruples $\Phi = (Z, W, \varphi; g)$ to denote explicit framed correspondences.

(II) Two explicit framed correspondences $(Z, W, \varphi; g)$ and $(Z', W', \varphi'; g')$ of level *m* are said to be *equivalent* if Z = Z' and there exists an étale neighborhood W'' of Z in $W \times_{\mathbb{A}_U^m} W'$ such that $\varphi \circ pr$ agrees with $\varphi' \circ pr'$ and the morphism $g \circ pr$ agrees with $g' \circ pr'$ on W''.

(III) A framed correspondence of level m from U to Y/(Y-S) is the equivalence class of an explicit framed correspondence of level m from U to Y/(Y-S). We write $Fr_m(U,Y/(Y-S))$ to denote the set of framed correspondences of level m from U to Y/(Y-S). We regard it as a pointed set whose distinguished point is the class $0_{Y/(Y-S),m}$ of the explicit correspondence $(Z, W, \varphi; g)$ with $W = \emptyset$.

(IV) If S = Y then the pointed set $Fr_m(U, Y/(Y - S))$ coincides with the pointed set $Fr_m(U, Y)$ of framed correspondences of level *m* from *U* to *Y*.

Definition 2.6. A *framed presheaf* \mathscr{F} on Sm/k is a contravariant functor from $Fr_+(k)$ to the category of sets. A *framed functor* \mathscr{F} on Sm/k is a contravariant functor from $Fr_+(k)$ to the category of pointed sets such that $\mathscr{F}(\emptyset) = pt$ and $\mathscr{F}(X \sqcup Y) = \mathscr{F}(X) \times \mathscr{F}(Y)$.

A framed Nisnevich sheaf on Sm/k is a framed presheaf \mathscr{F} such that its restriction to Sm/k is a Nisnevich sheaf.

Note that the representable presheaves on $Fr_+(k)$ are not framed functors.

Construction 2.7. We set $Fr_+(-,Y/(Y-S)) := \bigvee_{m \ge 0} Fr_m(-,Y/(Y-S))$ and define the structure of a framed presheaf on it as follows. Let *X*, *Y* and *S* be *k*-smooth schemes and let

$$\Psi = (Z', \mathbb{A}^k \times V \xleftarrow{(a, k)} W', \psi_1, \psi_2, \dots, \psi_k; g: W' \to U) \in Fr_k(V, U)$$

be an explicit correspondence of level k from V to U. Suppose

$$\Phi = (Z, \mathbb{A}^m \times U \xleftarrow{(\beta,\pi)} W, \varphi_1, \varphi_2, \dots, \varphi_m; g' : W \to Y) \in Fr_m(U, Y/(Y-S))$$

is an explicit correspondence of level *m* from *U* to Y/(Y-S). We define $\Psi^*(\Phi)$ as an explicit correspondence of level k + m from *V* to Y/(Y-S) as

$$(Z \times_U Z', \mathbb{A}^{k+m} \times V \xleftarrow{(\alpha, p, \pi)} W' \times_U W, \psi_1, \psi_2, \dots, \psi_k, \varphi_1, \varphi_2, \dots, \varphi_m, g' \circ pr_W) \in Fr_{k+m}(V, Y/(Y-S)).$$

Clearly, the pullback operation $(\Psi, \Phi) \mapsto \Psi^*(\Phi)$ of explicit correspondences respects the equivalence relation on them. We get a pairing

$$Fr_k(V,U) \times Fr_m(U,Y/(Y-S)) \rightarrow Fr_{k+m}(V,Y/(Y-S))$$
 (2)

making $Fr_+(-,Y/(Y-S))$ a $Fr_+(k)$ -presheaf.

(... 0 -1)

Denote by

$$\sigma_{Y/(Y-S)}: Fr_m(U, Y/(Y-S)) \to Fr_{m+1}(U, Y/(Y-S))$$
(3)

a map, which takes $\Phi = (Z, W, \varphi; g)$ to $(Z \times \{0\}, W \times \mathbb{A}^1, \varphi \circ pr_W, pr_{\mathbb{A}^1}; g)$.

Following Voevodsky [22] we give the following

Definition 2.8. We shall refer to the set

$$Fr(U,Y/(Y-S)) :=$$

$$= \operatorname{colim}(Fr_0(U, Y/(Y-S)) \xrightarrow{\sigma_{Y/(Y-S)}} Fr_1(U, Y/(Y-S)) \xrightarrow{\sigma_{Y/(Y-S)}} Fr_2(U, Y/(Y-S)) \cdots)$$

as the set of stable framed correspondences from U to Y/(Y-S).

Remark 2.9. It is straightforward to check that for any framed correspondences $\Psi \in Fr_n(U', U)$ and $\Phi \in Fr_m(U, Y/(Y - S))$ one has $\sigma_{Y/(Y-S)}(\Psi^*(\Phi)) = \Psi^*(\sigma_{Y/(Y-S)}(\Phi))$. This shows that the assignment $U \mapsto Fr(U, Y/(Y - S))$ from Definition 2.8 is *a framed presheaf*.

For a scheme X we let Et/X denote the category of schemes separated and étale over X.

Theorem 2.10 (Voevodsky [22]). Let X be a k-smooth scheme. Then for any scheme Y the functor $U \mapsto Fr_n(U,Y)$ from Et/X to Sets_• is a sheaf in the etale topology.


The proof of the latter theorem given in [22] yields the following

Corollary 2.11. Given $Y \in Sm/k$ and any closed subset S in Y, the presheaf $Fr_n(-,Y/(Y-S))$ on Sm/k is a pointed Nisnevich sheaf. Also, the framed presheaf Fr(-,Y/(Y-S)) is a framed Nisnevich sheaf.

3. THE VOEVODSKY LEMMA

In this section we discuss the Voevodsky lemma computing framed correspondences in terms of morphisms of associated Nisnevich sheaves. It is crucial in our analysis. Corollary 3.3 and a sketch of its proof was communicated to us by A. Suslin. It very much helped the authors in understanding Voevodsky's notes [22].

Construction 3.1. Given an explicit framed correspondence $\alpha = (Z, W, g : W \to Y)$ from *X* to Y/(Y - S) of level zero, consider an elementary distinguished square of the form



where $\rho: W \to X$ is an étale neighborhood of *Z*. Let $q: Y \to F := Y/(Y - S)$ be the canonical morphism of Nisnevich sheaves. Take a morphism of sheaves $q \circ g: W \to F$ and a morphism of sheaves $c: X - Z \to F$ sending X - Z to the distinguished point of *F*. These two morphisms agree on W - Z. Thus there is a unique morphism of Nisnevich sheaves

$$s_{(Z,W,g)}: X \to F$$

such that $s_{(Z,W,g)} \circ in = c$ and $s_{(Z,W,g)} \circ \rho = q \circ g$. Clearly, the sheaf morphism $s_{(Z,W,g)}$ depends only on the equivalence class of (Z,W,g) in $Fr_0(U,Y/(Y-S))$. The assignment $(Z,W,g) \mapsto s_{(Z,W,g)}$ defines a map of pointed sets

$$a_{X,Y/(Y-S)}$$
: $Fr_0(X,Y/(Y-S)) \rightarrow Mor_{Shv}(X,Y/(Y-S))$

The map $a_{X,Y/(Y-S)}$ is natural in X with respect to morphisms of smooth varieties. Hence $a_{Y/(Y-S)}: Fr_0(-,Y/(Y-S)) \to \operatorname{Mor}_{Shv}(-,Y/(Y-S))$ is a morphism of presheaves on the category Sm/k. Using Corollary 2.11, the morphism

$$a_{Y/(Y-S)}: Fr_0(-,Y/(Y-S)) \rightarrow \operatorname{Mor}_{Shv}(-,Y/(Y-S))$$

is a morphism of Nisnevich sheaves on Sm/k.

Lemma 3.2 (Voevodsky's Lemma). Let Y be a k-smooth scheme and $S \subset Y$ be a closed subset. The morphism of pointed Nisnevich sheaves

$$a_{Y/(Y-S)}: Fr_0(-,Y/(Y-S)) \to \operatorname{Mor}_{Shv}(-,Y/(Y-S))$$
(4)

is an isomorphism.

Proof. Since $a_{Y/(Y-S)}$ is a morphism of Nisnevich sheaves, it suffices to check that for any essentially *k*-smooth local Henzelian *U* the map $a_{X,Y/(Y-S)}$ is a bijection. Let *U* be local essentially smooth Henselian with the closed point $u \in U$. Since *U* is local Henzelian the following holds: for any non-empty closed subset *Z* in *U* the henzelization U_Z^h of *U* at *Z* coincides with *U* itself. This shows that

$$Fr_0(U, Y/(Y-S)) - 0_0 = \{(Z, U, f : U \to Y) \mid Z = f^{-1}(S), Z \neq \emptyset\} = \{f : U \to Y \mid f(u) \in S\}.$$

Here 0_0 is the distinguished point in $Fr_0(U, Y/(Y - S))$. The map $a_{U,Y/(Y-S)}$ takes a triple (Z, U, f) to the morphism $q \circ f : U \to Y/(Y - S)$, where $q : Y \to Y/(Y - S)$ is the quotient map.

Let $Y_u(U) \subset Y(U)$ be the subset of *U*-points of *Y* consisting of $g \in Y(U)$ with $g(u) \in S$. Then the map $Y(U) = \operatorname{Mor}_{Shv}(U,Y) \to \operatorname{Mor}_{Shv_{\bullet}}(U,Y/(Y-S))$ taking $f \in Y(U)$ to $q \circ f$ identifies $Y_u(U)$ with the subset $\operatorname{Mor}_{Shv_{\bullet}}(U,Y/(Y-S)) - *$. In fact,

$$Mor_{Shv}(U, Y/(Y-S)) = (Y_u(U) \sqcup (Y-S)(U))/(Y-S)(U)) = Y_u(U) \sqcup *,$$
(5)

where * is a singleton. Hence the map

$$a_{U,Y/(Y-S)}|_{Fr_0(U,Y/(Y-S))-\{0_0\}}: Fr_0(U,Y/(Y-S))-\{0_0\} \to \operatorname{Mor}_{Shv_{\bullet}}(U,Y/(Y-S)) \setminus *$$

is a bijection. Thus the map $a_{U,Y/(Y-S)}$ is a bijection, too.

Corollary 3.3. Let Y be a k-smooth scheme and $S \subset Y$ be a closed subset. Let X be a k-smooth variety and $B \subset X$ its closed subset. Suppose $Fr_0(X/B,Y/(Y-S))$ is the subset of $Fr_0(X,Y/(Y-S))$ consisting of framed correspondences (Z,W,g) with $Z \cap B = \emptyset$. Then the map of pointed sets

$$a_{X/B,Y/(Y-S)}$$
: $Fr_0(X/B,Y/(Y-S)) \rightarrow \operatorname{Mor}_{Shv_{\bullet}}(X/B,Y/(Y-S))$

is a bijection.

Proof. Consider a commutative diagram

$$Fr_{0}(X/B,Y/(Y-S)) \xrightarrow{a_{X/B,Y/(Y-S)}} \operatorname{Mor}_{Shv_{\bullet}}(X/B,Y/(Y-S)) \xrightarrow{in} r^{*}$$

$$Fr_{0}(X,Y/(Y-S)) \xrightarrow{a_{X,Y/(Y-S)}} \operatorname{Mor}_{Shv}(X,Y/(Y-S))$$

where r^* is induced by the quotient map $r: X \to X/(X - S)$. Since r is an epimorphism the map r^* is injective. The map *in* is an inclusion by the definition of $Fr_0(X/B, Y/(Y - S))$. By Lemma 3.2 the map $a_{X,Y/(Y-S)}$ is bijective. Thus the map $a_{X/B,Y/(Y-S)}$ is injective. It remains to check its surjectivity.

Let $g: X \to Y/(Y - S)$ be a Nisnevich sheaf morphism. It is in $Mor_{Shv_{\bullet}}(X/B, Y/(Y - S))$ if and only if g(B) is the distinguished point * in Y/(Y - S).

Let $g: X \to Y/(Y - S)$ be a Nisnevich sheaf morphism such that g(B) = *. We claim that g is in the image of $a_{X/B,Y/(Y-S)}$. By Lemma 3.2 there is an explicit framed correspondence $(Z, W, \tilde{g}: W \to Y)$ from X to Y/(Y - S) such that $g = a_{X,Y/(Y-S)}((Z, W, \tilde{g}: W \to Y))$. The latter equality means that the morphism g is unique such that g(X - Z) = * and $q \circ \tilde{g} = g \circ \rho$. If g(B) = *, then $B \cap Z = \emptyset$. Indeed, if b is a closed point of the closed subset $B \cap Z$ then $* \neq \tilde{g}(b) \in S$. On the other hand, $\tilde{g}(b) = g(b) = *$. We see that $B \cap Z = \emptyset$, $(Z, W, \tilde{g}: W \to Y)$ is in $Fr_0(X/B, Y/(Y - S))$, and $g = a_{X/B,Y/(Y - S)}((Z, W, \tilde{g}: W \to Y))$ as claimed. \Box

Remark 3.4. Let n > 0 be an integer. Let $B_n \subset (\mathbb{P}^1)^n$ be a closed subset which is the union of all subsets of the form $\mathbb{P}^1 \times ... \times \{\infty\} \times ... \times \mathbb{P}^1$. Set $B_0 = \{\infty\}$. For any $X, Y \in Sm/k$ and any $n \ge 0$ the inclusion

$$Fr_n(X,Y) \subset Fr_0(X \times (\mathbb{P}^1)^n / X \times B_n, Y \times \mathbb{A}^n / Y \times (\mathbb{A}^n - \{0\}))$$

is an equality. It suffices to check that any element $(Z, W, f : W \to Y \times \mathbb{A}^n)$ from $Fr_0(X \times (\mathbb{P}^1)^n / X \times B_n, Y \times \mathbb{A}^n / Y \times (\mathbb{A}^n - \{0\}))$ is contained in $Fr_n(X, Y)$. Since $Z \cap (X \times B_n) = \emptyset$, it follows that $Z \subset X \times \mathbb{A}^n$. Since Z is closed in $X \times (\mathbb{P}^1)^n$, then Z is projective over X. Since Z is also affine over X, it is finite over X. Giving a morphism $f : W \to Y \times \mathbb{A}^n$ is the same as giving n functions $\varphi_1, ..., \varphi_n$ and a morphism $g : W \to Y$. The condition $Z = f^{-1}(Y \times \{0\})$ is equivalent to that of $Z = \{\varphi_1 = ... = \varphi_n = 0\}$. The desired equality is checked.

The preceding remark and Corollary 3.3 imply the following

Proposition 3.5 (Voevodsky). *For any* $X, Y \in Sm/k$ *and any* $n \ge 0$ *the map*

$$a_{n,X,Y} = a_{X \times (\mathbb{P}^1)^n / X \times B_n, Y \times \mathbb{A}^n / Y \times (\mathbb{A}^n - \{0\})} : Fr_n(X,Y) \to \\ \to \operatorname{Hom}_{Shv_{\bullet}^{\operatorname{nis}}(Sm/k)}(X_+ \wedge (\mathbb{P}^1, \infty)^{\wedge n}, Y_+ \wedge (\mathbb{A}^1 / (\mathbb{A}^1 - 0))^n) = \\ = \operatorname{Hom}_{Shv_{\bullet}^{\operatorname{nis}}(Sm/k)}(X_+ \wedge (\mathbb{P}^1, \infty)^{\wedge n}, Y_+ \wedge T^n).$$

is a bijection. In what follows we shall write $\mathbb{P}^{\wedge n}$ for $(\mathbb{P}^1, \infty)^{\wedge n}$ and $\operatorname{Hom}(X_+ \wedge \mathbb{P}^{\wedge n}, Y_+ \wedge T^n)$ instead of $\operatorname{Hom}_{Shv_{\bullet}^{\operatorname{vis}}(Sm/k)}(X_+ \wedge \mathbb{P}^{\wedge n}, Y_+ \wedge T^n)$. We shall also write $\mathscr{F}r_n(X, Y)$ to denote $\operatorname{Hom}(X_+ \wedge \mathbb{P}^{\wedge n}, Y_+ \wedge T^n)$.

Consider two categories $Fr_+(k)$ and $\mathscr{F}r_+(k)$, where the objects in both categories are those of Sm/k. The category $Fr_+(k)$ is defined in 2.3. The morphisms between X and Y in $\mathscr{F}r_+(k)$ are defined as $\bigvee_{n \ge 0} \mathscr{F}r_n(X,Y)$. The composition is defined as follows. Given two morphisms $\alpha : X_+ \wedge \mathbb{P}^{\wedge m} \to Y_+ \wedge T^m$ and $\beta : Y_+ \wedge \mathbb{P}^{\wedge n} \to S_+ \wedge T^n$, define a morphism $\beta \circ \alpha \in \mathscr{F}r_{m+n}(X,S)$ as the composite

$$X_+ \wedge \mathbb{P}^{\wedge m} \wedge \mathbb{P}^{\wedge n} \xrightarrow{\alpha \wedge \mathrm{id}} Y_+ \wedge T^m \wedge \mathbb{P}^{\wedge n} \cong T^m \wedge Y_+ \wedge \mathbb{P}^{\wedge n} \xrightarrow{\mathrm{id} \wedge \beta} T^m \wedge Y_+ \wedge T^n \cong Y_+ \wedge T^m \wedge T^n.$$

It is straightforward to check commutativity of the diagram

$$\begin{array}{c|c} Fr_m(X,Y) \times Fr_m(Y,S) & \stackrel{\circ}{\longrightarrow} Fr_{m+n}(X,S) \\ a \times a & & a \\ \mathscr{F}r_m(X,Y) \times \mathscr{F}r_n(Y,S) & \stackrel{\circ}{\longrightarrow} \mathscr{F}r_{m+n}(X,S) \end{array}$$

These observations imply the following

Corollary 3.6. *The functor*

:
$$Fr_+(k) \to \mathscr{F}r_+(k)$$

а

is an isomorphism of categories.

It is also worth to make the following

Remark 3.7. One has that

$$Fr_n(X,Y/(Y-S)) \subset Fr_0(X \times (\mathbb{P}^1)^n/X \times B_n, Y \times \mathbb{A}^n/(Y \times \mathbb{A}^n - S \times \{0\}))$$

is an equality. It suffices to check that any element $(Z, W, f : W \to Y \times \mathbb{A}^n)$ from $Fr_0(X \times (\mathbb{P}^1)^n / X \times B_n, Y \times \mathbb{A}^n / (Y \times \mathbb{A}^n - S \times \{0\}))$ is contained in $Fr_n(X, Y(Y - S))$. Since $Z \cap (X \times \mathbb{A}^n)$

 B_n) = \emptyset , then $Z \subset X \times \mathbb{A}^n$. Since *Z* is closed in $X \times (\mathbb{P}^1)^n$, then *Z* is projective over *X*. Since *Z* is affine over *X*, it is also finite over *X*. Giving a morphism $f: W \to Y \times \mathbb{A}^n = \mathbb{A}^n \times Y$ is the same as giving *n* functions $\varphi_1, ..., \varphi_n$ and a morphism $g: W \to Y$. The condition $Z = f^{-1}(S \times \{0\})$ is equivalent to that of $Z = \{\varphi_1 = ... = \varphi_n = 0\} \cap g^{-1}(S)$. The equality is checked.

The previous remark and Corollary 3.3 imply the following

Proposition 3.8 (Voevodsky). *For any* $X, Y \in Sm/k$ *and any* $n \ge 0$ *, the map*

$$a_{n,X,Y/(Y-S)}: Fr_n(X,Y/(Y-S)) \to \operatorname{Hom}_{Shv_{\bullet}}(X_+ \wedge \mathbb{P}^{\wedge n}, Y \times \mathbb{A}^n/(Y \times \mathbb{A}^n - S \times \{0\})) =$$

$$= \operatorname{Hom}_{Shv_{\bullet}}(X_{+} \wedge \mathbb{P}^{\wedge n}, Y/(Y-S) \wedge T^{n}).$$

is a bijection. For brevity, we shall write $\mathscr{F}r_n(X,Y/(Y-S))$ to denote $\operatorname{Hom}_{Shv_{\bullet}}(X_+ \wedge \mathbb{P}^{\wedge n},Y/(Y-S) \wedge T^n)$.

Remark 3.9. For any pointed Nisnevich sheaf \mathscr{F} , the presheaf $\mathscr{F}r_+(\mathscr{F}) := \bigvee_{n \ge 0} \operatorname{Hom}(\mathbb{P}^{\wedge n}, \mathscr{F} \wedge T^n)$ is a framed presheaf. Indeed, define for any $U, X \in Sm/k$ and any m, n a map of pointed sets

$$\operatorname{Hom}(U_+ \wedge \mathbb{P}^{\wedge m}, X_+ \wedge T^m) \times \operatorname{Hom}(X_+ \wedge \mathbb{P}^{\wedge n}, \mathscr{F} \wedge T^n) \to \operatorname{Hom}(U_+ \wedge \mathbb{P}^{\wedge (m+n)}, \mathscr{F} \wedge T^{m+n})$$

If $\alpha : U_+ \wedge \mathbb{P}^{\wedge m} \to X_+ \wedge T^m$ and $s : X_+ \wedge \mathbb{P}^{\wedge n} \to \mathscr{F} \wedge T^n$ are morphisms of pointed Nisnevich sheaves, then we define $\alpha^*(s)$ as the composite morphism

$$U_+ \wedge \mathbb{P}^{\wedge m} \wedge \mathbb{P}^{\wedge n} \xrightarrow{\alpha \wedge \mathrm{id}} X_+ \wedge T^m \wedge \mathbb{P}^{\wedge n} \cong T^m \wedge X_+ \wedge \mathbb{P}^{\wedge n} \xrightarrow{\mathrm{id} \wedge s} T^m \wedge \mathscr{F} \wedge T^n \cong \mathscr{F} \wedge T^m \wedge T^n.$$

By Corollary 3.6 the categories $Fr_+(k)$ and $\mathscr{F}r_+(k)$ are isomorphic. The category isomorphism makes $\mathscr{F}r_+(\mathscr{F})$ a framed presheaf. In the special case when $\mathscr{F} = Y/(Y-S)$ the bijections $a_{n,X,Y/(Y-S)}$ induce *isomorphisms of framed presheaves* $a_{Y/(Y-S)} : Fr_+(-,Y/(Y-S)) \to \mathscr{F}r_+(-,Y/(Y-S))$, where $Fr_+(-,Y/(Y-S)) := \bigvee_{n \ge 0} Fr_n(-,Y/(Y-S))$.

Definition 3.10. For a pointed Nisnevich sheaf \mathscr{F} set

$$\mathscr{F}r(-,\mathscr{F}) = \operatorname{colim}(\mathscr{F} \xrightarrow{\sigma} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge 1}, \mathscr{F} \wedge T) \xrightarrow{\sigma} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge 2}, \mathscr{F} \wedge T^2) \xrightarrow{\sigma} \cdots),$$

where $\sigma(\Phi: U_+ \wedge \mathbb{P}^{\wedge n} \to \mathscr{F} \wedge T^n) = (U_+ \wedge \mathbb{P}^{\wedge n+1} \xrightarrow{\Phi \wedge 1_{\mathbb{P}^{\wedge 1}}} \mathscr{F} \wedge T^n \wedge \mathbb{P}^{\wedge 1} \xrightarrow{1 \wedge \sigma} \mathscr{F} \wedge T^{n+1})$. Observe that $\mathscr{F}r(-,\mathscr{F})$ is a framed Nisnevich sheaf.

In the special case $\mathscr{F} = Y/(Y-S)$ one has that

$$a_{Y/(Y-S)} = \operatorname{colim}_n a_{n,Y/(Y-S)} : Fr(-,Y/(Y-S)) \to \mathscr{F}r(-,Y/(Y-S)).$$
(6)

is an isomorphism of framed sheaves.

In what follows we shall identify the isomorphic framed sheaves Fr(-,Y/(Y-S)) and $\mathscr{F}r(-,Y/(Y-S))$. If we write Fr(-,Y/(Y-S)) then we use the geometric description of the sheaf. In turn, the use of $\mathscr{F}r(-,Y/(Y-S))$ will mostly refer to the equivalent categorical description. The reader should always keep in mind the equivalent desriptions of both framed sheaves thanks to Voevodsky's Lemma.

4. MOTIVIC VERSION OF SEGAL'S THEOREM

After collecting necessary facts about framed correspondences in previous sections, we can formulate the main computational result of the paper. It is reminiscent of Segal's theorem computing the suspension spectrum $\sum_{S^1}^{\infty} X$ of a topological space X as the Segal spectrum of an associated Γ -space $B\Sigma_X$ (see [19, Section 3] for more details). The motivic counterpart of the

Segal theorem computes the suspension \mathbb{P}^1 -spectrum $\Sigma_{\mathbb{P}^1}^{\infty}X_+$ of a smooth algebraic variety X in terms of associated motivic spaces with framed correspondences. In a certain sense the theory of framed correspondences gives rise to an infinite \mathbb{P}^1 -loop space machine. In order to formulate the theorem, we need some preparations.

In Section 2 we introduced Nisnevich sheaves $Fr_s(-,Y/Y-S)$ and Fr(-,Y/Y-S). In the special case when $Y = X \times \mathbb{A}^n$ and $S = X \times 0$ we shall write $Fr_s(-,X \times T^n)$ and $Fr(-,X \times T^n)$ to denote the Nisnevich sheaves $Fr_s(-,X \times \mathbb{A}^n/(X \times \mathbb{A}^n - X \times 0))$ and $Fr(-,X \times \mathbb{A}^n/(X \times \mathbb{A}^n - X \times 0))$ respectively. Recall that an element of $Fr_s(-,X \times T^n)$ can be written as a tuple $(Z,W,(\varphi_1,...,\varphi_s;\psi_1,...,\psi_n):W \to \mathbb{A}^{s+n};g:W \to X)$ such that the support $Z = Z(\varphi_1,...,\varphi_s) \cap Z(\psi_1,...,\psi_n) = Z(\varphi_1,...,\varphi_s,\psi_1,...,\psi_n)$.

For $X \in Sm/k$ and integers $s, n \ge 0$ consider a morphism of pointed Nisnevich sheaves

$$\sigma_{s,n}: Fr_s(-, X \times T^n) \to \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, Fr_s(-, X \times T^{n+1})).$$
(7)

It takes $(Z, W, \varphi_1, ..., \varphi_s; \psi_1, ..., \psi_n; g)$ to $(Z \times 0, W \times \mathbb{A}^1, \varphi_1, ..., \varphi_s; \psi_1, ..., \psi_n, pr_{\mathbb{A}^1}; g)$. On the other hand, we have canonical maps (3)

$$\sigma_{s,X\times T^n}: Fr_s(-,X\times T^n) \to Fr_{s+1}(-,X\times T^n)$$
(8)

taking $(Z, W, \varphi_1, ..., \varphi_s; \psi_1, ..., \psi_n; g)$ to $(Z \times 0, W \times \mathbb{A}^1, \varphi_1, ..., \varphi_s, pr_{\mathbb{A}^1}; \psi_1, ..., \psi_n; g)$. We have a commutative diagram

$$\begin{array}{c} Fr_{s}(-,X\times T^{n}) \xrightarrow{\sigma_{s,n}} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1},Fr_{s}(-,X\times T^{n+1})) \\ \sigma_{s,X\times T^{n}} \downarrow & \downarrow^{(\sigma_{s,X\times T^{n+1}})_{*}} \\ Fr_{s+1}(-,X\times T^{n}) \xrightarrow{\sigma_{s+1,n}} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1},Fr_{s+1}(-,X\times T^{n+1})). \end{array}$$

Passing to colimits in the s-direction, we get a morphism of pointed sheaves

$$\sigma_n: Fr(-, X \times T^n) \to \underline{\mathrm{Hom}}(\mathbb{P}^{\wedge 1}, Fr(-, X \times T^{n+1})).$$

We can form a \mathbb{P}^1 -spectrum

$$Fr_{\mathbb{P}^{\Lambda_1},T}(X) = (Fr(-,X), Fr(-,X \times T), Fr(-,X \times T^2), ...)$$
(9)

with structure morphisms given by σ_n -s.

We can now take the Suslin complex of each motivic space of the spectrum $Fr_{\mathbb{P}^{\wedge 1},T}(X)$ to form a \mathbb{P}^1 -spectrum

$$M_{\mathbb{P}^{\wedge 1}}(X) = (C_*Fr(-,X), C_*Fr(-,X \times T), C_*Fr(-,X \times T^2), ...)$$

with structure maps defined by $C_*(\sigma_n)$ -s.

There is a canonical morphism of \mathbb{P}^1 -spectra

$$\varkappa: \Sigma^{\infty}_{\mathbb{P}^1} X_+ o M_{\mathbb{P}^{\wedge 1}}(X)$$

given by the section $id_X \in Fr_0(X, X)$ (recall that a morphism from the suspension spectrum of a variety $X \in Sm/k$ to any other spectrum is fully determined by a section of the zeroth motivic space of the spectrum at X).

By [13, 2.7] the category of simplicial Nisnevich sheaves on Sm/k has the injective local model structure with cofibrations monomorphisms and local weak equivalences. Take a fibrant replacement $C_*Fr(-, X \times T^n)_f$ of every motivic space $C_*Fr(-, X \times T^n)$ within the injective local model structure. We then arrive at a \mathbb{P}^1 -spectrum

$$M_{\mathbb{P}^{\wedge 1}}(X)_f = (C_*Fr(-,X)_f, C_*Fr(-,X imes T)_f, C_*Fr(-,X imes T^2)_f,...).$$

Notice that $M_{\mathbb{P}^{\wedge 1}}(X)_f$ is a fibrant replacement of the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(X)$ within the level injective local model structure of \mathbb{P}^1 -spectra.

If the characteristic of the base field is different from 2, then we shall prove (see Theorem 4.1) that $M_{\mathbb{P}^{\wedge 1}}(X)_f$ is a positively fibrant \mathbb{P}^1 -spectrum. In turn, if char k = 2 then we are able to prove a similar result only "after inverting 2". Precisely, there are explicitly constructed motivic spaces $C_*Fr(-, X \times T^n)\{1/2\}$, n > 0, such that the canonical morphism of spaces

$$\tau: C_*Fr(-, X \times T^n) \to C_*Fr(-, X \times T^n) \{1/2\}$$

induces an isomorphism of sheaves $\pi_*^{nis}(\tau) \otimes \mathbb{Z}[1/2]$ (see Proposition 6.7).

We can form a \mathbb{P}^1 -spectrum

$$M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\} := (C_*Fr(-,X), C_*Fr(-,X\times T)\{1/2\}, C_*Fr(-,X\times T^2)\{1/2\}, \dots)$$

with structure maps defined as for $M_{\mathbb{P}^{\wedge 1}}(X)$. The reader may have observed that we do not modify the zeroth space $C_*Fr(-,X)$. The reason is that we only deal with spaces in positive degrees in the proof of Theorem 4.1. Take a fibrant replacement $C_*Fr(-,X \times T^n)\{1/2\}_f$ of every motivic space $C_*Fr(-,X \times T^n)\{1/2\}$ within the injective local model structure. We then arrive at a \mathbb{P}^1 -spectrum

$$M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\}_f = (C_*Fr(-,X)_f, C_*Fr(-,X\times T)\{1/2\}_f, C_*Fr(-,X\times T^2)\{1/2\}_f, \ldots).$$

Notice that $M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\}_f$ is a fibrant replacement of the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\}$ within the level injective local model structure of \mathbb{P}^1 -spectra.

Let

$$\varkappa_f: \Sigma^{\infty}_{\mathbb{P}^1} X_+ \to M_{\mathbb{P}^{\wedge 1}}(X) \to M_{\mathbb{P}^{\wedge 1}}(X)_f$$

denote the composite morphism. We also have a morphism of spectra

$$\tau_f \circ \varkappa_f : \Sigma^{\infty}_{\mathbb{P}^1} X_+ \to M_{\mathbb{P}^{\wedge 1}}(X)_f \to M_{\mathbb{P}^{\wedge 1}}(X) \{1/2\}_f.$$

A morphism of \mathbb{P}^1 -spectra is called a 2^{-1} -stable motivic equivalence if it becomes an isomorphism in the category SH(k)[1/2].

A motivic counterpart of the Segal theorem says:

Theorem 4.1. Let k be an infinite perfect field. Then the following statements are true:

- (1) The morphism $\varkappa_f : \Sigma_{\mathbb{P}^1}^{\infty} X_+ \to M_{\mathbb{P}^{\wedge 1}}(X)_f$ is a stable motivic equivalence of \mathbb{P}^1 -spectra and the morphism $\tau_f \circ \varkappa_f : \Sigma_{\mathbb{P}^1}^{\infty} X_+ \to M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\}_f$ is a 2^{-1} -stable motivic equivalence of \mathbb{P}^1 -spectra.
- (2) If char $k \neq 2$ (respectively k is of any characteristic) then the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(X)_f$ (respectively $M_{\mathbb{P}^{\wedge 1}}(X)\{1/2\}_f$) is a motivically fibrant Ω -spectrum in posisitive degrees. This means that for every positive integer n > 0 each motivic space $C_*(Fr(-,X \times T^n))_f$ (respectively $C_*(Fr(-,X \times T^n))\{1/2\}_f$) is motivically fibrant in the Morel–Voevodsky [17] motivic model category of simplicial Nisnevich sheaves and the structure map

$$C_*(Fr(-,X\times T^n))_f\to \Omega_{\mathbb{P}^1}(C_*(Fr(-,X\times T^{n+1}))_f)$$

(respectively $C_*(Fr(-,X \times T^n))\{1/2\}_f \rightarrow \Omega_{\mathbb{P}^1}(C_*(Fr(-,X \times T^{n+1}))\{1/2\}_f))$ is a weak equivalence schemewise.

We shall also extend Theorem 4.1 to directed colimits of simplicial schemes (see Theorem 10.1). The next two corollaries are immediate consequences of the preceding theorem.

Corollary 4.2. Let k be an infinite perfect field of characteristic different from 2. Then for any positive integer m > 0 the natural morphism of \mathbb{P}^1 -spectra

$$\varkappa_f: \Sigma_{\mathbb{P}^1}^{\infty}(X_+ \wedge \mathbb{P}^{\wedge m}) \to M_{\mathbb{P}^{\wedge 1}}(X \times T^m)_f := (C_*Fr(-, X \times T^m)_f, C_*Fr(-, X \times T^{m+1})_f, C_*Fr(-, X \times T^{m+2})_f, \dots)$$

is a fibrant replacement of the suspension \mathbb{P}^1 -spectrum $\Sigma_{\mathbb{P}^1}^{\infty}(X_+ \wedge \mathbb{P}^{\wedge m})$ in the stable motivic model structure of \mathbb{P}^1 -spectra in the sense of Jardine [14].

Given a \mathbb{P}^1 -spectrum E, let \mathscr{E} be an Ω -spectrum stably equivalent to E. By $\Omega^{\infty}_{\mathbb{P}^1}(E)$ we mean the zeroth motivic space \mathscr{E}_0 . If $E = \Sigma^{\infty}_{\mathbb{P}^1} \mathscr{X}$ is the suspension \mathbb{P}^1 -spectrum of a pointed motivic space \mathscr{X} , we shall write $\Omega^{\infty}_{\mathbb{P}^1} \Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X})$ to denote $\Omega^{\infty}_{\mathbb{P}^1}(E)$.

Corollary 4.3. Let k be an infinite perfect field of characteristic different from 2. Then for any positive integer m > 0 the natural morphism of motivic spaces

$$C_*(Fr(X \times T^m)) \to \Omega^{\infty}_{\mathbb{P}^1} \Sigma^{\infty}_{\mathbb{P}^1}(X_+ \wedge \mathbb{P}^{\wedge m})$$

is a stalkwise weak equivalence for the Nisnevich topology. Particularly, for any field extension K/k the natural morphism of simplicial sets

$$Fr(\Delta_{K}^{ullet},X imes T^{m}) o \Omega_{\mathbb{P}^{1}}^{\infty}\Sigma_{\mathbb{P}^{1}}^{\infty}(X_{+}\wedge\mathbb{P}^{\wedge m})(K)$$

is a weak equivalence.

The proof of Theorem 4.1 is lengthy and is postponed. Although it states something for motivic spaces, the main strategy to prove it is to use the machinery of framed motives introduced and studied in this paper. By definition, the framed motive of a variety or a sheaf is a S^1 spectrum of simplicial Nisnevich sheaves, and hence may have nothing to do with Theorem 4.1 at the first glance. *But this is not the case! It is the theory of framed motives that allows to prove Theorem 4.1*.

The proof also depends on a theorem of [7] about homotopy invariant presheaves with framed correspondences and further two papers [1, 8], in which the Cancellation Theorem for framed motives of algebraic varieties is proved and framed motives of relative motivic spheres are computed.

5. FRAMED MOTIVES

As we have mentioned above, framed motives give the main technical tool to prove Theorem 4.1. Before introducing them, we fix the following useful

General Framework. Let (\mathcal{V}, \otimes) be a closed symmetric monoidal category and \mathscr{C} is a bicomplete category which is tensored and cotensored over \mathscr{V} . Then for every $V \in \mathscr{V}$ and $X \in \mathscr{C}$ there are defined objects $V \otimes X$, $X \otimes V$, $\underline{\text{Hom}}(V, X)$ of \mathscr{C} . They are all functorial in V and X. Moreover, for every morphism $u : V \to V'$ in \mathscr{V} the square

is commutative.

As an important example, \mathscr{V} is the category $(sShv_{\bullet}(Sm/k), \wedge)$ of Nisnevich sheaves of pointed simplicial sets and \mathscr{C} is either $sShv_{\bullet}(Sm/k)$ or the category of S^1 -spectra of simplicial Nisnevich sheaves.

Another example is the category $(Fr_0(k), \times, pt)$ and the category $Fr_+(k)$ The functor

$$Fr_+(k) \times Fr_0(k) \to Fr_+(k)$$

takes (X, Y) to $X \times Y$.

Let Γ^{op} be the category of finite pointed sets and pointed maps. Its skeleton has objects $n^+ = \{0, 1, ..., n\}$. We shall also regard each finite pointed set as a pointed smooth scheme. For example, we identify n^+ with the pointed scheme $(\sqcup_1^n \operatorname{Spec} k)_+$ with the distinguished point + corresponding to $0 \in n^+$. Note that $0^+ = \emptyset_+$. A Γ -space is a covariant functor from Γ^{op} to the category of simplicial sets taking 0^+ to a one point simplicial set. A morphism of Γ -spaces is a natural transformation of functors. A Γ -space X is called *special* if the map $X((k+l)^+)$: $X(k^+) \times X(l^+)$ induced by the projections from $(k+l)^+ \cong k^+ \vee l^+$ to k^+ and l^+ is a weak equivalence for all k and l. X is called *very special* if it is special and the monoid $\pi_0(X(1^+))$ is a group.

In what follows we shall regard Γ^{op} as a full subcategory in $sShv_{\bullet}(Sm/k)$ by means of the identification $K \in \Gamma^{\text{op}}$ with the pointed scheme $(\operatorname{Spec} k \sqcup \ldots \sqcup \operatorname{Spec} k)_+$, where the coproduct is indexed by the non-based elements in K.

By the General Framework above, for every $\mathscr{F}, \mathscr{G} \in sShv_{\bullet}(Sm/k)$ the association

$$K \in \Gamma^{\mathrm{op}} \mapsto \mathrm{Hom}_{sShv_{\bullet}(Sm/k)}(\mathscr{F}, \mathscr{G} \wedge K)$$

gives rise to a Γ -space, where the right hand side is regarded as a discrete simplicial set. In particular, if $\mathscr{F} = X_+ \wedge \mathbb{P}^{\wedge n}$ and $\mathscr{G} = \mathscr{H} \wedge T^n$ with $X \in Sm/k$, $\mathscr{H} \in sShv_{\bullet}(Sm/k)$, we have that the association

$$K \in \Gamma^{\mathrm{op}} \mapsto \mathscr{F}r_n(X, \mathscr{H} \wedge K) := \mathrm{Hom}_{sShv_{\bullet}(Sm/k)}(X_+ \wedge \mathbb{P}^{\wedge n}, \mathscr{H} \wedge T^n \wedge K)$$

is a Γ -space. Taking the colimit over *n*, we get that

$$K \in \Gamma^{\mathrm{op}} \mapsto \mathscr{F}r(X, \mathscr{H} \wedge K) = \operatorname{colim}_n(\mathscr{F}r_n(X, \mathscr{H} \wedge K))$$

is a Γ -space as well.

Using the geometric description of framed correspondences for $\mathscr{H} = Y_+$, $Y \in Sm/k$, the Γ -spaces $K \in \Gamma^{\text{op}} \mapsto \mathscr{F}r_n(X, \mathscr{H} \wedge K)$ and $K \in \Gamma^{\text{op}} \mapsto \mathscr{F}r(X, \mathscr{H} \wedge K)$ can equivalently be defined as

$$K \in \Gamma^{\mathrm{op}} \mapsto Fr_n(X, Y \otimes K)$$
 and $Fr(X, Y \otimes K)$

respectively. Here $Y \otimes K := Y \sqcup ... \sqcup Y$ with the coproduct indexed by the non-based elements in *K*. Observe that $\emptyset \otimes K = \emptyset$ and $X \otimes * = \emptyset$. These Γ -spaces are functorial in *X* and *Y* in framed correspondences of level zero. The second Γ -space is furthermore a framed functor in *X*.

Definition 5.1. We define a category $SmOp(Fr_0(k))$, which will be often used in our constructions. Its objects are pairs (X,U), where $X \in Sm/k$ and $U \subset X$ is its open subset. A morphism between (X,U) and (X',U') in $SmOp(Fr_0(k))$ is a morphism $f \in Fr_0(X,X')$ such that $f(U) \subset U'$. We shall also identify $X \in Sm/k$ with the pair $(X, \emptyset) \in SmOp(Fr_0(k))$.

The category $SmOp(Fr_0(k))$ is symmetric monoidal with the monoidal product \land given by

$$(X,U) \land (Y,V) := (X \times Y, X \times V \cup U \times Y).$$

The point *pt* is its monoidal unit. Also, by $(X,U) \sqcup (Y,V)$ we shall mean $(X \sqcup Y, U \sqcup V)$.

Let $\Delta^{op}SmOp(Fr_0(k))$ be the category of simplicial objects in $SmOp(Fr_0(k))$. There is an obvious functor $spc : SmOp(Fr_0(k)) \rightarrow Shv_{\bullet}(Sm/k)$ sending an object $(X,U) \in SmOp(Fr_0(k))$ to the Nisnevich sheaf X/U. Observe that this functor is a strict symmetric monoidal functor. It induces a functor

$$spc: \Delta^{op}SmOp(Fr_0(k)) \rightarrow sShv_{\bullet}(Sm/k),$$

taking an object $[n] \mapsto (Y_n, U_n)$ to the simplicial Nisnevich sheaf $[n] \mapsto (Y_n/U_n)$. Given $Y \in SmOp(Fr_0(k))$ there is a Γ -space

$$K \in \Gamma^{\mathrm{op}} \mapsto Fr(X, Y \otimes K) := \mathscr{F}r(X, spc(Y) \otimes K).$$

Notice that the right hand side has an explicit geometric description thanks to Voevodsky's lemma.

Definition 5.2. (1) The framed motive $\mathcal{M}_{fr}(\mathcal{G})$ of a pointed Nisnevich simplicial sheaf \mathcal{G} is the Segal S^1 -spectrum $(C_*\mathscr{F}r(-,\mathscr{G}), C_*\mathscr{F}r(-,\mathscr{G} \wedge S^1), C_*\mathscr{F}r(-,\mathscr{G} \wedge S^2), \ldots)$ associated with the Γ -space $K \in \Gamma^{\mathrm{op}} \mapsto C_*\mathscr{F}r(-,\mathscr{G} \wedge K) = \mathscr{F}r(\Delta^{\bullet}_+ \wedge -, \mathscr{G} \wedge K)$. More precisely, each structure map

$$C_*\mathscr{F}r(-,\mathscr{G}\wedge S^m)\wedge S^1\to C_*\mathscr{F}r(-,\mathscr{G}\wedge S^{m+1})$$

is given as follows. For any r and m, it coincides termwise with the natural morphisms

$$\bigvee \mathscr{F}r(\Delta^r_+ \wedge -, \mathscr{G} \wedge S^m) \to \mathscr{F}r(\Delta^r_+ \wedge -, \bigvee (\mathscr{G} \wedge S^m)),$$

where coproducts are indexed by non-basepoint elements of $S_n^1 = n^+$.

(2) The *framed motive* $M_{fr}(Y)$ of $Y \in \Delta^{\text{op}}SmOp(Fr_0(k))$ is the framed motive $\mathscr{M}_{fr}(spc(Y))$. It is the Segal S^1 -spectrum $(C_*Fr(-,Y), C_*Fr(-,Y \otimes S^1), C_*Fr(-,Y \otimes S^2), \ldots)$ associated with the Γ -space $K \in \Gamma^{\text{op}} \mapsto C_*Fr(-,Y \otimes K) = Fr(\Delta^{\bullet} \times -, Y \otimes K)$.

(3) In particular, by the *framed motive* $M_{fr}(Y)$ of a smooth algebraic variety $Y \in Sm/k$ we mean the framed motive of $(Y, \emptyset) \in SmOp(Fr_0(k))$.

Remark 5.3. (1) We should point out that it is framed motives $M_{fr}(Y^{\bullet})$ of simplicial *k*-varieties which are used to construct the functor $M_{fr}: H_{\mathbb{A}^1}(k) \to SH_{S^1}(k)$ in Section 11.

(2) Framed motives of pointed Nisnevich simplicial sheaves are not suitable for constructing such a functor. Particularly, we do not expect that the functor $\mathscr{G} \mapsto \mathscr{M}_{fr}(\mathscr{G})$ from Definition 5.2(1) preserves motivic equivalences. Therefore we do not expect that for a general motivic space \mathscr{G} the value of the functor M_{fr} on \mathscr{G} constructed in Section 11 has stable motivic homotopy type of $\mathscr{M}_{fr}(\mathscr{G})$ from Definition 5.2(1).

(3) It is for this reason that the main result of [8] stating that the natural motivic equivalence $X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n} \to X \times T^n$ of motivic spaces induces for any $n \ge 1$ a motivic equivalence (and even a level Nisnevich local weak equivalence) $M_{fr}(X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \simeq \mathscr{M}_{fr}(X_+ \wedge T^n)$ of S^1 -spectra is not obvious at all. In other words, the framed motive of the sheaf $X_+ \wedge T^n$ is computed as the framed motive of the associated simplicial scheme. This result is necessary to prove Theorem 4.1. It is also of independent interest.

(4) More generally, we can raise a problem asking for which motivic spaces \mathscr{G} the framed motive $\mathscr{M}_{fr}(\mathscr{G})$ from Definition 5.2(1) is isomorphic in $SH_{S^1}(k)$ to its image under the functor $M_{fr}: H_{\mathbb{A}^1}(k) \to SH_{S^1}(k)$ constructed in Section 11.

(5) However, the functor $M_{fr} : \Delta^{\text{op}}(Fr_0(k)) \to Sp_{S^1}(k)$ does preserve motivic equivalences (see Corollary 11.6).

(6) Framed motives of the form $M_{fr}(Y)$ with $Y \in \Delta^{op} SmOp(Fr_0(k))$ are of great utility to prove Theorem 4.1.

The framed motive $M_{fr}(Y)$ or $\mathcal{M}_{fr}(\mathcal{G})$ is a symmetric semistable S^1 -spectrum, because it is the value of the Γ -space $C_*Fr(-,Y)$ or $C_*\mathcal{F}r(-,\mathcal{G})$ at the sphere spectrum $\mathbb{S} = (S^0, S^1, S^2, ...)$. Since we also deal with framed spectra/spaces whose presheaves of homotopy groups are presheaves of $\mathbb{Z}[1/2]$ -modules, we have to introduce "framed motives with 1/2-coefficients". For this consider $2 \in \pi_1(S^1) = [S^1, Ex^{\infty}S^1] = \mathbb{Z}$ (here square brackets refer to the maps up to simplicial homotopy). There is the smallest integer k > 0 such that any representative of $2 \in \pi_1(S^1)$ factors as

$$S^1 \to \operatorname{sd}^k S^1 \to Ex^{\infty}S^1.$$

Here $\operatorname{sd}^k S^1$ stands for the *k*th subdivision of S^1 . If there is no likelihood of confusion, we denote the left arrow by $2: S^1 \to \operatorname{sd}^k S^1$ and fix a representative in its homotopy class from now on. We also denote the finite pointed simplicial set $\operatorname{sd}^k S^1$ by \mathscr{S}^1 . We also set $\mathscr{S}^n := \mathscr{S}^1 \wedge \stackrel{n}{\cdots} \wedge \mathscr{S}^1$.

Let $\mathbb{S}[1/2]$ be the spectrum $(S^0, \mathscr{S}^1, \mathscr{S}^2, \ldots)$. Its structure maps are defined as $\mathscr{S}^n \wedge S^1 \xrightarrow{\mathrm{id} \wedge 2} \mathscr{S}^n \wedge \mathscr{S}^1 = \mathscr{S}^{n+1}$.

Definition 5.4. (1) The framed motive with 1/2-coefficients $\mathcal{M}_{fr}(\mathcal{G})\{1/2\}$ of a pointed Nisnevich simplicial sheaf \mathcal{G} is the value of the Γ -space $C_* \mathscr{F}r(-,\mathcal{G})$ at the spectrum $\mathbb{S}[1/2]$. Explicitly, it is the symmetric S^1 -spectrum $(C_* \mathscr{F}r(-,\mathcal{G}), C_* \mathscr{F}r(-,\mathcal{G} \land \mathscr{S}^1), C_* \mathscr{F}r(-,\mathcal{G} \land \mathscr{S}^2), \ldots)$ with each structure map given by the composition

$$C_*\mathscr{F}r(-,\mathscr{G}\wedge\mathscr{S}^m)\wedge S^1\to C_*\mathscr{F}r(-,\mathscr{G}\wedge\mathscr{S}^m\wedge S^1)\xrightarrow{2_*}C_*\mathscr{F}r(-,\mathscr{G}\wedge\mathscr{S}^{m+1}).$$

(2) The framed motive with 1/2-coefficients $M_{fr}(Y)\{1/2\}$ of $Y \in \Delta^{op}SmOp(Fr_0(k))$ is the framed motive $\mathcal{M}_{fr}(spc(Y))\{1/2\}$. It is the symmetric S^1 -spectrum $(C_*Fr(-,Y), C_*Fr(-,Y \otimes \mathcal{S}^1), C_*Fr(-,Y \otimes \mathcal{S}^2), \ldots)$ with structure maps defined as above.

(3) In particular, by the *framed motive* $M_{fr}(Y)\{1/2\}$ of a smooth algebraic variety $Y \in Sm/k$ we mean the framed motive with 1/2-coefficients of $(Y, \emptyset) \in SmOp(Fr_0(k))$.

Our next goal is to show that the framed motive $M_{fr}(Y)$ of $Y \in \Delta^{op}SmOp(Fr_0(k))$ is a positively fibrant Ω -spectrum. To this end we need to prove the "Additivity Theorem".

6. Additivity Theorem

In this section we prove the Additivity Theorem. It is reminiscent of the Additivity Theorem in algebraic *K*-theory. We shall use it to produce special Γ -spaces in the sense of Segal [19] for associated motivic spaces with framed correspondences. In particular, Segal's machine then implies that the framed motive of a variety or, more generally, $Y \in \Delta^{\text{op}} SmOp(Fr_0(k))$ is a positively fibrant S^1 -spectrum.

Following [22] for any $X \in Sm/k$ denote by *m* the explicit correspondence from *X* to $X \sqcup X$ with $U = (\mathbb{A}^1 - \{0\} \sqcup \mathbb{A}^1 - \{1\})_X$, $\varphi = (t-1) \sqcup t$ where $t : \mathbb{A}^1_X \to X$ is the projection and $g : (\mathbb{A}^1 - \{0\} \sqcup \mathbb{A}^1 - \{1\})_X \to X \sqcup X$. For any framed functor \mathscr{F} it defines a map

$$\mathscr{F}(X) \times \mathscr{F}(X) = \mathscr{F}(X \sqcup X) \xrightarrow{m^*} \mathscr{F}(X).$$

Definition 6.1. Let \mathscr{F} and \mathscr{G} be two presheaves of sets on the category of *k*-smooth schemes and let $\varphi_0, \varphi_1 : \mathscr{F} \rightrightarrows \mathscr{G}$ be two morphisms. An \mathbb{A}^1 -homotopy between φ_0 and φ_1 is a morphism $H : \mathscr{F} \to \underline{\text{Hom}}(\mathbb{A}^1, \mathscr{G})$ such that $H_0 = \varphi_0$ and $H_1 = \varphi_1$. We write $\varphi_0 \sim \varphi_1$ if there is an \mathbb{A}^1 homotopy between φ_0 and φ_1 . We say that $\varphi_0, \varphi_n : \mathscr{F} \rightrightarrows \mathscr{G}$ are \mathbb{A}^1 -homotopic, if there is a chain of morphisms $\varphi_1, ..., \varphi_n$ such that $\varphi_i \sim \varphi_{i+1}$ for i = 0, 1, ..., n - 1. We now want to discuss matrices actions on framed correspondences and \mathbb{A}^1 -homotopies associated to them. Let *Y* be a *k*-smooth scheme and let $A \in GL_n(k)$ be a matrix. Then *A* defines an automorphism $\varphi_{A\perp Id_m} : Fr_{n+m}(-,Y) \to Fr_{n+m}(-,Y)$ of the presheaf $Fr_{n+m}(-,Y)$ in the following way. Given $W \in Sm/k$ and an *explicit framed correspondence of level n*

$$\Phi = (\mathbb{A}_X^{n+m} \stackrel{p}{\leftarrow} U, \varphi : U \to \mathbb{A}_k^{n+m}, g : U \to Y) \in Fr_{n+m}(X, Y),$$

set $\varphi_{A\perp Id_m}(\Phi) = ((A\perp Id_m) \circ p, (A\perp Id_m) \circ \varphi, g)$. Within the notation of Remark 2.2

$$\varphi_{A\perp Id_m}(Z,U,(\varphi_1,\varphi_2,\ldots,\varphi_{n+m}),g)) := ((A\perp Id_m)(Z),U,(A\perp Id_m)\circ(\varphi_1,\varphi_2,\ldots,\varphi_{n+m}),g),$$

where $A \perp Id_m$ is a linear automorphism of \mathbb{A}_k^{n+m} . In more details, if $(U, \mathbb{A}_X^{n+m} \xleftarrow{p} U, s : Z \to U)$ is an étale neighbourhood of Z in \mathbb{A}_X^{n+m} , then we take

$$(U, \mathbb{A}_X^{n+m} \xleftarrow{(A \perp Id_m) \circ p} U, s \circ ((A \perp Id_m)^{-1}|_{(A \perp Id_m)(Z)}) : (A \perp Id_m)(Z) \to U)$$

as an étale neighbourhood of $(A \perp Id_m)(Z)$ in \mathbb{A}_X^{n+m} . Clearly, $\sigma \circ \varphi_{A \perp Id_m} = \varphi_{A \perp Id_{m+1}} \circ \sigma$. Hence the maps $\varphi_{A \perp Id_m}$ give rise to a unique automorphism of *presheaves on* Sm/k

$$\varphi_A: Fr(-,Y) \to Fr(-,Y) \tag{11}$$

such that for any $m \ge 0$ one has $\varphi_A|_{Fr_{n+m}(-,Y)} = \varphi_{A \perp Id_m}$.

Definition 6.2. Let $A \in SL_n(k)$. Choose a matrix $A_s \in SL_n(k[s])$ such that $A_0 = id$ and $A_1 = A$. The matrices $A_s \perp Id_m \in SL_{n+m}(k[s])$, regarded as morphisms $\mathbb{A}^{n+m} \times \mathbb{A}^1 \to \mathbb{A}^{n+m}$, give rise to an \mathbb{A}^1 -homotopy *h* between the automorphisms *id* and φ_A of Fr(-,Y) as follows. Given $a = (Z, U, (\varphi_1, \varphi_2, ..., \varphi_{n+m}), g) \in Fr_n(-,Y)$, one sets

$$h(a) = (Z \times \mathbb{A}^1, U \times \mathbb{A}^1, (A_s \perp Id_m) \circ (\varphi \times id_{\mathbb{A}^1}), g \circ pr_U) \in Fr_n(W \times \mathbb{A}^1, Y).$$

In this way we get a morphism $h : Fr(-,Y) \to Fr(-\times \mathbb{A}^1,Y)$ such that $h_0 = id$ and $h_1 = \varphi_A$. We see that *h* is an \mathbb{A}^1 -homotopy between the identity and φ_A .

Definition 6.3. Let $\tau \in \Sigma_n$ be an even permutation regarded as a matrix in $SL_n(k)$. Let $A_s \in SL_n(k[s])$ be such that $A_0 = id$ and $A_1 = \tau$. Then the morphism *h* from Definition 6.2 defines an \mathbb{A}^1 -homotopy between the automorphisms φ_{id} and φ_{τ} of Fr(-,Y).

We are now in a position to prove the Additivity Theorem. The category $SmOp(Fr_0(k))$ was introduced in Definition 5.1. Given a simplicial object Y of $SmOp(Fr_0(k))$, by $C_*Fr(-,Y)$ we mean as usual the diagonal of the bisimplicial presheaf $(m,n) \mapsto Fr(\Delta^m \times -, Y_n)$.

Theorem 6.4 (Additivity). For any two simplicial objects Y_1, Y_2 in $SmOp(Fr_0(k))$, the natural map $\alpha : Fr(-, Y_1 \sqcup Y_2) \to Fr(-, Y_1) \times Fr(-, Y_2)$, given by the morphisms $id \sqcup \emptyset : Y_1 \sqcup Y_2 \to Y_1, \emptyset \sqcup id : Y_1 \sqcup Y_2 \to Y_2$, induces a map of simplicial framed presheaves

$$C_*(\alpha): C_*Fr(-,Y_1 \sqcup Y_2) \to C_*Fr(-,Y_1) \times C_*Fr(-,Y_2),$$

which is a schemewise weak equivalence.


Proof. Since the realization functor takes simplicial weak equivalences of simplicial sets to weak equivalences, it is enough to prove the theorem for any objects Y_1, Y_2 in $SmOp(Fr_0(k))$. Moreover, it is sufficient to prove that for any $X \in Sm/k$ and any finite unpointed simplicial set K, the map

$$C_*(\alpha)(X,K): [K,C_*Fr(X,Y_1\sqcup Y_2)] \to [K,C_*Fr(X,Y_1)\times C_*Fr(X,Y_2)]$$

between the Hom-sets in the homotopy category Ho(*sSets*) of unpointed simplicial sets is a bijection. Furthermore, for the simplicity of the exposition we shall assume that Y_1, Y_2 are just *k*-smooth varieties. Let $Y = Y_1 \sqcup Y_2$. Define a morphism

$$\beta$$
: $Fr(-,Y_1) \times Fr(-,Y_2) \rightarrow Fr(-,Y)$

of presheaves on Sm/k by the following commutative diagram:



Here $i_{\varepsilon}: Y_{\varepsilon} \to Y$, $\varepsilon = 1, 2$, is the corresponding embedding.

We claim that $(\beta \alpha)|_{Fr_{2n}(-,Y)}$ is \mathbb{A}^1 -homotopic to the inclusion $in_{2n} : Fr_{2n}(-,Y) \to Fr(-,Y)$ and $(\alpha \beta)|_{Fr_{2n}(-,Y_1) \times Fr_{2n}(-,Y_2)}$ is \mathbb{A}^1 -homotopic to the inclusion

$$inc_{2n}^1 \times inc_{2n}^2 : Fr_{2n}(-,Y_1) \times Fr_{2n}(-,Y_2) \to Fr(-,Y_1) \times Fr(-,Y_2).$$

The first of these \mathbb{A}^1 -homotopies will imply that $C_*(\beta) \circ C_*(\alpha)|_{C_*(Fr_{2n}(-,Y))}$ is simplicially homotopic to the inclusion $C_*(in_{2n})$, because $C_*(-)$ converts \mathbb{A}^1 -homotopies into simplicial ones. The second of these \mathbb{A}^1 -homotopies will imply that $(C_*(\alpha) \circ C_*(\beta))|_{C_*(Fr_{2n}(-,Y_1) \times C_*(Fr_{2n}(-,Y_1)))}$ is simplicially homotopic to the inclusion $C_*(inc_{2n}^1 \times inc_{2n}^2)$. It will follow that for any $X \in Sm/k$ and any finite simplicial set K the map $C_*(\alpha)(X, K)$ is bijective. Indeed, one should use the fact that the functor [K, -]: Ho(*sSets*) \rightarrow Ho(*sSets*) commutes with sequential colimits whenever K is finite. It therefore remains to prove the claim.

Firstly, let us focus on the morphism $\alpha\beta$. The map $\alpha\beta$ is of the form

$$\rho_1 \times \rho_2 : Fr(X, Y_1) \times Fr(X, Y_2) \to Fr(X, Y_1) \times Fr(X, Y_2)$$

Here ρ_1 takes a framed correspondence $(Z_1, W_1, \underline{\phi}^{(1)}; g_1)$ of level *n* to the framed correspondence $(0 \times Z_1, \mathbb{A}^1 \times W_1, t_1, \underline{\phi}^{(1)}; g_1)$ of level n+1, ρ_2 takes a framed correspondence $(Z_2, W_2, \underline{\phi}^{(2)}; g_2)$ of level *n* to the framed correspondence $(1 \times Z_2, \mathbb{A}^1 \times W_2, t_0 - 1, \underline{\phi}^{(2)}; g_2)$ of level n+1. We first observe that the morphism ρ_2 is \mathbb{A}^1 -homotopic to the morphism ρ_2^0 : $Fr(-,Y_2) \to Fr(-,Y_2)$ taking a framed correspondence $(Z_2, W_2, \underline{\phi}^{(2)}; g_2)$ of level n to the framed correspondence $(0 \times Z_2, \mathbb{A}^1 \times W_2, t_0, \underline{\phi}^{(2)}; g_2)$ of level n+1. To see this, send a framed correspondence $(Z_2, W_2, \underline{\phi}^{(2)}; g_2)$ of level n to the framed correspondence

$$(\Delta \times Z_2, \mathbb{A}^1 \times \mathbb{A}^1 \times W_2, t_0 - \lambda, \underline{\varphi}^{(2)}; g_2)$$

of level n + 1. Evaluating the latter framed correspondence at $\lambda = 1$, we get $\rho_2(Z_2, W_2, \underline{\phi}^{(2)}; g_2)$. Evaluating the same framed correspondence at $\lambda = 0$, we get $(0 \times Z_2, \mathbb{A}^1 \times W_2, t_0, \underline{\phi}^{(2)}; g_2)$.

Let n > 0 be an *even* integer and let $\tau \in \Sigma_{n+1}$ be the even permutation $(n+1, 1, \overline{2}, ..., n)$. Let h_1 denote the associated \mathbb{A}^1 -homotopy from Definition 6.3 between the automorphisms φ_{id} and φ_{τ} of $Fr(-, Y_1)$. Then $h_1 \circ inc_n^1$ is an \mathbb{A}^1 -homotopy between inc_n^1 and $\varphi_{\tau} \circ inc_n^1 = \rho_1|_{Fr_n(-,Y_1)}$: $Fr_n(-,Y_1) \to Fr(-,Y_1)$. Let h_2 be the associated \mathbb{A}^1 -homotopy from Definition 6.3 between the automorphisms φ_{id} and $\varphi_{\tau} \circ inc_n^2$ is an \mathbb{A}^1 -homotopy between inc_n^2 and $\varphi_{\tau} \circ inc_n^2$ and $\varphi_{\tau} \circ inc_n^2$ is an \mathbb{A}^1 -homotopy between inc_n^2 and $\varphi_{\tau} \circ inc_n^2$ of $Fr(-,Y_2)$.

 $inc_n^2 = \rho_2^0|_{Fr_n(-,Y_2)} : Fr_n(-,Y_2) \to Fr(-,Y_2)$. Thus $(\alpha\beta)|_{Fr_{2n}(-,Y_1) \times Fr_{2n}(-,Y_2)}$ is \mathbb{A}^1 -homotopic to the inclusion $inc_{2n}^1 \times inc_{2n}^2$.

Next, let us focus on the morphism $\beta \alpha$. Since $Y = Y_1 \sqcup Y_2$ then every framed correspondence of level *n* from *X* to *Y* is of the form $a = (Z_1 \sqcup Z_2, W_1 \sqcup W_2, \varphi^{(1)} \sqcup \varphi^{(2)}; g_1 \sqcup g_2)$. One has,

$$\beta \alpha(a) = (0 \times Z_1 \sqcup 1 \times Z_2, \mathbb{A}^1 \times W_1 \sqcup \mathbb{A}^1 \times W_2), (t_0, \underline{\varphi}^{(1)}) \sqcup (t_0 - 1, \underline{\varphi}^{(2)}); g_1 \sqcup g_2).$$

Firstly, the morphism $\beta \alpha$ is \mathbb{A}^1 -homotopic to the morphism $\rho^0 : Fr(-,Y) \to Fr(-,Y)$ taking a framed correspondence $(Z, W, \underline{\varphi}; g)$ to the framed correspondence $(0 \times Z, \mathbb{A}^1 \times W, t_0, \underline{\varphi}; g_2)$ of level n+1. To see this, send a framed correspondence $a = (Z_1 \sqcup Z_2, W_1 \sqcup W_2, \underline{\varphi}^{(1)} \sqcup \underline{\varphi}^{(2)}; g_1 \sqcup g_2)$ of level n to the framed correspondence of level n+1

$$(\mathbb{A}^1 \times Z_1 \sqcup \Delta \times Z_2, \mathbb{A}^1 \times \mathbb{A}^1 \times (W_1 \sqcup W_2), (t_0, \underline{\varphi}^{(1)}) \sqcup ((t_0 - \lambda), \underline{\varphi}^{(2)}); g_1 \sqcup g_2).$$

Evaluating the latter framed correspondence of level n + 1 at $\lambda = 1$, we get $\beta \alpha(a)$. Evaluating the same framed correspondence $\lambda = 0$, we get $\rho^0(a)$. Furthermore, using the associated homotopy from Definition 6.3, we see that $\rho^0|_{Fr_{2n}(-,Y)}$ is \mathbb{A}^1 -homotopic to the inclusion in_{2n} . Hence $\beta \alpha|_{Fr_{2n}(-,Y)}$ is \mathbb{A}^1 -homotopic to the inclusion in_{2n} . \square

Now Additivity Theorem 6.4 together with the Segal machine [19] imply the following

Theorem 6.5. Let $Y \in \Delta^{\operatorname{op}} SmOp(Fr_0(k))$. Then the Γ -space $K \in \Gamma^{\operatorname{op}} \mapsto C_*Fr(-, Y \otimes K)$ is sectionwise special. As a consequence, the framed motive $M_{fr}(Y)$ of Y is sectionwise a positively fibrant Ω -spectrum, which is sectionwise (respectively locally in the Nisnevich topology) an Ω -spectrum whenever the motivic space $C_*Fr(-,Y)$ is sectionwise (respectively locally in the Nisnevich topology) connected.

Remark 6.6. Whenever we say that $M_{fr}(Y)$ is a (positively) fibrant Ω -spectrum we tacitly assume that each of its spaces $C_*Fr(-,Y \otimes S^n)$ is replaced with $Ex^{\infty}(C_*Fr(-,Y \otimes S^n))$, where Ex^{∞} refers to Kan's complex. The spaces $Ex^{\infty}(C_*Fr(-,Y \otimes S^n))$ are then spaces with framed correspondences and sectionwise fibrant simplicial sets. A detailed description of the spaces will be given in Section 12. We can equally take any naive sectionwise fibrant resolution functor in the category of spaces with framed correspondences (which exist by using standard arguments) in place of Ex^{∞} .

It is worth to mention that the latter theorem is a kind of the "Cancellation Theorem for framed motives in the S^1 -direction" (whose meaning will become clear in the proof of Theorem 4.1(2)). One should also stress the motivic spaces $C_*Fr(-,X \times T^n)_f$ from Theorem 4.1 are zero spaces of sheaves of S^1 -spectra $M_{fr}(X \times T^n)_f$ (these are level local Nisnevich replacements of the framed motives $M_{fr}(X \times T^n)$). So each space $C_*Fr(-,X \times T^n)_f$ is part of the S^1 -spectrum $M_{fr}(X \times T^n)_f$. If we can prove that each S^1 -spectrum $M_{fr}(X \times T^n)_f$, n > 0, is motivically fibrant over a field of characteristic different from 2, then each space $C_*Fr(-,X \times T^n)_f$ becomes motivically fibrant (what is claimed in Theorem 4.1). In turn, with a little extra work the same is true for spaces $C_*Fr(-,X \times T^n)\{1/2\}_f$ we are going to define now.

Suppose Y is a simplicial object in $SmOp(Fr_0(k))$, then $M_{fr}(Y)\{1/2\}$ is not a positive Ω spectrum in contrast to $M_{fr}(Y)$. The problem is that each structure map $C_*Fr(Y \otimes \mathscr{S}^n) \to \Omega C_*Fr(Y \otimes \mathscr{S}^{n+1})$ of the spectrum $M_{fr}(Y)\{1/2\}$ factors as

$$C_*Fr(Y\otimes\mathscr{S}^n)\to \Omega C_*Fr(Y\otimes\mathscr{S}^n\otimes S^1)\xrightarrow{2_*}\Omega C_*Fr(Y\otimes\mathscr{S}^{n+1}).$$

The left map is a weak equivalence of spaces for all n > 0, but the right arrow is such only "after inverting 2" explained below.

It is important to observe that the composite map

$$C_*Fr(Y\otimes\mathscr{S}^n\otimes S^1)\xrightarrow{2_*}C_*Fr(Y\otimes\mathscr{S}^{n+1})\xrightarrow{\iota^*}C_*Fr(Y\otimes\mathscr{S}^n\otimes S^1)$$

where t^* is a weak equivalence induced by the "last vertex weak equivalence" $t: \mathscr{S}^1 = \operatorname{sd}^k S^1 \to S^1$, is such that it is equal to the homomorphism $2 \times \operatorname{id}_{\pi_*(C_*Fr(Y \otimes \mathscr{S}^n \otimes S^1))}$ on presheaves of homotopy groups. This easily follows from the fact that these presheaves of homotopy groups are the same with the stable homotopy groups of their Segal S^1 -spectra and basic facts of Γ -spaces.

In particular, there is a canonical isomorphism of presheaves of stable homotopy groups

$$\pi_*(M_{fr}(Y)\{1/2\}) \cong \pi_*(M_{fr}(Y)) \otimes \mathbb{Z}[1/2]$$

and hence the canonical map of spectra $\mu: M_{fr}(Y) \to M_{fr}(Y)\{1/2\}$ is a "2⁻¹-stable weak equivalence" in the sense that $\pi_*(\mu) \otimes \mathbb{Z}[1/2]$ is an isomorphism.

The inversion of the map 2_* above is achieved if we take the space

$$C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\} := \operatorname{colim}_k \Omega^k C_*Fr(Y \otimes \mathscr{S}^{n+k}).$$

It follows from above that each map of motivic spaces

$$\tau: C_*Fr(Y \otimes \mathscr{S}^n) \to C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\}, \quad n > 0,$$

induces an isomorphism on homotopy presheaves after tensoring with $\mathbb{Z}[1/2]$ and the homotopy presheaves of $C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\}$ are already presheaves of $\mathbb{Z}[1/2]$ -modules. In turn, if $C_*Fr(Y)$ is locally connected, say if $Y = X \times T^n$ with $X \in Sm/k$ and n > 0 (see [8, A.1]), then $\tau : C_*Fr(Y) \to C_*Fr(Y)\{1/2\}$ induces isomorphisms on sheaves of homotopy groups with $\mathbb{Z}[1/2]$ -coefficients $\pi^{nis}_*(\tau) \otimes \mathbb{Z}[1/2]$ (for this we use Theorem 6.5). Moreover,

$$\pi_*(C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\}) = \pi_{*-n}(M_{fr}(Y)\{1/2\}) \cong \pi_{*-n}(M_{fr}(Y)) \otimes \mathbb{Z}[1/2].$$

We shall also consider a spectrum

$$\widetilde{M}_{fr}(Y)\{1/2\} := \Omega^{\infty} M_{fr}(Y)\{1/2\} = (C_*Fr(Y)\{1/2\}, C_*Fr(Y \otimes \mathscr{S}^1)\{1/2\}, \ldots).$$

Then the natural morphism of spectra $\ell: M_{fr}(Y)\{1/2\} \to \widetilde{M}_{fr}(Y)\{1/2\}$ is a stable equivalence by construction and $\widetilde{M}_{fr}(Y)\{1/2\}$ becomes an Ω -spectrum. By $\widetilde{M}_{fr}(Y)\{1/2\}_f$ we shall mean a fibrant replacement of the spectrum $\widetilde{M}_{fr}(Y)\{1/2\}$ in the level injective Nisnevich local model structure of spectra. Its spaces are of the form $CFr(Y \otimes \mathscr{S}^n)\{1/2\}_f$, $n \ge 0$. More generally, for every simplicial sheaf \mathscr{G} we shall write $\widetilde{M}_{fr}(\mathscr{G})\{1/2\}$ to denote the spectrum $\Omega^{\infty}\widetilde{M}_{fr}(\mathscr{G})\{1/2\}$. Its spaces will be denoted by $C_*\mathscr{F}r(\mathscr{G} \otimes \mathscr{S}^n)\{1/2\}$, $n \ge 0$.

We document the above arguments as follows.

Proposition 6.7. Suppose Y is a simplicial object in $SmOp(Fr_0(k))$, then the presheaves of homotopy groups of $\widetilde{M}_{fr}(Y)\{1/2\}$ and $C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\}$, $n \ge 0$, are presheaves of $\mathbb{Z}[1/2]$ -modules and the canonical morphisms

$$\ell \circ \mu: M_{fr}(Y) \to M_{fr}(Y)\{1/2\}, \quad \tau: C_*Fr(Y \otimes \mathscr{S}^n) \to C_*Fr(Y \otimes \mathscr{S}^n)\{1/2\}, \quad n > 0,$$

induce isomorphisms on presheaves of homotopy groups with $\mathbb{Z}[1/2]$ -coefficients $\pi_*(\ell \circ \mu) \otimes \mathbb{Z}[1/2]$ and $\pi_*(\tau) \otimes \mathbb{Z}[1/2]$ respectively. Furthermore, if $C_*Fr(Y)$ is locally connected (e.g. if $Y = X \times T^n$ with $X \in Sm/k$ and n > 0), then $\tau : C_*Fr(Y) \to C_*Fr(Y)\{1/2\}$ induces isomorphisms on sheaves of homotopy groups with $\mathbb{Z}[1/2]$ -coefficients $\pi_*^{nis}(\tau) \otimes \mathbb{Z}[1/2]$.

If we can prove that each S^1 -spectrum $M_{fr}(X \times T^n)$ or $\widetilde{M}_{fr}(X \times T^n)\{1/2\}_f$, n > 0, is motivically fibrant, then each space $C_*Fr(-,X \times T^n)_f$ or $C_*Fr(-,X \times T^n)\{1/2\}_f$ becomes motivically fibrant (what is claimed in Theorem 4.1). Therefore our next goal is to investigate this kind of fibrant motivic spaces coming from relevant S^1 -spectra in more detail.

7. FIBRANT MOTIVIC SPACES GENERATED BY S^1 -SPECTRA

Let $sShv_{\bullet}(Sm/k)$ denote the category of Nisnevich sheaves of pointed simplicial sets. It has the injective model structure [13] in which cofibrations are the monomorphisms and weak equivalences are stalkwise weak equivalences of simplicial sets. The category of S^1 -spectra $Sp_{S^1}(sShv_{\bullet}(Sm/k))$ associated with $sShv_{\bullet}(Sm/k)$ will also be called the *category of ordinary* S^1 -spectra of simplicial Nisnevich sheaves. It has level and stable model structures (the standard references here are [11, 14]). In this section we describe a class of motivic spaces coming from ordinary S^1 -spectra of simplicial Nisnevich sheaves, which are fibrant in the motivic model category $sShv_{\bullet}(Sm/k)_{mot}$ of Morel–Voevodsky [17]. This class occurs in our analysis. Recall that $sShv_{\bullet}(Sm/k)_{mot}$ is obtained from $sShv_{\bullet}(Sm/k)$ by Bousfield localization with respect to the projections $p: X \times \mathbb{A}^1 \to X, X \in Sm/k$. As above, the level/stable model category of S^1 -spectra associated with $sShv_{\bullet}(Sm/k)_{mot}$ will also be called the *level/stable injective model category of* S^1 -spectra.

Proposition 7.1. Let E be an S^1 -spectrum in the category of simplicial Nisnevich sheaves such that each space $E_n \in sShv_{\bullet}(Sm/k)$ of the spectrum is fibrant in $sShv_{\bullet}(Sm/k)$. Suppose E is sectionwise an Ω -spectrum in the category of ordinary S^1 -spectra of pointed simplicial sets. Suppose E is locally (-1)-connected in the Nisnevich topology. Finally suppose that for any integer n, the Nisnevich sheaf $\pi_n^{nis}(E)$ is strictly homotopy invariant. Then the following statements are true:

- (1) every motivic space E_n of E is motivically fibrant;
- (2) *E* is fibrant in the stable injective motivic model structure of S^1 -spectra.

Proof. (1). Since *E* is sectionwise an Ω -spectrum, then every E_n , $n \ge 0$, is sectionwise fibrant. Therefore it suffices to prove that E_n is \mathbb{A}^1 -local. So we have to check that for any smooth variety *X* the projection $p: X \times \mathbb{A}^1 \to X$ induces a weak equivalence of simplicial sets $p^*: E_n(X) \to E_n(X \times \mathbb{A}^1)$. Since *E* is sectionwise an Ω -spectrum it suffices to check that the pull-back map $p^*: E(X) \to E(X \times \mathbb{A}^1)$ is a stable equivalence of ordinary S^1 -spectra. So it is sufficient to verify that for any integer *r* the map $p^*: \pi_r(E(X)) \to \pi_r(E(X \times \mathbb{A}^1))$ is an isomorphism. Consider two convergent spectral sequences

$$H^p_{Nis}(X, \pi^{\operatorname{nis}}_q(E)) \Rightarrow \pi_{q-p}(E(X)) \quad \text{and} \quad H^p_{Nis}(X \times \mathbb{A}^1, \pi^{\operatorname{nis}}_q(E)) \Rightarrow \pi_{q-p}(E(X \times \mathbb{A}^1)).$$

The projection p induces a pull-back morphism between these two spectral sequences. This morphism is an isomorphism on the second page, because each Nisnevich sheaf $\pi_n^{nis}(E)$ is strictly homotopy invariant by assumption. Hence $p^* : \pi_r(E(X)) \to \pi_r(X \times \mathbb{A}^1)$ is an isomorphism. Assertion (2) easily follows from assertion (1).

Below we shall need the following

Lemma 7.2. Let *F* be a radditive framed presheaf of Abelian groups. Then the associated sheaf in the Nisnevich topology has a unique structure of a framed presheaf such that the map $F \rightarrow F_{nis}$ is a map of framed presheaves.

Proof. This is proved in [22, 4.5].

Remark 7.3. We should stress that [22, 4.5] used in the proof of the preceding lemma is not true unless *F* is radditive.

Corollary 7.4. Let k be an infinite perfect field and E be an S¹-spectrum of simplicial Nisnevich sheaves with framed correspondences. Suppose E is locally an Ω -spectrum in the Nisnevich topology. Suppose it is (-1)-connected locally in the Nisnevich topology. Finally suppose that if char $k \neq 2$ (respectively char k = 2) then for any integer n, the Nisnevich presheaf $\pi_n(E)$ is homotopy invariant, quasi-stable and radditive (respectively a homotopy invariant, quasi-stable and radditive presheaf of $\mathbb{Z}[1/2]$ -modules). Let $E \to E^f$ be a fibrant replacement of E in the level injective model structure of ordinary sheaves of S¹-spectra. Then the following statements are true:

- (1) each motivic space E_n^f is motivically fibrant;
- (2) the spectrum E^{f} is fibrant in the stable injective motivic model category of S^{1} -spectra.

Proof. The Nisnevich presheaf $\pi_n(E)$ is a radditive framed presheaf. Hence the associated Nisnevich sheaf $\pi_n^{nis}(E)$ is equipped with a unique structure of a framed presheaf such that the canonical morphism $\pi_n(E) \to \pi_n^{nis}(E)$ is a morphism of framed presheaves by Lemma 7.2. By [7, 1.1] the Nisnevich sheaf $\pi_n^{nis}(E)$ is strictly homotopy invariant, hence so is the Nisnevich sheaf $\pi_n^{nis}(E^f)$ of E^f . Our statement now follows from the previous proposition.

Corollary 7.5. Let k be an infinite perfect field and Y be a simplicial object in $SmOp(Fr_0(k))$. Suppose the simplicial Nisnevich sheaf $C_*Fr(Y)$ is locally connected in the Nisnevich topology. Let $M_{fr}(Y) \rightarrow M_{fr}(Y)_f$ and $\widetilde{M}_{fr}(Y)\{1/2\} \rightarrow \widetilde{M}_{fr}(Y)\{1/2\}_f$ be fibrant replacements in the level injective model structure of ordinary sheaves of S^1 -spectra. Then:

- (1) if char $k \neq 2$ (respectively of any characteristic) then $M_{fr}(Y)_f$ (respectively $M_{fr}(Y)\{1/2\}_f$) is fibrant in the stable injective motivic model category of S^1 -spectra;
- (2) if char $k \neq 2$ (respectively of any characteristic) then for any $n \ge 0$ and any fibrant relpacement $C_*(Fr(-,Y \otimes S^n)) \to C_*(Fr(-,Y \otimes S^n))_f$ in $sShv_{\bullet}(Sm/k)$ (respectively $C_*(Fr(-,Y \otimes \mathscr{S}^n))\{1/2\} \to C_*(Fr(-,Y \otimes \mathscr{S}^n))\{1/2\}_f$), the space $C_*(Fr(-,Y \otimes S^n))_f$ (respectively $C_*(Fr(-,Y \otimes \mathscr{S}^n))\{1/2\}_f$) is fibrant in $sShv_{\bullet}(Sm/k)_{mot}$.

Proof. Suppose char $k \neq 2$. The zeroth space $C_*Fr(Y)$ of the framed spectrum $M_{fr}(Y)$ is locally connected. Hence the framed spectrum $M_{fr}(Y)$ is locally an Ω -spectrum by the Segal machine [19] and Theorem 6.5. The presheaves $\pi_n(M_{fr}(Y))$ are homotopy invariant, quasistable and radditive framed presheaves. Corollary 7.4 implies assertion (1). To prove the second one, note that any two fibrant replacements of $C_*Fr(Y)$ in $sShv_{\bullet}(Sm/k)$ are sectionwise weakly equivalent. Hence Corollary 7.4(1) implies the second assertion. The statement for $\widetilde{M}_{fr}(Y)\{1/2\}_f$ and $C_*(Fr(-,Y \otimes \mathscr{S}^n))\{1/2\}_f$ is proved in a similar fashion if we use the fact that the presheaves of (stable) homotopy groups of $\widetilde{M}_{fr}(Y)\{1/2\}$ and $C_*(Fr(-,Y \otimes \mathscr{S}^n))\{1/2\}$ are homotopy invariant, quasi-stable and radditive framed presheaves of $\mathbb{Z}[1/2]$ -modules (see Proposition 6.7 as well).

Under the notation of the preceding corollary we can now prove the following

Corollary 7.6. Let k be an infinite perfect field. If char $k \neq 2$ then the following statements are *true:*

(1) For any integer n > 0, the S¹-spectrum $M_{fr}(X \times T^n)_f$ is motivically fibrant and the motivic space $C_*Fr(X \times T^n)_f$ is motivically fibrant.

- (2) For any integer $n \ge 0$, the S^1 -spectrum $M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)_f$ is motivically fibrant and the motivic space $C_*(Fr(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1))_f$ is motivically fibrant. (3) For any integer $n \ge 0$, the S^1 -spectrum $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))_f$ is motivically fibrant
- and the motivic space $C_*Fr(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))_f$ is motivically fibrant.

In turn, in the case of any characteristic of k the following statements are true:

- (1) For any integer n > 0, the S¹-spectrum $\widetilde{M}_{fr}(X \times T^n)\{1/2\}_f$ is motivically fibrant and the motivic space $C_*Fr(X \times T^n)\{1/2\}_f$ is motivically fibrant.
- (2) For any integer $n \ge 0$, the S^1 -spectrum $\widetilde{M}_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)\{1/2\}_f$ is motivically
- (1) for any integer n ≥ 0, the S¹-spectrum M_{fr}(X × Tⁿ × (A¹//G_m)){1/2}_f is motivically fibrant.
 (3) For any integer n ≥ 0, the S¹-spectrum M_{fr}(X × Tⁿ × (A¹//G_m)){1/2}_f is motivically fibrant and the motivic space C_{*}Fr(X × Tⁿ × (A¹//G_m)){1/2}_f is motivically fibrant.

Proof. Suppose char $k \neq 2$. By [8, A.1] the spaces $C_*Fr(X \times T^n)$, $C_*Fr(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))$ of the corollary are locally connected in the Nisnevich topology. The space $C_*(Fr(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes$ S^{1}) is, moreover, sectionwise connected. Now our assertions follow from Corollary 7.5. In the case of any characteristic of k the second part of our statement is proved in a similar fashion.

We should stress that the previous corollary is of great utility in the proof of Theorem 4.1.

8. COMPARING FRAMED MOTIVES

One of the key properties of framed motives is that they convert motivic equivalences between certain motivic spaces to Nisnevich local weak equivalences. Some of such motivic equivalences are discussed in this section. Its main result, Theorem 8.2, is an essential step in proving Theorem 4.1. We start with preparations.

Every category \mathscr{A} with coproducts and zero object 0 has a natural action of finite pointed sets. For example, $\mathscr{A} = Fr_0(k)$ or, more generally, $\mathscr{A} = SmOp(Fr_0(k))$. Precisely, if $A \in \mathscr{A}$ and (K,*) is a finite pointes set, then we set $A \otimes K := A \sqcup \ldots \sqcup A$, where the coproduct is taken over non-base elements of K. Clearly, $A \otimes K$ is functorial in A and K. Note that $A \otimes * = 0$ and $0 \otimes K = 0.$

This action is extended to an action of finite pointed simplicial sets on the category $\Delta^{op} \mathscr{A}$ of simplicial objects in \mathscr{A} . Let (I, 1) denote the pointed simplicial set $\Delta[1]$ with basepoint 1. The *cone* of $A \in \mathscr{A}$ is the simplicial object $A \otimes I$ in \mathscr{A} . There is a natural morphism $i_0 : A \to A \otimes I$ in $\Delta^{\mathrm{op}} \mathscr{A}$. Given a morphism $f: A \to B$ in \mathscr{A} , denote by $B//_f A$ a simplicial object in \mathscr{A} which is obtained from the pushout in $\Delta^{\mathrm{op}} \mathscr{A}$ of the diagram

$$B \xleftarrow{f} A \xleftarrow{\iota_0} A \otimes I$$

We can think of B/fA as a cone of f. In practice if A is a subobject of B, we shall also write B/A to denote the simplicial object $B/I_{1}A$ in \mathscr{A} with $\iota: A \to B$ the inclusion. We have a sequence of simplicial objects in \mathscr{A}

$$A \xrightarrow{f} B \to B / / f A \to A \otimes S^1.$$

In practice, this sequence is a typical "triangle" of an associated triangulateed category (see, e.g., the proof of Theorem 8.2).

Notation 8.1. (1) In the particular example when $\mathscr{A} = Fr_0(k)$ and (X, x) is a pointed smooth variety, we shall write $X^{\wedge 1}$ to denote the cone X//x of the inclusion $x \hookrightarrow X$. The most common example is $\mathbb{G}_m^{\wedge 1}$ given by the pointed scheme ($\mathbb{G}_m, 1$). Regarding $\Delta^{\text{op}} Fr_0(k)$ as a full subcategory of the symmetric monoidal category $\Delta^{\text{op}} SmOp(Fr_0(k))$, we can take the *n*th monoidal power of X//x for every n > 0, which we shall denote by $X^{\wedge n}$. The most common example is $\mathbb{G}_m^{\wedge n}$.

(2) If X is an open subset of $Y \in Sm/k$, we shall denote by $(Y//X)^{\wedge n}$ the *n*th monoidal power of $Y//X \in \Delta^{\text{op}} Fr_0(k)$. The most common example will be $(\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}$.

If $\mathscr{A} = SmOp(Fr_0(k))$ then the symmetric monoidal product on $SmOp(Fr_0(k))$ defined above gives rise to a natural pairing


$$SmOp(Fr_0(k)) \times \Delta^{op}Fr_0(k) \rightarrow \Delta^{op}SmOp(Fr_0(k)).$$

Composing it with the framed motive functor, we get a functor

$$M_{fr}: SmOp(Fr_0(k)) \times \Delta^{op}Fr_0(k) \to Sp_{S^1}(sShv_{\bullet}(Sm/k)).$$

Taking pairings of $(X \times \mathbb{A}^n, X \times \mathbb{A}^n - X \times 0) \in SmOp(Fr_0(k))$ with $\mathbb{A}^1//\mathbb{G}_m \in \Delta^{op}Fr_0(k)$ and $\mathbb{G}_m^{\wedge 1} \otimes S^1 \in \Delta^{op}Fr_0(k)$, we get the framed motives $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))$ and $M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)$ respectively.

Consider a commutative diagram in $\Delta^{op} Fr_0(k)$



It induces a morphism of framed motives

$$\beta_*\alpha_*: M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m)) \to M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1), \quad n \ge 0,$$

and

$$\beta_*\{1/2\}\alpha_*\{1/2\}: M_{fr}(X\times T^n\times (\mathbb{A}^1//\mathbb{G}_m))\{1/2\} \to M_{fr}(X\times T^n\times \mathbb{G}_m^{\wedge 1}\otimes S^1)\{1/2\}, \quad n \ge 0.$$

The main result of this section is as follows.

Theorem 8.2. The morphism $\beta_*\alpha_*$ (respectively $\beta_*\{1/2\}\alpha_*\{1/2\}$) is a stable Nisnevich local weak equivalence of S^1 -spectra whenever the base field k is of characteristic different from 2 (respectively of any characteristic).

We postpone the proof of the theorem. It requires the language of "linear framed motives".

Definition 8.3. Let *X* and *Y* be smooth schemes. Denote by

- ◇ $\mathbb{Z}Fr_n(X,Y) := \mathbb{Z}[Fr_n(X,Y)] = \mathbb{Z}[Fr_n(X,Y)]/\mathbb{Z} \cdot 0_n$, i.e the free abelian group generated by the set $Fr_n(X,Y)$ modulo $\mathbb{Z} \cdot 0_n$;
- $\diamond \mathbb{Z}F_n(X,Y) := \mathbb{Z}Fr_n(X,Y)/A$, where *A* is a subgroup generated by the elementts

$$(Z \sqcup Z', U, (\varphi_1, \varphi_2, \dots, \varphi_n), g) - (Z, U \setminus Z', (\varphi_1, \varphi_2, \dots, \varphi_n)|_{U \setminus Z'}, g|_{U \setminus Z'}) - (Z', U \setminus Z, (\varphi_1, \varphi_2, \dots, \varphi_n)|_{U \setminus Z}, g|_{U \setminus Z}).$$

We shall also refer to the latter relation as the *additivity property for supports*. In other words, it says that a framed correspondence in $\mathbb{Z}F_n(X,Y)$ whose support is a disjoint union $Z \sqcup Z'$ equals the sum of the framed correspondences with supports Z and Z' respectively. Note that $\mathbb{Z}F_n(X,Y)$ is $\mathbb{Z}[Fr_n(X,Y)]$ modulo the subgroup generated by the elements as above, because $0_n = 0_n + 0_n$ in this quotient group, hence 0_n equals zero. Indeed, it is enough to observe that the support of 0_n equals $\emptyset \sqcup \emptyset$ and then apply the above relation to this support.

The elements of $\mathbb{Z}F_n(X,Y)$ are called *linear framed correspondences of level n* or just *linear framed correspondences*. It is worth to mention that $\mathbb{Z}F_n(X,Y)$ is a free abelian group generated by the elements of $Fr_n(X,Y)$ with connected support.

Denote by $\mathbb{Z}F_*(k)$ an additive category whose objects are those of Sm/k with Hom-groups defined as

$$\operatorname{Hom}_{\mathbb{Z}F_*(k)}(X,Y) = \bigoplus_{n \ge 0} \mathbb{Z}F_n(X,Y).$$

The composition is induced by the composition in the category $Fr_*(k)$.

There is a functor $Sm/k \to \mathbb{Z}F_*(k)$ which is the identity on objects and which takes a regular morphism $f: X \to Y$ to the linear framed correspondence $1 \cdot (X, X \times \mathbb{A}^0, pr_{\mathbb{A}^0}, f \circ pr_X) \in \mathbb{Z}F_0(k)$.

Definition 8.4. Let X, Y, S and T be schemes. The external product from Definition 2.4 induces a unique external product

$$\mathbb{Z}F_n(X,Y) \times \mathbb{Z}F_m(S,T) \xrightarrow{-\boxtimes -} \mathbb{Z}F_{n+m}(X \times S, Y \times T)$$

such that for any elements $a \in Fr_n(X,Y)$ and $b \in Fr_m(S,T)$ one has $1 \cdot a \boxtimes 1 \cdot b = 1 \cdot (a \boxtimes b) \in \mathbb{Z}F_{n+m}(X \times S, Y \times T)$.

For the constant morphism $c: \mathbb{A}^1 \to pt$, we set

$$\Sigma := - \boxtimes 1 \cdot (\{0\}, \mathbb{A}^1, t, c) \colon \mathbb{Z}F_n(X, Y) \to \mathbb{Z}F_{n+1}(X, Y)$$

and refer to it as the suspension.

Definition 8.5. For any *k*-smooth variety *Y* there is a presheaf $\mathbb{Z}F_*(-,Y)$ on the category $\mathbb{Z}F_*(k)$ represented by *Y*. We also have a $\mathbb{Z}F_*(k)$ -presheaf

$$\mathbb{Z}F(-,Y) := \operatorname{colim}(\mathbb{Z}F_0(-,Y) \xrightarrow{\Sigma} \mathbb{Z}F_1(-,Y) \xrightarrow{\Sigma} \cdots \xrightarrow{\Sigma} \mathbb{Z}F_n(-,Y) \xrightarrow{\Sigma} \cdots).$$

For a *k*-smooth variety *X*, the elements of $\mathbb{Z}F(X,Y)$ are also called *stable linear framed correspondences*. Stable linear framed correspondences *do not form* morphisms of a category.

Remark 8.6. For any X, Y in Sm/k one has $\mathbb{Z}F_*(-, X \sqcup Y) = \mathbb{Z}F_*(-, X) \oplus \mathbb{Z}F_*(-, Y)$ and $\mathbb{Z}F(-, X \sqcup Y) = \mathbb{Z}F(-, X) \oplus \mathbb{Z}F(-, Y)$.

For every $(Y, Y - S) \in SmOp(Fr_0(k))$, $\mathbb{Z}F_*(k)$ -presheaves $\mathbb{Z}F_*(-, Y/Y - S)$ and $\mathbb{Z}F(-, Y/Y - S)$ are defined in a similar fashion. Namely, each $\mathbb{Z}F_n(X, Y/Y - S)$ is the free abelian group generated by the elements of $Fr_n(X, Y/Y - S)$ with connected support. Then we set $\mathbb{Z}F_*(X, Y/Y - S) = \bigoplus_{n \ge 0} \mathbb{Z}F_n(X, Y/Y - S)$ and $\mathbb{Z}F(X, Y/Y - S)$ is obtained from $\mathbb{Z}F_*(X, Y/Y - S)$ by stabilization in the Σ -direction.

For every $Y \in \Delta^{\text{op}} SmOp(Fr_0(k))$ there is a Γ -space

$$(K,*) \in \Gamma^{\mathrm{op}} \mapsto \mathbb{Z}F(-, (Y/Y - S) \otimes K).$$

Definition 8.7. The *linear framed motive* $LM_{fr}(Y)$ of $Y \in \Delta^{\text{op}}SmOp(Fr_0(k))$ is the Segal S^1 -spectrum $(C_*\mathbb{Z}F(-,Y), C_*\mathbb{Z}F(-,Y \otimes S^1), C_*\mathbb{Z}F(-,Y \otimes S^2), \ldots)$ of spaces in $sShv_{\bullet}(Sm/k)$ associated with the Γ -space $K \in \Gamma^{\text{op}} \mapsto C_*\mathbb{Z}F(-,Y \otimes K) = \mathbb{Z}F(\Delta^{\bullet} \times -, Y \otimes K)$.

The linear framed motive $LM_{fr}(Y)\{1/2\}$ of $Y \in \Delta^{op}SmOp(Fr_0(k))$ with 1/2-coefficients is the S^1 -spectrum $(C_*\mathbb{Z}F(-,Y), C_*\mathbb{Z}F(-,Y\otimes \mathscr{S}^1), C_*\mathbb{Z}F(-,Y\otimes \mathscr{S}^2), ...)$ of motivic spaces in $sShv_{\bullet}(Sm/k)$. Equivalently, it is the value of the Γ -space $K \in \Gamma^{op} \mapsto C_*\mathbb{Z}F(-,Y\otimes K)$ at the spectrum S[1/2].

Note that $LM_{fr}(Y)$ is the Eilenberg–Mac Lane spectrum associated with the complex of Nisnevich sheaves $C_*\mathbb{Z}F(-,Y)$ (we often identify simplicial abelian groups with their normalized complexes by the Dold–Kan correspondence). Therefore $\pi_*(LM_{fr}(Y)) = H_*(C_*\mathbb{Z}F(-,Y))$.

The following lemma is proved similar to Proposition 6.7.

Lemma 8.8. Suppose Y is a simplicial object in $SmOp(Fr_0(k))$, then the presheaves of homotopy groups of $LM_{fr}(Y)\{1/2\}$ are presheaves of $\mathbb{Z}[1/2]$ -modules and the canonical morphism

$$\mu: LM_{fr}(Y) \to LM_{fr}(Y) \{1/2\}$$

induces isomorphisms on presheaves of homotopy groups with $\mathbb{Z}[1/2]$ -coefficients $\pi_*(\mu) \otimes \mathbb{Z}[1/2]$.

Denote by $LM_{fr}(Y)[1/2]$ the S¹-spectrum

$$(C_*\mathbb{Z}F(-,Y)\otimes\mathbb{Z}[1/2], C_*\mathbb{Z}F(-,Y\otimes S^1)\otimes\mathbb{Z}[1/2], C_*\mathbb{Z}F(-,Y\otimes S^2)\otimes\mathbb{Z}[1/2], \ldots)$$

It is the Segal S^1 -spectrum associated with the Γ -space $K \in \Gamma^{\text{op}} \mapsto C_* \mathbb{Z}F(-, Y \otimes K) \otimes \mathbb{Z}[1/2] := \mathbb{Z}F(\Delta^{\bullet} \times -, Y \otimes K) \otimes \mathbb{Z}[1/2]$. By the very construction, $\pi_*(LM_{fr}(Y)[1/2]) = \pi_*(LM_{fr}(Y)) \otimes \mathbb{Z}[1/2]$. We also denote by $\widetilde{LM}_{fr}(Y)\{1/2\} := \Omega^{\infty}LM_{fr}(Y)\{1/2\}$. The canonical morphism $\zeta : LM_{fr}(Y)\{1/2\} \to \widetilde{LM}_{fr}(Y)\{1/2\}$ is a stable equivalence. Notice that $LM_{fr}(Y)[1/2], \widetilde{LM}_{fr}(Y)\{1/2\}$ are sectionwise Ω -spectra.

Corollary 8.9. The composite map $LM_{fr}(Y) \xrightarrow{\mu} LM_{fr}(Y) \{1/2\} \xrightarrow{\zeta} \widetilde{LM}_{fr}(Y) \{1/2\}$ factors as

$$LM_{fr}(Y) \xrightarrow{\rho} LM_{fr}(Y)[1/2] \xrightarrow{\xi} \widetilde{LM}_{fr}(Y)\{1/2\}$$

and ξ is a level weak equivalence of spectra.

Proof of Theorem 8.2. Suppose char $k \neq 2$. Since the spectra $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))$ and $M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)$ are sectionwise connected, $\beta_* \alpha_*$ is a stable Nisnevich local equivalence of spectra if and only if so is the induced map on homology

$$\beta_*\alpha_*:\mathbb{Z}M_{fr}(X\times T^n\times (\mathbb{A}^1//\mathbb{G}_m))\to \mathbb{Z}M_{fr}(X\times T^n\times \mathbb{G}_m^{\wedge 1}\otimes S^1).$$

Here both linear spectra are defined by taking free Abelian groups of every entry of $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))$ and $M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)$ respectively. By [8, 1.2] the latter arrow is schemewise stably equivalent to the map of linear framed motives

$$\beta_*\alpha_*: LM_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m)) \to LM_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1).$$

We see that the map of the theorem is a stable Nisnevich local equivalence of spectra if and only if the morphism of complexes of Nisnevich sheaves

$$\beta_*\alpha_*: C_*\mathbb{Z}F(-,X\times T^n\times (\mathbb{A}^1//\mathbb{G}_m))\to C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{G}_m^{\wedge 1}\otimes S^1)$$

is a quasi-isomorphism.

Since $C_*\mathbb{Z}F(-,Y_1 \sqcup Y_2) = C_*\mathbb{Z}F(-,Y_1) \oplus C_*\mathbb{Z}F(-,Y_2)$ for any $Y_1, Y_2 \in \Delta^{\text{op}}SmOp(Fr_0(k))$, the digram (12) induces a commutative diagram of triangles of complexes of Nisnevich sheaves

Firstly, we claim that α is a schemewise quasi-isomorphism of complexes of presheaves. Indeed, we have a map of two triangles of complexes of presheaves

We see that the left square of the diagram is Mayer–Vietoris, hence α is a schemewise quasiisomorphisms of complexes of presheaves.

Secondly, we claim that β is a local quasi-isomorphism of complexes of Nisnevich sheaves. This is equivalent to showing that the complex $C_*\mathbb{Z}F(-,X \times T^n \times \mathbb{A}^{\wedge 1})$ of diagram (13) is locally quasi-isomorphic to zero. To prove the latter, consider a map of two triangles of complexes of sheaves

$$C_*\mathbb{Z}F(-,X\times T^n\times pt) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times pt) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times pt) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times pt) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \longrightarrow C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^1) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^n) \xrightarrow{+} C_*\mathbb{Z}F(-,X\times T^n\times \mathbb{A}^n)$$

The cohomology sheaves of the lower right complex are homotopy invariant and quasi-stable framed presheaves. By [7, 1.1] these cohomology sheaves are strictly homotopy invariant. The terms of this complex are contractible sheaves. Now the proof of [20, 1.10.2] yields the local acyclicity of the complex. It follows that $C_*\mathbb{Z}F(-, X \times T^n \times \mathbb{A}^{\wedge 1})$ is locally acyclic, and hence β is locally a quasi-isomorphism.

The proof that $\beta_*\{1/2\}\alpha_*\{1/2\}$ is a stable Nisnevich local weak equivalence repeats the above proof word for word if we notice that homology of framed motives with 1/2-coefficients is computed as $LM_{fr}(Y)[1/2]$ (see Lemma 8.8 and Corollary 8.9 as well). This completes the proof of the theorem.

In fact, the proof of Theorem 8.2 also shows the following fact.

Corollary 8.10. Suppose char $k \neq 2$ (respectively k is of any characteristic). Then for every $n \geq 0$ and $X \in Sm/k$, the natural maps $M_{fr}(X \times T^n \times \mathbb{A}^1) \to M_{fr}(X \times T^n)$ and $LM_{fr}(X \times T^n \times \mathbb{A}^1) \to LM_{fr}(X \times T^n)$ (respectively $M_{fr}(X \times T^n \times \mathbb{A}^1)\{1/2\} \to M_{fr}(X \times T^n)\{1/2\}$ and $LM_{fr}(X \times T^n \times \mathbb{A}^1)\{1/2\} \to LM_{fr}(X \times T^n)\{1/2\}$) are stable local weak equivalences of S^1 -spectra.

Let us take the *n*th power $(\beta \alpha)^{\wedge n} : (\mathbb{A}^1 / /\mathbb{G}_m)^{\wedge n} \to (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n}$ of the morphism $\beta \alpha$ in the symmetric monoidal category $\Delta^{\text{op}} SmOp(Fr_0(k))$. Below we shall also need the following

Corollary 8.11. Suppose chark $\neq 2$ (respectively k is of any characteristic). For every $n \ge 1$ and $X \in Sm/k$, the map $(\beta \alpha)^{\wedge n}_* : C_*Fr(-, X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \to C_*Fr(-, X \times (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n})$

(respectively $(\beta \alpha)^{\wedge n}_*: C_*Fr(-, X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})\{1/2\} \to C_*Fr(-, X \times (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n})\{1/2\})$ is a local Nisnevich weak equivalence of motivic spaces.

Proof. Suppose char $k \neq 2$. The space $C_*Fr(-, X \times (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n})$ is plainly sectionwise connected. By [8, A.1] the space $C_*Fr(-, X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})$ is locally connected. Therefore $M_{fr}(X \times (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n})$ is sectionwise an Ω -spectrum and $M_{fr}(X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})$ is locally an Ω -spectrum by Theorem 6.5. Therefore our assertion would follow if we showed that the map $(\beta \alpha)_*^{\wedge n} : M_{fr}(X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \to M_{fr}(X \times (\mathbb{G}_m^{\wedge 1} \otimes S^1)^{\wedge n})$ is a local Nisnevich weak equivalence of S^1 -spectra. The latter follows by using induction in n, Theorem 8.2 and the fact that the realization of Nisnevich local weak equivalences is a local Nisnevich weak equivalence. If k is of any characteristic then our proof is similar to the proof above.

We finish the section by proving the following useful

Theorem 8.12. Suppose char $k \neq 2$ (respectively k is of any characteristic). For every $n \ge 0$ and every elementary Nisnevich square



the square of S¹-spectra

(repectively the square

of framed motives with 1/2-coefficients) is homotopy cartesian locally in the Nisnevich topology. The same is also true for linear framed motives and linear framed motives with 1/2-coefficients.

Proof. Suppose char $k \neq 2$. Since we deal with connected S^1 -spectra, the proof of Theorem 8.2 shows that our statement suffuces to verify for linear framed motives. It follows from the proof of [22, 4.4] that the sequence of presheaves

$$0 \to \mathbb{Z}F_s(-, U' \times T^n) \to \mathbb{Z}F_s(-, U \times T^n) \oplus \mathbb{Z}F_s(-, X' \times T^n) \to \mathbb{Z}F_s(-, X \times T^n) \to 0$$

is locally exact for every $s \ge 0$. Passing to the colimit over *s*, the sequence of presheaves

$$0 \to \mathbb{Z}F(-, U' \times T^n) \to \mathbb{Z}F(-, U \times T^n) \oplus \mathbb{Z}F(-, X' \times T^n) \to \mathbb{Z}F(-, X \times T^n) \to 0$$

is locally exact as well. It follows that the sequence of Eilenberg–Mac Lane spectra $EM(\mathbb{Z}F(-,U'\times T^n)) \rightarrow EM(\mathbb{Z}F(-,U\times T^n)) \times EM(\mathbb{Z}F(-,X'\times T^n)) \rightarrow EM(\mathbb{Z}F(-,X\times T^n))$

is locally a homotopy fibre sequence. It follows that the sequence

$$LM_{fr}(U' \times T^n) \to LM_{fr}(U \times T^n) \times LM_{fr}(X' \times T^n) \to LM_{fr}(X \times T^n)$$
(14)

is a homotopy fibre sequence in the motivic model structure of S^1 -spectra, and hence so is the sequence

$$LM_{fr}(U' \times T^n)_f \to LM_{fr}(U \times T^n)_f \times LM_{fr}(X' \times T^n)_f \to LM_{fr}(X \times T^n)_f,$$

where "f" as in Corollary 7.4. It follows from Corollary 7.4 that the latter sequence is a sequence of fibrant objects in the stable injective motivic model structure of S^1 -spectra. Therefore this sequence is also locally a homotopy fibre sequence, and hence so is sequence (14). If k is of any characteristic then our proof is similar to the proof above (recall that homology of framed motives with 1/2-coefficients is computed as linear framed motives tensored with $\mathbb{Z}[1/2]$).

9. PROOF OF THEOREM 4.1

In this section we prove Theorem 4.1. We first give a proof for the second statement of the theorem followed by the first statement.

9.1. Proof of Theorem 4.1(2)

Recall that $A \in sShv_{\bullet}(Sm/k)$ is *finitely presentable* if the functor $Hom_{sShv_{\bullet}(Sm/k)}(A, -)$ preserves directed colimits. Using the General Framework on p. 14, for every $A, L \in sShv_{\bullet}(Sm/k)$ with A finitely presentable there are canonical morphisms in $sShv_{\bullet}(Sm/k)$

$$C_*\mathscr{F}r(L) \to \underline{\operatorname{Hom}}(A, C_*\mathscr{F}r(L \wedge A)), \quad C_*\mathscr{F}r(L)\{1/2\} \to \underline{\operatorname{Hom}}(A, C_*\mathscr{F}r(L \wedge A)\{1/2\})$$

as well as canonical morphisms of ordinary S^1 -spectra of simplicial Nisnevich sheaves

$$\mathscr{M}_{fr}(L) \to \underline{\mathrm{Hom}}(A, \mathscr{M}_{fr}(L \wedge A)), \quad \widetilde{\mathscr{M}}_{fr}(L)\{1/2\} \to \underline{\mathrm{Hom}}(A, \widetilde{\mathscr{M}}_{fr}(L \wedge A)\{1/2\}).$$
(15)

They induce the corresponding morphisms of spaces and S^1 -spectra respectively

$$a_A : C_* \mathscr{F} r(L)_f \to \underline{\operatorname{Hom}}(A, C_* \mathscr{F} r(L \wedge A)_f), \quad a_A \{1/2\} : C_* \mathscr{F} r(L) \{1/2\}_f \to \underline{\operatorname{Hom}}(A, C_* \mathscr{F} r(L \wedge A) \{1/2\}_f)$$
and

and

$$\alpha_{A}: \mathscr{M}_{fr}(L)_{f} \to \underline{\mathrm{Hom}}(A, \mathscr{M}_{fr}(L \wedge A)_{f}), \quad \alpha_{A}\{1/2\}: \mathscr{M}_{fr}(L)\{1/2\}_{f} \to \underline{\mathrm{Hom}}(A, \mathscr{M}_{fr}(L \wedge A)\{1/2\}_{f})$$

where $C_* \mathscr{F}r(L)_f, C_* \mathscr{F}r(L)\{1/2\}_f$ (respectively $\mathscr{M}_{fr}(L)_f, \widetilde{\mathscr{M}}_{fr}(L)\{1/2\}_f$) are Nisnevich local fibrant replacements of $C_* \mathscr{F} r(L), C_* \mathscr{F} r(L) \{1/2\}$ (respectively a level Nisnevich local fibrant replacement of $\mathcal{M}_{fr}(L), \mathcal{M}_{fr}(L)\{1/2\}$ in the category of ordinary S¹-spectra).

Lemma 9.1. Suppose $u: A \to B$ is a motivic weak equivalence in $sShv_{\bullet}(Sm/k)$ between finitely presentable objects such that the induced map $u_* : \mathscr{M}_{fr}(L \wedge A) \to \mathscr{M}_{fr}(L \wedge B)$ is a stable Nisnevich local weak equivalence of spectra. Suppose $\mathcal{M}_{fr}(L)_f, \mathcal{M}_{fr}(L \wedge A)_f, \mathcal{M}_{fr}(L \wedge B)_f$ are all motivically fibrant S¹-spectra. Then $\alpha_A : \mathscr{M}_{fr}(L)_f \to \operatorname{Hom}(A, \mathscr{M}_{fr}(L \wedge A)_f)$ is a sectionwise stable equivalence if and only if so is $\alpha_B : \mathscr{M}_{fr}(L)_f \to \operatorname{Hom}(B, \mathscr{M}_{fr}(L \wedge B)_f).$

Suppose $u: A \rightarrow B$ is a motivic weak equivalence in $sShv_{\bullet}(Sm/k)$ between finitely presentable objects such that $u\{1/2\}_*$: $\mathscr{M}_{fr}(L \wedge A)\{1/2\} \to \mathscr{M}_{fr}(L \wedge B)\{1/2\}$ is a stable Nisnevich local weak equivalence of spectra. Suppose $\widetilde{\mathcal{M}}_{fr}(L)\{1/2\}_f, \widetilde{\mathcal{M}}_{fr}(L \wedge A)\{1/2\}_f, \widetilde{\mathcal{M}}_{fr}(L \wedge B)\{1/2\}_f$ are all motivically fibrant S^1 -spectra. Then $\alpha\{1/2\}_A : \widetilde{\mathcal{M}}_{fr}(L)\{1/2\}_f \to \underline{\mathrm{Hom}}(A, \widetilde{\mathcal{M}}_{fr}(L \wedge \mathbb{Z}))$ A) $\{1/2\}_f$ is a sectionwise stable equivalence if and only if so is $\alpha\{1/2\}_B : \widetilde{\mathcal{M}}_{fr}(L)\{1/2\}_f \to \mathcal{M}_{fr}(L)\{1/2\}_f$ <u>Hom</u> $(B, \mathscr{M}_{fr}(L \wedge B)\{1/2\}_f)$.

Proof. Commutative square (10) of the General Framework gives rise to a commutative square

By assumption, $\mathscr{M}_{fr}(L \wedge A)_f, \mathscr{M}_{fr}(L \wedge B)_f$ are motivically fibrant S^1 -spectra, hence u^* is sectionwise stable equivalence. Since $\mathscr{M}_{fr}(L \wedge A) \to \mathscr{M}_{fr}(L \wedge B)$ is a stable Nisnevich local weak equivalence of spectra, it follows that $\mathscr{M}_{fr}(L \wedge A)_f \to \mathscr{M}_{fr}(L \wedge B)_f$ is a sectionwise stable weak equivalence of spectra. We see that the right vertical arrow u_* of the square is a sectionwise stable weak equivalence of spectra. Our statement now follows from the two-out-three property for weak equivalences (the statement for $\alpha\{1/2\}_A, \alpha\{1/2\}_B$ repeats the above arguments word for word).

Corollary 9.2. (1) Under the assumptions of Lemma 9.1 the map of spaces $a_A : C_* \mathscr{F}r(L)_f \to \underline{\operatorname{Hom}}(A, C_* \mathscr{F}r(L \wedge A)_f)$ is a sectionwise weak equivalence if and only if so is $a_B : C_* \mathscr{F}r(L)_f \to \underline{\operatorname{Hom}}(B, C_* \mathscr{F}r(L \wedge B)_f)$.

(2) Under the assumptions of Lemma 9.1 the map of spaces $a\{1/2\}_A : C_* \mathscr{F}r(L)\{1/2\}_f \rightarrow \underline{\operatorname{Hom}}(A, C_* \mathscr{F}r(L \wedge A)\{1/2\}_f)$ is a sectionwise weak equivalence if and only if so is $a\{1/2\}_B : C_* \mathscr{F}r(L)\{1/2\}_f \rightarrow \underline{\operatorname{Hom}}(B, C_* \mathscr{F}r(L \wedge B)\{1/2\}_f)$.

Lemma 9.3. (1) Suppose $u : A \to B$ is a motivic weak equivalence in $sShv_{\bullet}(Sm/k)$ between finitely presentable objects. Suppose $\mathscr{M}_{fr}(L)_f, \mathscr{M}_{fr}(L \wedge B)_f$ are motivically fibrant S^1 -spectra. Then the composite map of spaces $C_*\mathscr{F}r(L)_f \xrightarrow{a_B} \underline{Hom}(B, C_*\mathscr{F}r(L \wedge B)_f) \xrightarrow{u^*} \underline{Hom}(A, C_*\mathscr{F}r(L \wedge B)_f)$ is a sectionwise weak equivalence if and only if $\alpha_B : \mathscr{M}_{fr}(L)_f \to \underline{Hom}(B, \mathscr{M}_{fr}(L \wedge B)_f)$ is a sectionwise stable equivalence of spectra.

(2) Suppose $u : A \to B$ is a motivic weak equivalence in $sShv_{\bullet}(Sm/k)$ between finitely presentable objects. Suppose $\widetilde{\mathcal{M}}_{fr}(L)\{1/2\}_f, \widetilde{\mathcal{M}}_{fr}(L \wedge B)\{1/2\}_f$ are motivically fibrant S^1 -spectra. Then the composite map of spaces

$$C_*\mathscr{F}r(L)\{1/2\}_f \xrightarrow{a\{1/2\}_B} \underline{\operatorname{Hom}}(B, C_*\mathscr{F}r(L \wedge B)\{1/2\}_f) \xrightarrow{u^*} \underline{\operatorname{Hom}}(A, C_*\mathscr{F}r(L \wedge B)\{1/2\}_f)$$

is a sectionwise weak equivalence if and only if

$$\alpha\{1/2\}_B: \mathscr{M}_{fr}(L)\{1/2\}_f \to \underline{\mathrm{Hom}}(B, \mathscr{M}_{fr}(L \wedge B)\{1/2\}_f)$$

is a sectionwise stable equivalence of spectra.

Proof. Our assumptions on spectra imply $C_*\mathscr{F}r(L)_f, C_*\mathscr{F}r(L \wedge B)_f$ are motivically fibrant spaces. It follows that $C_*\mathscr{F}r(L)_f \xrightarrow{a_B} \operatorname{Hom}(B, C_*\mathscr{F}r(L \wedge B)_f) \xrightarrow{u^*} \operatorname{Hom}(A, C_*\mathscr{F}r(L \wedge B)_f)$ is a sectionwise weak equivalence if and only if so is $C_*\mathscr{F}r(L)_f \xrightarrow{a_B} \operatorname{Hom}(B, C_*\mathscr{F}r(L \wedge B)_f)$, because $u: A \to B$ is a motivic weak equivalence (recall that all spaces in the Morel–Voevodsy's model category $sShv_{\bullet}(Sm/k)_{mot}$ are cofibrant).

Again because of our assumptions on spectra we have that $C_*\mathscr{F}r(L)_f \xrightarrow{a_B} \operatorname{Hom}(B, C_*\mathscr{F}r(L \wedge B)_f)$ is a sectionwise weak equivalence if and only if $\alpha_B : \mathscr{M}_{fr}(L)_f \to \operatorname{Hom}(B, \mathscr{M}_{fr}(L \wedge B)_f)$ is a sectionwise stable equivalence of spectra. The corresponding proof for maps $a\{1/2\}_B$, $\alpha\{1/2\}_B$ is similar to that above.

We are now in a position to prove the second statement of Theorem 4.1.

Proof of Theorem 4.1(2). By Corollary 7.6 for any integer n > 0, the S^1 -spectrum $M_{fr}(X \times T^n)_f$ is motivically fibrant and the motivic space $C_*Fr(X \times T^n)_f$ is motivically fibrant. Let $u : \mathbb{P}^{\wedge 1} = (\mathbb{P}^1, \infty) \to T$ be the canonical motivic weak equivalence in $sShv_{\bullet}(Sm/k)$. It is also given by the framed correspondence of level one $(\{0\}, \mathbb{A}^1, t) \in Fr_1(pt, pt)$. By Lemma 9.3 the map

$$C_*Fr(X \times T^n)_f \to \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*Fr(X \times T^{n+1})_f)$$

is a sectionwise weak equivalence if and only if $\alpha_T : M_{fr}(X \times T^n)_f \to \underline{\text{Hom}}(T, M_{fr}(X \times T^{n+1})_f)$ is a sectionwise stable equivalence of spectra.

Consider the zigzag of motivic weak equivalences

$$T \xleftarrow{\sim} (-, \mathbb{A}^1 / / \mathbb{G}_m)_+ \xrightarrow{\sim} (-, \mathbb{G}_m^{\wedge 1} \otimes S^1)_+,$$

where the right arrow is induced by $\beta \alpha$ of diagram (12). By Corollary 7.6 for any integer $n \ge 0$, the S^1 -spectra $M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)_f, M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))_f$ are motivically fibrant and $C_*(Fr(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1))_f, C_*Fr(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m))_f$ are motivically fibrant spaces.

By [8, 8.1] $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m)) \to M_{fr}(X \times T^{n+1})$ is a stable Nisnevich local weak equivalence of spectra. By Theorem 8.2 $M_{fr}(X \times T^n \times (\mathbb{A}^1//\mathbb{G}_m)) \to M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)$ is a stable Nisnevich local weak equivalence of spectra. By Lemma 9.1 $\alpha_T : M_{fr}(X \times T^n)_f \to \underline{\mathrm{Hom}}(T, M_{fr}(X \times T^{n+1})_f)$ is a sectionwise stable equivalence of spectra if and only if so is the map of spectra $\alpha_{\mathbb{G}_m^{\wedge 1} \otimes S^1} : M_{fr}(X \times T^n)_f \to \underline{\mathrm{Hom}}((\mathbb{G}_m^{\wedge 1} \otimes S^1)_+, M_{fr}(X \times T^n \times \mathbb{G}_m^{\wedge 1} \otimes S^1)_f).$

Consider a commutative diagram

with $M_{fr}(X \times (\mathbb{A}^1//\mathbb{G}_m)^n)_f$ a stable Nisnevich local fibrant replacement of ordinary spectra and $(\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}$ from Notation 8.1. It follows from [8, 1.1] that the left vertical arrow is a sectionwise stable weak equivalence of spectra, hence so is the right vertical arrow. We see that the lower arrow is a sectionwise stable weak equivalence of spectra if and only if so is the upper one. But the upper arrow is a sectionwise stable weak equivalence of spectra by the Cancellation Theorem for framed motives of algebraic varieties [1, Theorem A] and Theorem 6.5. The proof for spaces $C_*Fr(X \times T^n)\{1/2\}_f$ repeats the above proof word for word.

The proof of Theorem 4.1(2) and Corollary 8.11 also implies the following

Corollary 9.4. For any $n \ge 1$, the map $C_*Fr(-,X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \to C_*Fr(-,X \times T^n)$ (respectively $C_*Fr(-,X \times (\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})\{1/2\} \to C_*Fr(-,X \times T^n)\{1/2\}$) is a local Nisnevich weak equivalence of motivic spaces whenever char $k \ne 2$ (respectively k is of any characteristic).

9.2. Proof of Theorem 4.1(1)

In this section we finish the proof of Theorem 4.1. It remains to show part (1) of the theorem. Denote by $\mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ the category of \mathbb{P}^1 -spectra with \mathbb{P}^1 pointed at ∞ . We shall work with the injective stable motivic model structure on $\mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ (see [14] for details). The weak equivalences in this model category will be referred to as *stable equivalences*.

We define the *fake suspension functor* $\Sigma_{\mathbb{P}^1}^{\ell} : \mathbf{Spt}^{\mathbb{P}^1}(Sm/k) \to \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ by $(\Sigma_{\mathbb{P}^1}^{\ell}\mathscr{Z})_n = \mathscr{Z}_n \wedge \mathbb{P}^1$ and structure maps

$$(\Sigma_{\mathbb{P}^1}^{\ell}\mathscr{Z}_n) \wedge \mathbb{P}^1 \xrightarrow{\sigma_n \wedge \mathbb{P}^1} \Sigma_{\mathbb{P}^1}^{\ell} \mathscr{Z}_{n+1}$$

where σ_n is a structure map of \mathscr{Z} . The fake suspension functor is left adjoint to the *fake loops* functor $\Omega_{\mathbb{P}^1}^{\ell}$: **Spt**^{\mathbb{P}^1}(*Sm*/*k*) \to **Spt**^{\mathbb{P}^1}(*Sm*/*k*) defined by $(\Omega_{\mathbb{P}^1}^{\ell}\mathscr{Z})_n = \Omega_{\mathbb{P}^1}\mathscr{Z}_n = \underline{\mathrm{Hom}}(\mathbb{P}^1, \mathscr{Z}_n)$ and structure maps adjoint to

$$\Omega_{\mathbb{P}^1}\mathscr{Z}_n \xrightarrow{\Omega_{\mathbb{P}^1} \widetilde{\sigma}_n} \Omega_{\mathbb{P}^1}(\Omega_{\mathbb{P}^1}\mathscr{Z}_{n+1}),$$

where $\tilde{\sigma}_n$ is adjoint to the structure map σ_n of \mathscr{Z} .

Define the *shift functors* $t : \mathbf{Spt}^{\mathbb{P}^1}(Sm/k) \to \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ and $s : \mathbf{Spt}^{\mathbb{P}^1}(Sm/k) \to \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ by $(s\mathscr{Z})_n = \mathscr{Z}_{n+1}$ and $(t\mathscr{Z})_n = \mathscr{Z}_{n-1}, (t\mathscr{Z})_0 = pt$, with the evident structure maps. Note that t is left adjoint to s.

Define Θ : $\mathbf{Spt}^{\mathbb{P}^1}(Sm/k) \to \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ to be the functor $s \circ \Omega_{\mathbb{P}^1}^{\ell}$, where *s* is the shift functor. Then we have a natural map $\iota_{\mathscr{Z}} : \mathscr{Z} \to \Theta \mathscr{Z}$, and we define

$$\Theta^{\infty}\mathscr{Z} = \operatorname{colim}(\mathscr{Z} \xrightarrow{\iota_{\mathscr{Z}}} \Theta \mathscr{Z} \xrightarrow{\Theta\iota_{\mathscr{Z}}} \Theta^{2} \mathscr{Z} \xrightarrow{\Theta^{2}\iota_{\mathscr{Z}}} \cdots \xrightarrow{\Theta^{n-1}\iota_{\mathscr{Z}}} \Theta^{n} \mathscr{Z} \xrightarrow{\Theta^{n}\iota_{\mathscr{Z}}} \cdots).$$
(16)

Set $\eta_{\mathscr{Z}}: \mathscr{Z} \to \Theta^{\infty} \mathscr{Z}$ to be the obvious natural transformation.

Lemma 9.5. For every \mathbb{P}^1 -spectrum \mathscr{L} the natural map $\eta_{\mathscr{Z}} : \mathscr{Z} \to \Theta^{\infty} \mathscr{Z}$ is a stable motivic weak equivalence.

Proof. The assertion would follow from [11, 4.11] as soon as we find a weakly finitely generated model structure on pointed simplicial presheaves $sPre_{\bullet}(Sm/k)$ in the sense of [3] such that its model category of \mathbb{P}^1 -spectra is Quillen equivalent to the injective stable motivic model structure of Jardine [14]. Such a model structure on $sPre_{\bullet}(Sm/k)$ is the flasque motivic model structure of Isaksen [12]. The fact that it is weakly finitely generated follows from [12, 3.10, 4.9, 5.1] and [5, 2.2].

We are now in a position to prove Theorem 4.1(1).

Proof of Theorem 4.1(1). We shall construct the stable equivalence of the theorem not only for smooth algebraic varieties but also for all pointed motivic spaces. This is of independent interest. Let $\mathscr{X} \in sPre_{\bullet}(Sm/k)$ be a pointed motivic space. Consider its suspension spectrum

$$\Sigma^{\infty}_{\mathbb{P}^1}\mathscr{X} = (\mathscr{X}, \mathscr{X} \wedge \mathbb{P}^1, \mathscr{X} \wedge \mathbb{P}^{\wedge 2}, \ldots)$$

We set

$$\Sigma^{\infty}_{\mathbb{P}^1 T} \mathscr{X} = (\mathscr{X}, \mathscr{X} \wedge T, \mathscr{X} \wedge T^2, \ldots)$$

be the \mathbb{P}^1 -spectrum with structure maps defined by $(\mathscr{X} \wedge T^n) \wedge \mathbb{P}^1 \xrightarrow{\text{id} \wedge \sigma} \mathscr{X} \wedge T^{n+1}$, where $\sigma : \mathbb{P}^1 \to T$ is the canonical motivic equivalence of sheaves.

Since smash product of a motivic weak equivalence and a motivic space is again a motivic weak equivalence, we get a level motivic equivalence of spectra

$$\sigma: \Sigma^{\infty}_{\mathbb{P}^1}\mathscr{X} \to \Sigma^{\infty}_{\mathbb{P}^1,T}\mathscr{X}$$

Let us take the Nisnevich sheaf $(\mathscr{X} \wedge T^n)_{nis}$ of each space $\mathscr{X} \wedge T^n$ of $\Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}$. Then we get a \mathbb{P}^1 -spectrum $\Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}_{nis}$ and a level local weak equivalence of spectra

$$\nu: \Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X} \to \Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}_{\mathrm{nis}}$$

By Lemma 9.5 the natural map of spectra

$$\eta: \Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}_{\text{nis}} \to \Theta^{\infty} \Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}_{\text{nis}}$$

is a stable equivalence.

Evaluation of the spectrum $\sum_{\mathbb{P}^1,T}^{\infty} \mathscr{X}_{nis}$ at a smooth scheme U has entries

 $(\operatorname{colim}_{n}(\operatorname{Hom}_{Shv_{\bullet}(Sm/k)}(U_{+} \wedge \mathbb{P}^{\wedge n}, \mathscr{X}_{\operatorname{nis}} \wedge T^{n})), \operatorname{colim}_{n}(\operatorname{Hom}_{Shv_{\bullet}(Sm/k)}(U_{+} \wedge \mathbb{P}^{\wedge n}, \mathscr{X}_{\operatorname{nis}} \wedge T^{n+1})), \ldots),$

where each smash product $\mathscr{X}_{nis} \wedge T^n$ is taken in the category of pointed simplicial sheaves.

So the spectrum $\Theta^{\infty} \Sigma^{\infty}_{\mathbb{P}^1,T} \mathscr{X}_{nis}$ can be written as

$$(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}),\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T),\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T^{2}),\ldots)$$

If we apply the functor C_* to the spectrum $\Theta^{\infty} \Sigma_{\mathbb{P}^1}^{\infty} \mathscr{X}_{nis}$ levelwise, we get a spectrum

$$(C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}})), C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T)), C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T^2)), \ldots).$$

By [17, 2.3.8] the natural map of spectra

$$\delta: \Theta^{\infty} \Sigma^{\infty}_{\mathbb{P}^1, T} \mathscr{X}_{\operatorname{nis}} \to C_*(\Theta^{\infty} \Sigma^{\infty}_{\mathbb{P}^1, T} \mathscr{X}_{\operatorname{nis}})$$

is a level motivic weak equivalence.

Taking a Nisnevich local fibrant replacement $C_*(\mathscr{F}r(-,\mathscr{X}_{nis} \wedge T^n))_f$ of every motivic space $C_*(\mathscr{F}r(-,\mathscr{X}_{nis} \wedge T^n))$, we get a spectrum

$$C_*(\Theta^{\infty}\Sigma^{\infty}_{\mathbb{P}^1,T}\mathscr{X}_{\mathrm{nis}})_f := (C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}))_f, C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T))_f, C_*(\mathscr{F}r(-,\mathscr{X}_{\mathrm{nis}}\wedge T^2))_f, \ldots)$$

Denote by

$$\rho := \delta \circ \eta \circ \nu \circ \sigma : \Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X}) \to C_*(\Theta^{\infty}\Sigma^{\infty}_{\mathbb{P}^1,T}\mathscr{X}_{\mathrm{nis}})_f.$$

Then ρ is a stable motivic weak equivalence of \mathbb{P}^1 -spectra. Moreover, if \mathscr{X} is represented by a smooth scheme X (or, more generally, by a directed colimit of simplicial smooth schemes from $\Delta^{\operatorname{op}}Fr_0(k)$), then $C_*(\Theta^{\infty}\Sigma_{\mathbb{P}^1,T}^{\infty}\mathscr{X}_{\operatorname{nis}})_f$ is naturally isomorphic in SH(k) to the spectrum $M_{\mathbb{P}^{\Lambda}1}(\mathscr{X})_f$ such that the composite isomorphism

$$\Sigma^{\infty}_{\mathbb{P}^{1}}(\mathscr{X}) \xrightarrow{\rho} C_{*}(\Theta^{\infty}\Sigma^{\infty}_{\mathbb{P}^{1},T}\mathscr{X}_{\mathrm{nis}})_{f} \xrightarrow{\cong} M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_{f}$$

equals \varkappa_f of the theorem.

Moreover, the canonical morphism

$$M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_f \to M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})\{1/2\}_f$$

is a 2^{-1} -stable motivic equivalence of \mathbb{P}^1 -spectra by Proposition 6.7. This finishes the proof of part (1) of Theorem 4.1.

10. Computing infinite \mathbb{P}^1 -loop spaces

Let $\Delta^{\text{op}} Fr_0(k)$ be the full subcategory of $sShv_{\bullet}(Sm/k)$ consisting of directed colimits of objects from $\Delta^{\text{op}} Fr_0(k)$. Recall that $\Delta^{\text{op}} Fr_0(k)$ can be regarded as a full subcategory of $sShv_{\bullet}(Sm/k)$ sending $X \in Fr_0(k)$ to $X_+ \in Shv_{\bullet}(Sm/k)$.

In order to compute $\Omega_{\mathbb{P}^1}^{\infty} \Sigma_{\mathbb{P}^1}^{\infty}(\mathscr{F})$ of any motivic space $\mathscr{F} \in sShv_{\bullet}(Sm/k)$ (see Theorem 10.5), we need the following extension of Theorem 4.1 to objects of $\Delta^{op} Fr_0(k)$:

Theorem 10.1. Let k be an infinite perfect field and Y an object of $\Delta^{op} Fr_0(k)$. Then the following statements are true:

- (1) The morphism $\varkappa_f : \Sigma_{\mathbb{P}^1}^{\infty} Y_+ \to M_{\mathbb{P}^{\wedge 1}}(Y)_f$ is a stable motivic equivalence of \mathbb{P}^1 -spectra and the morphism $\tau_f \circ \varkappa_f : \Sigma_{\mathbb{P}^1}^{\infty} Y_+ \to M_{\mathbb{P}^{\wedge 1}}(Y)\{1/2\}_f$ is a 2^{-1} -stable motivic equivalence of \mathbb{P}^1 -spectra.
- (2) If char $k \neq 2$ (respectively k is of any characteristic) then the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(Y)_f$ (respectively $M_{\mathbb{P}^{\wedge 1}}(Y)\{1/2\}_f$) is a motivically fibrant Ω -spectrum in posisitive degrees. This means that for every positive integer n > 0 each motivic space $C_*(Fr(-, Y \times T^n))_f$ (respectively $C_*(Fr(-, Y \times T^n))\{1/2\}_f$) is motivically fibrant in the Morel–Voevodsky [17] motivic model category of simplicial Nisnevich sheaves and the structure map

$$C_*(Fr(-,Y\times T^n))_f\to \Omega_{\mathbb{P}^1}(C_*(Fr(-,Y\times T^{n+1}))_f)$$

(respectively $C_*(Fr(-,Y \times T^n))\{1/2\}_f \rightarrow \Omega_{\mathbb{P}^1}(C_*(Fr(-,Y \times T^{n+1}))\{1/2\}_f))$ is a weak equivalence schemewise.

Proof. The first statement of the theorem can be proved similar to Theorem 4.1(1) for any $\overrightarrow{F} \in \Delta^{\operatorname{op}} Fr_0(k)$. Without loss of generality it is enough to prove the second statement of the theorem for $Y \in \Delta^{\operatorname{op}} Fr_0(k)$. Indeed, we use the facts that the functor $M_{\mathbb{P}^{\wedge 1}}(-)$ respects directed colimits and directed colimits of locally fibrant objects are Nisnevich excisive (even more: they are fibrant in the local flasque model structure of sheaves in the sense of [12, 4.6]).

We first observe that each space $C_*(Fr(-,Y \times T^n))$, n > 0, is locally connected, because it is the geometric realization of a simplicial locally connected *H*-space $[k \in \Delta^{\text{op}} \mapsto C_*(Fr(-,Y_k \times T^n))]$ and $\pi_0^{\text{nis}}(C_*(Fr(-,Y \times T^n))) = 0$ by [9, 7.1]. Now Corollary 7.6 is true if we replace *X* by *Y* in it. Indeed, its proof relies on connectedness of the corresponding spaces which we have just verified and on Corollary 7.5. As a result, for every positive integer n > 0 each motivic space $C_*(Fr(-,Y \times T^n))_f$ (respectively $C_*(Fr(-,Y \times T^n))\{1/2\}_f$) is motivically fibrant in the Morel–Voevodsky [17] motivic model category of simplicial Nisnevich sheaves.

In order to show that the structure map

$$C_*(Fr(-,Y\times T^n))_f\to \Omega_{\mathbb{P}^1}(C_*(Fr(-,Y\times T^{n+1}))_f), \quad n>0$$

(respectively $C_*(Fr(-, Y \times T^n))\{1/2\}_f \to \Omega_{\mathbb{P}^1}(C_*(Fr(-, Y \times T^{n+1}))\{1/2\}_f))$ is a weak equivalence schemewise, we use Corollary 7.6 (replacing *X* by *Y* in it) and the proof of Theorem 4.1(2) (in there we also use the fact that the geometric realization of a simplicial stable local equivalence of *S*¹-spectra is a stable local equivalence) to say that this is equivalent to showing that the map

$$M_{fr}(Y \times (\mathbb{A}^1//\mathbb{G}_m)^n)_f \to \Omega_{\mathbb{G}_m^{\wedge 1}} \Omega_{S^1}(M_{fr}(Y \times (\mathbb{A}^1//\mathbb{G}_m)^n \otimes \mathbb{G}_m^{\wedge 1} \otimes S^1)_f)$$

(respectively the map $\widetilde{M}_{fr}(Y \times (\mathbb{A}^1//\mathbb{G}_m)^n)\{1/2\}_f \to \Omega_{\mathbb{G}_m^{\wedge 1}}\Omega_{S^1}(\widetilde{M}_{fr}(Y \times (\mathbb{A}^1//\mathbb{G}_m)^n \otimes \mathbb{G}_m^{\wedge 1} \otimes S^1)\{1/2\}_f))$ is a schemewise stable weak equivalence of spectra. But the latter follows from the Cancellation Theorem for framed motives [1].

Corollary 10.2. Let k be an infinite perfect field. If char $k \neq 2$ (respectively k is of any characteristic), n > 0 and $\varphi : \mathscr{X} \to \mathscr{Y}$ is a map between spaces from $\Delta^{\operatorname{op}}Fr_0(k)$ such that $\Sigma_{\mathbb{P}^1}^{\infty}(\varphi)$ is an isomorphism in SH(k) (respectively in SH(k)[1/2]) then the induced map $\varphi_* : C_*Fr(-, \mathscr{X} \times T^n) \to C_*Fr(-, \mathscr{Y} \times T^n)$ (respectively $\varphi_* : C_*Fr(-, \mathscr{X} \times T^n) \{1/2\} \to C_*Fr(-, \mathscr{Y} \times T^n)\{1/2\}$) is a local weak equivalence of motivic spaces.

In the situation when $\mathscr{X} \in \Delta^{\text{op}} Fr_0(k)$ is such that the space $C_*Fr(-,\mathscr{X})$ is locally connected we arrive at the following result:

Theorem 10.3. Let k be an infinite perfect field of characteristic different from 2. Suppose $\mathscr{X} \in \Delta^{\operatorname{op}} Fr_0(k)$ is such that the space $C_*Fr(-,\mathscr{X})$ is locally connected. Then $C_*Fr(-,\mathscr{X})$ is an \mathbb{A}^1 -local space and there is an isomorphism

$$\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X})\cong C_*Fr(-,\mathscr{X})$$

in $H_{\mathbb{A}^1}(k)$.

Proof. The fact that $C_*Fr(-,\mathscr{X})$ is an \mathbb{A}^1 -local space is proved similar to Corollary 7.5(2). The theorem would follow if we showed that the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_f$ is motivically fibrant, because the suspension spectrum $\Sigma_{\mathbb{P}^1}^{\infty}\mathscr{X}$ is stably equivalent to $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_f$ by Theorem 10.1(1). Now the fact that $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_f$ is motivically fibrant repeats the proof of Theorem 10.1(2) word for word.

The proof of the preceding theorem shows the following

Corollary 10.4. Under the assumptions of Theorem 10.3 the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})_f$ is motivically fibrant.

By [2, 3.1] $sShv_{\bullet}(Sm/k)$ has the projective motivic model structure in which generating cofibrations are given by

$$X_+ \wedge \partial \Delta^n_+ \to X_+ \wedge \Delta^n_+, \quad X \in Sm/k, \quad n \ge 0.$$

Equivalently, this family can be regarded as a family in $\Delta^{op}Fr_0(k)$ of the arrows $X \otimes \partial \Delta^n \to X \otimes \Delta^n$, $n \ge 0$.

Let $\mathscr{X} \mapsto \mathscr{X}^c$ be the cofibrant replacement functor in $sShv_{\bullet}(Sm/k)$ with respect to the projective model structure. Then $\mathscr{X}^c \in \Delta^{\operatorname{op}}Fr_0(k)$, and hence Theorem 10.1 is applicable to it. It also follows from Corollary 10.2 that each functor

$$C_*Fr(-,(-)^c \times T^n) : \mathscr{X} \in sShv_{\bullet}(Sm/k) \mapsto C_*Fr(-,\mathscr{X}^c \times T^n) \in sShv_{\bullet}(Sm/k), \quad n \ge 1,$$

(respectively $\mathscr{X} \mapsto C_*Fr(-, \mathscr{X}^c \times T^n)\{1/2\}$) takes motivic weak equivalences to local weak equivalences whenever char $k \neq 2$ (respectively k is of any characteristic). Thus we get functors

$$C_*Fr(-,(-)^c \times T^n), C_*Fr(-,(-)^c \times T^n)\{1/2\} : H_{\mathbb{A}^1}(k) \to H_{nis}(k), \quad n \ge 1,$$

where $H_{nis}(k)$ stands for the homotopy category of $sShv_{\bullet}(Sm/k)$ equipped with the local injective model structure.

Denote by $\Omega_{\mathbb{P}^1}^{\infty} \Sigma_{\mathbb{P}^1}^{\infty}(H_{\text{nis}}(k))$ the full subcategory of $H_{\text{nis}}(k)$ consisting of the infinite \mathbb{P}^1 -loop spaces. Also, denote by $\Sigma_{\mathbb{P}^1,1/2}^{\infty} : H_{\mathbb{A}^1}(k) \to SH(k)[1/2]$ the composite functor

$$H_{\mathbb{A}^1}(k) \xrightarrow{\Sigma_{\mathbb{P}^1}^{\infty}} SH(k) \to SH(k)[1/2],$$

where the right arrow is the standard localising functor inverting 2. The above arguments together with Theorem 10.1 imply the following

Theorem 10.5. *Let k be an infinite perfect field. Then the following statements are true for every field of characteristic different from 2:*

- (1) The functor $C_*Fr(-,(-)^c \times T^n)_f : H_{\mathbb{A}^1}(k) \to H_{\text{nis}}(k), n \ge 1$, lands in $\Omega^{\infty}_{\mathbb{P}^1} \Sigma^{\infty}_{\mathbb{P}^1}(H_{\text{nis}}(k))$.
- (2) For every $n \ge 1$ and $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the space $C_*Fr(-, \mathscr{X}^c \times T^n)_f$ has motivic homotopy type of $\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X} \wedge T^n)$. In particular, the functor $\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1} \circ (-\wedge T^n)$: $H_{\mathbb{A}^1}(k) \to \Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(H_{nis}(k))$ is isomorphic to the functor $\mathscr{X} \mapsto C_*Fr(-, \mathscr{X}^c \times T^n)_f$.

- (3) For every $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the space $\Omega_{\mathbb{P}^1}(C_*Fr(-,\mathscr{X}^c \times T)_f)$ has motivic homotopy type of $\Omega_{\mathbb{P}^1}^{\infty} \Sigma_{\mathbb{P}^1}^{\infty}(\mathscr{X})$. In particular, the functor $\Omega_{\mathbb{P}^1}^{\infty} \Sigma_{\mathbb{P}^1}^{\infty} : H_{\mathbb{A}^1}(k) \to \Omega_{\mathbb{P}^1}^{\infty} \Sigma_{\mathbb{P}^1}^{\infty}(H_{\text{nis}}(k))$ is isomorphic to the functor $\mathscr{X} \mapsto \Omega_{\mathbb{P}^1}(C_*Fr(-, \mathscr{X}^c \times T)_f)$.
- (4) The functor $\Sigma_{\mathbb{P}^1}^{\infty}: H_{\mathbb{A}^1}(k) \to SH(k)$ is isomorphic to the functor $\mathscr{X} \mapsto M_{\mathbb{P}^{\wedge 1}}(\mathscr{X}^c)_f$.

In turn, if k is of any characteristic, then:

- (1') The functor $C_*Fr(-,(-)^c \times T^n)\{1/2\}_f : H_{\mathbb{A}^1}(k) \to H_{\text{nis}}(k)$ lands in $\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(H_{\text{nis}}(k))$ for every $n \ge 1$.
- (2') For every $n \ge 1$ and $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the space $C_*Fr(-, \mathscr{X}^c \times T^n)\{1/2\}_f$ has motivic homotopy type of $\Omega^{\infty}_{\mathbb{P}^1}(\Sigma^{\infty}_{\mathbb{P}^1,1/2}(\mathscr{X}\wedge T^n))$. In particular, the functor $\Omega^{\infty}_{\mathbb{P}^1}(\Sigma^{\infty}_{\mathbb{P}^1,1/2}\circ(-\wedge T^n))$ $T^{n})): H_{\mathbb{A}^{1}}(k) \to \Omega^{\infty}_{\mathbb{P}^{1}}\Sigma^{\infty}_{\mathbb{P}^{1}}(H_{\text{nis}}(k)) \text{ is isomorphic to the functor } \mathscr{X} \mapsto C_{*}Fr(-,\mathscr{X}^{c} \times \mathbb{C})$ T^n){1/2}_f.
- (3') For every $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the space $\Omega_{\mathbb{P}^1}(C_*Fr(-,\mathscr{X}^c \times T)\{1/2\}_f)$ has motivic homotopy type of $\Omega^{\infty}_{\mathbb{P}^1}(\Sigma^{\infty}_{\mathbb{P}^1,1/2}(\mathscr{X}))$. In particular, the functor $\Omega^{\infty}_{\mathbb{P}^1}(\Sigma^{\infty}_{\mathbb{P}^1,1/2}): H_{\mathbb{A}^1}(k) \to \mathbb{C}^{\infty}$ $\Omega^{\infty}_{\mathbb{P}^{1}}\Sigma^{\infty}_{\mathbb{P}^{1}}(H_{\mathrm{nis}}(k)) \text{ is isomorphic to the functor } \mathscr{X} \mapsto \Omega_{\mathbb{P}^{1}}(C_{*}Fr(-,\mathscr{X}^{c} \times T)\{1/2\}_{f}).$ $(4') \Sigma^{\infty}_{\mathbb{P}^{1},1/2} : H_{\mathbb{A}^{1}}(k) \to SH(k)[1/2] \text{ is isomorphic to the functor } \mathscr{X} \mapsto M_{\mathbb{P}^{\Lambda}}(\mathscr{X}^{c})\{1/2\}_{f}.$

Corollary 10.6. Let k be an infinite perfect field. If char $k \neq 2$ (respectively k is of any characteristic) and n > 0 then for any map of motivic spaces $\varphi : \mathscr{Y} \to \mathscr{Z}$ such that $\Sigma_{\mathbb{D}^1}^{\infty}(\varphi)$ is an isomorphism in SH(k) (respectively in SH(k)[1/2]) the induced map $\varphi_* : C_*Fr(-, \mathscr{Y}^c \times T^n) \to \mathcal{Y}^c$ $C_*Fr(-,\mathscr{Z}^c \times T^n)$ (respectively $\varphi_* : C_*Fr(-,\mathscr{Y}^c \times T^n)\{1/2\} \to C_*Fr(-,\mathscr{Z}^c \times T^n)\{1/2\}$ is a local weak equivalence.

Remark 10.7. In fact, assertion (3) of Theorem 10.5 is true for any local group-like completion of the motivic space $C_*Fr(-,\mathscr{X}^c)_f$, functorial in \mathscr{X} . It follows from the Additivity Theorem that the π_0 -sheaf of $C_*Fr(-,\mathscr{X}^c)$ is a sheaf of Abelian monoids. One of such local group-like completion functors is given in the assertion. Another local group-like completion is given by $\mathscr{X} \mapsto \Omega_{S^1}(C_*Fr(-,\mathscr{X}^c \otimes S^1)_f)$. In particular, there is a motivic weak equivalence of spaces

$$\Omega^{\infty}_{\mathbb{P}^1} \Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X}) \simeq \Omega_{S^1}(C_* Fr(-, \mathscr{X}^c \otimes S^1)_f)$$

for any $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ and any infinite perfect field with char $k \neq 2$.

In turn, if $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ is such that the space $C_*Fr(-,\mathscr{X}^c)$ is locally connected we get the following result:

Theorem 10.8. Let k be an infinite perfect field of characteristic different from 2 and let $\mathscr{X} \in$ $sShv_{\bullet}(Sm/k)$ be such that the space $C_*Fr(-,\mathscr{X}^c)$ is locally connected. Then $C_*Fr(-,\mathscr{X}^c)$ is an \mathbb{A}^1 -local space and there is an isomorphism

$$\Omega^{\infty}_{\mathbb{P}^1}\Sigma^{\infty}_{\mathbb{P}^1}(\mathscr{X})\cong C_*Fr(-,\mathscr{X}^c)$$

in $H_{\mathbb{A}^1}(k)$.

Proof. Since $\mathscr{X}^c \in \Delta^{\operatorname{op}} Fr_0(k)$, the theorem follows from Theorem 10.3.

The proof of the preceding theorem shows the following

Corollary 10.9. Under the assumptions of Theorem 10.8 the \mathbb{P}^1 -spectrum $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X}^c)_f$ is motivically fibrant. Moreover, $\Sigma_{\mathbb{P}^1}^{\infty} \mathscr{X}$ is isomorphic to $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X}^c)_f$ in SH(k).

11. FURTHER APPLICATIONS OF FRAMED MOTIVES

Having applied the macinery of framed motives to prove Theorem 4.1, we want to give further applications. One of the applications computes the suspension bispectrum $\sum_{S^1}^{\infty} \sum_{G}^{\infty} X_+$ of a smooth algebraic variety X in terms of twisted framed motives of X. Another important application will be purely topological. It will compute the classical sphere spectrum $\sum_{S^1}^{\infty} S^0$ as the framed motive $M_{fr}(pt)(pt)$ of the point $pt = \operatorname{Spec} k$ evaluated at the point whenever the base field k is algebraically closed of characteristic zero.

Denote by \mathbb{G} the cone $(\mathbb{G}_m)_+//pt_+$ of the embedding $pt_+ \stackrel{1}{\hookrightarrow} (\mathbb{G}_m)_+$ in the category of pointed simplicial presheaves $sPre_{\bullet}(Sm/k)$. It is termwise equal to

$$(-, \mathbb{G}_m)_+, (-, \mathbb{G}_m)_+ \lor (-, pt)_+, (-, \mathbb{G}_m)_+ \lor (-, pt)_+ \lor (-, pt)_+, \dots$$

Moreover, its sheafification equals $(\mathbb{G}_m^{\wedge 1})_+$, which is termwise equal to

$$(-,\mathbb{G}_m)_+,(-,\mathbb{G}_m\sqcup pt)_+,(-,\mathbb{G}_m\sqcup pt\sqcup pt)_+,\ldots$$

The sheafification is represented in the category $\Delta^{op}(Fr_0(k))$ by the object $\mathbb{G}_m^{\wedge 1}$ (see Notation 8.1), which is termwise equal to

$$\mathbb{G}_m, \mathbb{G}_m \sqcup pt, \mathbb{G}_m \sqcup pt \sqcup pt, \ldots$$

One of the models for Morel–Voevodsky's SH(k) can be defined in terms of (S^1, \mathbb{G}) -bispectra (see, e.g., [14]). The main (S^1, \mathbb{G}) -bispectrum we work with is given by the sequence of framed motives $M_{fr}^{\mathbb{G}}(X) = (M_{fr}(X), M_{fr}(X \times \mathbb{G}_m^{\wedge 1}), M_{fr}(X \times \mathbb{G}_m^{\wedge 2}), \ldots), \quad X \in Sm/k$, where simplicial objects $\mathbb{G}_m^{\wedge n} \in \Delta^{\operatorname{op}}(Fr_0(k))$ are defined in Notation 8.1. Each structure map is defined as the composition

$$M_{fr}(X \times \mathbb{G}_m^{\wedge n}) \to \underline{\mathrm{Hom}}((\mathbb{G}_m^{\wedge 1})_+, M_{fr}(X \times \mathbb{G}_m^{\wedge n+1})) \to \underline{\mathrm{Hom}}(\mathbb{G}, M_{fr}(X \times \mathbb{G}_m^{\wedge n+1})),$$

where the left map is defined as (15). We shall also call $M_{fr}(X \times \mathbb{G}_m^{\wedge n})$ the *n*-twisted framed motive of X, and write $M_{fr}(X)(n)$ to denote $M_{fr}(X \times \mathbb{G}_m^{\wedge n})$. So we can write for brevity

$$M_{fr}^{\mathbb{G}}(X) = (M_{fr}(X), M_{fr}(X)(1), M_{fr}(X)(2), \ldots),$$

to denote the (S^1, \mathbb{G}) -bispectrum $M_{fr}^{\mathbb{G}}(X)$.

In turn, we can similarly define a bispectrum

$$\widetilde{M}_{fr}^{\mathbb{G}}(X)\{1/2\} = (\widetilde{M}_{fr}(X)\{1/2\}, \widetilde{M}_{fr}(X \times \mathbb{G}_m^{\wedge 1})\{1/2\}, \widetilde{M}_{fr}(X \times \mathbb{G}_m^{\wedge 2})\{1/2\}, \ldots), \quad X \in Sm/k.$$

It is worth to mention that the bispectrum $M_{fr}^{\mathbb{G}}(X)$ is constructed in the same way as the bispectrum

$$M_{\mathbb{K}}^{\mathbb{G}}(X) = (M_{\mathbb{K}}(X), M_{\mathbb{K}}(X)(1), M_{\mathbb{K}}(X)(2), \ldots), \quad X \in Sm/k,$$

where each S^1 -spectrum $M_{\mathbb{K}}(X)(n) = M_{\mathbb{K}}(X \times \mathbb{G}_m^{\wedge n})$ is the *n*-twisted *K*-motive of *X* in the sense of [6]. It was shown in [6] that $M_{\mathbb{K}}^{\mathbb{G}}(pt)$ represents the bispectrum $f_0(KGL)$.

The main result of this section is as follows.

Theorem 11.1. Let k be an infinite perfect field. If char $k \neq 2$ (respectively k is of any characteristic) then for any $X \in Sm/k$, the canonical map of bispectra $\sum_{S^1}^{\infty} \Sigma_{\mathbb{G}}^{\infty} X_+ \to M_{fr}^{\mathbb{G}}(X)$ (respectively $\sum_{S^1}^{\infty} \Sigma_{\mathbb{G}}^{\infty} X_+ \to \widetilde{M}_{fr}^{\mathbb{G}}(X)\{1/2\}$) is a stable motivic weak equivalence (respectively 2⁻¹-stable motivic weak equivalence). *Proof.* If char $k \neq 2$ (respectively k is of any characteristic) then it is enough to prove that on bigraded presheaves $\pi_{*,*}^{\mathbb{A}^1}(\gamma)$ (respectively $\pi_{*,*}^{\mathbb{A}^1}(\gamma) \otimes \mathbb{Z}[1/2]$) is an isomorphism. We shall prove the theorem for the case when char $k \neq 2$, because the other case is proved in a similar fashion. Every bispectrum yields a $S^1 \wedge \mathbb{G}$ -spectrum by taking diagonal. In order to avoid massive notation, we prove the theorem for the case X = pt. The same proof works for any $X \in Sm/k$. Let $r \ge 0$, $s, n \ge 1$ be integers with $s + n \ge 0$. There is a commutative diagram in the homotopy category $H_{\mathbb{A}^1}(k)$ of pointed motivic spaces

$$\begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, (S^{1} \wedge \mathbb{G})^{n} \end{bmatrix} \xrightarrow{(1_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, C_{*}Fr((S^{1} \otimes \mathbb{G}_{m}^{\wedge 1})^{\wedge n}) \end{bmatrix} \\ \uparrow (u^{\wedge n})_{*} & u_{*}^{*} \uparrow \\ \begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, (\mathbb{A}_{+}^{1}//(\mathbb{G}_{m})_{+})^{n} \end{bmatrix} \xrightarrow{(2_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, C_{*}Fr((\mathbb{A}^{1}//\mathbb{G}_{m})^{\wedge n}) \end{bmatrix} \\ \downarrow (v^{\wedge n})_{*} & v_{*}^{*} \downarrow \\ \begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, T^{n} \end{bmatrix} \xrightarrow{(3_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge (S^{1} \wedge \mathbb{G})^{s+n}, C_{*}Fr(T^{n}) \end{bmatrix} \\ \downarrow (u^{\wedge (s+n)})^{*} & (u^{\wedge (s+n)})^{*} \downarrow \\ \begin{bmatrix} S^{r} \wedge U_{+} \wedge (\mathbb{A}_{+}^{1}//(\mathbb{G}_{m})_{+})^{s+n}, T^{n} \end{bmatrix} \xrightarrow{(4_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge (\mathbb{A}_{+}^{1}//(\mathbb{G}_{m})_{+})^{s+n}, C_{*}Fr(T^{n}) \end{bmatrix} \\ \uparrow (v^{\wedge (s+n)})^{*} & (v^{\wedge (s+n)})^{*} \uparrow \\ \begin{bmatrix} S^{r} \wedge U_{+} \wedge T^{s+n}, T^{n} \end{bmatrix} \xrightarrow{(5_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge T^{s+n}, C_{*}Fr(T^{n}) \end{bmatrix} \\ \downarrow (\sigma^{\wedge (s+n)})^{*} & (\sigma^{\wedge (s+n)})^{*} \downarrow \\ \begin{bmatrix} S^{r} \wedge U_{+} \wedge \mathbb{P}^{\wedge s+n}, T^{n} \end{bmatrix} \xrightarrow{(6_{n})} \begin{bmatrix} S^{r} \wedge U_{+} \wedge \mathbb{P}^{\wedge s+n}, C_{*}Fr(T^{n}) \end{bmatrix}$$

In this diagram all left vertical arrows are bijections, since the natural morphisms $u : \mathbb{A}^1_+ / / (\mathbb{G}_m)_+ \to S^1 \wedge \mathbb{G}, v : \mathbb{A}^1_+ / / (\mathbb{G}_m)_+ \to T$ and $\sigma : \mathbb{P}^{\wedge 1} \to T$ are motivic equivalences. All right vertical arrows are bijections, since the morphisms

$$C_*Fr((\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \xrightarrow{u_*^n} C_*Fr((S^1 \otimes \mathbb{G}_m^{\wedge 1})^{\wedge n}), \ C_*Fr((\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}) \xrightarrow{v_*^n} C_*Fr(T^n)$$

are local equivalences by Corollaries 8.11 and 9.4.

Fit now each of twelve vertices of the diagram into a direct colimit over *n* as follows. For vertices on the right hand side we form direct colimits with respect to the *T*-spectum with entries $\{C_*Fr(T^n)\}$, with respect to the $\mathbb{A}^1_+//(\mathbb{G}_m)_+$ -spectrum with entries $\{C_*Fr((\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})\}$ as well as with respect to the $S^1 \wedge \mathbb{G}$ -spectrum with entries $\{C_*Fr((S^1 \otimes \mathbb{G}_m^{\wedge 1})^{\wedge n})\}$. Likewise, for vertices on the right hand side we form direct colimits with respect to the *T*-spectum with entries $\{T^n\}$, with respect to the $\mathbb{A}^1_+//(\mathbb{G}_m)_+$ -spectum with entries $\{(\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}\}$, with respect to the $\mathbb{A}^1_+//(\mathbb{G}_m)_+$ -spectum with entries $\{(\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}\}$, with respect to the $S^1 \wedge \mathbb{G}$ -spectrum with entries $\{(S^1 \wedge \mathbb{G})^{\wedge n}\}$.

The family of morphisms $\{(6_n)\}$ forms a morphism on the direct colimit, since it corresponds to the *T*-spectra morphism $\{T^n\} \to \{C_*Fr(T^n)\}$. For i = 5,4,3 the families of morphisms $\{(i_n)\}$ form morphisms on the direct colimits by the same reason. The family of morphisms $\{(1_n)\}$ forms a morphism on the direct colimit, since it corresponds to the $S^1 \wedge \mathbb{G}$ -spectra morphism $\{(S^1 \wedge \mathbb{G})^{\wedge n}\} \to \{C_*Fr((S^1 \otimes \mathbb{G}_m^{\wedge 1})^{\wedge n})\}$. Finally, the family of morphisms $\{(2_n)\}$ forms a morphism on the direct colimit, since it corresponds to the $\mathbb{A}^1_+//(\mathbb{G}_m)_+$ -spectra morphism $\{(\mathbb{A}^1//\mathbb{G}_m)^{\wedge n}\} \to \{C_*Fr((\mathbb{A}^1//\mathbb{G}_m)^{\wedge n})\}$. In a similar fashion for each vertical map the corresponding family of arrows forms a morphism on the direct colimits. In this way we get a commutative diagram consisting of twelve direct colimits and morphisms between them. We also get a commutative diagram consisting of twelve groups and homomorphisms between them. In that diagram of groups all the vertical arrows are isomorphisms as mentioned above. The bottom arrow is an isomorphism by Theorem 4.1, hence so is the top arrow. We conclude that for $r \ge 0$ the map of presheaves $\pi_{r+2s,s}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0)) \xrightarrow{\gamma_*} \pi_{r+2s,s}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))$ is an isomorphism for any integer *s*. In another words, the map γ_* is an isomorphisms on presheaves $\pi_{a,b}^{\mathbb{A}^1}$ with $2a - b \ge 0$. In particular, for any $U \in Sm/k$ and any t > 0 the map

$$\pi_{2a,a}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0))(U\times\mathbb{G}_m^{\times t})\xrightarrow{\gamma_*}\pi_{2a,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U\times\mathbb{G}_m^{\times t})$$

is an isomorphism. Note that $\pi_{2a,a}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0))(U_+ \wedge \mathbb{G}_m^{\wedge t})$ is a canonical direct summand of the group $\pi_{2a,a}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0))(U \times \mathbb{G}_m^{\times t})$, and the group $\pi_{2a,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U \wedge \mathbb{G}_m^{\wedge t})$ is a canonical direct summand of the group $\pi_{2a,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U \times \mathbb{G}_m^{\times t})$. Hence the map

$$\pi_{2a-t,a}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0))(U) = \pi_{2a,a}^{\mathbb{A}^1}(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^1}^{\infty}(S^0))(U_+ \wedge \mathbb{G}_m^{\wedge t}) \xrightarrow{\gamma_*} \pi_{2a,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U_+ \wedge \mathbb{G}_m^{\wedge t}) = \pi_{2a-t,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U) \xrightarrow{\gamma_*} \pi_{2a,a}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt))(U_+ \wedge \mathbb{G}_m^{\wedge t}) \xrightarrow{\gamma_*} \pi_{2a,a}^{\mathbb{G}^1}(M_{fr}^{\mathbb{G}}(pt))(U_+ \wedge \mathbb{G}_m^{\mathbb{G}^1}(pt))(U_+ \wedge \mathbb{G}_m^{\mathbb{G}^1$$

is an isomorphism too. Thus, the map γ_* is an isomorphism on presheaves $\pi_{a,b}^{\mathbb{A}^1}$ with 2a - b < 0, whence the theorem.

Corollary 11.2. Let k be an infinite perfect field and X be smooth. If char $k \neq 2$ (respectively k is of any characteristic) then $\pi_{-n,-n}^{\mathbb{A}^1}((\Sigma_{S^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}X_+)(pt))$ (respectively $\pi_{-n,-n}^{\mathbb{A}^1}((\Sigma_{S^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}X_+)(pt)) \otimes \mathbb{Z}[1/2]$), $n \geq 0$, is the Grothendieck group of the commutative monoid $\pi_0(C_*Fr(pt, X \times \mathbb{G}_m^{\wedge n}))$ (respectively the Abelian group $\pi_0(C_*Fr(pt, X \times \mathbb{G}_m^{\wedge n})\{1/2\})$).

Corollary 11.3. Let k be an infinite perfect field and let X be smooth. If char $k \neq 2$ (respectively k is of any characteristic) then $\pi_{-n,-n}^{\mathbb{A}^1}(\Sigma_{\mathbb{S}^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}X_+)(pt)$

$$\pi_{-n,-n}^{\mathbb{A}^1}(\Sigma^{\infty}_{\mathbb{S}^1}\Sigma^{\infty}_{\mathbb{G}}X_+)(pt) = H_0(\mathbb{Z}F(\Delta^{\bullet}_k, X\times \mathbb{G}_m^{\wedge n})), \quad n \ge 0$$

(respectively $\pi_{-n,-n}^{\mathbb{A}^1}(\Sigma_{S^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}X_+)(pt)\otimes\mathbb{Z}[1/2] = H_0(\mathbb{Z}F(\Delta_k^{\bullet}, X\times\mathbb{G}_m^{\wedge n}))\otimes\mathbb{Z}[1/2])$. In particular, $\pi_{-n,-n}^{\mathbb{A}^1}(\Sigma_{S^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}S^0)(pt) = H_0(\mathbb{Z}F(\Delta_k^{\bullet}, \mathbb{G}_m^{\wedge n})) = K_n^{MW}(k)$ if $n \ge 0$ and char k = 0.

Proof. The fact that $H_0(\mathbb{Z}F(\Delta_k^{\bullet}, \mathbb{G}_m^{\wedge n})) = K_n^{MW}(k), n \ge 0$, has been proved by Neshitov in [18] for fields of characteristic zero. Thus the statement recovers the celebrated theorem of Morel [16] for Milnor–Witt *K*-theory.

It is also useful to have Theorem 11.1 for simplicial schemes or, more generally, for objects in $\Delta^{op} Fr_0(k)$ (cf. Theorem 10.1).

Theorem 11.4. Let k be an infinite perfect field. If char $k \neq 2$ (respectively k is of any characteristic) then for any $Y \in \Delta^{\operatorname{op}} Fr_0(k)$, the canonical map of bispectra $\Sigma^{\infty}_{S^1} \Sigma^{\infty}_{\mathbb{G}} Y_+ \to M^{\mathbb{G}}_{fr}(Y)$ (respectively $\Sigma^{\infty}_{S^1} \Sigma^{\infty}_{\mathbb{G}} Y_+ \to \widetilde{M}^{\mathbb{G}}_{fr}(Y)\{1/2\}$) is a stable motivic weak equivalence (respectively 2⁻¹-stable motivic weak equivalence).

Proof. If we use Theorem 10.1, our proof repeats that of Theorem 11.1 word for word. \Box

Here is an application of the preceding theorem.

Theorem 11.5. Suppose the base field k is perfect infinite with char $k \neq 2$ (respectively k is of any characteristic). For any $Y \in \Delta^{\operatorname{op}} Fr_0(k)$ one has a canonical isomorphism

$$SH(k)(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}X_{+},\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}Y_{+}[n]) = SH_{S^{1}}^{\operatorname{nis}}(k)(\Sigma_{S^{1}}^{\infty}X_{+},M_{fr}(Y)[n]), \quad n \ge 0.$$

(respectively $SH(k)[1/2](\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}X_{+}, \Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}Y_{+}[n]) = SH_{S^{1}}^{\text{nis}}(k)(\Sigma_{S^{1}}^{\infty}X_{+}, M_{fr}(Y)\{1/2\}[n]))$, where $SH_{S^{1}}^{\text{nis}}(k)$ is the ordinary stable homotopy category of Nisnevich sheaves of S^{1} -spectra.

Proof. Suppose the base field k is perfect infinite with char $k \neq 2$. Consider a bispectrum

$$M_{fr}^{\mathbb{G}}(Y)_f = (M_{fr}(Y)_f, M_{fr}(Y \times \mathbb{G}_m^{\wedge 1})_f, M_{fr}(Y \times \mathbb{G}_m^{\wedge 2})_f, \ldots)$$

obtained from $M_{fr}^{\mathbb{G}}(Y)$ by taking Nisnevich local stable fibrant replacements at each level. It is shown similar to [1, Theorem B] that $M_{fr}^{\mathbb{G}}(Y)_f$ is a motivically fibrant bispectrum.

Theorem 11.5 implies an isomorphism

$$SH(k)(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}X_{+},\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}Y_{+}[n]) = SH(k)(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}X_{+},M_{fr}^{\mathbb{G}}(Y)_{f}[n]), \quad n \ge 0$$

But

$$SH(k)(\Sigma_{\mathbb{G}}^{\infty}\Sigma_{S^{1}}^{\infty}X_{+}, M_{fr}^{\mathbb{G}}(Y)_{f}[n]) \cong SH_{S^{1}}(k)(\Sigma_{S^{1}}^{\infty}X_{+}, M_{fr}(Y)_{f}[n]) \cong$$
$$SH_{S^{1}}^{\operatorname{nis}}(k)(\Sigma_{S^{1}}^{\infty}X_{+}, M_{fr}(Y)_{f}[n]) \cong SH_{S^{1}}^{\operatorname{nis}}(k)(\Sigma_{S^{1}}^{\infty}X_{+}, M_{fr}(Y)[n]).$$

If k is of any characteristic then we can likewise show an isomorphism

$$SH(k)[1/2](\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}X_+,\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}Y_+[n])=SH^{\operatorname{nis}}_{S^1}(k)(\Sigma^{\infty}_{S^1}X_+,M_{fr}(Y)\{1/2\}[n]).$$

It remains to observe that $M_{fr}(Y)\{1/2\}$ is stably equivalent to $\widetilde{M}_{fr}(Y)\{1/2\}$.

Corollary 11.6. Suppose the base field k is perfect infinite with char $k \neq 2$ (respectively k is of any characteristic). For any morphism $\varphi: Y \to Z$ in $\Delta^{\operatorname{op}}Fr_0(k)$ such that $\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}(\varphi)$ is an isomorphism in SH(k) (respectively in SH(k)[1/2]), the morphism of framed motives $M_{fr}(\varphi)$: $M_{fr}(Y) \to M_{fr}(Z)$ (respectively $M_{fr}(\varphi)\{1/2\}: M_{fr}(Y)\{1/2\} \to M_{fr}(Z)\{1/2\}$) is a local stable equivalence of S^1 -spectra.

Let $\mathscr{X} \mapsto \mathscr{X}^c$ be the cofibrant replacement functor in $sShv_{\bullet}(Sm/k)$ (see p. 36). Then \mathscr{X}^c is in $\Delta^{\operatorname{op}}Fr_0(k)$, and hence Theorems 11.4-11.5 are applicable to it. It also follows from Corollary 11.6 that each functor

$$M_{fr}((-)^c): \mathscr{X} \in sShv_{\bullet}(Sm/k) \mapsto M_{fr}(\mathscr{X}^c) \in Sp_{S^1}(sShv_{\bullet}(Sm/k))$$

(respectively $\mathscr{X} \mapsto M_{fr}(\mathscr{X}^c)\{1/2\}$) takes motivic weak equivalences to stable local weak equivalences whenever char $k \neq 2$ (respectively k is of any characteristic). Thus we get functors

$$M_{fr}((-)^c), M_{fr}((-)^c)\{1/2\} : H_{\mathbb{A}^1}(k) \to SH_{S^1}^{nis}(k)$$

where $SH_{S^1}^{nis}(k)$ stands for the homotopy category of $Sp_{S^1}(sShv_{\bullet}(Sm/k))$ equipped with the stable local injective model structure.

In a similar fashion we define functors

$$M_{fr}^{\mathbb{G}}((-)^{c}), M_{fr}^{\mathbb{G}}((-)^{c})\{1/2\} : H_{\mathbb{A}^{1}}(k) \to SH(k).$$

Explicitly, they take a motivic space \mathscr{X} to bispectra $M_{fr}^{\mathbb{G}}(\mathscr{X}^c), M_{fr}^{\mathbb{G}}(\mathscr{X}^c) \{1/2\}$. In fact, we shall extend both functors to SH(k) in Section 12.

Denote by $\Omega^{\infty}_{\mathbb{G}}(SH^{\operatorname{nis}}_{S^1}(k))$ the full subcategory of $SH^{\operatorname{nis}}_{S^1}(k)$ consisting of the infinite \mathbb{G} -loop spectra. Also, denote by $\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1,1/2}: H_{\mathbb{A}^1}(k) \to SH(k)[1/2]$ the composite functor

$$H_{\mathbb{A}^1}(k) \xrightarrow{\Sigma_{\mathbb{G}}^{\infty} \Sigma_{S^1}^{\infty}} SH(k) \to SH(k)[1/2],$$

where the right arrow is the standard localising functor inverting 2. The above arguments together with Theorems 11.4-11.5 and Corollary 11.6 imply the following **Theorem 11.7.** Let k be an infinite perfect field. Then the following statements are true for every field of characteristic different from 2:

- (1) The functor $M_{fr}((-)^c)_f : H_{\mathbb{A}^1}(k) \to SH_{S^1}^{nis}(k)$ lands in $\Omega^{\infty}_{\mathbb{G}}(SH_{S^1}^{nis}(k))$, where "f" refers to the stable local fibrant replacement of S^1 -spectra.
- (2) For every $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the spectrum $M_{fr}(\mathscr{X}^c)_f$ has stable motivic homotopy type of $\Omega^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{\mathbb{S}^1}(\mathscr{X})$. In particular, the functor $\Omega^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{\mathbb{S}^1}: H_{\mathbb{A}^1}(k) \to \Omega^{\infty}_{\mathbb{G}}(SH^{\operatorname{nis}}_{S^1}(k))$ is isomorphic to the functor $\mathscr{X} \mapsto M_{fr}(\mathscr{X}^c)_f$.
- (3) The functor $\Sigma^{\infty}_{\mathbb{G}_{r}}\Sigma^{\infty}_{S^{1}}: H_{\mathbb{A}^{1}}(k) \to SH(k)$ is isomorphic to the functor $\mathscr{X} \mapsto M^{\mathbb{G}}_{fr}(\mathscr{X}^{c})$.

In turn, if k is of any characteristic, then:

- (1') The functor $M_{fr}((-)^c)\{1/2\}_f: H_{\mathbb{A}^1}(k) \to SH_{S^1}^{nis}(k)$ lands in $\Omega^{\infty}_{\mathbb{G}}(SH_{S^1}^{nis}(k))$, where "f" refers to the stable local fibrant replacement of S^1 -spectra.
- (2') For every $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ the spectrum $M_{fr}(\mathscr{X}^{c})\{1/2\}_{f}$ has stable motivic homotopy type of $\Omega^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{S^{1},1/2}(\mathscr{X})$. In particular, the functor $\Omega^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{\mathbb{G}} \Sigma^{\infty}_{S^{1},1/2} : H_{\mathbb{A}^{1}}(k) \to \mathbb{C}^{1}_{\mathbb{G}}$ $\Omega^{\infty}_{\mathbb{G}}(SH^{\operatorname{nis}}_{S^{1}}(k)) \text{ is isomorphic to the functor } \mathscr{X} \mapsto M_{fr}(\mathscr{X}^{c})\{1/2\}_{f}.$ (3') $\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^{1},1/2} : H_{\mathbb{A}^{1}}(k) \to SH(k) \text{ is isomorphic to the functor } \mathscr{X} \mapsto M^{\mathbb{G}}_{fr}(\mathscr{X}^{c})\{1/2\}.$

Corollary 11.8. Let k be an infinite perfect field. If char $k \neq 2$ (respectively k is of any characteristic) then for any map of motivic spaces $\varphi : \mathscr{Y} \to \mathscr{Z}$ such that $\Sigma_{S^1}^{\infty} \Sigma_{\mathbb{G}}^{\infty}(\varphi)$ is an isomorphism in SH(k) (respectively in SH(k)[1/2]) the induced map $\varphi_* : M_{fr}(\mathscr{Y}^c) \to M_{fr}(\mathscr{Z}^c)$ (respectively $\varphi_*: M_{fr}(\mathscr{Y}^c)\{1/2\} \to M_{fr}(\mathscr{Z}^c)\{1/2\})$ is a local stable equivalence.

We finish the section with a topological application of framed motives. It gives an explicit model for the classical sphere spectrum $\sum_{s_1}^{\infty} S^0$.

Theorem 11.9. Let k be an algebraically closed field of characteristic zero, with embedding $k \hookrightarrow \mathbb{C}$. Then the framed motive $M_{fr}(pt)(pt)$ of the point $pt = \operatorname{Spec} k$ evaluated at pt has stable homotopy type of the classical sphere spectrum $\sum_{s1}^{\infty} S^0$. In particular, $C_*Fr(pt, pt \otimes S^1)$ has homotopy type of $\Omega_{S^1}^{\infty} \Sigma_{S^1}^{\infty} S^1$.

Proof. By a theorem of Levine [15] the functor $c: SH \to SH(k)$, induced by the functor

$$sSets_{\bullet} \rightarrow sPre_{\bullet}(Sm/k)$$

sending a pointed simplicial set to the constant presheaf on Sm/k, is fully faithful. The functor c comes from a left Quillen functor (see the proof of [15, 6.5]). Its right Quillen functor from bispectra to ordinary S¹-specra takes a bispectrum $E = (E_0, E_1, ...)$ to $E_0(pt)$. Moreover, c induces an isomorphism $\pi_n(E) \to \pi_{n,0}^{\mathbb{A}^1}(c(E))$ for all spectra *E*.

Consider $M_{fr}^{\mathbb{G}}(pt)$. By Theorem 11.1 the canonical morphism

$$\Sigma^{\infty}_{S^1} \Sigma^{\infty}_{\mathbb{G}} S^0 = c(\Sigma^{\infty}_{S^1} S^0) \to M^{\mathbb{G}}_{fr}(pt)$$

is a motivic stable equivalence of bispectra. Consider a bispectrum

$$M_{fr}^{\mathbb{G}}(pt)_f = (M_{fr}(pt)_f, M_{fr}(\mathbb{G}_m^{\wedge 1})_f, M_{fr}(\mathbb{G}_m^{\wedge 2})_f, \dots)$$

obtained from $M_{fr}^{\mathbb{G}}(pt)$ by taking Nisnevich local stable fibrant replacements at each level. By [1, Theorem B] $M_{fr}^{\mathbb{G}}(pt)_f$ is a motivically fibrant bispectrum, and hence a fibrant replacement of $\Sigma_{S^1}^{\infty} \Sigma_{\mathbb{G}}^{\infty} S^0.$

It follows that $M_{fr}(pt)_f(pt)$ is a fibrant replacement of $\Sigma_{S^1}^{\infty}S^0$, because each homomorphism $\pi_n(\Sigma_{S^1}^{\infty}S^0) \to \pi_{n,0}^{\mathbb{A}^1}(c(\Sigma_{S^1}^{\infty}S^0)) \xrightarrow{\cong} \pi_{n,0}^{\mathbb{A}^1}(M_{fr}^{\mathbb{G}}(pt)_f) = \pi_n(M_{fr}(pt)_f(pt))$ is an isomorphism. It remains to observe that $M_{fr}(pt)_f(pt)$ is stably equivalent to $M_{fr}(pt)(pt)$.

12. The BIG FRAMED MOTIVE FUNCTOR \mathcal{M}_{fr}^{b}

In Section 11 we have computed the functor $\Sigma_{S^1}^{\infty}\Sigma_{\mathbb{G}}^{\infty}: H_{\mathbb{A}^1}(k) \to SH(k)$ as the functor $M_{fr}^{\mathbb{G}}: H_{\mathbb{A}^1}(k) \to SH(k)$. We extend the latter functor to bispectra below, but first we start with preparations. We assume in this section that the base field k is infinite perfect with char $k \neq 2$.

We first give an explicit description of $Ex^{\infty}(C_*\mathscr{F}r(\mathscr{X}))$ for every space $\mathscr{X} \in sShv_{\bullet}(Sm/k)$, where Ex^{∞} is the Kan complex. Voevodsky [21, Section 3] defined a realization functor from simplicial sets to Nisnevich sheaves $|-|: sSets \to Shv_{nis}(Sm/k)$ such that $|\Delta[n]| = \Delta^n$, where $\Delta[n]$ is the standard *n*-simplex. Under this notation the cosimplicial scheme Δ^{\bullet} equals $|\Delta[\bullet]|$. For every $\ell \ge 0$ denote by $sd^{\ell}\Delta^{\bullet}$ the cosimplicial Nisnevich sheaf $|sd^{\ell}\Delta[\bullet]|$. Under this notation we then have a canonical isomorphism of motivic spaces

$$Ex^{\ell}(C_*\mathscr{F}r(\mathscr{X})) = \mathscr{F}r(|sd^{\ell}\Delta[\bullet]|_+ \wedge -, \mathscr{X}).$$

It follows that $Ex^{\ell}(C_*\mathscr{F}r(\mathscr{X}))$ is a space with framed correspondences as well as the space

$$Ex^{\infty}(C_*\mathscr{F}r(\mathscr{X})) = \operatorname{colim}_{\ell} Ex^{\ell}(C_*\mathscr{F}r(\mathscr{X})).$$

Then $Ex^{\infty}(C_*\mathscr{F}r(\mathscr{X}))$ is a sectionwise fibrant pointed simplicial set and the canonical map $C_*\mathscr{F}r(\mathscr{X}) \to Ex^{\infty}(C_*\mathscr{F}r(\mathscr{X}))$ is a sectionwise weak equivalence.

For brevity we denote by **G** the sheafification of \mathbb{G} . It equals the simplicial sheaf $(\mathbb{G}_m^{\wedge 1})_+ \in sShv_{\bullet}(Sm/k)$ (see Section 11). Given a (S^1, \mathbf{G}) -bispectrum *E* of simplicial Nisnevich sheaves, replace it by a stably cofibrant bispectrum E^c in the stable projective motivic model structure of bispectra associated with the projective motivic structure on $sShv_{\bullet}(Sm/k)$ in the sense of [2] (we shall with this model structure throughout the section). Then each (i, j)-entry $E_{i,j}^c$ of E^c belongs

to
$$\Delta^{\text{op}} Fr_0(k)$$
. We set

$$M_{fr}^{\mathbb{G}}(E)_{i,j} := Ex^{\infty}(C_*Fr(E_{i,j}^c)), \quad i, j \ge 0$$

The structure maps in the S^1 - and **G**-direction

$$Ex^{\infty}(C_*Fr(E_{i,j}^c)) \to \underline{\operatorname{Hom}}(S^1, Ex^{\infty}(C_*Fr(E_{i+1,j}^c))), Ex^{\infty}(C_*Fr(E_{i,j}^c)) \to \underline{\operatorname{Hom}}(\mathbf{G}, Ex^{\infty}(C_*Fr(E_{i,j+1}^c))))$$

are obviously induced by the structure maps u_v , u_h of E^c (see [14, p. 488] for the relevant definitions on bispectra). Precisely, they are compositions

$$Ex^{\infty}(C_*Fr(E^c_{i,j})) \to \underline{\operatorname{Hom}}(S^1, Ex^{\infty}(C_*Fr(E^c_{i,j} \otimes S^1))) \xrightarrow{u_{v_{\tau}}} \underline{\operatorname{Hom}}(S^1, Ex^{\infty}(C_*Fr(E^c_{i+1,j})))$$

and

$$Ex^{\infty}(C_*Fr(E_{i,j}^c)) \to \underline{\operatorname{Hom}}(\mathbf{G}, Ex^{\infty}(C_*Fr(E_{i,j}^c \otimes \mathbb{G}_m^{\wedge 1}))) \xrightarrow{u_h} \underline{\operatorname{Hom}}(\mathbf{G}, Ex^{\infty}(C_*Fr(E_{i,j+1}^c)))$$

respectively.

For brevity we drop Ex^{∞} from notation and tacitly assume below that all spaces like $C_*Fr(E_{i,j}^c)$ are sectionwise fibrant with framed correspondences. We then have a canonical morphism of bispectra

$$\zeta: E^c \to M^{\mathbb{G}}_{fr}(E).$$

Clearly, ζ is functorial in *E*.

Denote by $SH^{fr}(k)$ the full subcategory of SH(k) consisting of framed bispectra, i.e. those bispectra \mathscr{E} such that each space $\mathscr{E}_{i,j}$ is a space with framed correspondences and the structure maps $\mathscr{E}_{i,j} \to \underline{\operatorname{Hom}}(S^1, \mathscr{E}_{i+1,j}), \mathscr{E}_{i,j} \to \underline{\operatorname{Hom}}(\mathbf{G}, \mathscr{E}_{i,j+1})$ preserve framed correspondences.

Theorem 12.1. For every (S^1, \mathbf{G}) -bispectrum of simplicial Nisnevich sheaves E the morphism of bispectra $\zeta : E^c \to M_{fr}^{\mathbb{G}}(E)$ is a stable motivic equivalence. In particular, E is isomorphic to $M_{fr}^{\mathbb{G}}(E)$ in SH(k) by the zigzag of equivalences $E \leftarrow E^c \to M_{fr}^{\mathbb{G}}(E)$ and $M_{fr}^{\mathbb{G}}$ induces an equivalence of categories $M_{fr}^{\mathbb{G}} : SH(k) \xrightarrow{\simeq} SH^{fr}(k)$.

Proof. Let $diag(E^c)$, $diag(M_{fr}^{\mathbb{G}}(E))$ be the diagonal $S^1 \wedge \mathbf{G}$ -spectra. They consist of motivic spaces $(E_{0,0}^c, E_{1,1}^c, \ldots)$ and $(C_*Fr(E_{0,0}^c), C_*Fr(E_{1,1}^c), \ldots)$ respectively. It suffices to show that $diag(\zeta): diag(E^c) \rightarrow diag(M_{fr}^{\mathbb{G}}(E))$ is a stable motivic equivalence of $S^1 \wedge \mathbf{G}$ -spectra.

By [14, p. 496] $diag(E^c)$ has a natural filtration

$$diag(E^c) = \operatorname{colim}_n L_n(diag(E^c)),$$

where $L_n(diag(E^c))$ is the spectrum

$$E_{0,0}^c, E_{1,1}^c, \ldots, E_{n,n}^c, E_{n,n}^c \wedge (S^1 \wedge \mathbf{G}), E_{n,n}^c \wedge (S^1 \wedge \mathbf{G})^2, \ldots$$

In turn, $diag(M_{fr}^{\mathbb{G}}(E))$ has a natural filtration

$$diag(M_{fr}^{\mathbb{G}}(E)) = \operatorname{colim}_{n} M_{fr}^{S^{1} \wedge \mathbb{G}}(L_{n}(diag(E^{c}))),$$

where $M_{fr}^{S^1 \wedge \mathbb{G}}(L_n(diag(E^c)))$ is the spectrum

 $C_*Fr(E_{0,0}^c), C_*Fr(E_{1,1}^c), \dots, C_*Fr(E_{n,n}^c), C_*Fr(E_{n,n}^c \otimes (S^1 \otimes \mathbb{G}_m^{\wedge 1})), C_*Fr(E_{n,n}^c \otimes (S^1 \otimes \mathbb{G}_m^{\wedge 1})^2), \dots$

It follows from Theorem 11.4 that each morphism $L_n(diag(E^c)) \to M_{fr}^{S^1 \land \mathbb{G}}(L_n(diag(E^c)))$ is a stable motivic equivalence, and hence so is $diag(\zeta)$.

Observe that the composite functor

$$H_{\mathbb{A}^1}(k) \xrightarrow{\Sigma^{\infty}_{\mathbb{G}}\Sigma^{\infty}_{S^1}} SH(k) \xrightarrow{M^{\mathbb{G}}_{fr}} SH(k)$$

is isomorphic to the functor $M_{fr}^{\mathbb{G}}$ of Theorem 11.7(3). Thus the functor of Theorem 12.1 extends that of Theorem 11.7(3).

Next, denote by \mathscr{M}_{fr}^{b} the functor taking a bispectrum E to $(\Theta_{\mathbb{G}}^{\infty} \circ \Theta_{S^{1}}^{\infty} \circ M_{fr}^{\mathbb{G}})(E)$ and refer to it as the *big framed motive functor*. Here $\Theta_{S^{1}}^{\infty}$ applies to each S^{1} -spectrum of the bispectrum $M_{fr}^{\mathbb{G}}(E)$ similar to the formula (16) followed by $\Theta_{\mathbb{G}}^{\infty}$ which applies to the bispectrum $(\Theta_{S^{1}}^{\infty} \circ M_{fr}^{\mathbb{G}})(E)$ in the **G**-direction similar to (16).

Theorem 12.2. The following statements are true for every bispectrum E:

- (1) the natural map $\mu: M^{\mathbb{G}}_{fr}(E) \to \mathscr{M}^b_{fr}(E)$ is a stable motivic equivalence of bispectra;
- (2) for any $i, j \ge 0$ the space with framed correspondences $\mathscr{M}_{fr}^{b}(E)_{i,j}$ is \mathbb{A}^{1} -local as an ordinary motivic space;
- (3) the bispectrum $\mathscr{M}_{fr}^{b}(E)^{f}$, obtained from $\mathscr{M}_{fr}^{b}(E)$ by taking Nisnevich local replacements $\mathscr{M}_{fr}^{b}(E)_{i,j}^{f}$ in all entries, is motivically fibrant.

Proof. (1). Since the projective motivic model structure on $sShv_{\bullet}(Sm/k)$ is weakly finitely generated in the sense of [3], our assertion is proved similar to that of Lemma 9.5.

(2). Write $E^c = (E_{*,0}^c, E_{*,1}^c, ...)$ as a collection of S^1 -spectra. Then $M_{fr}^{\mathbb{G}}(E)$ is a collection of S^1 -spectra $(C_*Fr(E_{*,0}^c), C_*Fr(E_{*,1}^c), ...)$ and

$$\Theta_{S^1}^{\infty} M_{fr}^{\mathbb{G}}(E) = (\Theta_{S^1}^{\infty}(C_*Fr(E_{*,0}^c)), \Theta_{S^1}^{\infty}(C_*Fr(E_{*,1}^c)), \ldots).$$

By construction, the (i, j)-entry equals

$$\Theta_{S^1}^{\infty} M_{fr}^{\mathbb{G}}(E)_{i,j} := \operatorname{colim}(C_* Fr(E_{i,j}^c) \to \underline{\operatorname{Hom}}(S^1, C_* Fr(E_{i+1,j}^c)) \to \underline{\operatorname{Hom}}(S^2, C_* Fr(E_{i+2,j}^c)) \to \cdots).$$

In each weight j, the S¹-spectrum $E_{*,j}^c$ has a natural filtration $E_{*,j}^c = \operatorname{colim}_n L_n(E_{*,j}^c)$, where $L_n(E_{*,j}^c)$ is the spectrum

$$E_{0,j}^{c}, E_{1,j}^{c}, \ldots, E_{n,j}^{c}, E_{n,j}^{c} \wedge S^{1}, E_{n,j}^{c} \wedge S^{2}, \ldots$$

Then $C_*Fr(E_{*,j}^c) = C_*Fr(\operatorname{colim}_n L_n(E_{*,j}^c)) = \operatorname{colim}_n C_*Fr(L_n(E_{*,j}^c))$, where $C_*Fr(L_n(E_{*,j}^c))$ is the spectrum

$$C_*Fr(E_{0,j}^c), C_*Fr(E_{1,j}^c), \dots, C_*Fr(E_{n,j}^c), C_*Fr(E_{n,j}^c \otimes S^1), C_*Fr(E_{n,j}^c \otimes S^2), \dots$$

Since each space $C_*Fr(E_{n,j}^c \otimes S^\ell)$, $\ell > 0$, is \mathbb{A}^1 -local, then so is the space $\underline{\operatorname{Hom}}(S^m, C_*Fr(E_{n,j}^c \otimes S^\ell))$, where $m \ge 0$. Therefore each space of $\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j}^c)) = \Theta_{S^1}^{\infty}(\operatorname{colim}_n C_*Fr(L_n(E_{*,j}^c))) = \operatorname{colim}_n \Theta_{S^1}^{\infty}(C_*Fr(L_n(E_{*,j}^c)))$ is \mathbb{A}^1 -local, because so is each $\Theta_{S^1}^{\infty}(C_*Fr(L_n(E_{*,j}^c)))$. We use here the fact that a directed colimit of Nisnevich excisive spaces is Nisnevich excisive to conclude that a directed colimit of \mathbb{A}^1 -local spaces is \mathbb{A}^1 -local. We see that each $\Theta_{S^1}^{\infty}M_{fr}^{\mathbb{G}}(E)_{i,j}$ is \mathbb{A}^1 -local. If we take a Nisnevich local fibrant resolution $\Theta_{S^1}^{\infty}M_{fr}^{\mathbb{G}}(E)_{*,j}^f$ of the bispectrum $\Theta_{S^1}^{\infty}M_{fr}^{\mathbb{G}}(E)^f$ is motivically fibrant in the local stable model structure of S^1 -spectra. Indeed, this follows from the fact that the structure maps of the S^1 -spectrum $\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j}^c))$ are isomorphisms and that the functor $\operatorname{Hom}(S^1, -)$ preserves local equivalences.

Next, the S¹-spectrum in each weight j of $\mathscr{M}_{fr}^{b}(E)$ equals by definition

$$\operatorname{colim}(\Theta_{S^{1}}^{\infty}(C_{*}Fr(E_{*,j}^{c})) \to \operatorname{\underline{Hom}}(\mathbf{G}, \Theta_{S^{1}}^{\infty}(C_{*}Fr(E_{*,j+1}^{c}))) \to \operatorname{\underline{Hom}}(\mathbf{G}^{\wedge 2}, \Theta_{S^{1}}^{\infty}(C_{*}Fr(E_{*,j+2}^{c}))) \to \cdots)$$

We have shown above that $(\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j}^c)))_f$ is a motivically fibrant S^1 -spectrum. We claim that

$$\underline{\operatorname{Hom}}(\mathbf{G}, \Theta_{S^1}^{\infty}(C_*Fr(E_{*,j+1}^c))) \to \underline{\operatorname{Hom}}(\mathbf{G}, (\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j+1}^c)))_f)$$

is a levelwise local equivalence. In this case it will follow that each space $\mathscr{M}_{fr}^{b}(E)_{i,j}$ is \mathbb{A}^{1} -local.

Since both spectra are sectionwise Ω -spectra, it suffices to prove that this arrow is a stable local equivalence. The presheaves of stable homotopy groups of the left spectrum are \mathbb{A}^1 -invariant quasi-stable radditive with framed correspondences (see [6, Introduction]. Therefore our claim follows from the following

Sublemma. Let \mathscr{X} be an \mathbb{A}^1 -local motivic S^1 -spectrum whose presheaves of stable homotopy groups are homotopy invariant quasi-stable radditive presheaves with framed correspondences (see [7] for the definition of such presheaves). Suppose \mathscr{X}^f is a local stable fibrant replacement of \mathscr{X} . Then the map of spectra $\operatorname{Hom}(\mathbf{G}, \mathscr{X}) \to \operatorname{Hom}(\mathbf{G}, \mathscr{X}^f)$ is a local stable equivalence.

Proof. First let us compute $\pi^{nis}_*(\underline{\text{Hom}}((\mathbb{G}_m)_+, \mathscr{X}^f))$. We have $\mathscr{X}^f = \text{hocolim}_{n \to -\infty} \mathscr{X}^f_{\geq n}$, where $\mathscr{X}_{\geq n}$ is the naive *n*th truncation of \mathscr{X} in $Sp_{S^1}(sShv_\bullet(Sm/k))$. \mathscr{X} has homotopy invariant, quasistable radditive presheaves with framed correspondences of stable homotopy groups $\pi_*(\mathscr{X})$. By [7, 1.1] the Nisnevich sheaves $\pi^{nis}_*(\mathscr{X}_{\geq n})$ are strictly homotopy invariant. If $\mathscr{X}^f_{\geq n}$ is a stable local replacement of $\mathscr{X}_{\geq n}$, it follows from Proposition 7.1 that $\mathscr{X}^f_{\geq n}$ is motivically fibrant, hence $\mathscr{X}_{\geq n}$ is \mathbb{A}^1 -local in $SH_{S^1}(k)$. Since

$$\underline{\operatorname{Hom}}((\mathbb{G}_m)_+,\mathscr{X}^f) = \operatorname{hocolim}_{n \to -\infty} \underline{\operatorname{Hom}}((\mathbb{G}_m)_+,\mathscr{X}^f_{\geq n})$$

 $\pi^{\text{nis}}_*(\underline{\text{Hom}}((\mathbb{G}_m)_+,\mathscr{X}^f)) = \operatorname{colim}_{n \to -\infty} \pi^{\text{nis}}_*(\underline{\text{Hom}}((\mathbb{G}_m)_+,\mathscr{X}^f_{\geq n})).$ Consider the Brown–Gersten convergent spectral sequence

$$H^p_{\mathrm{nis}}(U \times \mathbb{G}_m, \pi^{\mathrm{nis}}_q(\mathscr{X}_{\geqslant n})) \Longrightarrow \pi_{q-p}(\mathscr{X}^f_{\geqslant n})(U \times \mathbb{G}_m), \quad U \in Sm/k.$$

It follows from [7, 16.10-11] that each presheaf $U \mapsto H^p_{nis}(U \times \mathbb{G}_m, \pi^{nis}_q(\mathscr{X}_{\geq n}))$ is \mathbb{A}^1 -invariant quasi-stable radditive with framed correspondences. If U is a smooth local Henzelian scheme then by [7, 2.15(3')] there is an embedding

$$H^p_{\mathrm{nis}}(U \times \mathbb{G}_m, \pi^{\mathrm{nis}}_q(\mathscr{X}_{\geq n})) \hookrightarrow H^p_{\mathrm{nis}}(\mathbb{G}_{m,k(U)}, \pi^{\mathrm{nis}}_q(\mathscr{X}_{\geq n})),$$

where k(U) is the function field on U. By the Sublemma in [6, Appendix A] we have that $H^p_{\text{nis}}(\mathbb{G}_{m,k(U)}, \pi^{\text{nis}}_q(\mathscr{X}_{\geq n})) = 0$ for p > 0, and hence $H^p_{\text{nis}}(U \times \mathbb{G}_m, \pi^{\text{nis}}_q(\mathscr{X}_{\geq n}^f)) = 0$ for p > 0. We can conclude that

$$\pi_n^{\operatorname{nis}}(\operatorname{\underline{Hom}}((\mathbb{G}_m)_+,\mathscr{X}^f))=\pi_n^{\operatorname{nis}}(\mathscr{X}^f)(\mathbb{G}_m\times-).$$

It also follows that

(

$$\pi_n^{\operatorname{nis}}(\operatorname{\underline{Hom}}(\mathbf{G},\mathscr{X}^f)) = (\pi_n^{\operatorname{nis}}(\mathscr{X}^f))_{-1} = (\pi_n^{\operatorname{nis}}(\mathscr{X}^f))_{-1}$$

It remains to show that the morphism of \mathbb{A}^1 -invariant radditive quasi-stable framed sheaves

$$(\pi_n(\operatorname{\underline{Hom}}(\mathbf{G},\mathscr{X})))^{\operatorname{nis}} = (\pi_n(\mathscr{X}))^{\operatorname{nis}}_{-1} \to (\pi_n^{\operatorname{nis}}(\mathscr{X}))_{-1}$$

is an isomorphism. Using [7, 2.15(3')] it suffices to check that it is an isomorphism for every field extension K/k. The homomorphism of Abelian groups

$$(\pi_n(\mathscr{X}))_{-1}^{\mathrm{nis}}(K) = (\pi_n(\mathscr{X}))_{-1}(K) \to (\pi_n^{\mathrm{nis}}(\mathscr{X}))_{-1}(K)$$

is an isomorphism, because for any \mathbb{A}^1 -invariant radditive quasi-stable framed presheaf of Abelian groups \mathscr{F} and every open $V \subset \mathbb{A}^1_K$, one has $\mathscr{F}(V) = \mathscr{F}^{nis}(V)$ (see the proof of [7, 2.1]). \Box

(3). By the previous assertion each space $\mathscr{M}_{fr}^{b}(E)_{i,j}$ is \mathbb{A}^{1} -local, and hence $\mathscr{M}_{fr}^{b}(E)_{i,j}^{f}$ is a motivically fibrant space. The proof of the assertion shows that $\mathscr{M}_{fr}^{b}(E)_{i,j}^{f}$ can be computed as the *i*th space of the the motivically fibrant S^{1} -spectrum

$$\operatorname{colim}(\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j}^c))^f \to \Omega_{\mathbf{G}}(\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j+1}^c))^f) \to \Omega_{\mathbf{G}^{\wedge 2}}(\Theta_{S^1}^{\infty}(C_*Fr(E_{*,j+2}^c))^f) \to \cdots).$$

Moreover, this colimit is the S^1 -spectrum in weight j of the bispectrum $\mathcal{M}_{fr}^b(E)^f$. By [11, 4.12] such a bispectrum must be motivically fibrant. This completes the proof.

Definition 12.3. (1) Given a (S^1, \mathbf{G}) -bispectrum E and $i, j \ge 0$, denote by $\underline{\pi}_{i,j}^{fr}(E)$ the graded Nisnevich sheaf $\bigoplus_{n\ge 0} \pi_n^{\text{nis}}(\mathscr{M}_{fr}^b(E)_{i,j})$ and refer to such sheaves as *framed sheaves of* E.

(2) Denote by $SH_{nis}^{fr}(k)$ the full subcategory of $SH^{fr}(k)$ consisting of framed (S^1, \mathbf{G}) -bispectra whose motivic spaces are \mathbb{A}^1 -local as ordinary motivic spaces.

Theorem 12.2 implies the following

Theorem 12.4. *The following statements are true:*

- (1) Framed sheaves detect stable weak equivalences of bispectra. Namely, a map of bispectra $f: E \to E'$ is a stable motivic equivalence if and only if $\underline{\pi}_{i,j}^{fr}(E) \xrightarrow{f_*} \underline{\pi}_{i,j}^{fr}(E)$ is an isomorphism of sheaves for all $i, j \ge 0$.
- (2) For every bispectrum E the bispectrum $\mathscr{M}_{fr}^b(E)$ belongs to $SH_{nis}^{fr}(k)$. Moreover, the functor $\mathscr{M}_{fr}^b: SH(k) \to SH_{nis}^{fr}(k)$ is an equivalence of categories and there is a natural isomorphism of endofunctors on SH(k):

$$\mathrm{id} \to \iota \circ \mathscr{M}_{fr}^{\mathfrak{o}},$$

where $\iota: SH_{nis}^{fr}(k) \to SH(k)$ is the natural inclusion.

In other words, the preceding theorem says that the functor \mathscr{M}_{fr}^b converts classical Morel– Voevodsky's stable motivic homotopy theory SH(k) into an equivalent local homotopy theory of \mathbb{A}^1 -local framed bispectra from $SH_{nis}^{fr}(k)$. The main ingredient of this equivalent local homotopy theory is framed motivic spaces of the form $C_*Fr(-,Y)$ with $Y \in \Delta^{op}Fr_0(k)$ a simplicial scheme as well as their framed motives $M_{fr}(Y)$.

We document this as follows.

Theorem 12.5. The Morel–Voevodsky stable motivic homotopy category SH(k) can be defined as follows. Its objects are (S^1, \mathbf{G}) -bispectra and morphisms between two bispectra E, E' are given by the set $\pi_0(E^c, \mathscr{M}^b_{fr}(E')^f)$ of ordinary morphisms between bispectra E^c and $\mathscr{M}^b_{fr}(E')^f$ modulo the naive homotopy. In particular, $SH(k)(\Sigma^{\infty}_{S^1}\Sigma^{\infty}_{\mathbb{G}}X_+, E') = \pi_0(\mathscr{M}^b_{fr}(E')^f_{0,0}(X))$ for any $X \in Sm/k$.

To conclude the section, the interested reader will easily construct the "framed functor with 1/2-coefficients" $\mathcal{M}_{fr}^b\{1/2\}$ and prove the same theorems from this section for SH(k)[1/2] in any characteristic of the base field k.

13. Framed \mathbb{P}^1 -spectra

In Theorem 10.5 we have shown that the suspension spectrum $\Sigma_{\mathbb{P}^1}^{\infty} \mathscr{X}$ of a motivic space $\mathscr{X} \in sShv_{\bullet}(Sm/k)$ is naturally equivalent to the spectrum

$$M_{\mathbb{P}^{\wedge 1}}(\mathscr{X}^{c}) = (C_{*}\mathscr{F}r(-,\mathscr{X}^{c}), C_{*}\mathscr{F}r(-,\mathscr{X}^{c}\wedge T), C_{*}\mathscr{F}r(-,\mathscr{X}^{c}\wedge T^{2}), \ldots).$$

It induces a functor of homotopy categories

$$M_{\mathbb{P}^{\wedge 1}}: H_{\mathbb{A}^1}(k) \to SH(k),$$

which is equivalent to $\Sigma_{\mathbb{P}^1}^{\infty}$ (see Theorem 10.5). Observe that $M_{\mathbb{P}^{\wedge 1}}$ lands in the full subcategory of \mathbb{P}^1 -spectra, denote it by $SH^{fr}(k)$, whose motivic spaces are \mathbb{A}^1 -invariant with framed correspondences. This section will show the reader how to get naturally framed \mathbb{P}^1 -spectra. Another approach for bispectra was illustrated in the previous section.

The purpose of this final section is to construct an equivalence of categories

$$SH(k) \xrightarrow{\cong} SH^{fr}(k)$$

(see Theorem 13.4). Moreover, we shall prove that every spectrum is isomorphic in SH(k) to a framed Ω -spectrum from $SH^{fr}(k)$.

Throughout the section k is any field and by $\mathbf{Spt}^{\mathbb{P}^1}(\mathbf{Sm}/k)$ (respectively $\mathbf{Spt}^T(\mathbf{Sm}/k)$) we mean the category of \mathbb{P}^1 -spectra (respectively T-spectra) associated with simplicial Nisnevich sheaves. We shall consider the level and stable flasque model structures on $\mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ or **Spt**^T (Sm/k) in the sense of [12]. An advantage of the model structures is that a filtered colimit of fibrant objects is again fibrant [12, 5.3, 6.7] (in fact, both model structures are weakly finitely generated in the sense of [3]). A fibrant \mathbb{P}^1 -spectrum with respect to the stable motivic model structure will also be referred to as an Ω -spectrum.

The motivic equivalence $\sigma: \mathbb{P}^{\wedge 1} \to T$ induces an adjoint pair

$$f: \mathbf{Spt}^{\mathbb{P}^1}(Sm/k) \rightleftharpoons \mathbf{Spt}^T(Sm/k): g,$$

where g is the forgetful functor. When proving Theorem 4.1(1) the reader may have observed that we first replaced the suspension spectrum $\Sigma_{\mathbb{P}^1}^{\infty} \mathscr{X}$ by the suspension *T*-spectrum $\Sigma_T^{\infty} \mathscr{X} =$ $f(\Sigma_{\mathbb{P}^1}^{\infty}\mathscr{X})$ and then applied Θ^{∞} to the \mathbb{P}^1 -spectrum $\Sigma_{\mathbb{P}^1,T}^{\infty}\mathscr{X} := gf(\Sigma_{\mathbb{P}^1}^{\infty}\mathscr{X}) = g(\Sigma_T^{\infty}\mathscr{X})$ in order to get framed correspondences. We see that framed correspondences are obtained from the T-spectrum $\Sigma_T^{\infty} \mathscr{X}$.

We want to extend this construction to spectra. Suppose $\mathscr{E} \in \mathbf{Spt}^T(Sm/k)$. By [14, p. 496] $\mathscr{E} = (\mathscr{E}_0, \mathscr{E}_1, \ldots)$ has a natural filtration

$$\mathscr{E} = \operatorname{colim}_n L_n \mathscr{E},$$

where $L_n \mathscr{E}$ is the spectrum

$$\mathscr{E}_0, \mathscr{E}_1, \ldots, \mathscr{E}_n, \mathscr{E}_n \wedge T, \mathscr{E}_n \wedge T^2, \ldots$$

Denote by $L_n^{\mathbb{P}^1} \mathscr{E}$ the spectrum $g(L_n \mathscr{E})$. By Lemma 9.5 the natural map of spectra

$$D_n: L_n^{\mathbb{P}^1}\mathscr{E} \to \Theta^{\infty} L_n^{\mathbb{P}^1}\mathscr{E}$$

is a stable equivalence. There is an isomorphism of spectra

 $\iota_n : \Theta^{\infty} L_n^{\mathbb{P}^1} \mathscr{E} \xrightarrow{\cong} \Theta^{\infty} L_{n,fr}^{\mathbb{P}^1} \mathscr{E} := (\underline{\operatorname{Hom}}(\mathbb{P}^{\wedge n}, \mathscr{F}r(\mathscr{E}_n)), \dots, \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, \mathscr{F}r(\mathscr{E}_n)), \mathscr{F}r(\mathscr{E}_n), \mathscr{F}r(\mathscr{E}_n \wedge T), \dots).$ If we apply the Suslin complex functor C_* levelwise, we get a spectrum

 $C_*\Theta^{\infty}L_{n,fr}^{\mathbb{P}^1}\mathscr{E} := (\underline{\operatorname{Hom}}(\mathbb{P}^{\wedge n}, C_*\mathscr{F}r(\mathscr{E}_n)), \ldots, \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_n)), C_*\mathscr{F}r(\mathscr{E}_n), C_*\mathscr{F}r(\mathscr{E}_n \wedge T), \ldots).$

By [17, 2.3.8] the natural map of spectra

$$\delta_n: \Theta^{\infty} L_{n,fr}^{\mathbb{P}^1} \mathscr{E} \to C_* \Theta^{\infty} L_{n,fr}^{\mathbb{P}^1} \mathscr{E}$$

is a level motivic weak equivalence.

Passing to the colimit over *n* we get that the composite map of spectra

$$\operatorname{colim}_n(\delta_n\iota_n\eta_n):g(\mathscr{E})=\operatorname{colim}_nL_n^{\mathbb{P}^1}\mathscr{E}\to\operatorname{colim}_nC_*\Theta^{\infty}L_{n,fr}^{\mathbb{P}^1}\mathscr{E}$$

is a stable motivic weak equivalence. We use here the fact that a sequential colimit of stable motivic weak equivalences is a stable motivic weak equivalence by [14, 3.12].

Denote by $\Theta^{\infty}C_*\mathscr{F}r(\mathscr{E})$ the spectrum

 $\operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, C_{*}\mathscr{F}r(\mathscr{E}_{n})), \operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, C_{*}\mathscr{F}r(\mathscr{E}_{1+n})), \operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, C_{*}\mathscr{F}r(\mathscr{E}_{2+n})), \ldots$

Also, denote by $C_*\mathscr{F}r(\mathscr{E})$ the spectrum $(C_*\mathscr{F}r(\mathscr{E}_0), C_*\mathscr{F}r(\mathscr{E}_1), C_*\mathscr{F}r(\mathscr{E}_2), \ldots)$. Each structure map $C_*\mathscr{F}r(\mathscr{E}_k) \to \underline{\mathrm{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_{k+1})), k \ge 0$, is given by the natural composition

$$C_*\mathscr{F}r(\mathscr{E}_k) \to \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_k \wedge \mathbb{P}^{\wedge 1})) \xrightarrow{\sigma_*} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_k \wedge T)) \xrightarrow{(\varepsilon_k)_*} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_{k+1})) \xrightarrow{\sigma_*} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_{k+1}))$$

where $\varepsilon_k : \mathscr{E}_k \wedge T \to \mathscr{E}_{k+1}$ is the structure map of \mathscr{E} . Observe that for any $\mathscr{X} \in \Delta^{\operatorname{op}} \overrightarrow{Fr_0}(k)$ the spectrum $M_{\mathbb{P}^{\wedge 1}}(\mathscr{X})$ is isomorphic to $C_*\mathscr{F}r(\Sigma_{\mathbb{P}^1}^{\infty} {}_T(\mathscr{X}))$ in SH(k).

The spectrum $\operatorname{colim}_n C_* \Theta^{\infty} L_{n,fr}^{\mathbb{P}^1} \mathscr{E}$ is naturally isomorphic to the spectrum $\Theta^{\infty} C_* \mathscr{F} r(\mathscr{E})$ and the stable motivic equivalence

$$\alpha: g(\mathscr{E}) \to \Theta^{\infty}C_*\mathscr{F}r(\mathscr{E})$$

factors as $g(\mathscr{E}) \xrightarrow{\beta} C_* \mathscr{F}r(\mathscr{E}) \xrightarrow{\gamma} \Theta^{\infty} C_* \mathscr{F}r(\mathscr{E})$. Here γ equals the stable motivic equivalence of Lemma 9.5 and each $\beta_n : \mathscr{E}_n \to C_* \mathscr{F}r(\mathscr{E}_n)$ is the obvious map of motivic spaces. The twoout-of-three property implies β is a stable motivic equivalence. It is plainly functorial in $\mathscr{E} \in$ **Spt**^{*T*}(*Sm/k*).

Theorem 13.1. For every *T*-spectrum \mathscr{E} there is a natural stable motivic equivalence of spectra $\beta: g(\mathscr{E}) \to C_* \mathscr{F}r(\mathscr{E})$, functorial in \mathscr{E} . Moreover, a morphism of spectra $u: \mathscr{E} \to \mathscr{E}'$ is a stable motivic equivalence if and only if so is $C_* \mathscr{F}r(u): C_* \mathscr{F}r(\mathscr{E}) \to C_* \mathscr{F}r(\mathscr{E}')$. In particular, we have a functor

$$C_*\mathscr{F}r: \operatorname{Ho}(\operatorname{\mathbf{Spt}}^T(Sm/k)) \to \operatorname{Ho}(\operatorname{\mathbf{Spt}}^{\mathbb{P}^n}(Sm/k)),$$

which is an equivalence of categories.

Proof. The first statement has already been verified above. The second statement follows from the first statement and the fact that $u : \mathscr{E} \to \mathscr{E}'$ is a stable motivic equivalence if and only if so is $g(u) : g(\mathscr{E}) \to g(\mathscr{E}')$ (see [14, p. 477]). The functor

$$C_*\mathscr{F}r: \operatorname{Ho}(\operatorname{\mathbf{Spt}}^T(Sm/k)) \to \operatorname{Ho}(\operatorname{\mathbf{Spt}}^{\mathbb{P}^1}(Sm/k)),$$

is an equivalence of categories, because so is g and $\beta : g(\mathscr{E}) \to C_*\mathscr{F}r(\mathscr{E})$ is a stable motivic equivalence, functorial in \mathscr{E} .

Lemma 13.2. For every *T*-spectrum \mathscr{E} , each space of the \mathbb{P}^1 -spectrum $\Theta^{\infty} \mathscr{F}r(\mathscr{E})$ (respectively $\Theta^{\infty}C_*\mathscr{F}r(\mathscr{E})$) is a motivic space with framed correspondences.

Proof. $\Theta^{\infty} \mathscr{F} r(\mathscr{E})$ is the spectrum

$$\operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, \mathscr{F}r(\mathscr{E}_{n})), \operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, \mathscr{F}r(\mathscr{E}_{1+n})), \operatorname{colim}_{n} \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, \mathscr{F}r(\mathscr{E}_{2+n})), \dots$$

Given a sheaf *F* and s > 0, we claim that $\underline{\text{Hom}}(\mathbb{P}^{\wedge s}, \mathscr{F}r(F))$ is a sheaf with framed correspondences (the internal Hom is taken in the category of pointed Nisnevich sheaves). To see this, define for any $U, X \in Sm/k$ and any m, n a map of pointed sets

$$\operatorname{Hom}(U_+ \wedge \mathbb{P}^{\wedge m}, X_+ \wedge T^m) \wedge \operatorname{Hom}(X_+ \wedge \mathbb{P}^{\wedge s+n}, F \wedge T^n) \to \operatorname{Hom}(U_+ \wedge \mathbb{P}^{\wedge (s+m+n)}, F \wedge T^{m+n}).$$

If $\alpha : U_+ \wedge \mathbb{P}^{\wedge m} \to X_+ \wedge T^m$ and $v : X_+ \wedge \mathbb{P}^{\wedge s+n} \to F \wedge T^n$ are morphisms of pointed Nisnevich sheaves, then we define $\alpha^*(v)$ as the composite morphism

$$U_{+} \wedge \mathbb{P}^{\wedge s} \wedge \mathbb{P}^{\wedge m} \wedge \mathbb{P}^{\wedge n} \cong U_{+} \wedge \mathbb{P}^{\wedge m} \wedge \mathbb{P}^{\wedge s+n} \xrightarrow{\alpha \wedge \mathrm{id}} X_{+} \wedge T^{m} \wedge \mathbb{P}^{\wedge s+n} \cong$$
$$\cong T^{m} \wedge X_{+} \wedge \mathbb{P}^{\wedge s+n} \xrightarrow{\mathrm{id} \wedge \nu} T^{m} \wedge F \wedge T^{n} \cong F \wedge T^{m} \wedge T^{n}.$$

Passing to the colimit over *n*, we get that $\underline{\text{Hom}}(\mathbb{P}^{\wedge s}, \mathscr{F}r(F))$ is a sheaf with framed correspondences as claimed.

Next, the composite morphism

$$\underline{\operatorname{Hom}}(\mathbb{P}^{\wedge s}, \mathscr{F}r(\mathscr{E}_{k})) \to \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge s+1}, \mathscr{F}r(\mathscr{E}_{k} \wedge T)) \xrightarrow{(\varepsilon_{k})_{*}} \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge s+1}, \mathscr{F}r(\mathscr{E}_{k+1})), \quad s, k \ge 0,$$

is plainly a morphism of framed Nisnevich sheaves. Recall that the left arrow is induced by the
map taking $(v: X_{+} \wedge \mathbb{P}^{\wedge s+n} \to \mathscr{E}_{k} \wedge T^{n}) \in \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge s}, \mathscr{F}r_{n}(\mathscr{E}_{k}))$ to the composition

$$(X_{+} \wedge \mathbb{P}^{\wedge s+1+n} \cong X_{+} \wedge \mathbb{P}^{\wedge s+n} \wedge \mathbb{P}^{\wedge 1} \xrightarrow{\nu \wedge \sigma} \mathscr{E}_{k} \wedge T^{n} \wedge T \cong \mathscr{E}_{k} \wedge T \wedge T^{n}) \in \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge s+1}, \mathscr{F}r_{n}(\mathscr{E}_{k} \wedge T)).$$

It follows that each motivic space $(\Theta^{\infty} \mathscr{F}r(\mathscr{E}))_k = \operatorname{colim}_n \operatorname{\underline{Hom}}(\mathbb{P}^{\wedge n}, \mathscr{F}r(\mathscr{E}_{k+n}))$ of the spectrum $\Theta^{\infty} \mathscr{F}r(\mathscr{E})$ has framed correspondences. Obviously, the same is true for the spectrum $\Theta^{\infty}C_*\mathscr{F}r(\mathscr{E})$.

Suppose $\mathscr{E} \in \mathbf{Spt}^T(Sm/k)$. Then there are isomorphisms of \mathbb{P}^1 -spectra

$$\Theta^{\infty}(g(\mathscr{E})) = \Theta^{\infty}(\operatorname{colim}_{n} L_{n}^{\mathbb{P}^{1}} \mathscr{E}) \cong \operatorname{colim}_{n} \Theta^{\infty}(L_{n}^{\mathbb{P}^{1}} \mathscr{E})) \cong \Theta^{\infty} \mathscr{F}r(\mathscr{E}).$$

If \mathscr{E} is levelwise motivically fibrant it follows that $\Theta^{\infty} \mathscr{F}r(\mathscr{E})$ is an Ω -spectrum, because so is $\Theta^{\infty}(g(\mathscr{E}))$ by [11, 4.6]. Using Lemma 13.2, we have shown therefore the following

Lemma 13.3. For every levelwise fibrant *T*-spectrum \mathscr{E} , the spectrum $\Theta^{\infty} \mathscr{F}r(\mathscr{E}) \in \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ is an Ω -spectrum with framed correspondences.

We are now in a position to prove that SH(k) is canonically equivalent to its full subcategory $SH^{fr}(k)$ of framed spectra, i.e. those spectra whose motivic spaces are \mathbb{A}^1 -invariant with framed correspondences.

Theorem 13.4. The functor $C_* \mathscr{F}r : \mathbf{Spt}^T(Sm/k) \to \mathbf{Spt}^{\mathbb{P}^1}(Sm/k)$ takes a spectrum \mathscr{E} to a framed spectrum $C_* \mathscr{F}r(\mathscr{E})$. The composite functor

$$SH(k) = \operatorname{Ho}(\operatorname{\mathbf{Spt}}^{\mathbb{P}^1}(Sm/k)) \xrightarrow{f} \operatorname{Ho}(\operatorname{\mathbf{Spt}}^T(Sm/k)) \xrightarrow{C_*\mathscr{F}r} \operatorname{Ho}(\operatorname{\mathbf{Spt}}^{\mathbb{P}^1}(Sm/k)),$$

induces an equivalence of categories $SH(k) \xrightarrow{\sim} SH^{fr}(k)$. Moreover, every \mathbb{P}^1 -spectrum is isomorphic in SH(k) to a framed Ω -spectrum.

Proof. By construction, $C_* \mathscr{F}r(\mathscr{E})$ is an \mathbb{A}^1 -invariant with framed correspondences. The next statement follows from Theorem 13.1 and the fact that f is an equivalence of categories (see [14, 2.13]). To show that every \mathbb{P}^1 -spectrum E is isomorphic in SH(k) to a framed Ω -spectrum, let E^c be a cofibrant replacement of E and $f(E^c)_{lf} \in \mathbf{Spt}^T(Sm/k)$ be a level fibrant replacement of the T-spectrum $f(E^c)$. Let Θ^{∞}_T denote the stabilization functor in the category of T-spectra. Since the flasque motivic model structure on spaces is weakly finitely generated (this follows from [12, 3.10]), the T-spectrum $\Theta^{\infty}_T(f(E^c)_{lf})$ is fibrant by [11, 4.6] and the morphism $f(E^c)_{lf} \to \Theta^{\infty}_T(f(E^c)_{lf})$ is a stable equivalence by [11, 4.11]. The canonical arrow $g(\Theta^{\infty}_T(f(E^c)_{lf})) \to \Theta^{\infty}(gf(E^c)_{lf})$ is a level weak equivalence by [14, p. 477], and hence the composite morphism

$$E^{c} \to gf(E^{c}) \to g(\Theta^{\infty}_{T}(f(E^{c})_{lf})) \to \Theta^{\infty}(gf(E^{c})_{lf})$$

is a stable weak equivalence, where the left arrow is the adjunction unit morphism. The \mathbb{P}^1 -spectrum $\Theta^{\infty}(gf(E^c)_{lf})$ is a framed Ω -spectrum by Lemma 13.3, so the zigzag

$$E \leftarrow E^c \to \Theta^{\infty}(gf(E^c)_{lf})$$

gives an isomorphism in SH(k).

The preceding theorem says that we can define the stable motivic homotopy theory as framed \mathbb{P}^1 -spectra or even framed Ω -spectra. However, such framed Ω -spectra are hardly amenable for explicit calculations in general, because they require levelwise motivically fibrant replacements by construction. Instead, the main point of this paper is to show that whenever the base field *k* is infinite perfect with char $k \neq 2$, one can nevertheless construct *exlicit* fibrant Ω -spectra using framed correspondences of Voevodsky and the machinery of framed motives introduced and developed in this paper.

Namely, suppose $\mathscr{E} = (\mathscr{E}_0, \mathscr{E}_1, \ldots) \in \mathbf{Spt}^T(Sm/k)$ is such that its framed spectrum $C_*\mathscr{F}r(\mathscr{E})$ satisfies the following conditions:

- (1) each space $C_*\mathscr{F}r(\mathscr{E}_n)$, $n \ge 0$, is locally connected;
- (2) each space C_{*} ℱr(ℰ_n), n ≥ 0, is σ-invariant (i.e. it takes the framed correspondence σ_X = (X × 0, X × Å¹, pr_{Å¹}, pr_X) of level one to a weak equivalence of simplicial sets for every X ∈ Sm/k);
- (3) each structure morphism induces a motivic equivalence

$$C_*\mathscr{F}r(\mathscr{E}_n)_f \to \underline{\operatorname{Hom}}(\mathbb{P}^{\wedge 1}, C_*\mathscr{F}r(\mathscr{E}_{n+1})_f), \quad n \ge 0,$$

where the subscript "f" refers to a local fibrant replacement.

Then the spectrum $C_*\mathscr{F}r(\mathscr{E})_f := (C_*\mathscr{F}r(\mathscr{E}_0)_f, C_*\mathscr{F}r(\mathscr{E}_1)_f, ...)$ is an Ω -spectrum stably equivalent to \mathscr{E} . In particular, if a motivic space \mathscr{X} is such that its suspension *T*-spectrum satisfies (1) - (3), then $C_*\mathscr{F}r(\mathscr{X})$ is locally equivalent to the space $\Omega_{\mathbb{P}^1}^{\infty}\Sigma_{\mathbb{P}^1}^{\infty}\mathscr{X}$ (see p. 14 for the definition of the latter space). The hardest condition in practice is condition (3), where the machinery of framed motives works in its full capacity.

14. CONCLUDING REMARKS

To finish the paper, we should mention that Voevodsky [22] also defines another type of framed correspondences $Fr_*^{rat}(U,X)$, in which regular functions on etale neighborhoods of supports are replaced by *rational* functions. Additivity Theorem 6.4 is also true for the motivic space

$$U \in Sm/k \mapsto C_*Fr^{rat}(U,X) \in \mathbb{S}_{\bullet}$$
 $X \in Sm/k$.

It follows from [22] that $C_*Fr^{rat}(U,X)$ is a group-like space, and hence the S¹-spectrum

$$M_{fr}^{rat}(X) = (C_*Fr^{rat}(-,X), C_*Fr^{rat}(-,X \otimes S^1), C_*Fr^{rat}(-,X \otimes S^2), \dots)$$

is an Ω -spectrum by the Segal machine [19]. We call it the *rational framed motive of* $X \in Sm/k$.

Conjecture 1. The natural map of S^1 -spectra $M_{fr}(X) \to M_{fr}^{rat}(X)$ is a local stable weak equivalence for all $X \in Sm/k$.

Basing on the technique developed in the previous sections, it is quite standard to develop homotopy theory of rational framed motives. If the conjecture is true, one can then construct explicit motivic fibrant replacements for spectra/bispectra of the form $\sum_{\mathbb{P}^1}^{\infty} X_+$ and $\sum_{S^1}^{\infty} \sum_{\mathbb{G}}^{\infty} X_+$ respectively in terms of spaces $C_*Fr^{rat}(-,X)$.

REFERENCES

- A. Ananyevskiy, G. Garkusha, I. Panin, Cancellation theorem for framed motives of algebraic varieties, preprint arXiv:1601.06642.
- [2] B. Blander, Local projective model structures on simplicial presheaves, K-Theory 24(3) (2001), 283-301.
- [3] B. I. Dundas, O. Röndigs, P. A. Østvær, Enriched functors and stable homotopy theory, Doc. Math. 8 (2003), 409-488.
- [4] B. I. Dundas, M. Levine, P. A. Østvær, O. Röndigs, V. Voevodsky, Motivic homotopy theory, Universitext, Springer-Verlag, Berlin, 2007.
- [5] G. Garkusha, I. Panin, K-motives of algebraic varieties, Homology, Homotopy Appl. 14(2) (2012), 211-264.
- [6] G. Garkusha, I. Panin, On the motivic spectral sequence, J. Inst. Math. Jussieu 17(1) (2018), 137-170.
- [7] G. Garkusha, I. Panin, Homotopy invariant presheaves with framed transfers, preprint arXiv:1504.00884.
- [8] G. Garkusha, A. Neshitov, I. Panin, Framed motives of relative motivic spheres, preprint arXiv:1604.02732.
- [9] D. Grayson, Weight filtrations via commuting automorphisms, K-Theory 9 (1995), 139-172.
- [10] A. Grothendieck, Éléments de géométrie algébrique. IV, Étude locale des schémas et des morphismes de schémas, Quatrième partie, Publ. Math. IHES 32 (1967), 5-361.

- [11] M. Hovey, Spectra and symmetric spectra in general model categories, J. Pure Appl. Algebra, 165(1) (2001), 63-127.
- [12] D. Isaksen, Flasque model structures for simplicial presheaves, K-Theory 36 (2005), 371-395.
- [13] J. F. Jardine, Simplicial presheaves, J. Pure Appl. Algebra 47 (1987), 35-87.
- [14] J. F. Jardine, Motivic symmetric spectra, Doc. Math. 5 (2000), 445-552.
- [15] M. Levine, A comparison of motivic and classical stable homotopy theories, J. Topology 7 (2014), 327-362.
- [16] F. Morel, \mathbb{A}^1 -Algebraic topology over a field, Lecture Notes in Mathematics, No. 2052, Springer-Verlag, 2012.
- [17] F. Morel, V. Voevodsky, A¹-homotopy theory of schemes, Publ. Math. IHES 90 (1999), 45-143.
- [18] A. Neshitov, Framed correspondences and the Milnor–Witt K-theory, J. Inst. Math. Jussieu, published online, 2016.
- [19] G. Segal, Categories and cohomology theories, Topology 13 (1974), 293-312.
- [20] A. Suslin, V. Voevodsky, Bloch-Kato conjecture and motivic cohomology with finite coefficients, The Arithmetic and Geometry of Algebraic Cycles (Banff, AB, 1998), NATO Sci. Ser. C Math. Phys. Sci., Vol. 548, Kluwer Acad. Publ., Dordrecht (2000), pp. 117-189.
- [21] V. Voevodsky, \mathbb{A}^1 -Homotopy theory, Doc. Math., Extra Vol. ICM 1998(1), 417-442.
- [22] V. Voevodsky, Notes on framed correspondences, unpublished, 2001. Also available at math.ias.edu/vladimir/files/framed.pdf

DEPARTMENT OF MATHEMATICS, SWANSEA UNIVERSITY, SINGLETON PARK, SWANSEA SA2 8PP, UNITED KINGDOM

E-mail address: g.garkusha@swansea.ac.uk

ST. PETERSBURG BRANCH OF V. A. STEKLOV MATHEMATICAL INSTITUTE, FONTANKA 27, 191023 ST. PETERSBURG, RUSSIA

ST. PETERSBURG STATE UNIVERSITY, DEPARTMENT OF MATHEMATICS AND MECHANICS, UNIVERSITET-SKY PROSPEKT, 28, 198504, PETERHOF, ST. PETERSBURG, RUSSIA

E-mail address: paniniv@gmail.com