

# Size corrected significance tests in seemingly unrelated regressions with autocorrelated errors

Karavias, Yiannis; Tzavalis, Elias; Symeonides, Spyridon

DOI:

[10.1515/jtse-2015-0014](https://doi.org/10.1515/jtse-2015-0014)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Karavias, Y, Tzavalis, E & Symeonides, S 2017, 'Size corrected significance tests in seemingly unrelated regressions with autocorrelated errors', *Journal of Time Series Econometrics*, vol. 9, no. 1.  
<https://doi.org/10.1515/jtse-2015-0014>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Checked 26/07/2016

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Size corrected significance tests in Seemingly Unrelated Regressions with autocorrelated errors

Spyridon D. Symeonides\*    Yiannis Karavias<sup>†</sup>    Elias Tzavalis<sup>‡</sup>

September 8, 2015

## Abstract

Refined asymptotic methods are used to produce degrees-of-freedom-adjusted Edgeworth and Cornish-Fisher size corrections of the  $t$  and  $F$  testing procedures for the parameters of a S.U.R. model with serially correlated errors. The corrected tests follow the Student- $t$  and  $F$  distributions, respectively, with an approximation error of order  $O(\tau^3)$ , where  $\tau = 1/\sqrt{T}$  and  $T$  is the number of time observations. Monte Carlo simulations provide evidence that the size corrections suggested hereby have better finite sample properties, compared to the asymptotic testing procedures (either standard or Edgeworth corrected), which do not adjust for the degrees of freedom.

*Key words:* Linear regression; S.U.R. models; stochastic expansions; asymptotic approximations; AR(1) errors.

*JEL classification:* C10, C12, D24.

---

\*Department of Economics, University of Ioannina, Ioannina 455 00, Greece. *Email:* ssymeoni@cc.uoi.gr

<sup>†</sup>Corresponding Author, School of Economics, University of Birmingham. *Email:* i.karavias@bham.ac.uk

<sup>‡</sup>Department of Economics, Athens University of Economics and Business, 76 Patission Str., 10434 Athens, Greece. *Email:* etzavalis@aueb.gr

# 1 Introduction

The use of refined asymptotic techniques can considerably improve the finite-sample performance of testing procedures in applied econometric research (see, e.g., Ullah (2004), for a survey). These techniques involve the use of Edgeworth expansions which effectively provide higher-order asymptotic approximations of the finite-sample distributions of well known econometric test statistics (see Magdalinos and Symeonides (1995), Magee (1985), Rothenberg (1984b), Symeonides et al. (2007), *inter alia*). In finite samples, there are considerable discrepancies between the actual (sample) and nominal size of many standard testing procedures, employed in econometric literature. These discrepancies are found to be very severe, especially for the generalized linear regression model with a non-scalar covariance matrix of the error terms estimated by the feasible generalized least squares (FGLS), or maximum likelihood (see, e.g., Kiviet and Phillips (1996), Ullah (2004)).

Despite the substantial amount of work on refined asymptotic bias expansions of alternative estimators for the linear regression model or simultaneous equations systems (see, e.g., Iglesias and Phillips (2010, 2012), Kiviet and Phillips (1996), Kiviet et al. (1995), Phillips (2000, 2007), *inter alia*), there are only a few papers applying these methods to conventional tests, like the  $F$  and  $t$ . Rothenberg (1984b, 1988) used Edgeworth expansions in terms of the chi-square and normal distributions to derive general formulae of corrected critical values of the Wald (or  $F$ ) and  $t$  tests, respectively.

In this paper, we derive size corrections of the  $t$  and  $F$  tests for the system of Seemingly Unrelated Regression (S.U.R.) equations with first-order autoregressive error terms, introduced by Parks (1967). The oversizedness of these tests in finite samples can be attributed to two sources: (i) the non-zero cross-correlations of the error terms of the S.U.R. equations, and (ii) the specific dynamic structure of these error terms, i.e., the existence of serial correlation (with possibly distinct autocorrelation coefficients) across the S.U.R. equations.

Since the Edgeworth expansions are not well-defined distribution functions and they may assign negative ‘probabilities’ to the tails of the approximated distributions, the paper suggests using the Cornish-Fisher expansion of the tests

rather than the Edgeworth expansion of their distribution functions (see Cornish and Fisher (1937), Fisher and Cornish (1960), Hill and Davis (1968), Magdalinos (1985), Ogasawara (2012), *inter alia*). The above suggested corrections are asymptotically equivalent, but there are arguments—both theoretical and practical—in favour of the Cornish-Fisher correction: First, the Cornish-Fisher corrected test statistics are theoretically superior because they are proper random variables and their distributions have well-behaved tails; second, since they do not require the calculation of new critical values, they can be readily implemented in applied research based on the publicly available tables of standard distributions.

The paper proposes the use of degrees-of-freedom-adjusted Edgeworth corrected critical values and Cornish-Fisher corrected statistics of the  $t$  and  $F$  tests when the S.U.R. model with serially correlated errors is estimated using the Parks' estimator (see Parks (1967)). These corrections follow the Student- $t$  and  $F$  distributions, respectively, with an approximation error of order  $O(\tau^3)$ , where  $\tau = 1/\sqrt{T}$  and  $T$  is the number of time observations of the sample. The use of degree-of-freedom-adjusted forms of the above tests lead to approximations that are 'locally exact' (see Magdalinos (1985)), which means that the approximate distributions reduce to the exact ones, when the model is sufficiently simplified. These approximations are found to improve the small-sample performance of the tests (see Magdalinos and Symeonides (1995), Symeonides et al. (2007)). To our knowledge, this is the first attempt in the literature to develop analytic size corrected testing procedures for the S.U.R. model with serially correlated errors.

The analytic size corrections suggested by the paper take into account the magnitude of the various nuisance parameters, as well as the way in which they influence the elements of the disturbance covariance matrix. They can be implemented separately to correct for the non-zero cross-correlations of the error terms, or their serial correlation effects, or the combination of the above.

The paper is organised as follows. Section 2 provides some preliminary notations. Section 3 presents the S.U.R. model and the assumptions needed in our expansions. Analytic formulae for the locally exact Edgeworth and Cornish-Fisher second-order size corrections of the  $t$  and  $F$  test statistics are derived

in Section 4. Section 5 conducts out a Monte Carlo simulation evaluating the performance of the suggested corrected tests. Finally, Section 6 concludes the paper. Proofs of the results of the paper are given in the Appendix.

## 2 Preliminary notation

Throughout the paper, we use the  $tr$ ,  $vec$ ,  $\otimes$ , and matrix differentiation notation as defined in Dhrymes (1978, pages 518–540), and for any two indices  $i$  and  $j$ , we denote Kronecker's delta as  $\delta_{ij}$ . Moreover, any  $(n \times m)$  matrix  $L$  with elements  $l_{ij}$  is denoted as

$$L = [(l_{ij})_{i=1, \dots, n; j=1, \dots, m}],$$

with obvious modifications for vectors and square matrices. If  $l_{ij}$  are  $(n_i \times m_j)$  matrices, then  $L$  is the  $(\sum n_i \times \sum m_j)$  partitioned matrix with submatrices  $l_{ij}$ . The following matrices:

$$P_X = X(X'X)^{-1}X', \quad \bar{P}_X = I - P_X = I - X(X'X)^{-1}X'$$

denote the orthogonal projectors into the spaces spanned by the columns of the matrix  $X$  and its orthogonal complement, respectively. Finally, for any stochastic quantity (scalar, vector, or matrix) we use the symbol  $\mathcal{E}(\cdot)$  to denote the expectation operator.

## 3 The model

Consider a S.U.R. system of  $M$  contemporaneously correlated regression equations of the form

$$y_\mu = X_\mu \beta_\mu + u_\mu \quad (\mu = 1, \dots, M), \quad (1)$$

where  $y_\mu$  are  $(T \times 1)$  vectors of observations on the dependent variables,  $X_\mu$  are  $(T \times n_\mu)$  matrices of observations on sets of  $n_\mu$  non-stochastic regressors,  $\beta_\mu$  are  $(n_\mu \times 1)$  vectors of parameters to be estimated and  $u_\mu$  are  $(T \times 1)$  vectors of non-observable serially correlated stochastic error terms of the  $\mu$ -th equation, defined as  $u_{t\mu}$  ( $t = 1, \dots, T$ ). These terms are generated by the following stationary first-order autoregressive (AR(1)) process:

$$u_{t\mu} = \rho_\mu u_{(t-1)\mu} + \varepsilon_{t\mu}, \quad -1 < \rho_\mu < 1 \quad (t = 1, \dots, T; \mu = 1, \dots, M), \quad (2)$$

where  $\varepsilon_{t\mu}$  are normally distributed innovations. For any two indices  $\mu, \mu' = 1, \dots, M$ , we have  $\mathcal{E}(\varepsilon_{t\mu}) = 0$ , for all  $t$ . Moreover, for  $t \neq 1$  or  $t' \neq 1$ , the covariance between two innovations  $\varepsilon_{t\mu}$  and  $\varepsilon_{t'\mu'}$  is given as  $\mathcal{E}(\varepsilon_{t\mu}\varepsilon_{t'\mu'}) = \delta_{tt'}\sigma_{\mu\mu'}$ . For  $t = t' = 1$  and  $\mu, \mu' = 1, \dots, M$ ,  $\mathcal{E}(\varepsilon_{t\mu}\varepsilon_{t'\mu'})$  becomes

$$\mathcal{E}(\varepsilon_{1\mu}\varepsilon_{1\mu'}) = \sigma_{\mu\mu'}(1 - \rho_\mu^2)^{1/2}(1 - \rho_{\mu'}^2)^{1/2}/(1 - \rho_\mu\rho_{\mu'}) \quad (3)$$

(see Parks (1967, pages 507–508)). In addition to assumption  $\rho_\mu \in (-1, 1)$ , stationarity of AR(1) processes (2) implies the following relationships on the initial conditions of the error terms of the S.U.R. equations:

$$u_{1\mu} = (1 - \rho_\mu^2)^{-1/2}\varepsilon_{1\mu} \quad (t = 1; \mu = 1, \dots, M). \quad (4)$$

These relationships imply that, for all  $t = 1, \dots, T$  and  $\mu, \mu' = 1, \dots, M$ , the error terms  $u_{t\mu}$  satisfy the following conditions:

$$\mathcal{E}(u_{t\mu}) = 0, \quad \mathcal{E}(u_{t\mu}^2) = \sigma_{\mu\mu}/(1 - \rho_\mu^2), \quad \mathcal{E}(u_{t\mu}u_{t\mu'}) = \sigma_{\mu\mu'}/(1 - \rho_\mu\rho_{\mu'}). \quad (5)$$

Let  $n = \sum_{\mu=1}^M n_\mu$ , and define the  $(MT \times 1)$  vectors  $y$  and  $u$ , the  $(n \times 1)$  vector  $\beta$  and the  $(MT \times n)$  block diagonal matrix  $X$  as follows:

$$\begin{aligned} y &= [(y_\mu)_{\mu=1, \dots, M}], \quad u = [(u_\mu)_{\mu=1, \dots, M}], \\ \beta &= [(\beta_\mu)_{\mu=1, \dots, M}], \\ X &= [(\delta_{\mu\mu'}X_\mu)_{\mu, \mu'=1, \dots, M}]. \end{aligned} \quad (6)$$

Then, the system of equations (1) can be written in a matrix form as follows:

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_M \end{bmatrix} = \begin{bmatrix} X_1 & 0 & \cdots & 0 \\ 0 & X_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & X_M \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_M \end{bmatrix} + \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_M \end{bmatrix}, \quad (7)$$

or more compactly as

$$y = X\beta + u. \quad (8)$$

To derive size corrected significance tests for the elements of the vector  $\beta$ , the above representations of the S.U.R. system will be written in an autocorrelation-free form, after applying appropriate transformations on  $y$ ,  $X$  and  $u$ . Following

Parks (1967), define the  $(T \times T)$  matrices  $P_\mu$  and  $R^{\mu\mu'}$  as follows:

$$P_\mu = \begin{bmatrix} (1 - \rho_\mu^2)^{-\frac{1}{2}} & 0 & 0 & \cdots & 0 \\ (1 - \rho_\mu^2)^{-\frac{1}{2}} \rho_\mu & 1 & 0 & \cdots & 0 \\ (1 - \rho_\mu^2)^{-\frac{1}{2}} \rho_\mu^2 & \rho_\mu & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ (1 - \rho_\mu^2)^{-\frac{1}{2}} \rho_\mu^{T-1} & \rho_\mu^{T-2} & \rho_\mu^{T-3} & \cdots & 1 \end{bmatrix}, \quad R^{\mu\mu'} = P_\mu^{-1} P_{\mu'}^{-1}, \quad (9)$$

and the following  $(MT \times MT)$  block diagonal matrix

$$P = [(\delta_{\mu\mu'} P_\mu)_{\mu, \mu'=1, \dots, M}]. \quad (10)$$

Then, (2) implies that the  $(T \times 1)$  random vectors  $u_\mu$  can be written as

$$u_\mu = P_\mu \varepsilon_\mu \quad (\mu = 1, \dots, M), \quad (11)$$

where  $\varepsilon_\mu$  are  $(T \times 1)$  random vectors with non-autocorrelated elements  $\varepsilon_{t\mu}$ , i.e.,

$$\varepsilon_\mu = [(\varepsilon_{t\mu})_{t=1, \dots, T; \mu=1, \dots, M}]. \quad (12)$$

As in (11), consider the  $(T \times 1)$  vectors  $y_{\mu*}$  and  $(T \times n_\mu)$  matrices  $X_{\mu*}$ , with non-autocorrelated elements, satisfying the following relations:

$$y_{\mu*} = P_\mu^{-1} y_\mu, \quad X_{\mu*} = P_\mu^{-1} X_\mu, \quad (13)$$

and define the  $(MT \times 1)$  vector  $y_*$  and  $(MT \times n)$  block diagonal matrix  $X_*$  as follows:

$$y_* = [(y_{\mu*})_{\mu=1, \dots, M}], \quad X_* = [(\delta_{\mu\mu'} X_{\mu*})_{\mu, \mu'=1, \dots, M}]. \quad (14)$$

Then, premultiplying the  $\mu$ -th equation of (7) by  $P_\mu^{-1}$ , we can derive the following S.U.R. model with non-autocorrelated error terms:

$$\begin{bmatrix} y_{1*} \\ y_{2*} \\ \vdots \\ y_{M*} \end{bmatrix} = \begin{bmatrix} X_{1*} & 0 & \cdots & 0 \\ 0 & X_{2*} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & X_{M*} \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_M \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_M \end{bmatrix} \quad (15)$$

(see Zellner (1962, 1963), Zellner and Huang (1962), Zellner and Theil (1962)).

In more compact form, this model can be written as

$$y_* = X_* \beta + \varepsilon, \quad (16)$$

where  $y_* = P^{-1}y$ ,  $X_* = P^{-1}X$  and  $\varepsilon = P^{-1}u$ . The above representation of the S.U.R. system implies that the  $(MT \times 1)$  error vector  $u$  in (8) is normally distributed with mean and variance-covariance matrix given as follows:

$$\mathcal{E}(u) = 0, \quad \mathcal{E}(uu') = \Omega^{-1} = P\mathcal{E}(\varepsilon\varepsilon')P' = P(\Sigma \otimes I_T)P', \quad (17)$$

where

$$\Sigma = [(\sigma_{\mu\mu'})_{\mu,\mu'=1, \dots, M}]. \quad (18)$$

The last relationship implies that

$$\Omega = P'^{-1}(\Sigma^{-1} \otimes I_T)P^{-1} \quad (19)$$

is a function of the  $((M + M^2) \times 1)$  parameter vector  $\gamma = (\varrho', \varsigma')'$ , where  $\varrho = (\rho_1, \dots, \rho_M)'$  is the  $(M \times 1)$  vector of autocorrelation coefficients in (2) and the  $(M^2 \times 1)$  vector  $\varsigma = \text{vec}(\Sigma^{-1}) \in \mathcal{L} = \mathbb{R}^{M^2} - \mathfrak{U}$ , where  $\mathfrak{U}$  is the subspace of  $\mathbb{R}^{M^2}$  in which  $\Sigma$  is not positive definite. After defining the composite index

$$(\mu\mu') = \mu + M(\mu' - 1) \quad ((\mu\mu') = 1, \dots, M^2), \quad (20)$$

for any two indices  $\mu, \mu' = 1, \dots, M$ , it can be easily seen that the  $(\mu\mu')$ -th element of vector  $\varsigma$ , denoted as  $\varsigma_{(\mu\mu')}$ , is actually the  $(\mu, \mu')$ -th element of matrix  $\Sigma^{-1}$ , denoted as  $\sigma^{\mu\mu'}$ .

The system of equations (16) (or (15)) can be seen as the vectorization outcome of the following form of the S.U.R. model of  $M$  equations:

$$Y_* = ZB + E, \quad (21)$$

where  $Y_*$  and  $E$  are  $(T \times M)$  random matrices defined as

$$y_* = \text{vec}(Y_*), \quad \varepsilon = \text{vec}(E), \quad (22)$$

respectively, where the rows of matrix  $E$  are  $\mathcal{N}_M(0, \Sigma)$  random vectors and  $B$  is a  $(K \times M)$  matrix whose columns, denoted as  $b_\mu$ , are defined as

$$b_\mu = \Psi_\mu \beta_\mu \quad (\mu = 1, \dots, M), \quad (23)$$

where  $\Psi_\mu$  are  $(K \times n_\mu)$  known submatrices of the  $(MK \times n)$  block diagonal matrix

$$\Psi = [(\delta_{\mu\mu'} \Psi_\mu)_{\mu,\mu'=1, \dots, M}]. \quad (24)$$



Finally,  $Z$  is a  $(T \times K)$  matrix with non-autocorrelated columns, defined by the following relationship:

$$\begin{aligned} X_* &= [(\delta_{\mu\mu'} X_{\mu*})_{\mu,\mu'=1, \dots, M}] = [(\delta_{\mu\mu'} Z \Psi_\mu)_{\mu,\mu'=1, \dots, M}] \\ &= [(\delta_{\mu\mu'} Z)_{\mu,\mu'=1, \dots, M}] [(\delta_{\mu\mu'} \Psi_\mu)_{\mu,\mu'=1, \dots, M}] \\ &= (I_M \otimes Z) \Psi. \end{aligned} \quad (25)$$

The above representation of the S.U.R. model, given by (21), will facilitate the expansions needed in our derivations of the size corrected tests suggested in the paper.

### 3.1 Assumptions

To carry out our expansions, it would be theoretically convenient to introduce a reparameterization of the error covariance matrix of model (8) as follows:

$$y = X\beta + \sigma u, \quad \sigma > 0, \quad u \sim \mathcal{N}_{MT}(0, \Omega^{-1}), \quad (26)$$

assuming that parameter  $\sigma^2$  can be estimated separately from the rest terms of the covariance matrix  $\Omega^{-1}$  of vector  $u$ .<sup>1</sup>

For the derivation of our size corrected tests, we need to make a number of assumptions on the elements of matrix  $\Omega$ , which is the inverse of the variance-covariance matrix of the error vector  $u$ . To this end, we denote as  $\Omega_i$ ,  $\Omega_{ij}$ , etc., the  $(MT \times MT)$  matrices of first-, second- and higher-order derivatives, respectively, of the elements of matrix  $\Omega$  with respect to the elements of the  $((M + M^2) \times 1)$  vector of nuisance parameters  $\gamma = (\varrho', \varsigma')'$ . For any estimator of  $\gamma$ , define the  $((1 + M + M^2) \times 1)$  vector  $\delta$ , with elements

$$\delta_0 = \frac{\hat{\sigma}^2 - 1}{\tau}, \quad \delta_{\rho_\mu} = \frac{\hat{\rho}_\mu - \rho_\mu}{\tau}, \quad \delta_{\varsigma_{(\mu\mu')}} = \frac{\hat{\varsigma}_{(\mu\mu')} - \varsigma_{(\mu\mu')}}{\tau}, \quad (27)$$

where  $\mu = 1, \dots, M$ ,  $(\mu\mu') = 1, \dots, M^2$  and  $\tau = 1/\sqrt{T}$  is the ‘asymptotic

---

<sup>1</sup>The nuisance parameters  $\sigma$  and  $\gamma$  can be simultaneously identified under the restriction  $\sigma = 1$ , which implies that the estimate of matrix  $\Sigma$ , denoted as  $\hat{\Sigma}$ , is accurate, up to a multiplicative factor. This is not true in samples with small time dimension. A convenient method to estimate  $\sigma$  is through the following feasible generalized least squares (GL) estimator

$$\hat{\sigma}_{GL} = \left[ (y - X\hat{\beta})' \left( \hat{P}_{GL}^{-1} (\hat{\Sigma}_{GL}^{-1} \otimes I_T) \hat{P}_{GL}^{-1} \right) (y - X\hat{\beta}) / (MT - n) \right]^{1/2},$$

where  $\hat{\beta}$  is the feasible GL estimator based on any consistent estimators of  $\Sigma^{-1}$  and  $P^{-1}$ .

scale' of our second-order stochastic expansions. Then, our size corrected tests can be derived based on the following assumption.

**Assumption 1:**

- (i) The elements of matrices  $\Omega$  and  $\Omega^{-1}$  are bounded for all  $T$ , all vectors  $\varrho$  with elements  $\rho_\mu \in (-1, 1)$ , and all vectors  $\varsigma \in \mathcal{L}$ . Moreover, the following matrices:

$$A = X'\Omega X/T, \quad F = X'X/T, \quad \Gamma = Z'Z/T \quad (28)$$

converge to non-singular limits, as  $T \rightarrow \infty$ .

- (ii) Up to the fourth order, the partial derivatives of the elements of  $\Omega$  with respect to the elements of  $\varrho$  and  $\varsigma$ , are bounded for all  $T$ , all vectors  $\varrho$  with elements in the interval  $(-1, 1)$ , and all vectors  $\varsigma \in \mathcal{L}$ .
- (iii) The estimators  $\hat{\varrho}$  and  $\hat{\varsigma}$  are even functions of  $u$ , and they are functionally unrelated to the parameter vector  $\beta$ . As a result, they can be written as functions of  $X$ ,  $Z$ , and  $u$  only.
- (iv) The vector of nuisance parameters  $\delta$  admits a stochastic expansion of the form

$$\begin{aligned} \delta &= \left[ \delta_0, [(\delta_{\rho_\mu})_{\mu=1, \dots, M}]', [(\delta_{\varsigma_{(\mu\mu')}})_{(\mu\mu')=1, \dots, M^2}]' \right]' \\ &= d_1 + \tau d_2 + \omega(\tau^2), \end{aligned} \quad (29)$$

where the order of magnitude  $\omega(\cdot)$ , defined in the Appendix, has the same operational properties as order  $O(\cdot)$ . Moreover, the expectations

$$\mathcal{E}(d_1 d_1'), \quad \mathcal{E}(\sqrt{T}d_1 + d_2) \quad (30)$$

exist and have finite limits, as  $T \rightarrow \infty$ .

The first two conditions of Assumption 1 imply that the following matrices:

$$A_i = X'\Omega_i X/T, \quad A_{ij} = X'\Omega_{ij} X/T, \quad A_{ij}^* = X'\Omega_i \Omega^{-1} \Omega_j X/T \quad (31)$$

are bounded. Thus, according to Magdalinos (1992), the Taylor series expansion of  $\beta$  constitutes a stochastic expansion. Since the vectors of nuisance parameters  $\varrho$  and  $\varsigma$  are functionally unrelated to  $\beta$ , condition (iii) of Assumption 1 is satisfied for a wide class of estimators  $\hat{\varrho}$  and  $\hat{\varsigma}$ , including the maximum likelihood

estimators and the simple or iterative estimators based on the regression residuals (see Breusch (1980), Rothenberg (1984a)). Note that we *need not* assume that estimators  $\hat{\varrho}$  and  $\hat{\varsigma}$  are asymptotically efficient.

Further, conditions (i)–(iv) of Assumption 1 should be satisfied by all the estimators of vectors  $\varrho$  and  $\varsigma$ , considered in the paper. The estimators of the elements of vector  $\varrho$ , i.e.,  $\rho_\mu$  ( $\mu = 1, \dots, M$ ) include the following: the least squares (LS), Durbin-Watson (DW), generalized least squares (GL), Prais-Winsten (PW) and maximum likelihood (ML).<sup>2</sup> The elements of vector  $\varsigma = \text{vec}(\Sigma^{-1})$  can be estimated by

$$\hat{\varsigma} = \text{vec} \left[ (Y_* - Z\hat{B})'(Y_* - Z\hat{B})/T \right]^{-1}, \quad (32)$$

where  $\hat{B}$  is any consistent estimator of the matrix of parameters  $B$  of regression model (21). Consistent estimators of  $B$  include the unrestricted and restricted least squares (denoted as UL and RL, respectively), the simple and iterative generalized least squares (denoted as GL and IG, respectively) and the maximum likelihood (ML) estimators.<sup>3</sup>

To present the expansions suggested in the paper, expectations  $\mathcal{E}(d_1 d_1')$  and

---

<sup>2</sup>The closed forms of these estimators of  $\rho_\mu$ , for all  $\mu$ , are given as follows:

(i) LS:

$$\tilde{\rho}_\mu = \sum_{t=2}^T \tilde{u}_{t\mu} \tilde{u}_{(t-1)\mu} / \sum_{t=1}^T \tilde{u}_{t\mu}^2,$$

where  $\tilde{u}_{t\mu}$  are the LS residuals of regression model (1).

(ii) DW:

$$\hat{\rho}_\mu^{(DW)} = 1 - (DW/2),$$

where the  $DW$  is the Durbin-Watson statistic.

(iii) GL:

$$\hat{\rho}_\mu = \sum_{t=2}^T \hat{u}_{t\mu} \hat{u}_{(t-1)\mu} / \sum_{t=1}^T \hat{u}_{t\mu}^2,$$

where  $\hat{u}_{t\mu}$  denote the GL estimates of  $u_{t\mu}$ , based on the autocorrelation-correction of regression model (1), for all  $\mu$ , using any asymptotically efficient estimator of  $\rho_\mu$ .

(iv) PW: This estimator of  $\rho_\mu$ , denoted as  $\hat{\rho}_\mu^{(PW)}$ , together with the PW estimator of  $\beta$ , denoted as  $\hat{\beta}_\mu^{(PW)}$ , minimize the sum of squared GL residuals (Prais and Winsten (1954)).

(v) ML: This estimator, denoted as  $\hat{\rho}_\mu^{(ML)}$ , satisfies a cubic equation with coefficients defined in terms of the ML residuals (Beach and MacKinnon (1978)).

<sup>3</sup>The closed forms of these estimators of  $B$  are given as follows:

(i) UL:

$$\hat{B}_{(UL)} = (Z'Z)^{-1}Z'Y_*.$$

$\mathcal{E}(\sqrt{T}d_1 + d_2)$  will be defined as follows:

$$\lim_{T \rightarrow \infty} \mathcal{E}(d_1 d_1') = \begin{bmatrix} \lambda_0 & \lambda'_\varrho & \lambda'_\varsigma \\ \lambda_\varrho & \Lambda_\varrho & \Lambda'_{\varrho\varsigma} \\ \lambda_\varsigma & \Lambda_{\varrho\varsigma} & \Lambda_\varsigma \end{bmatrix} \quad \text{and} \quad \lim_{T \rightarrow \infty} \mathcal{E}(\sqrt{T}d_1 + d_2) = \begin{bmatrix} \kappa_0 \\ \kappa_\varrho \\ \kappa_\varsigma \end{bmatrix}, \quad (33)$$

respectively, where  $\lambda_0$  and  $\kappa_0$  are scalars,  $\lambda_\varrho$  and  $\kappa_\varrho$  are  $(M \times 1)$  vectors,  $\lambda_\varsigma$  and  $\kappa_\varsigma$  are  $(M^2 \times 1)$  vectors,  $\Lambda_\varrho$  is a  $(M \times M)$  matrix,  $\Lambda_\varsigma$  is a  $(M^2 \times M^2)$  matrix and  $\Lambda_{\varrho\varsigma}$  is a  $(M^2 \times M)$  matrix. The following partitions of the above matrix and vector will be of use in the paper:

$$\begin{bmatrix} \lambda_0 & \lambda' \\ \lambda & \Lambda \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} \kappa_0 \\ \kappa \end{bmatrix}, \quad (34)$$

where

$$\Lambda = \begin{bmatrix} \Lambda_\varrho & \Lambda'_{\varrho\varsigma} \\ \Lambda_{\varrho\varsigma} & \Lambda_\varsigma \end{bmatrix}, \quad \lambda = \begin{bmatrix} \lambda_\varrho \\ \lambda_\varsigma \end{bmatrix} \quad \text{and} \quad \kappa = \begin{bmatrix} \kappa_\varrho \\ \kappa_\varsigma \end{bmatrix}, \quad (35)$$

where  $\Lambda$  is a  $((M + M^2) \times (M + M^2))$  matrix, and  $\lambda$  and  $\kappa$  are  $((M + M^2) \times 1)$  vectors. The elements of the vectors and matrices in (33), (34) and (35) can be interpreted as ‘measures’ of the accuracy of the expansions of estimators  $\hat{\sigma}^2$ ,  $\hat{\rho}_\mu$  and  $\hat{\varsigma}_{(\mu\mu')}$  around the true values of the corresponding parameters.

## 4 Size corrected test statistics

In this section, we derive size corrected  $t$ , Wald and  $F$  test statistics, as well as the second-order approximations of their distributions based on the conditions of Assumption 1. The versions of the test statistics which adjust for the degrees

---

(ii) RL:

$$vec(\hat{B}_{(RL)}) = \Psi(X'_* X_*)^{-1} X'_* y_*.$$

(iii) GL:

$$vec(\hat{B}_{(GL)}) = \Psi \left[ X'_* (\hat{\Sigma}_I^{-1} \otimes I_T) X_* \right]^{-1} X'_* (\hat{\Sigma}_I^{-1} \otimes I_T) y_*,$$

where  $\hat{\Sigma}_I^{-1}$  is the UL or RL estimator of  $\Sigma$ .

(iv) IG: This estimator, denoted as  $\hat{B}_{(IG)}$ , is computed by iterative implementation of the GL estimator.

(v) ML: This estimator, denoted as  $\hat{B}_{(ML)}$ , can be computed by iterating the GL estimation process up to convergence (Dhrymes (1971)).

of freedom, namely the Student- $t$  and  $F$ , are locally exact. That is, if the vector of parameters  $\gamma = (\varrho', \varsigma')'$  is known to belong to a ball of radius  $\vartheta$ , then the approximate distributions of these test statistics become exact, as  $\vartheta \rightarrow 0$ .

The analytic size corrections developed in this section can provide size corrections to either the non-zero cross-correlations of the error terms or their serial correlation effects. The part of the size corrections corresponding to the serial correlation effects constitutes an extension of the results in Magdalinos and Symeonides (1995) to the multiple equation framework. On the other hand, the part of the size corrections due to the non-zero cross-correlations constitutes a completely genuine contribution to the literature, which can be readily implemented to correct the size of the  $t$  and  $F$  tests in the standard Zellner's S.U.R. model (see Zellner (1962)) alone.

#### 4.1 The $t$ test

Let the elements of the  $(n \times 1)$  vector  $e$  and scalar  $e_0$  be known quantities. Testing any null hypothesis of the form

$$H_0 : e' \beta = e_0 \quad (36)$$

against its one-sided alternatives, can be based upon the following  $t$  statistic:

$$t = (e' \beta - e_0) / \left[ \hat{\sigma}^2 e' (X' \hat{\Omega} X)^{-1} e \right]^{1/2}, \quad (37)$$

which is adjusted for the degrees of freedom of the Student- $t$  distribution.

For the derivation of the suggested asymptotic expansions, we define the  $((M + M^2) \times 1)$  vector  $l$  and the  $((M + M^2) \times (M + M^2))$  matrix  $L$  as follows:

$$l = \left[ [(l_{\rho_\mu})_{\mu=1, \dots, M}]', [(l_{\varsigma_{(\mu\mu')}})_{(\mu\mu')=1, \dots, M^2}]' \right]', \quad (38)$$

$$L = \begin{bmatrix} [(l_{\rho_\mu \rho_{\mu'}})_{\mu, \mu'=1, \dots, M}] & [(l_{\rho_\mu \varsigma_{(\nu\nu')}})_{\substack{\mu=1, \dots, M; \\ (\nu\nu')=1, \dots, M^2}}] \\ [(l_{\varsigma_{(\nu\nu')} \rho_\mu})_{\substack{(\nu\nu')=1, \dots, M^2; \\ \mu=1, \dots, M}}] & [(l_{\varsigma_{(\mu\mu') \varsigma_{(\nu\nu')}}})_{\substack{(\mu\mu')=1, \dots, M^2; \\ (\nu\nu')=1, \dots, M^2}}] \end{bmatrix}, \quad (39)$$

where the elements of vector  $l$  and matrix  $L$  are defined below:

$$\begin{aligned} l_{\rho_\mu} &= h' G A_{\rho_\mu} G h, & l_{\varsigma_{(\mu\mu')}} &= h' G A_{\varsigma_{(\mu\mu')}} G h, \\ l_{\rho_\mu \rho_{\mu'}} &= h' G C_{\rho_\mu \rho_{\mu'}} G h, & l_{\rho_\mu \varsigma_{(\nu\nu')}} &= h' G C_{\rho_\mu \varsigma_{(\nu\nu')}} G h, \\ l_{\varsigma_{(\nu\nu')} \rho_\mu} &= h' G C_{\varsigma_{(\nu\nu')} \rho_\mu} G h, & l_{\varsigma_{(\mu\mu') \varsigma_{(\nu\nu')}}} &= h' G C_{\varsigma_{(\mu\mu') \varsigma_{(\nu\nu')}}} G h, \end{aligned} \quad (40)$$

where  $G = A^{-1} = (X'\Omega X/T)^{-1}$  is a  $(n \times n)$  matrix,  $h = e/(e'Ge)^{1/2}$  is a  $(n \times 1)$  vector and

$$\begin{aligned} C_{\rho_\mu \rho_{\mu'}} &= A_{\rho_\mu \rho_{\mu'}}^* - 2A_{\rho_\mu} G A_{\rho_{\mu'}} + A_{\rho_\mu \rho_{\mu'}}/2, \\ C_{\rho_\mu \varsigma_{(\nu\nu')}} &= A_{\rho_\mu \varsigma_{(\nu\nu')}}^* - 2A_{\rho_\mu} G A_{\varsigma_{(\nu\nu')}} + A_{\rho_\mu \varsigma_{(\nu\nu')}}/2, \\ C_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} &= A_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}}^* - 2A_{\varsigma_{(\mu\mu')}} G A_{\varsigma_{(\nu\nu')}} + A_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}}/2, \end{aligned} \quad (41)$$

with obvious modifications for  $C_{\varsigma_{(\nu\nu')} \rho_\mu}$ .

The next two theorems give alternative Edgeworth approximations of the distribution function of the  $t$  statistic, given in (37), in terms of the normal and Student- $t$  distributions, respectively.

**Theorem 1.** *The distribution function of the  $t$  statistic (37), under the null hypothesis (36), admits the Edgeworth expansion*

$$\Pr\{t \leq x\} = I(x) - \frac{\tau^2}{2} \left[ (p_1 + \frac{1}{2}) + (p_2 + \frac{1}{2}) x^2 \right] xi(x) + O(\tau^3), \quad (42)$$

where  $I(\cdot)$  and  $i(\cdot)$  are the standard normal distribution and density functions, respectively, and scalars  $p_1$  and  $p_2$  can be calculated as follows:

$$p_1 = \text{tr}(\Lambda L) + \frac{l'\Lambda l}{4} + l'(\kappa + \frac{\lambda}{2}) - \kappa_0 + \frac{\lambda_0 - 2}{4}, \quad p_2 = \frac{l'\Lambda l - 2l'\lambda + \lambda_0 - 2}{4}. \quad (43)$$

Analytic formulae for the computation of scalars  $\lambda_0$ ,  $\kappa_0$ , and the elements of  $\lambda$ ,  $\kappa$ ,  $\Lambda$ ,  $l$  and  $L$  are given in the Appendix (see Lemmas A.15 and A.17).

Instead of using the Edgeworth expansion (42), we can approximate the distribution function of the  $t$  statistic in terms of the Student- $t$  distribution as follows:

**Theorem 2.** *The distribution function of the  $t$  statistic (37), under the null hypothesis (36), admits the Edgeworth expansion*

$$\Pr\{t \leq x\} = I_{MT-n}(x) - \frac{\tau^2}{2} [p_1 + p_2 x^2] xi_{MT-n}(x) + O(\tau^3), \quad (44)$$

where  $I_{MT-n}(\cdot)$  and  $i_{MT-n}(\cdot)$  are the Student- $t$  distribution and density functions, respectively, with  $MT - n$  degrees of freedom, and scalars  $p_1$  and  $p_2$  are defined in (43).

Theorem 1 implies that we can calculate the Edgeworth corrected  $\alpha\%$  critical value of the  $t$  statistic (37) as

$$n_\alpha^* = n_\alpha + \frac{\tau^2}{2} \left[ (p_1 + \frac{1}{2}) + (p_2 + \frac{1}{2}) n_\alpha^2 \right] n_\alpha, \quad (45)$$

based on the  $\alpha\%$  significant point of the standard normal distribution, denoted as  $n_\alpha$ . Similarly, based on Theorem 2, we can calculate the Edgeworth corrected  $\alpha\%$  critical value of the  $t$  statistic (37) as

$$t_\alpha^* = t_\alpha + \frac{\tau^2}{2} [p_1 + p_2 t_\alpha^2] t_\alpha, \quad (46)$$

using the  $\alpha\%$  significant point of the Student- $t$  distribution, denoted as  $t_\alpha$ .

The Edgeworth approximation employed by Theorems 1 and 2 to obtain the size corrected critical values  $n_\alpha^*$  and  $t_\alpha^*$  is not a proper distribution function, as it may assign negative ‘probabilities’ in the tails of the approximate distribution. To overcome this problem, we can use a Cornish-Fisher expansion. This corrects the test statistics of interest, instead of their critical values. The Cornish-Fisher expansion is simply the inversion of the Edgeworth correction of the critical values and, thus, it is expected to have very similar properties around the mean of the approximate distribution. However, at the tails of this distribution, which are important for inference, the properties of the Cornish-Fisher expansion are different. In fact, the Cornish-Fisher size corrected statistics constitute random variables with well-behaved tails, and thus they *do not* assign negative ‘probabilities’ at the tails of their distributions.

The Cornish-Fisher corrected  $t$  statistic for testing null hypothesis (36) is given in the following theorem.

**Theorem 3.** *The Cornish-Fisher size corrected  $t$  statistic*

$$t_* = t - \frac{\tau^2}{2} [p_1 + p_2 t^2] t \quad (47)$$

*is distributed, under the null hypothesis (36), as a Student- $t$  random variable with  $MT - n$  degrees of freedom, with an approximation error of order  $O(\tau^3)$ .*

The Cornish-Fisher size corrected  $t$  statistic  $t_*$ , given by equation (47), can be readily used, in practice, to test null hypothesis (36) against its one-sided alternatives. This can be done by using the standard tables of the Student- $t$  distribution with  $MT - n$  degrees of freedom.

## 4.2 The Wald and $F$ tests

Let  $H$  be a  $(m \times n)$  matrix of rank  $m$  with known elements and  $h_0$  be a known  $(m \times 1)$  vector. Testing any null hypothesis of the form

$$H_0 : H\beta = h_0 \quad (48)$$

against all possible alternatives, can be based upon the Wald statistic

$$w = (H\hat{\beta} - h_0)' \left[ H(X'\hat{\Omega}X)^{-1}H' \right]^{-1} (H\hat{\beta} - h_0)/\hat{\sigma}^2, \quad (49)$$

or the familiar  $F$  statistic

$$F = (H\hat{\beta} - h_0)' \left[ H(X'\hat{\Omega}X)^{-1}H' \right]^{-1} (H\hat{\beta} - h_0)/m\hat{\sigma}^2, \quad (50)$$

which is adjusted for the degrees of freedom of the  $F$  distribution.

For the derivation of the suggested asymptotic expansions, we define the  $(n \times n)$  matrix

$$Q = H'(HGH')^{-1}H, \quad (51)$$

and we partition the  $(n \times n)$  matrices  $G = A^{-1} = (X'\Omega X/T)^{-1}$  and  $\Xi = GQG$  and the  $(n \times 1)$  vector  $h$  as follows:

$$G = [(G_{ij})_{i,j=1, \dots, M}], \quad \Xi = [(\Xi_{ij})_{i,j=1, \dots, M}], \quad h = [(h_i)_{i=1, \dots, M}], \quad (52)$$

where  $G_{ij}$  and  $\Xi_{ij}$  are the  $(i, j)$ -th  $(n_i \times n_j)$  submatrices of  $G$  and  $\Xi$ , respectively, and  $h_i = e_i/(e'Ge)^{1/2}$  is the  $i$ -th  $(n_i \times 1)$  subvector of  $h$ , where  $e_i$  is the corresponding  $i$ -th  $(n_i \times 1)$  subvector of the  $(n \times 1)$  vector  $e$ .

Next, define the  $((M + M^2) \times 1)$  vector  $c$ , and the  $((M + M^2) \times (M + M^2))$  matrices  $C$  and  $D_*$  as follows:

$$c = \left[ [(c_{\rho_\mu})_{\mu=1, \dots, M}]', [(c_{\varsigma_{(\mu\mu')}})_{(\mu\mu')=1, \dots, M^2}]' \right]', \quad (53)$$

$$C = \begin{bmatrix} [(c_{\rho_\mu \rho_{\mu'}})_{\mu, \mu'=1, \dots, M}] & [(c_{\rho_\mu \varsigma_{(\nu\nu')}})_{\substack{\mu=1, \dots, M; \\ (\nu\nu')=1, \dots, M^2}}] \\ [(c_{\varsigma_{(\nu\nu')} \rho_\mu})_{\substack{(\nu\nu')=1, \dots, M^2; \\ \mu=1, \dots, M}}] & [(c_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}})_{\substack{(\mu\mu')=1, \dots, M^2; \\ (\nu\nu')=1, \dots, M^2}}] \end{bmatrix} \quad (54)$$

and

$$D_* = \begin{bmatrix} [(d_{\rho_\mu \rho_{\mu'}})_{\mu, \mu'=1, \dots, M}] & [(d_{\rho_\mu \varsigma_{(\nu\nu')}})_{\substack{\mu=1, \dots, M; \\ (\nu\nu')=1, \dots, M^2}}] \\ [(d_{\varsigma_{(\nu\nu')} \rho_\mu})_{\substack{(\nu\nu')=1, \dots, M^2; \\ \mu=1, \dots, M}}] & [(d_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}})_{\substack{(\mu\mu')=1, \dots, M^2; \\ (\nu\nu')=1, \dots, M^2}}] \end{bmatrix}, \quad (55)$$



where the elements of vector  $c$  and matrices  $C$  and  $D_*$  are defined as follows:

$$\begin{aligned}
c_{\rho_\mu} &= \text{tr}(A_{\rho_\mu} \Xi), \quad c_{\rho_\mu \rho_{\mu'}} = \text{tr}(C_{\rho_\mu \rho_{\mu'}} \Xi), \\
c_{\rho_\mu \varsigma_{(\nu\nu')}} &= \text{tr}(C_{\rho_\mu \varsigma_{(\nu\nu')}} \Xi), \\
c_{\varsigma_{(\mu\mu')}} &= \text{tr}(A_{\varsigma_{(\mu\mu')}} \Xi), \quad c_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} = \text{tr}(C_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} \Xi), \\
d_{\rho_\mu \rho_{\mu'}} &= \text{tr}(D_{*\rho_\mu \rho_{\mu'}} \Xi), \quad d_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} = \text{tr}(D_{*\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} \Xi), \\
d_{\rho_\mu \varsigma_{(\nu\nu')}} &= \text{tr}(D_{*\rho_\mu \varsigma_{(\nu\nu')}} \Xi),
\end{aligned} \tag{56}$$

where

$$\begin{aligned}
D_{*\rho_\mu \rho_{\mu'}} &= \frac{A_{\rho_\mu} \Xi A_{\rho_{\mu'}}}{2}, \quad D_{*\rho_\mu \varsigma_{(\nu\nu')}} = \frac{A_{\rho_\mu} \Xi A_{\varsigma_{(\nu\nu')}}}{2}, \\
D_{*\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} &= \frac{A_{\varsigma_{(\mu\mu')}} \Xi A_{\varsigma_{(\nu\nu')}}}{2},
\end{aligned} \tag{57}$$

with obvious modifications for  $c_{\varsigma_{(\nu\nu')} \rho_\mu}$ ,  $d_{\varsigma_{(\nu\nu')} \rho_\mu}$  and  $D_{*\varsigma_{(\nu\nu')} \rho_\mu}$ .

The next two theorems give Edgeworth approximations of the distribution functions of the Wald ( $w$ ) and  $F$  statistics, given by (49) and (50), respectively.

**Theorem 4.** *The distribution function of the Wald statistic (49), under the null hypothesis (48), admits the Edgeworth expansion*

$$\Pr\{w \leq x\} = F_m(x) - \tau^2 [\xi_1 + (\xi_2/(m+2))x] \frac{x}{m} f_m(x) + O(\tau^3), \tag{58}$$

where  $F_m(\cdot)$  and  $f_m(\cdot)$  are the chi-square distribution and density functions, respectively, and scalars  $\xi_1$  and  $\xi_2$  can be calculated as follows:

$$\xi_1 = \text{tr}[\Lambda(C + D_*)] - c' \Lambda c / 4 + c' \kappa + m[c' \lambda / 2 - \kappa_0 - (m-2)\lambda_0 / 4], \tag{59}$$

$$\xi_2 = \text{tr}(\Lambda D_*) + [c' \Lambda c - (m+2)(2c' \lambda - m\lambda_0)] / 4.$$

Analytic formulae for the computation of scalars  $\lambda_0$  and  $\kappa_0$ , and the elements of  $\lambda$ ,  $\kappa$ ,  $\Lambda$ ,  $c$ ,  $C$  and  $D_*$  are given in the Appendix (see Lemmas A.16 and A.17).

Instead of using the Wald statistic (49) and the Edgeworth expansion of its distribution, given in (58), we can use the  $F$  statistic, given by (50), and approximate its distribution function in terms of the  $F$  distribution as follows:

**Theorem 5.** *The distribution function of the  $F$  statistic (50), under null hypothesis (48), admits the Edgeworth expansion*

$$\Pr\{F \leq x\} = F_{MT-n}^m(x) - \tau^2 [q_1 + q_2 x] x f_{MT-n}^m(x) + O(\tau^3), \quad (60)$$

where  $F_{MT-n}^m(\cdot)$  and  $f_{MT-n}^m(\cdot)$  are the  $F$  distribution and density functions, respectively, with  $m$  and  $MT - n$  degrees of freedom, and scalars  $q_1$  and  $q_2$  can be calculated as follows:

$$q_1 = \xi_1/m + (m - 2)/2, \quad q_2 = \xi_2/(m + 2) - m/2, \quad (61)$$

where scalars  $\xi_1$  and  $\xi_2$  are defined in (59).

Theorem 4 implies that the Edgeworth corrected  $\alpha\%$  critical value of the Wald statistic (49) is given as

$$\chi_\alpha^* = \chi_\alpha + \tau^2 \left[ \frac{\xi_1}{m} + \frac{\xi_2}{m(m + 2)} \chi_\alpha \right] \chi_\alpha, \quad (62)$$

based on the  $\alpha\%$  significant point of the chi-square distribution, denoted as  $\chi_\alpha$ . Theorem 5 enables us to calculate the Edgeworth corrected  $\alpha\%$  critical value of  $F$  statistic (50) as

$$F_\alpha^* = F_\alpha + \tau^2 [q_1 + q_2 F_\alpha] F_\alpha, \quad (63)$$

based on the  $\alpha\%$  significant point of the  $F$  distribution, denoted as  $F_\alpha$ .

The Cornish-Fisher size corrected  $F$  statistic for testing null hypothesis (48) is given in the next theorem.

**Theorem 6.** *The Cornish-Fisher size corrected  $F$  statistic*

$$F_* = F - \tau^2 [q_1 + q_2 F] F \quad (64)$$

*is distributed, under null hypothesis (48), as an  $F$  random variable with  $m$  and  $MT - n$  degrees of freedom, with an approximation error of order  $O(\tau^3)$ .*

Unlike the Edgeworth approximation, the Cornish-Fisher corrected  $F$  statistic, denoted as  $F_*$  in equation (64), is a proper random variable and it does not assign negative ‘probabilities’ in the tails of its distribution. Thus, the Cornish-Fisher corrected  $F$  statistic can be readily implemented, in applied research, to test null hypothesis (48). This can be done by using the standard tables of the  $F$  distribution, with  $m$  and  $MT - n$  degrees of freedom.

## 5 Monte-Carlo simulations

In this section, we evaluate the small-sample performance of the size corrected tests suggested in the previous section, compared to their corresponding standard (first-order asymptotic approximation) versions. To this end, we rely on a Monte Carlo simulation based on 5000 iterations and we consider small-samples of  $T = 15, 20, 40$  observations.

In our simulation, we consider the S.U.R. model of  $M = 2$  seemingly unrelated equations (see, e.g., Zellner (1962)), i.e.,

$$\begin{aligned} y_{t,1} &= \beta_{0,1} + \beta_{1,1}x_{t1,1} + \beta_{2,1}x_{t2,1} + u_{t,1} \\ y_{t,2} &= \beta_{0,2} + \beta_{1,2}x_{t1,2} + \beta_{2,2}x_{t2,2} + u_{t,2} \end{aligned} \quad (t = 1, \dots, T), \quad (65)$$

where the error terms,  $u_{t,1}$  and  $u_{t,2}$ , are contemporaneously correlated with covariance  $\sigma_{12}$ . Both of these error terms follow AR(1) process (2), with normally distributed innovations. The autoregressive coefficients of this process  $\rho_1$  and  $\rho_2$  are assumed to be equal, i.e.,  $\rho_1 = \rho_2 = \rho = \pm 0.5, \pm 0.8$ . To ensure stationarity of error terms  $u_{t,1}$  and  $u_{t,2}$ , conditions (3) are satisfied. For  $t = 0$ , these conditions require that

$$\begin{aligned} y_{0,1} &\sim \mathcal{N}(0, \sigma_{11}/(1 - \rho_1^2)) \quad \text{and} \quad \mathcal{E}(y_{0,1}y_{0,2}) = \sigma_{12} \frac{(1 - \rho_1^2)^{1/2}(1 - \rho_2^2)^{1/2}}{1 - \rho_1\rho_2} \\ y_{0,2} &\sim \mathcal{N}(0, \sigma_{22}/(1 - \rho_2^2)) \end{aligned}$$

In our analysis, we assume  $\sigma_{11} = \sigma_{22} = 1$  and we are focused on investigating the consequences of the different sign and magnitude of covariances  $\sigma_{12}$  on our tests, for the following cases:  $\sigma_{12} = \pm 0.5, \pm 0.75, \pm 0.9$ . Since  $\sigma_{11} = \sigma_{22} = 1$ ,  $\sigma_{12}$  is the correlation coefficient between  $u_{t,1}$  and  $u_{t,2}$ .

According to (15) (or (16)), the above S.U.R. model can be written in terms of the following transformed equations, with non-autocorrelated errors:

$$y_{1*} = X_{1*}\beta_1 + \varepsilon_1; \quad y_{2*} = X_{2*}\beta_2 + \varepsilon_2,$$

where  $y_{1*}$  and  $y_{2*}$  are  $(TX1)$  vectors of observations on the dependent variables, with  $P_\mu y_{\mu*} = y_\mu$ , for  $\mu = 1, 2$ , where  $P_\mu$  is defined by (9),  $X_{1*}$  and  $X_{2*}$  are  $(T \times 3)$  matrices of regressors, with  $P_\mu X_{\mu*} = X_\mu$  and  $\beta_1 = (\beta_{0,1}, \beta_{1,1}, \beta_{2,1})'$ ,  $\beta_2 = (\beta_{0,2}, \beta_{1,2}, \beta_{2,2})'$  are  $(3 \times 1)$  vectors of parameters, including the constant. In terms of the S.U.R. representation (21), the above equations can be written

as

$$Y_* = ZB + E,$$

where  $Y_*$  is a  $(T \times 2)$  matrix of observations on vectors  $y_{1*}$  and  $y_{2*}$ ,  $E$  is a  $(T \times 2)$  matrix whose rows are vectors of normally distributed innovations with variance-covariance  $\Sigma = [(\sigma_{\mu\mu'})_{\mu,\mu'=1,2}]$ ,  $B$  is a  $(3 \times 2)$ -dimension matrix whose columns,  $\beta_1$  and  $\beta_2$ , are vectors of parameters, and  $Z$  is a  $(T \times 6)$  matrix whose columns are vectors of possibly collinear variables defined as

$$\begin{aligned} z_{t1} &\equiv z_{t6} \equiv (1 - \rho^2)^{1/2} & (t = 1), \\ z_{t1} &\equiv z_{t6} \equiv (1 - \rho) & (t = 2, 3, \dots, T), \\ z_{tj} &= \alpha^{1/2}\zeta_{t1} + (1 - \alpha)^{1/2}\zeta_{tj} & (j = 2, 3, 4, 5), \end{aligned}$$

where  $\zeta_{tj}$  ( $j = 2, 3, 4, 5$ ) are  $\mathcal{N}(0, 1)$  random variables and  $\alpha$  stands for the common correlation coefficient between any two non-constant columns of  $Z$  (see also McDonald and Galarneau (1975)). This captures the same degree of multicollinearity between regressors  $x_{t1,\mu}$  and  $x_{t2,\mu}$  of S.U.R. model (65). In our simulation, we consider the following two values of the collinearity coefficient:  $\alpha = 0.5, 0.9$ . According to (25), submatrices  $X_{1*}$  and  $X_{2*}$  (collected in matrix  $X_*$ ) can be obtained from  $Z$  by assuming that submatrices  $\Psi_1$  and  $\Psi_2$ , of the block diagonal matrix  $\Psi$  are given as follows:

$$\Psi_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}; \quad \Psi_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

In all iterations of our simulation, the two equations of S.U.R. model (65) were estimated by LS. The residuals of these equations were used to compute the LS estimates of autoregressive coefficients  $\rho_1$  and  $\rho_2$ , denoted as  $\tilde{\rho}_1$  and  $\tilde{\rho}_2$ . Then, the transformed variables  $y_{1,\mu}^*$  and  $x_{tj,\mu}^*$ , for  $j = 0, 1, 2$  (where ‘0’ stands for the constant), are calculated as follows:

$$\begin{aligned} y_{1,\mu}^* &= (1 - \tilde{\rho}_\mu^2)^{1/2} y_{1,\mu} & x_{1j,\mu}^* &= (1 - \tilde{\rho}_\mu^2)^{1/2} x_{1j,\mu} & (t = 1), \\ y_{t,\mu}^* &= y_{t,\mu} - \tilde{\rho}_\mu y_{(t-1),\mu} & x_{tj,\mu}^* &= x_{tj,\mu} - \tilde{\rho}_\mu x_{(t-1)j,\mu} & (t \neq 1). \end{aligned} \tag{66}$$

These variables were then used to compute the feasible GL estimates of  $\beta_{j,\mu}$  ( $j = 0, 1, 2; \mu = 1, 2$ ), denoted as  $\hat{\beta}_{j,\mu}$ . The columns of matrix  $Z$  were obtained as  $z_1 = x_{0,1}^*$ ,  $z_2 = x_{1,1}^*$ ,  $z_3 = x_{2,1}^*$ ,  $z_6 = x_{0,2}^*$ ,  $z_4 = x_{1,2}^*$ ,  $z_5 = x_{2,2}^*$ , while the unrestricted estimates of matrix  $B$  were based on the GL estimates  $\hat{\beta}_{j,\mu}$ . The unrestricted estimates of the inverse covariance matrix  $\Sigma^{-1}$  were estimated based on (32) and the feasible GL estimate  $\hat{\sigma}_{GL}$  which is calculated by using the following formula:

$$\hat{\sigma}_{GL} = \left[ (y - X\hat{\beta})' \left( \hat{P}_I'^{-1} (\hat{\Sigma}_I^{-1} \otimes I_T) \hat{P}_I^{-1} \right) (y - X\hat{\beta}) / (MT - n) \right]^{1/2},$$

where  $I$  denotes any consistent estimators of matrices  $\Sigma^{-1}$  and  $P^{-1}$  (see Appendix), used to obtain a feasible GL estimator of  $\beta$ .

The results of our simulation are presented in Tables 1a, 1b and 2. The actual sizes of our size corrected tests of the following null hypothesis:

$$H_0 : \beta_{2,1} = 0, \tag{67}$$

against its one-sided alternatives, are reported in Tables 1a and 1b. In particular, Table 1a presents results against alternative  $H_A : \beta_{2,1} > 0$ , while Table 1b against  $H_A : \beta_{2,1} < 0$ . The table presents the actual sizes (i.e., the rejection probabilities) at the 5% significance level of the following: the standard normal and Student- $t$  tests (denoted as  $z$  and  $t$ , respectively), their finite-sample size corrected versions based on the Edgeworth corrected critical values of the standard normal and Student- $t$  distributions (denoted as  $E$ - $z$  and  $E$ - $t$ , respectively) and the Cornish-Fisher finite-sample size corrected Student- $t$  test (denoted as  $CF$ - $t$ ). Note that we do not examine the performance of the above  $t$  tests for the null hypothesis (67) against its two-sided alternatives, since this is a special case of the  $F$  test examined in Table 2.

Table 2 presents the actual sizes of our size corrected tests of the following joint null hypothesis on the slope coefficients of S.U.R. model (65), across its two equations:

$$H_0 : \beta_{1,1} = \beta_{2,1} = \beta_{1,2} = \beta_{2,2} = 0. \tag{68}$$

This is done against the alternative hypothesis that at least one of these coefficients are different from zero, i.e., at least one  $\beta_{j,\mu} \neq 0$  ( $j = 1, 2; \mu = 1, 2$ ). The table presents the actual sizes at the 5% significance level of the following: the

standard Wald (chi-square) and  $F$  tests (denoted as  $\chi^2$  and  $F$ , respectively), their finite-sample size corrected versions based on the Edgeworth corrected critical values of the chi-square and  $F$  distributions (denoted by  $E\text{-}\chi^2$  and  $E\text{-}F$ , respectively) and the Cornish-Fisher finite-sample size corrected  $F$  test (denoted as  $CF\text{-}F$ ).

Turning now into the discussion of the results of our simulation, Tables 1a and 1b clearly indicate that the size corrected tests have better size performance in all reported sample sizes ( $T = 15, 20, 40$ ), compared to the standard versions of these tests, based on first-order approximations. This is true for both the Edgeworth and Cornish-Fisher size corrections, and across all different values of  $\rho$ ,  $\sigma_{12}$  and  $\alpha$  examined.

Between the above different categories of size corrected tests, our results indicate that the  $CF\text{-}t$  test outperforms the  $E\text{-}z$  and  $E\text{-}t$  ones. This is true for almost all cases of  $\alpha$  and  $\sigma_{12}$  considered, if  $\rho$  takes large values, i.e.,  $\rho = \pm 0.8$ . The same is true for small samples ( $T = 15$  or  $20$ ) and  $\rho = \pm 0.5$ .

Regarding the chi-square and  $F$  tests, the results of Table 2 indicate that, in most of the cases examined, the size corrected versions of these tests, i.e.,  $E\text{-}\chi^2$ ,  $E\text{-}F$  and  $CF\text{-}F$ , perform better in small samples, compared to their standard versions. Between the Edgeworth and Cornish-Fisher size corrected versions of these tests (i.e.,  $E\text{-}F$  (or  $E\text{-}\chi^2$ ) and  $CF\text{-}F$ ), the latter is found to perform better than the former for all sample sizes considered, and across all values of  $\rho$ ,  $\sigma_{12}$  and  $\alpha$  examined. Notice that, for relatively large samples ( $T = 40$ ), the  $E\text{-}\chi^2$  test outperforms the degrees-of-freedom-adjusted  $E\text{-}F$  test. This suggests that, for the model considered in our simulation, samples of 40 observations seem to be large enough to induce the reduction of the magnitude of the degrees-of-freedom-adjusted Edgeworth size corrections.

Summing up, the results of our simulation clearly indicate that the finite-sample size corrected tests  $E\text{-}\chi^2$ ,  $E\text{-}F$  and  $CF\text{-}F$  can considerably improve the performance of the standard (uncorrected) tests in small samples. This happens even for very high levels of autocorrelation and/or cross-correlation between the error terms of the equations of the S.U.R. model. Another interesting conclusion that can be drawn from the results of this exercise is that the adjusted for the degrees of freedom versions of the tests perform better than their unadjusted

ones in most of the cases considered in our simulation. Note that this is also true for the standard (uncorrected) versions of the tests.

Table 1a:  $H_0 : \beta_{2,1} = 0$  against  $H_A : \beta_{2,1} > 0$  (Nominal size: 5%)

Test:			Actual sizes (%)																				
			$\rho = -0.8$					$\rho = -0.5$					$\rho = 0.5$					$\rho = 0.8$					
			$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$	
$\alpha$	$\sigma_{12}$	$T$																					
0.5	-0.90	15	14.6	10.2	13.8	10.4	8.0	11.8	8.0	11.0	8.4	7.4	11.9	8.5	11.1	8.9	8.1	14.9	11.0	13.9	11.2	9.9	
		20	12.4	8.6	11.8	8.8	7.3	9.7	6.7	9.1	6.9	6.6	10.5	7.5	9.6	7.7	7.5	12.9	9.5	12.4	9.9	8.8	
		40	9.0	7.2	8.7	7.3	7.0	6.9	5.3	6.6	5.4	5.3	7.4	5.9	7.3	6.1	6.0	9.8	7.7	9.5	7.9	7.5	
	-0.75	15	14.6	10.1	13.9	10.3	7.9	11.1	7.7	10.3	8.0	7.3	11.7	8.2	10.9	8.6	7.9	14.5	10.5	13.5	10.7	9.6	
		20	12.5	9.0	11.9	9.2	7.6	9.0	6.3	8.3	6.5	6.1	10.2	7.6	9.7	7.9	7.5	13.4	9.8	12.7	10.1	9.2	
		40	8.1	6.0	7.9	6.2	5.8	7.1	5.7	6.9	5.9	5.8	7.4	5.9	7.2	6.0	5.9	9.1	7.2	8.9	7.4	7.0	
	-0.50	15	14.8	10.2	14.0	10.4	7.7	10.4	7.4	9.7	7.6	7.0	11.4	8.1	10.6	8.5	7.9	14.2	10.6	13.5	10.8	9.6	
		20	12.4	7.8	11.8	9.0	7.3	9.0	6.6	8.4	6.8	6.6	9.4	6.9	9.0	7.2	6.8	12.9	9.5	12.3	9.7	8.6	
		40	8.5	6.5	8.3	6.7	6.4	7.0	5.4	6.8	5.6	5.5	7.4	6.1	7.2	6.2	6.1	9.2	7.1	9.0	7.3	7.1	
	0.50	15	14.0	9.7	13.2	9.9	7.7	10.5	7.2	9.8	7.4	6.9	11.5	8.1	10.6	8.5	7.9	14.9	10.7	14.0	11.1	9.9	
		20	11.9	8.1	11.4	8.3	6.8	8.7	6.4	8.3	6.6	6.3	10.3	7.7	9.7	8.0	7.5	13.3	10.2	12.8	10.3	9.3	
		40	8.1	6.3	7.9	6.4	6.2	6.8	5.4	6.5	5.5	5.4	7.1	5.7	6.9	5.9	5.7	9.0	6.9	8.8	7.0	6.8	
	0.75	15	14.7	10.2	14.0	10.4	8.0	11.5	8.0	10.5	8.2	7.5	12.2	8.5	11.3	8.9	8.3	13.8	10.3	13.1	10.5	9.3	
		20	12.2	8.8	11.6	8.9	7.4	9.3	6.7	8.8	6.9	6.5	10.2	7.3	9.6	7.7	7.3	12.5	9.4	11.9	9.6	8.6	
		40	8.8	6.8	8.6	6.9	6.5	7.2	5.9	7.0	6.1	6.0	7.5	5.9	7.2	6.1	6.0	9.2	7.0	8.9	7.2	6.8	
	0.90	15	13.8	9.7	13.0	9.8	7.5	11.2	7.7	10.3	8.0	7.3	12.2	8.7	11.6	9.0	8.4	15.0	11.0	14.1	11.2	10.1	
		20	12.9	9.0	12.4	9.2	7.7	9.4	6.6	8.7	6.8	6.3	10.0	7.3	9.4	7.5	7.2	12.9	9.5	12.3	9.8	8.8	
		40	9.1	6.9	8.7	7.1	6.7	7.0	5.4	6.8	5.6	5.4	7.2	5.7	7.0	5.8	5.7	9.4	7.3	9.2	7.5	7.2	
	0.9	-0.90	15	14.6	10.4	13.8	10.5	7.7	11.2	7.7	10.4	7.9	7.3	11.8	8.5	11.0	8.7	8.2	14.5	10.9	13.8	11.1	9.8
			20	12.7	9.3	12.2	9.5	7.8	9.8	6.8	9.2	7.1	6.7	10.4	7.6	9.9	7.8	7.5	13.2	10.0	12.8	10.2	9.3
			40	9.2	7.2	9.0	7.4	7.1	7.4	6.0	7.2	6.2	6.0	7.3	5.9	7.1	6.0	6.0	9.9	7.9	9.7	8.0	7.7
		-0.75	15	14.5	9.8	13.5	10.0	7.8	10.7	7.3	9.8	7.5	6.9	11.7	8.4	10.9	8.7	8.1	14.9	11.0	13.9	11.3	10.1
			20	11.9	8.3	11.4	8.5	7.0	9.9	7.2	9.4	7.4	7.0	9.7	6.9	9.1	7.2	6.8	13.0	9.9	12.5	10.1	9.0
			40	8.5	6.5	8.3	6.7	6.4	6.7	5.2	6.4	5.3	5.2	7.5	5.9	7.3	6.1	6.0	9.7	7.8	9.4	7.9	7.7
-0.50		15	14.2	9.6	13.3	9.8	7.3	10.8	7.4	9.9	7.6	7.1	11.7	8.3	10.8	8.6	8.2	14.5	10.9	13.6	11.1	9.7	
		20	11.5	8.0	11.0	8.2	6.8	9.3	6.8	8.8	7.1	6.7	10.2	7.4	9.6	7.7	7.2	12.6	9.7	11.9	9.9	9.0	
		40	9.0	7.0	8.8	7.2	6.8	7.1	5.8	6.9	5.9	5.8	7.3	5.7	6.9	5.9	5.7	8.9	6.8	8.7	7.0	6.7	
0.50		15	14.6	10.3	13.8	10.4	7.9	10.6	7.5	9.8	7.7	7.1	11.9	8.3	11.0	8.6	8.0	14.9	11.2	14.2	11.4	10.1	
		20	12.7	8.8	12.1	9.0	7.7	9.1	6.4	8.6	6.7	6.4	9.8	7.0	9.2	7.3	6.9	12.8	9.3	12.2	9.5	8.5	
		40	8.5	6.5	8.3	6.7	6.3	6.9	5.4	6.7	5.6	5.4	7.3	5.8	7.1	5.9	5.9	9.3	7.0	9.1	7.1	6.8	
0.75		15	14.0	9.6	13.2	9.7	7.3	10.7	7.3	9.9	7.5	6.9	11.6	8.0	10.6	8.3	7.8	14.1	10.3	13.3	10.5	9.5	
		20	12.2	8.8	11.7	9.0	7.4	9.3	6.5	8.7	6.8	6.4	9.8	7.0	9.2	7.4	6.9	12.8	9.7	12.3	9.9	8.9	
		40	8.5	6.3	8.2	6.5	6.2	7.2	5.8	7.0	6.0	5.9	7.7	5.9	7.4	6.1	6.0	9.2	7.0	8.9	7.1	6.9	
0.90		15	14.3	10.0	13.5	10.2	7.8	11.1	7.8	10.2	8.0	7.3	12.3	8.7	11.5	9.1	8.3	15.3	11.3	14.3	11.5	10.1	
		20	13.0	9.1	12.4	9.3	7.7	9.1	6.8	8.7	7.0	6.6	9.9	7.2	9.4	7.4	7.0	12.8	9.3	12.2	9.4	8.5	
		40	8.8	6.9	8.6	7.0	6.8	7.1	5.5	6.8	5.7	5.6	7.2	5.6	6.9	5.8	5.7	9.6	7.6	9.4	7.7	7.4	



Table 1b:  $H_0 : \beta_{2,1} = 0$  against  $H_A : \beta_{2,1} < 0$  (Nominal size: 5%)

Test:			Actual sizes (%)																			
			$\rho = -0.8$					$\rho = -0.5$					$\rho = 0.5$					$\rho = 0.8$				
$\alpha$	$\sigma_{12}$	$T$	$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$	$z$	$E-z$	$t$	$E-t$	$CF-t$
0.5	-0.90	15	15.0	10.5	14.1	10.6	8.2	11.2	7.7	10.3	8.0	7.2	12.0	8.5	11.1	8.8	8.2	14.7	10.9	13.9	11.1	9.9
		20	12.5	8.8	11.8	9.0	7.3	10.1	7.3	9.6	7.5	7.0	10.0	7.3	9.5	7.7	7.3	13.2	9.6	12.5	9.9	8.9
		40	8.8	6.9	8.6	7.0	6.5	7.5	5.9	7.3	6.0	6.0	7.2	5.8	7.1	6.0	5.9	9.1	7.1	8.8	7.2	6.8
	-0.75	15	14.4	10.0	13.5	10.1	8.1	11.2	7.9	10.4	8.1	7.4	11.7	8.1	10.9	8.5	7.9	14.7	10.7	13.8	11.0	9.8
		20	12.9	9.4	12.2	9.5	8.1	9.3	6.6	8.7	6.8	6.4	9.6	6.8	9.0	7.1	6.7	12.7	9.4	12.1	9.6	8.8
		40	8.7	6.8	8.4	6.9	6.7	7.3	6.0	7.1	6.1	6.1	7.6	5.9	7.3	6.2	6.0	8.9	7.1	8.7	7.2	6.9
	-0.50	15	14.5	10.2	13.7	10.3	7.6	10.7	7.4	10.0	7.7	7.2	11.7	8.1	10.8	8.4	7.8	14.6	10.9	13.7	11.1	9.9
		20	12.3	8.7	11.7	8.9	7.3	9.5	6.7	8.8	7.0	6.6	9.7	7.1	9.1	7.3	7.0	13.1	9.7	12.5	10.1	9.0
		40	7.9	6.1	7.5	6.2	5.9	7.1	5.9	6.8	6.0	5.9	6.8	5.5	6.6	5.7	5.5	9.0	7.0	8.8	7.2	6.9
	0.50	15	13.8	9.9	12.9	10.1	7.6	10.9	7.3	10.1	7.6	6.9	11.4	8.2	10.5	8.5	7.9	14.8	11.0	14.0	11.2	10.1
		20	12.1	8.3	11.5	8.5	6.8	9.1	6.2	8.5	6.4	6.1	9.8	7.1	9.1	7.4	7.0	13.0	9.6	12.4	9.8	8.8
		40	8.6	6.4	8.4	6.6	6.4	7.2	5.7	6.9	5.9	5.8	7.6	5.8	7.4	6.0	5.9	9.9	7.6	9.6	7.7	7.3
	0.75	15	14.5	10.0	13.6	10.2	7.8	11.4	7.8	10.5	8.0	7.3	11.6	8.5	10.9	8.8	8.2	14.2	10.6	13.4	10.9	9.7
		20	12.9	8.9	12.2	9.2	7.8	9.8	7.1	9.2	7.4	7.0	9.8	6.9	9.2	7.2	6.9	12.8	9.2	12.2	9.4	8.5
		40	8.9	6.6	8.5	6.8	6.5	7.0	5.5	6.8	5.6	5.6	7.2	5.8	7.0	6.0	5.9	9.4	7.3	9.1	7.4	7.1
0.90	15	14.1	10.1	13.2	10.2	8.2	11.2	7.9	10.4	8.1	7.4	11.9	8.2	11.0	8.5	7.8	14.6	10.7	13.8	11.0	9.7	
	20	12.3	8.6	11.7	8.7	7.3	9.6	6.8	9.1	7.0	6.7	9.8	7.2	9.3	7.4	7.1	13.5	10.0	12.8	10.3	9.4	
	40	8.0	6.3	7.8	6.5	6.1	6.9	5.5	6.8	5.7	5.6	7.2	5.8	7.0	6.0	5.9	9.6	7.7	9.3	7.9	7.5	
0.9	-0.90	15	14.4	10.0	13.7	10.1	7.7	11.6	8.2	10.9	8.5	7.6	12.0	8.5	11.3	8.8	8.2	15.4	11.2	14.5	11.5	10.3
		20	12.4	8.9	11.8	9.1	7.5	9.3	6.5	8.7	6.7	6.4	9.9	7.2	9.2	7.4	6.9	13.0	9.8	12.3	10.1	8.9
		40	8.7	6.1	8.5	6.8	6.4	7.1	5.6	6.8	5.8	5.7	7.3	5.8	7.1	6.0	5.9	9.7	7.7	9.4	7.8	7.6
	-0.75	15	14.5	10.4	13.8	10.5	8.2	11.0	7.6	10.2	7.9	7.4	11.7	8.4	11.0	8.7	8.1	14.7	10.6	13.9	10.9	9.4
		20	12.3	8.7	11.7	8.9	7.4	9.3	6.7	8.8	7.0	6.6	9.9	7.3	9.2	7.6	7.3	12.5	9.2	12.1	9.5	8.5
		40	8.7	6.5	8.5	6.7	6.3	6.9	5.6	6.7	5.8	5.7	7.0	5.6	6.8	5.8	5.7	9.1	7.1	8.7	7.2	7.0
	-0.50	15	14.7	9.8	13.5	10.0	7.7	10.6	7.3	9.8	7.6	7.0	11.6	8.2	10.8	8.5	7.9	14.3	10.6	13.4	10.8	9.4
		20	11.7	8.1	11.2	8.4	6.7	9.5	6.8	9.0	7.1	6.6	10.2	7.5	9.6	7.9	7.3	12.5	9.0	12.0	9.3	8.4
		40	8.9	6.8	8.7	7.0	6.6	6.9	5.6	6.7	5.7	5.6	6.7	5.4	6.4	5.6	5.5	9.0	6.9	8.7	7.1	6.8
	0.50	15	14.0	9.6	13.1	9.7	7.6	10.2	7.0	9.5	7.2	6.6	11.2	8.0	10.5	8.3	7.7	14.0	10.5	13.3	10.7	9.5
		20	11.5	8.2	11.0	8.3	7.0	9.6	6.9	9.0	7.3	6.8	9.9	7.3	9.4	7.6	7.2	12.5	9.2	12.0	9.5	8.7
		40	8.5	6.3	8.1	6.5	6.2	7.2	5.7	6.9	5.9	5.8	7.4	5.7	7.1	6.0	5.9	8.9	6.8	8.7	7.0	6.7
	0.75	15	14.2	9.9	13.3	10.0	7.6	11.4	7.9	10.6	8.2	7.3	12.0	8.6	11.2	8.8	8.3	14.5	10.7	13.7	10.9	9.8
		20	12.0	8.6	11.4	8.8	7.1	9.3	6.9	8.7	7.1	6.8	9.5	6.9	9.0	7.2	6.7	12.8	9.6	12.2	9.8	9.0
		40	8.4	6.4	8.2	6.6	6.3	7.3	5.8	7.1	6.0	5.9	10.0	5.7	6.8	5.8	5.7	9.2	7.3	9.0	7.5	7.1
0.90	15	15.3	10.5	14.4	10.6	8.2	11.3	7.9	10.4	8.2	7.6	11.2	7.8	10.4	8.1	7.5	15.3	11.5	14.5	11.7	10.3	
	20	13.0	9.2	12.4	9.3	7.7	9.4	6.7	8.7	7.0	6.5	10.6	8.0	10.1	8.3	7.8	13.2	9.9	12.5	10.2	9.2	
	40	9.1	7.0	8.8	7.2	6.9	7.1	5.6	6.8	5.8	5.6	7.1	5.9	7.0	6.1	6.0	10.3	7.9	10.1	8.1	7.7	

Table 2:  $H_0 : \beta_{1,1} = \beta_{2,1} = \beta_{1,2} = \beta_{2,2} = 0$  (Nominal size: 5%)

Test:			Actual sizes (%)														
			$\chi^2$	$E\text{-}\chi^2$	$F$	$E\text{-}F$	$CF\text{-}F$	$\chi^2$	$E\text{-}\chi^2$	$F$	$E\text{-}F$	$CF\text{-}F$	$\chi^2$	$E\text{-}\chi^2$	$F$	$E\text{-}F$	$CF\text{-}F$
$\alpha$	$\sigma_{12}$	$T$	$\rho = -0.8$					$\rho = -0.5$					$\rho = 0.5$				
0.5	-0.90	15	46.1	31.4	40.2	27.9	4.7	30.5	18.2	24.7	16.5	9.8	33.3	20.7	26.7	19.3	14.0
		20	38.2	25.0	33.6	23.6	6.4	22.7	13.2	18.6	12.9	9.6	26.1	15.6	21.6	15.2	11.9
		40	20.5	13.3	18.7	13.5	10.7	12.6	7.6	11.1	8.2	7.4	12.9	8.0	11.3	8.4	7.7
	-0.75	15	45.8	31.5	39.9	28.4	5.8	28.4	16.6	22.4	15.4	10.9	33.2	21.4	27.1	20.2	15.9
		20	36.7	24.2	32.3	22.6	7.9	22.4	12.9	18.3	12.8	10.2	25.4	15.9	21.3	15.6	13.0
		40	20.2	12.8	18.2	13.0	10.2	12.4	7.6	10.9	8.0	7.4	13.2	8.4	11.7	8.8	8.0
	-0.50	15	46.2	31.6	39.7	28.3	7.3	28.9	17.6	23.2	16.5	12.4	33.0	21.0	26.7	20.1	16.6
		20	36.0	23.1	31.6	21.7	8.9	21.1	12.1	17.1	12.1	10.0	23.1	14.2	19.1	14.2	12.2
		40	17.6	11.4	16.2	11.7	9.7	11.9	7.5	10.4	7.9	7.5	12.8	8.1	11.2	8.5	7.8
	0.50	15	45.8	31.1	39.8	28.1	7.6	29.2	17.5	23.3	16.4	12.4	32.6	20.6	26.3	19.6	16.2
		20	35.9	23.4	31.5	21.9	8.7	21.2	12.5	17.6	12.4	10.4	24.4	14.6	20.0	14.5	12.2
		40	18.3	11.4	16.4	11.6	9.4	12.1	7.6	10.7	8.1	7.6	13.2	8.1	11.5	8.5	8.0
	0.75	15	45.5	31.1	39.4	28.1	6.2	30.3	18.5	24.1	17.2	11.6	33.9	21.8	27.8	20.6	16.2
		20	36.9	24.0	32.3	22.7	8.2	22.6	13.5	18.5	13.3	10.7	24.9	15.4	20.6	15.2	12.4
		40	19.2	12.6	17.5	12.8	10.2	12.9	7.9	11.4	8.4	7.7	13.3	8.2	11.8	8.8	8.0
0.9	0.90	15	46.1	31.7	40.1	28.2	4.9	29.9	18.0	24.2	16.4	9.7	35.0	22.2	28.9	20.7	14.9
		20	37.8	24.5	33.3	23.0	7.2	23.1	13.2	18.9	12.9	9.6	25.0	15.5	20.7	15.2	12.2
		40	20.6	13.5	18.9	13.7	10.8	12.2	7.4	10.7	7.9	7.2	13.2	7.8	11.4	8.4	7.4
	-0.90	15	46.23	32.0	40.1	28.9	5.4	29.8	18.2	23.8	17.0	11.2	34.4	22.4	28.3	21.2	16.6
		20	38.2	25.3	33.8	23.8	7.3	22.9	13.4	19.1	13.3	10.7	26.1	15.9	21.8	15.8	13.2
		40	20.6	13.7	19.0	14.0	11.3	12.2	7.8	10.8	8.2	7.8	14.2	9.2	12.7	9.8	9.0
	-0.75	15	45.7	32.0	39.8	29.0	6.8	29.1	17.4	22.8	16.4	11.7	33.4	21.0	26.9	19.9	16.1
		20	36.9	24.9	32.8	23.5	8.6	21.2	11.9	17.2	11.8	9.7	24.6	15.1	20.5	15.2	12.9
		40	18.8	11.8	17.1	12.1	9.7	12.3	7.6	10.5	8.0	7.5	13.0	8.2	11.4	8.6	7.9
	-0.50	15	44.5	30.4	38.3	27.6	7.4	27.7	16.1	21.8	15.2	11.7	32.4	21.0	26.5	20.2	16.9
		20	36.1	23.5	31.5	22.2	8.5	20.7	12.0	16.8	11.9	9.8	24.1	14.9	20.5	14.9	12.7
		40	18.1	11.4	16.3	11.7	9.4	11.6	7.3	10.1	7.7	7.3	12.3	7.8	10.9	8.2	7.8
	0.50	15	44.9	30.7	38.6	27.3	7.1	28.1	17.0	21.9	16.0	12.3	32.2	20.7	26.3	20.0	16.8
		20	35.4	23.4	31.0	22.2	8.9	20.7	11.9	16.8	11.8	9.9	23.8	14.5	19.6	14.5	12.3
		40	18.4	11.8	16.8	12.1	9.7	11.9	7.5	10.4	8.1	7.5	12.3	7.9	10.7	8.4	7.8
	0.75	15	46.4	32.2	40.3	29.0	6.4	29.2	17.4	22.9	16.3	11.5	33.1	20.7	26.6	19.6	15.9
		20	37.2	24.8	32.8	23.4	8.7	22.0	12.8	17.7	12.7	10.5	25.2	15.4	21.0	15.4	13.0
		40	19.4	12.8	17.9	13.2	10.6	12.0	7.4	10.4	7.9	7.4	13.1	8.1	11.5	8.6	8.0
	0.90	15	46.8	31.9	40.2	28.5	4.9	30.4	18.3	24.5	17.1	11.6	34.4	21.8	28.0	20.6	15.7
		20	38.8	25.8	34.3	24.2	7.9	22.6	13.3	18.7	13.1	10.3	26.2	16.3	22.0	16.0	13.1
		40	20.5	13.4	18.5	13.8	11.0	12.9	8.3	11.4	8.8	8.1	13.1	8.3	11.6	8.8	8.0

## 6 Conclusions

In this paper, we have employed Edgeworth expansions of the standard normal (or Student- $t$ ) and chi-square (or  $F$ ) distributions to derive second-order size corrected testing procedures for the coefficient of the S.U.R. model with first-order autocorrelated errors. These procedures include (i) the Edgeworth corrected critical values of the well-known Wald (or  $F$ ) and  $t$  tests and (ii) the Cornish-Fisher corrected  $F$  and  $t$  test statistics. Since the standard  $F$  and  $t$  tests are adjusted for the degrees of freedom, they are locally exact, which means that their approximate distributions become exact when the model is sufficiently simplified.

The Edgeworth and Cornish-Fisher expansions, employed by the paper, are equivalent to each other, since the latter constitutes an inversion of the former. However, in practice, the use of the Cornish-Fisher corrected test statistics is recommended, since they are proper random variables with well-behaved distribution tails. The Edgeworth approximation, on the other hand, may assign negative ‘probabilities’ in the tails of the approximate distributions. Furthermore, the Cornish-Fisher size corrected tests can be easily implemented, in practice, using the standard tables of the Student- $t$  and the  $F$  distributions.

To evaluate the small-sample performance of the suggested tests, we have conducted a Monte Carlo simulation. The results of this simulation indicate that the size corrected  $t$  and  $F$  tests lead to substantial size improvements upon their standard versions, which assume first-order asymptotic approximations. This is true even for very small samples of 15 or 20 observations. Between the Edgeworth and Cornish-Fisher categories of the size corrected tests suggested in the paper, the second category is found to perform better than the first for almost all cases of serial and cross-equation correlation of the error terms of the S.U.R. model examined. This result is also robust across different degrees of multicollinearity between the explanatory variables of the model considered. In particular, both the  $t$  and  $F$  Cornish-Fisher size corrected tests are found to outperform their Edgeworth size corrected counterparts even when the degree of serial correlation of the error terms is very high. This is true even for a close-to-unity degree of correlation across the S.U.R equations.

## Appendix

In this appendix, we provide proofs of the main results of the paper. To prove these results, we rely on a number of lemmas. Some of them are given with sketchy proofs only for reasons of space. The complete proofs are available upon request. The presentation of our proofs is scheduled as follows: First, we provide some preliminary matrix-algebra results, needed for the calculation of the quantities in the stochastic expansions of all estimators and tests considered. Then, using these lemmas, we give the proofs of the theorems.

### Matrix-algebra results

Following Magdalinos (1992, page 344), let  $\mathcal{I}$  be a given set of indices which, without loss of generality, can be considered to belong to the open interval  $(0, 1)$ . For any collection of real-valued stochastic quantities (scalars, vectors, or matrices)  $Y_\tau$  ( $\tau \in \mathcal{I}$ ), we write  $Y_\tau = \omega(\tau^i)$ , if for any given  $n > 0$ , there exists a  $0 < \epsilon < \infty$  such that

$$\Pr \left[ \|Y_\tau / \tau^i\| > (-\ln \tau)^\epsilon \right] = o(\tau^n), \quad (\text{A.1})$$

as  $\tau \rightarrow 0$ , where the  $\|\cdot\|$  is the Euclidean norm. If (A.1) is valid for any  $n > 0$ , we write  $Y_\tau = \omega(\infty)$ . The use of this order of magnitude is motivated by the fact that, if two stochastic quantities differ by a quantity of order  $\omega(\tau^i)$ , then, under general conditions, the distribution function of the one provides an asymptotic approximation of the distribution function of the other, with an error of order  $O(\tau^i)$ . Furthermore, orders  $\omega(\cdot)$  and  $O(\cdot)$  have similar operational properties (Magdalinos (1992)).

Define the following  $(T \times T)$  matrices:  $D$  which is a band matrix whose  $(t, t')$ -th element is equal to 1 if  $|t - t'| = 1$  and 0 elsewhere,  $D_j$  whose  $(t, t')$ -th element is equal to 1 if  $t - t' = 1$  and 0 elsewhere,  $D_i$  whose  $(t, t')$ -th element is equal to 1 if  $t - t' = -1$  and 0 elsewhere. Also, define the following  $(T \times T)$  matrices:  $\Delta$  with 1 in  $(1, 1)$ -st and  $(T, T)$ -th positions and 0's elsewhere,  $\Delta_{11}$  with 1 in  $(1, 1)$ -st position and 0's elsewhere,  $\Delta_{TT}$  with 1 in  $(T, T)$ -th position and 0's elsewhere. Moreover, by using matrix  $P_\mu$  in (9), we can calculate  $(T \times T)$  matrices  $R_{ij}$  as follows:

$$R_{ij} = P_i P_j' = \frac{1}{1 - \rho_i \rho_j} \begin{bmatrix} 1 & \rho_j & \cdots & \rho_j^{T-1} \\ \rho_i & 1 & \cdots & \rho_j^{T-2} \\ \vdots & \vdots & & \vdots \\ \rho_i^{T-1} & \rho_i^{T-2} & \cdots & 1 \end{bmatrix}. \quad (\text{A.2})$$

Matrices  $R^{ij}$  help us to write the elements of matrix  $\Omega$  analytically. For these matrices and their derivatives the following two lemmas hold:

**Lemma A.1.** For matrix  $R^{ii}$ , which is the inverse of  $R_{ii}$ , the following result holds:

$$R^{ii} = P_i'^{-1} P_i^{-1} = (1 + \rho_i^2) I_T - \rho_i D - \rho_i^2 \Delta, \quad (\text{A.3})$$

where  $R^{ii} = R_{ii}^{-1}$  ( $\forall i$ ). Moreover, for matrix  $R^{ij}$ , the following result holds:

$$\begin{aligned} R^{ij} = P_i'^{-1} P_j^{-1} &= (1 + \rho_i \rho_j) I_T - \rho_i D_i - \rho_j D_j - \rho_i \rho_j \Delta_{TT} \\ &\quad + [(1 - \rho_i^2)^{1/2} (1 - \rho_j^2)^{1/2} - 1] \Delta_{11}. \end{aligned} \quad (\text{A.4})$$

Note that  $R^{ij}$  is not the inverse of  $R_{ij}$ , i.e.,  $R^{ij} \neq R_{ij}^{-1}$  ( $\forall i \neq j$ ).

**Proof of Lemma A.1.** For  $i = j$ , definition (A.2) implies that matrix  $R_{ii}$  is the exact analogue of the error covariance matrix in a single-equation regression model with autocorrelated errors. And it is well-known from the autocorrelation literature that the inverse matrix  $R^{ii} = R_{ii}^{-1}$  can be expressed in the form of (A.3). Further, (A.4) can be proved along the same lines, as a straightforward generalization for  $i \neq j$ .  $\square$

Define the  $(M \times M)$  matrix  $\Sigma^{-1} = [(\sigma^{\mu\mu'})_{\mu, \mu'=1, \dots, M}]$  and scalars:

$$\begin{aligned} a_{ij} &= (1 - \rho_i^2)^{1/2} (1 - \rho_j^2)^{1/2}, \\ \xi'_{(i)j} &= \partial a_{ij} / \partial \rho_i, \quad \xi''_{(i)j} = \partial^2 a_{ij} / \partial^2 \rho_i, \quad \xi''_{(i)(j)} = \partial^2 a_{ij} / \partial \rho_i \partial \rho_j, \\ R_{\rho_\mu}^{ij} &= \partial R^{ij} / \partial \rho_\mu, \quad R_{\rho_\mu \rho_{\mu'}}^{ij} = \partial^2 R^{ij} / \partial \rho_\mu \partial \rho_{\mu'}. \end{aligned} \quad (\text{A.5})$$

**Lemma A.2.** For the partial derivatives of matrix  $R^{ij}$  the following results hold:

$$\begin{aligned} R_{\rho_i}^{ii} &= 2\rho_i I_T - D - 2\rho_i \Delta, \quad R_{\rho_i \rho_i}^{ii} = 2(I_T - \Delta) \quad (\forall i), \\ R_{\rho_j}^{ii} &= R_{\rho_j \rho_j}^{ii} = R_{\rho_i \rho_j}^{ii} = 0 \quad (\forall i \neq j), \\ R_{\rho_i}^{ij} &= \rho_j I_T - D_i - \rho_j \Delta_{TT} + \xi'_{(i)j} \Delta_{11} \quad (\forall i, j), \\ R_{\rho_i \rho_i}^{ij} &= \xi''_{(i)j} \Delta_{11}, \quad R_{\rho_i \rho_j}^{ij} = I_T - \Delta_{TT} + \xi''_{(i)(j)} \Delta_{11} \quad (\forall i, j), \\ R_{\rho_\mu}^{ij} &= R_{\rho_\mu \rho_\mu}^{ij} = R_{\rho_\mu \rho_i}^{ij} = R_{\rho_\mu \rho_j}^{ij} = 0 \quad (\forall \mu \neq i \wedge \forall \mu \neq j), \end{aligned} \quad (\text{A.6})$$

with obvious modifications for  $R_{\rho_j}^{ij}$  and  $R_{\rho_j \rho_j}^{ij}$ . Further,

$$\begin{aligned} \xi'_{(i)j} &= -\rho_i (1 - \rho_i^2)^{-1/2} (1 - \rho_j^2)^{1/2} \quad (\forall i), \\ \xi''_{(i)j} &= -(1 - \rho_i^2)^{-3/2} (1 - \rho_j^2)^{1/2} \quad (\forall i), \\ \xi''_{(i)(j)} &= \rho_i \rho_j (1 - \rho_i^2)^{-1/2} (1 - \rho_j^2)^{-1/2} \quad (\forall i, j), \\ \frac{\partial a_{ij}}{\partial \rho_\mu} &= \frac{\partial^2 a_{ij}}{\partial \rho_\mu^2} = \frac{\partial^2 a_{ij}}{\partial \rho_\mu \partial \rho_i} = \frac{\partial^2 a_{ij}}{\partial \rho_\mu \partial \rho_j} = 0 \quad (\forall \mu \neq i \wedge \forall \mu \neq j). \end{aligned} \quad (\text{A.7})$$

**Proof of Lemma A.2.** To prove the results of the lemma, it suffices to calculate the first- and second-order derivatives of matrices  $R^{ii}$  and  $R^{ij}$ , defined in (A.3) and (A.4), respectively, and of scalars  $a_{ij}$ , defined in (A.5).  $\square$

**Lemma A.3.** For the elements of matrix  $\Omega$  the following results hold:

$$\begin{aligned} \sum_{k=1}^M \sigma_{ik} \sigma^{ki} &= \sum_{k=1}^M \sigma^{ik} \sigma_{ki} = 1, \\ \sum_{k=1}^M \sigma_{ik} \sigma^{kj} &= \sum_{k=1}^M \sigma^{ik} \sigma_{kj} = 0 \quad (\forall i \neq j), \\ \sum_{k=1}^M \sigma_{ik} \sigma^{ki} R_{ik} R^{ki} &= \sum_{k=1}^M \sigma^{ik} \sigma_{ki} R^{ik} R_{ki} = I_{TM}, \\ \sum_{k=1}^M \sigma_{ik} \sigma^{kj} R_{ik} R^{kj} &= \sum_{k=1}^M \sigma^{ik} \sigma_{kj} R^{ik} R_{kj} = 0 \quad (\forall i \neq j). \end{aligned} \quad (\text{A.8})$$

**Proof of Lemma A.3.** The results of the lemma can be proved by noticing that that

$$\Omega^{-1} = P(\Sigma \otimes I_T)P' = [(\sigma_{ij} R_{ij})_{i,j=1, \dots, M}] \Rightarrow \Omega = [(\sigma^{ij} R^{ij})_{i,j=1, \dots, M}], \quad (\text{A.9})$$

since  $P$  is block diagonal,  $\Sigma \Sigma^{-1} = \Sigma^{-1} \Sigma = I_M$  and  $\Omega \Omega^{-1} = \Omega^{-1} \Omega = I_{TM}$ .  $\square$

To derive the partial derivatives of  $\Omega$  with respect to nuisance parameters, given in the next lemma, we need the following definitions. For the composite index  $(ij) = 1, \dots, M^2$ , defined in (20), let  $\varsigma_{(ij)} = \sigma^{ij}$  be the elements of the  $(M^2 \times 1)$  vector  $\varsigma = \text{vec}(\Sigma^{-1})$ . Also, let  $\Delta_{\mu\mu'} = [(\delta_{\mu i} \delta_{j \mu'})_{i,j=1, \dots, M}]$  be a  $(M \times M)$  matrix with 1 in the  $(\mu, \mu')$ -th position and 0's elsewhere. Then, for all  $\mu, \mu', \nu$  and  $\nu'$ , we have

$$\frac{\partial}{\partial \varsigma_{(\mu\mu')}} (\Sigma^{-1} \otimes I_T) = \Delta_{\mu\mu'} \otimes I_T, \quad \frac{\partial^2}{\partial \varsigma_{(\mu\mu')} \partial \varsigma_{(\nu\nu')}} (\Sigma^{-1} \otimes I_T) = 0. \quad (\text{A.10})$$

**Lemma A.4.** The partial derivatives of  $\Omega$ , with respect to the elements of vectors  $\varrho$  and  $\varsigma$ , can be analytically written as follows:

$$\Omega_{\varsigma_{(\mu\mu')}} = [(\delta_{\mu i} \delta_{j \mu'} R^{\mu\mu'})_{i,j=1, \dots, M}], \quad \Omega_{\varsigma_{(\mu\mu')} \varsigma_{(\nu\nu')}} = 0, \quad (\text{A.11})$$

$$\begin{aligned} \Omega_{\rho_\mu} &= [(\delta_{\mu i} \sigma^{\mu j} R_{\rho_\mu}^{\mu j} + \delta_{j \mu} \sigma^{i \mu} R_{\rho_\mu}^{i \mu} - \delta_{\mu i} \delta_{j \mu} \sigma^{\mu \mu} R_{\rho_\mu}^{\mu \mu})_{i,j=1, \dots, M}], \\ \Omega_{\rho_\mu \rho_\mu} &= [(\delta_{\mu i} \sigma^{\mu j} R_{\rho_\mu \rho_\mu}^{\mu j} + \delta_{j \mu} \sigma^{i \mu} R_{\rho_\mu \rho_\mu}^{i \mu} - \delta_{\mu i} \delta_{j \mu} \sigma^{\mu \mu} R_{\rho_\mu \rho_\mu}^{\mu \mu})_{i,j=1, \dots, M}], \\ \Omega_{\rho_\mu \rho_{\mu'}} &= [(\delta_{\mu i} \delta_{j \mu'} \sigma^{\mu \mu'} R_{\rho_\mu \rho_{\mu'}}^{\mu \mu'} + \delta_{\mu' i} \delta_{j \mu} \sigma^{\mu' \mu} R_{\rho_\mu \rho_{\mu'}}^{\mu' \mu} \\ &\quad - \delta_{\mu i} \delta_{j \mu} \sigma^{\mu \mu} \delta_{\mu \mu'} R_{\rho_\mu \rho_{\mu'}}^{\mu \mu})_{i,j=1, \dots, M}], \end{aligned} \quad (\text{A.12})$$

$$\begin{aligned} \Omega_{\rho_\mu \varsigma_{(\nu\nu')}} &= [(\delta_{\nu i} \delta_{j \nu'} \delta_{\mu \nu} R_{\rho_\mu}^{\mu \nu'} + \delta_{\nu i} \delta_{j \nu'} \delta_{\nu' \mu} R_{\rho_\mu}^{\nu \mu} - \delta_{\nu i} \delta_{j \nu'} \delta_{\mu \nu} \delta_{\nu' \mu} R_{\rho_\mu}^{\mu \mu})_{i,j=1, \dots, M}] \\ &\Rightarrow \Omega_{\rho_\mu \varsigma_{(\nu\nu')}} = 0 \quad (\forall \nu \neq \mu \wedge \forall \nu' \neq \mu). \end{aligned} \quad (\text{A.13})$$

**Proof of Lemma A.4.** To prove the results of the lemma, we rely on the results of Lemmas A.2 and A.3 in order to calculate the first- and second-order derivatives of matrix  $\Omega$  with respect to the elements of vectors  $\varrho$  and  $\varsigma$ . The results in (A.11) come immediately from equations (A.9) and (A.10).

Some comments must be made for the derivation of the results in (A.12) and (A.13). Matrix  $\Omega_{\rho_\mu}$  can be calculated as the sum of three matrices. The first matrix has non-zero elements on its  $\mu$ -th row; the second matrix has its non-zero elements on its  $\mu$ -th column; and the third matrix, which has only one non-zero element at the  $(\mu, \mu)$ -th position, is subtracted to correct for the double-counting of the derivative of the element at the intersection of the  $\mu$ -th row with the  $\mu$ -th column of matrix  $\Omega$ . The elements of matrix  $\Omega_{\rho_\mu \rho_\mu}$  can be readily calculated by taking the derivatives of the elements of  $\Omega_{\rho_\mu}$  with respect to  $\rho_\mu$ .

On taking the derivatives of the elements of  $\Omega_{\rho_\mu}$  with respect to  $\rho_{\mu'}$ , we can calculate the elements of matrix  $\Omega_{\rho_\mu \rho_{\mu'}}$ . Note that matrix  $\Omega_{\rho_\mu \rho_{\mu'}}$  has its non-zero elements at its  $(\mu, \mu')$ -th and  $(\mu', \mu)$ -th positions. The subtracted third term corrects for the double-counting of the derivative of the  $(\mu, \mu)$ -th element of matrix  $\Omega$  in cases with  $\mu' = \mu$ . The third term is eliminated, as it should be, in cases with  $\mu' \neq \mu$ .  $\square$

To derive the elements of the product of matrices  $\Omega_i \Omega^{-1} \Omega_j$ , needed for the partial derivatives of matrix  $A$  (see Lemmas A.14 – A.17), we define the following matrices:

$$\begin{aligned}
W_{ij} = & \sigma^{i\mu} \sigma_{\mu\mu'} \sigma^{\mu'j} R_{\rho_\mu}^{i\mu} R_{\mu\mu'} R_{\rho_{\mu'}}^{\mu'j} \\
& + \delta_{\mu i} \left\{ \left[ \sum_{k=1}^M \sigma^{\mu k} \sigma_{k\mu'} R_{\rho_\mu}^{\mu k} R_{k\mu'} \right] - \sigma^{\mu\mu} \sigma_{\mu\mu'} R_{\rho_\mu}^{\mu\mu} R_{\mu\mu'} \right\} \sigma^{\mu'j} R_{\rho_{\mu'}}^{\mu'j} \\
& + \delta_{j\mu'} \sigma^{i\mu} R_{\rho_\mu}^{i\mu} \left\{ \left[ \sum_{r=1}^M \sigma_{\mu r} \sigma^{r\mu'} R_{\mu r} R_{\rho_{\mu'}}^{r\mu'} \right] - \sigma_{\mu\mu'} \sigma^{\mu'\mu'} R_{\mu\mu'} R_{\rho_{\mu'}}^{\mu'\mu'} \right\} \\
& + \delta_{\mu i} \delta_{j\mu'} \left\{ \sum_{k=1}^M \sum_{r=1}^M \sigma^{\mu k} \sigma_{k r} \sigma^{r\mu'} R_{\rho_\mu}^{\mu k} R_{k r} R_{\rho_{\mu'}}^{r\mu'} \right. \\
& - \left[ \sum_{k=1}^M \sigma^{\mu k} \sigma_{k\mu'} R_{\rho_\mu}^{\mu k} R_{k\mu'} \right] \sigma^{\mu'\mu'} R_{\rho_{\mu'}}^{\mu'\mu'} \\
& - \sigma^{\mu\mu} R_{\rho_\mu}^{\mu\mu} \left[ \sum_{r=1}^M \sigma_{\mu r} \sigma^{r\mu'} R_{\mu r} R_{\rho_{\mu'}}^{r\mu'} \right] \\
& \left. + \sigma^{\mu\mu} \sigma_{\mu\mu'} \sigma^{\mu'\mu'} R_{\rho_\mu}^{\mu\mu} R_{\mu\mu'} R_{\rho_{\mu'}}^{\mu'\mu'} \right\}, \tag{A.14}
\end{aligned}$$

$$\Omega_{\rho_\mu \rho_{\mu'}}^* = \Omega_{\rho_\mu} \Omega^{-1} \Omega_{\rho_{\mu'}}, \quad \Omega_{\varsigma_{(\mu\mu')}\varsigma_{(\nu\nu')}}^* = \Omega_{\varsigma_{(\mu\mu')}} \Omega^{-1} \Omega_{\varsigma_{(\nu\nu')}}, \tag{A.15}$$

$$\Omega_{\rho_\mu \varsigma_{(\nu\nu')}}^* = \Omega_{\rho_\mu} \Omega^{-1} \Omega_{\varsigma_{(\nu\nu')}} \quad \text{and} \quad \Omega_{\varsigma_{(\nu\nu')}\rho_\mu}^* = \Omega_{\varsigma_{(\nu\nu')}} \Omega^{-1} \Omega_{\rho_\mu}.$$

**Lemma A.5.** *The elements of matrices  $\Omega_{\rho_\mu \rho_{\mu'}}^*$ ,  $\Omega_{\varsigma_{(\mu\mu')\varsigma_{(\nu\nu')}}}^*$ ,  $\Omega_{\rho_\mu \varsigma_{(\nu\nu')}}^*$  and  $\Omega_{\varsigma_{(\nu\nu')\rho_\mu}}^*$  can be analytically written as follows:*

$$\begin{aligned}\Omega_{\rho_\mu \rho_{\mu'}}^* &= [(W_{ij})_{i,j=1, \dots, M}], \\ \Omega_{\varsigma_{(\mu\mu')\varsigma_{(\nu\nu')}}}^* &= [(\delta_{\mu i} \delta_{j\nu'} \sigma_{\mu' \nu} R^{\mu\nu'})_{i,j=1, \dots, M}], \\ \Omega_{\rho_\mu \varsigma_{(\nu\nu')}}^* &= \left[ \left( \left( \sum_{k=1}^M \sigma^{ik} \sigma_{k\nu} R_{\rho_\mu}^{ik} R_{k\nu} \right) \delta_{j\nu'} R^{\nu\nu'} \right)_{i,j=1, \dots, M} \right], \\ \Omega_{\varsigma_{(\nu\nu')\rho_\mu}}^* &= \left[ \left( \delta_{\nu i} R^{\nu\nu'} \left( \sum_{r=1}^M \sigma_{\nu' r} \sigma^{rj} R_{\nu' r} R_{\rho_\mu}^{rj} \right) \right)_{i,j=1, \dots, M} \right].\end{aligned}\tag{A.16}$$

**Proof of Lemma A.5.** The results in (A.16) can be easily proved by combining Lemma A.4 with equations (A.9) and (A.14).  $\square$

## Asymptotic expansions of estimators

In the next lemmas we derive useful asymptotic expansions for all estimators of matrix  $B$  and of the nuisance parameters considered in the paper. In each case, these estimators are indexed by  $I$  (see footnotes 2 and 3).

**Lemma A.6.** *All estimators  $\hat{B}_I$  ( $I = UL, RL, GL, IG, ML$ ) of matrix  $B$ , defined in (21), admit a stochastic expansion of the form*

$$\hat{B}_I = B + \tau B_1^I + \omega(\tau^2),\tag{A.17}$$

where

$$\begin{aligned}B_1^{UL} &= \sqrt{T}(Z'Z)^{-1}Z'E, \\ \text{vec}(B_1^{RL}) &= \sqrt{T}\Psi(X_*'X_*)^{-1}X_*'\varepsilon, \\ \text{vec}(B_1^{GL}) &= \text{vec}(B_1^{IG}) = \text{vec}(B_1^{ML}) \\ &= \sqrt{T}\Psi[X_*(\Sigma_I^{-1} \otimes I_T)X_*]^{-1}X_*(\Sigma_I^{-1} \otimes I_T)\varepsilon.\end{aligned}\tag{A.18}$$

**Proof of Lemma A.6.** The results of the lemma follow immediately from models (16) and (21), and the definitions of all estimators  $B_I$  considered (see footnote 3). Thus, since  $\tau = 1/\sqrt{T}$ , we can readily find that

$$\hat{B}_{UL} = (Z'Z)^{-1}Z'(ZB + E) = B + \tau[\sqrt{T}(Z'Z)^{-1}Z'E] = B + B_1^{UL}.\tag{A.19}$$

Similarly, since (23) implies that  $\text{vec}(B) = \Psi\beta$ , we can easily find that

$$\begin{aligned}\text{vec}(\hat{B}_{RL}) &= \Psi(X_*'X_*)^{-1}X_*(X_*\beta + \varepsilon) = \text{vec}(B) + \tau[\sqrt{T}\Psi(X_*'X_*)^{-1}X_*'\varepsilon] \\ &= \text{vec}(B) + \tau \text{vec}(B_1^{RL}).\end{aligned}\tag{A.20}$$



The result for estimator  $\hat{B}_{GL}$  can be proved according to (A.20), taking into account that  $\hat{\Sigma}_I = \Sigma + \omega(\tau)$ , for any consistent estimator  $\hat{\Sigma}_I$  of matrix  $\Sigma$ , indexed by  $I$ .  $\square$

Let  $\hat{E}_I$  be the residuals corresponding to the estimators  $\hat{B}_I$ . Then, the following lemma holds for the estimators  $\hat{\Sigma}_I$  and  $\hat{\Sigma}_I^{-1}$  of matrix  $\Sigma$  and its inverse, respectively, based on  $\hat{E}_I$ .

**Lemma A.7.** *All estimators  $\hat{\Sigma}_I$  ( $I = UL, RL, GL, IG, ML$ ) of matrix  $\Sigma$  admit a stochastic expansion of the form*

$$\hat{\Sigma}_I = \Sigma + \tau(\Sigma_1 + \tau\Sigma_2^I) + \omega(\tau^3), \quad (\text{A.21})$$

where

$$\Sigma_1 = \sqrt{T}(E'E/T - \Sigma), \quad \Sigma_2^I = (B_1^I - B_1^{UL})'\Gamma(B_1^I - B_1^{UL}) - E'P_Z E, \quad (\text{A.22})$$

$\Gamma$  is any conformable matrix and  $P_Z$  is the orthogonal projector spanned by the columns of matrix  $Z$ . Estimator  $\hat{\Sigma}_I^{-1}$  admits a stochastic expansion of the form

$$\hat{\Sigma}_I^{-1} = \Sigma^{-1} - \tau S_1 + \tau^2 S_2^I + \omega(\tau^3), \quad (\text{A.23})$$

where

$$S_1 = \Sigma^{-1}\Sigma_1\Sigma^{-1}, \quad S_2^I = \Sigma^{-1}(\Sigma_1\Sigma^{-1}\Sigma_1 - \Sigma_2^I)\Sigma^{-1}. \quad (\text{A.24})$$

**Proof of Lemma A.7.** By using model (21) and Lemma A.6 we find that

$$\hat{E}_I = Y_* - Z\hat{B}_I = ZB + E - Z(B + \tau B_1^I + \omega(\tau^2)) = E - \tau ZB_1^I + \omega(\tau^2). \quad (\text{A.25})$$

Moreover, from the definition of matrix  $\Gamma$  and (A.18) we find that

$$(B_1^I)'Z'E/\sqrt{T} = (B_1^I)'(Z'Z/T)(Z'Z/T)^{-1}Z'E/\sqrt{T} = (B_1^I)'\Gamma B_1^{UL}. \quad (\text{A.26})$$

Then, since  $\hat{\Sigma}_I = \hat{E}_I'\hat{E}_I/T$ , equations (A.22), (A.25) and (A.26) imply that

$$\begin{aligned} \hat{\Sigma}_I &= E'E/T + \tau^2 \left[ (B_1^I)'\Gamma B_1^I - (B_1^{UL})'\Gamma B_1^I - (B_1^I)'Z^\Gamma B_1^{UL} \right] + \omega(\tau^3) \\ &= \Sigma + \tau\sqrt{T}(E'E/T - \Sigma) \\ &\quad + \tau^2 \left[ (B_1^I - B_1^{UL})'\Gamma(B_1^I - B_1^{UL}) - E'P_Z E \right] + \omega(\tau^3), \end{aligned} \quad (\text{A.27})$$

which completes the proof of (A.21). To prove (A.23), it suffices to use (A.24) and equation (2.6) in (Magdalinos 1992, Corollary 1), which implies that

$$\begin{aligned} \hat{\Sigma}_I^{-1} &= \left[ \Sigma + \tau(\Sigma_1 + \tau\Sigma_2^I) + \omega(\tau^3) \right]^{-1} \\ &= \Sigma^{-1} - \tau\Sigma^{-1}(\Sigma_1 + \tau\Sigma_2^I)\Sigma^{-1} \\ &\quad + \tau^2\Sigma^{-1}(\Sigma_1 + \tau\Sigma_2^I)\Sigma^{-1}(\Sigma_1 + \tau\Sigma_2^I)\Sigma^{-1} + \omega(\tau^3). \end{aligned} \quad (\text{A.28})$$

$\square$

The stochastic expansion of estimator of vector  $\varsigma$ , denoted as  $\hat{\varsigma}_I$ , is given in the next lemma:

**Lemma A.8.** *All estimators  $\hat{\varsigma}_I = \text{vec}([\hat{E}_I' \hat{E}_I / T]^{-1})$  of vector  $\varsigma$ , indexed by  $I = UL, RL, GL, IG, ML$ , admit a stochastic expansion of the form*

$$\hat{\varsigma}_I = \varsigma - \tau \text{vec}(S_1) + \tau^2 \text{vec}(S_2^I) + \omega(\tau^3) \quad (\text{A.29})$$

and thus, the  $(M^2 \times 1)$  vector  $\delta_\varsigma = (\hat{\varsigma} - \varsigma)/\tau$ , with elements  $\delta_{\varsigma(\mu\mu')}$  defined in (27), admits a stochastic expansion of the form

$$\begin{aligned} \delta_\varsigma &= -\text{vec}(S_1) + \tau \text{vec}(S_2^I) + \omega(\tau^2) \\ &= d_{1\varsigma} + \tau d_{2\varsigma} + \omega(\tau^2), \end{aligned} \quad (\text{A.30})$$

which implies that

$$d_{1\varsigma} = -\text{vec}(S_1), \quad d_{2\varsigma} = \text{vec}(S_2^I). \quad (\text{A.31})$$

**Proof of Lemma A.8.** The proof follows immediately from equations (21), (29), (32) and (A.23).  $\square$

To derive the stochastic expansion of the estimators of  $\sigma$ , denoted as  $\hat{\sigma}_I$ , we define the following  $(M \times M)$  matrices, indexed by  $I$ :

$$\begin{aligned} \Delta_I &= \lim_{T \rightarrow \infty} T \mathcal{E}[(\hat{B}_I - \hat{B}_{UL})' \Gamma (\hat{B}_I - \hat{B}_{UL})] \\ &= \lim_{T \rightarrow \infty} \mathcal{E}[(B_1^I - B_1^{UL})' \Gamma (B_1^I - B_1^{UL})], \end{aligned} \quad (\text{A.32})$$

where  $\Gamma$  is any conformable matrix.

**Lemma A.9.** *All estimators  $\hat{\sigma}_I^2$  ( $I = UL, RL, GL, IG, ML$ ) of  $\sigma^2$  (see footnote 1) satisfy the relation*

$$\begin{aligned} \hat{\sigma}_I^2 &= \text{tr}(\hat{\Sigma}_I^{-1} \hat{\Sigma}_J) / (M - \tau^2 n) \\ &= \{M + \tau^2 \text{tr}[(S_2^I - S_2^J) \Sigma]\} / (M - \tau^2 n) + \omega(\tau^3). \end{aligned} \quad (\text{A.33})$$

The last equation implies that

$$\begin{aligned} (\hat{\sigma}_I^2 - 1)/\tau &= \{M/\tau + \tau \text{tr}[(S_2^I - S_2^J) \Sigma]\} / (M - \tau^2 n) - 1/\tau + \omega(\tau^2) \\ &= \tau \{\text{tr}[(S_2^I - S_2^J) \Sigma] + n\} / M + \omega(\tau^2), \end{aligned} \quad (\text{A.34})$$

i.e., scalar  $\delta_0$ , defined in (27), admits a stochastic expansion of the form

$$\delta_0 = \sigma_0 + \tau \sigma_1 + \omega(\tau^2), \quad (\text{A.35})$$

which in turn implies that

$$\sigma_0 = 0 \quad \text{and} \quad \sigma_1 = \{\text{tr}[(S_2^I - S_2^J) \Sigma] + n\} / M. \quad (\text{A.36})$$

**Proof of Lemma A.9.** To prove the lemma we rely on the following results (see (A.37) and (A.38)): Since the rows  $\varepsilon_t$  ( $t = 1, \dots, T$ ) of  $E$  are independent  $\mathcal{N}_M(0, \Sigma)$  random vectors, matrix  $E'E$  is a Wishart matrix with weight matrix  $\Sigma$  and  $T$  degrees of freedom, i.e.,  $E'E \sim \mathcal{W}_M(\Sigma, T)$  and  $\mathcal{E}(E'E) = T\Sigma$ . Then, it is easy to show that

$$\mathcal{E}(E'E\Sigma^{-1}E'E) = T(M + T + 1)\Sigma. \quad (\text{A.37})$$

Moreover, since  $E'E \sim \mathcal{W}_M(\Sigma, T)$  and  $P_Z$  is idempotent of rank  $K$ , it follows that matrix  $E'P_Z E \sim \mathcal{W}_M(\Sigma, K)$  and  $\mathcal{E}(E'P_Z E) = \text{tr}(P_Z)\Sigma = K\Sigma$ . Further,  $\mathcal{E}(\Sigma_1) = 0$ ,  $\mathcal{E}(\Sigma_1\Sigma^{-1}\Sigma_1) = (M + 1)\Sigma$  and

$$\mathcal{E}(S_1) = 0, \quad \mathcal{E}(S_2^I) = (M + K + 1)\Sigma^{-1} - \Sigma^{-1}\mathcal{E}[(B_1^I - B_1^{UL})'\Gamma(B_1^I - B_1^{UL})]\Sigma^{-1}. \quad (\text{A.38})$$

Let  $\hat{\varepsilon}_{GL} = \text{vec}(\hat{E}_{GL})$  be the  $GL$  residuals of regression equation (16). Then, the corresponding estimator of  $\Sigma$  is  $\hat{\Sigma}_J = \hat{E}_{GL}'\hat{E}_{GL}/T$ . Also, let  $\hat{\beta}_{GL}$  be the  $GL$  estimator of  $\beta$  in (16). Define the  $(M \times M)$  matrices  $M_I = \lim_{T \rightarrow \infty} \mathcal{E}(S_2^I)$  ( $I = UL, RL, GL, IG, ML$ ) and the  $(M^2 \times M^2)$  matrix  $N$  whose  $((ij), (kr))$ -th element is  $\nu_{(ij)(kr)} = \sigma_{ik}\sigma_{jr} + \sigma_{ir}\sigma_{jk}$  ( $i, j, k, r = 1, \dots, M$ ). Then, (A.32) and (A.38) imply that

$$M_I = (M + K + 1)\Sigma^{-1} - \Sigma^{-1}\Delta_I\Sigma^{-1} \quad (\text{A.39})$$

$$\Rightarrow \lim_{T \rightarrow \infty} T\mathcal{E}[(S_2^I - S_2^J)\Sigma] = (M_I - M_{GL})\Sigma = \Sigma^{-1}(\Delta_{GL} - \Delta_I), \quad (\text{A.40})$$

where

$$\begin{aligned} \Delta_{UL} &= 0, \\ \Delta_{RL} &= \left[ \left( \text{tr}(B_{ii}^{-1}B_{ij}B_{jj}^{-1}B_{ji}) - n_i - n_j + K \right) \sigma_{ij} \right]_{i,j=1, \dots, M}, \\ \Delta_{GL} &= \Delta_{IG} = \Delta_{ML} = K\Sigma - \left[ \left( \text{tr}(G_{ij}B_{ji}) \right) \right]_{i,j=1, \dots, M}. \end{aligned} \quad (\text{A.41})$$

Since  $E'E \sim \mathcal{W}_M(\Sigma, T)$  and  $\mathcal{E}(E'E) = T\Sigma$ , matrix  $W_* = \sqrt{T}\Sigma_1 = E'E - T\Sigma$ , with elements  $w_{ij}$ , is a Wishart matrix in deviations from its expected values. Following Zellner (1971, page 389, equation (B.58)), we find that

$$\mathcal{E}(w_{ij}w_{kr}) = T(\sigma_{ik}\sigma_{jr} + \sigma_{ir}\sigma_{jk}) = T\nu_{(ij)(kr)} \quad (\text{A.42})$$

$$\Rightarrow \lim_{T \rightarrow \infty} \mathcal{E}[(\text{vec}(S_1))(\text{vec}(S_1))'] = (\Sigma^{-1} \otimes \Sigma^{-1})N(\Sigma^{-1} \otimes \Sigma^{-1}). \quad (\text{A.43})$$

The proof of the lemma can be completed using the following relationship:

$$(M - \tau^2 n)^{-1} = M^{-1}(1 - \tau^2 n/M)^{-1} = (1 + \tau^2 n/M)/M + \omega(\tau^4). \quad (\text{A.44})$$

□

Before deriving the asymptotic expansion of the estimators of  $\rho_\mu$ , next we define the following  $(T \times T)$  matrices:

$$R_i^{\mu\mu} = R_{\rho_\mu}^{\mu\mu} + i\rho_\mu\Delta \quad (i = 1, 2), \quad V_\mu = [I - X_\mu(X_\mu' R^{\mu\mu} X_\mu)^{-1} X_\mu' R_{\mu\mu}] R^{\mu\mu}. \quad (\text{A.45})$$

The first assumption in Subsection 3.1 implies that matrices

$$B_{\mu\mu} = X_\mu' R^{\mu\mu} X_\mu / T \quad \text{and} \quad F_{\mu\mu} = X_\mu' X_\mu / T \quad (\text{A.46})$$

converge to non-singular matrices, as  $T \rightarrow \infty$ , and that matrices

$$X_\mu' \Delta X_\mu / T, \quad X_\mu' \Delta R_{\mu\mu} X_\mu / T, \quad X_\mu' R_{\mu\mu} \Delta X_\mu / T, \quad (\text{A.47})$$

$$X_\mu' \Delta R_{\mu\mu} \Delta X_\mu / T \quad \text{and} \quad \Theta_{\mu\mu} = X_\mu' R_{\mu\mu} X_\mu / T$$

are of order  $O(T^{-1})$ . All the above matrices help to derive expectations of products of quadratic forms of  $u$ , needed in the expansions of estimators of  $\rho_\mu$ . These are given in the next lemma:

**Lemma A.10.** *For quadratic forms of vector  $u$ , we have the following results:*

$$\begin{aligned} \mathcal{E}(u_\mu' R_2^{\mu\mu} u_\mu) &= \frac{2\rho_\mu \sigma_{\mu\mu}}{1 - \rho_\mu^2}, \\ \mathcal{E}(u_\mu' u_\mu u_\mu' R_2^{\mu\mu} u_\mu) &= -\frac{2T\rho_\mu \sigma_{\mu\mu}^2}{(1 - \rho_\mu^2)^2} + O(1), \\ \mathcal{E}(u_\mu' R_2^{\mu\mu} u_\mu u_\mu' R_2^{\mu\mu} u_\mu) &= \frac{4T\sigma_{\mu\mu}^2}{1 - \rho_\mu^2} + O(1), \\ \mathcal{E}(u_\mu' R_2^{\mu\mu} u_\mu u_{\mu'}' R_2^{\mu'\mu'} u_{\mu'}) &= \frac{4T\sigma_{\mu\mu} \sigma_{\mu'\mu'}}{1 - \rho_\mu \rho_{\mu'}} + O(1), \\ \mathcal{E}(u_\mu' \bar{P}_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu) &= \frac{\sigma_{\mu\mu}}{\rho_\mu} [n_\mu - \text{tr}(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu})] \\ &\quad + O(T^{-1}), \\ \mathcal{E}(u_\mu' \bar{P}_{X_\mu} R_2^{\mu\mu} \bar{P}_{X_\mu} u_\mu) &= \frac{\sigma_{\mu\mu}}{\rho_\mu} [2 [\rho_\mu^2 / (1 - \rho_\mu^2) - n_\mu] \\ &\quad + (1 - \rho_\mu^2) \text{tr}(F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \\ &\quad + \text{tr}(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu})] + O(T^{-1}), \\ \mathcal{E}(u_\mu' R^{\mu\mu} V_\mu P_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu) &= \frac{\sigma_{\mu\mu}}{\rho_\mu} [ [\text{tr}(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu}) - n_\mu] \\ &\quad + (1 - \rho_\mu^2) [\text{tr}(F_{\mu\mu} B_{\mu\mu}^{-1}) - \text{tr}(F_{\mu\mu}^{-1} \Theta_{\mu\mu})] ] \\ &\quad + O(T^{-1}). \end{aligned} \quad (\text{A.48})$$

**Proof of Lemma A.10.** We begin the proof by noticing that  $\text{tr}(R_{\mu\mu}) = T/(1 - \rho_\mu^2)$  and  $\text{tr}(R_1^{\mu\mu} R_{\mu\mu}) = 0$ . Next, we define  $r = \rho_\mu^2$ , which implies that  $|r| < 1$ . Then, using

the following results:

$$\sum_{i=1}^T r^i = \frac{r(1-r^T)}{1-r}, \quad \sum_{i=0}^T r^i = \frac{1-r^{T+1}}{1-r}, \quad (\text{A.49})$$

$$\sum_{i=1}^T ir^i = \sum_{i=0}^T ir^i = \frac{r[1-(T+1)r^T + Tr^{T+1}]}{(1-r)^2},$$

which hold for any  $0 \leq r < 1$ , we can readily calculate the following traces:

$$\begin{aligned} tr(R_2^{\mu\mu} R_{\mu\mu}) &= \frac{2\rho_\mu}{1-\rho_\mu^2}, \quad tr(\Delta R_{\mu\mu}) = \frac{2}{1-\rho_\mu^2}, \quad tr\{(\Delta R_{\mu\mu})^2\} = \frac{2(1+\rho_\mu^{2(T-1)})}{(1-\rho_\mu^2)^2}, \\ tr\{(\Delta R_{\mu\mu})^3\} &= \frac{2(1+3\rho_\mu^{2(T-1)})}{(1-\rho_\mu^2)^3}, \quad tr(\Delta R_{\mu\mu}^3) = \frac{2}{(1-\rho_\mu^2)^4} + O(T^{-1}), \\ tr(R_{\mu\mu} \Delta R_{\mu\mu}) &= \frac{2(1-\rho_\mu^{2T})}{(1-\rho_\mu^2)^3}, \\ tr\{R_{\mu\mu}(\Delta R_{\mu\mu})^2\} &= \frac{2}{(1-\rho_\mu^2)^3} \left[ T\rho_\mu^{2(T-1)} + \frac{1-\rho_\mu^{2T}}{1-\rho_\mu^2} \right], \quad (\text{A.50}) \\ tr(R_{\mu\mu}^2)/T &= \frac{1+\rho_\mu^2}{(1-\rho_\mu^2)^3} + O(T^{-1}), \quad tr(R_{\mu\mu}^3)/T = \frac{1+\rho_\mu^4}{(1-\rho_\mu^2)^5} + O(T^{-1}), \\ tr(\rho_\mu^2 \Delta R_{\mu\mu} \Delta R_{\mu\mu}) &= \frac{2\rho_\mu^2(1+\rho_\mu^{2(T-1)})}{(1-\rho_\mu^2)^2}, \\ tr(\rho_\mu R_1^{\mu\mu} R_{\mu\mu} \Delta R_{\mu\mu}) &= \frac{2}{1-\rho_\mu^2} + \frac{2(1-\rho_\mu^{2T})}{(1-\rho_\mu^2)^2}. \end{aligned}$$

Note that in calculating the traces in (A.50), terms of the form  $T^n \rho_\mu^{2T} \rightarrow 0$  since  $0 \leq \rho_\mu < 1$  and L' Hospital's rule implies that  $\lim_{T \rightarrow \infty} T^n \rho_\mu^{2T} = 0$ .

Then, by using definitions (A.45), (A.46) and (A.47), the results in (A.50) and a large amount of tedious algebra, we can compute the following traces:

$$\begin{aligned} tr(R_{\mu\mu} R_i^{\mu\mu} R_{\mu\mu}) &= -\frac{2T\rho_\mu}{(1-\rho_\mu^2)^2} + O(1), \quad tr\{(R_i^{\mu\mu} R_{\mu\mu})^2\} = \frac{2T}{1-\rho_\mu^2} + O(1), \\ tr\{R_{\mu\mu}(R_i^{\mu\mu} R_{\mu\mu})^2\} &= \frac{2T(2\rho_\mu^2-1)}{(1-\rho_\mu^2)^3} + O(1), \quad tr\{(R_i^{\mu\mu} R_{\mu\mu})^3\} = \frac{2T(2-3\rho_\mu^2)}{\rho_\mu(1-\rho_\mu^2)^2} + O(1), \\ tr(P_{X_\mu} R_i^{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ tr(B_{\mu\mu} F_{\mu\mu}^{-1}) - (1-\rho_\mu^2)n_\mu \right] + O(T^{-1}), \quad (\text{A.51}) \\ tr(P_{X_\mu} R_i^{\mu\mu} R_{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ n_\mu - (1-\rho_\mu^2)tr(F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right] + O(T^{-1}), \\ tr(P_{X_\mu} R_i^{\mu\mu} P_{X_\mu} R_{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ tr(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu}) - (1-\rho_\mu^2)tr(F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right] + O(T^{-1}), \end{aligned}$$

where  $i = 1, 2$ . Working similarly we can calculate the following traces:

$$\begin{aligned} tr(\bar{P}_{X_\mu} R_2^{\mu\mu} \bar{P}_{X_\mu} R_{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ \frac{2[\rho_\mu^2 - n_\mu(1-\rho_\mu^2)]}{1-\rho_\mu^2} + (1-\rho_\mu^2)tr(F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right. \\ &\quad \left. + tr(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right] + O(T^{-1}), \quad (\text{A.52}) \\ tr(\bar{P}_{X_\mu} R_2^{\mu\mu} P_{X_\mu} R_{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ n_\mu - tr(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right] + O(T^{-1}), \end{aligned}$$

and

$$\begin{aligned} \text{tr}(R^{\mu\mu} V_\mu P_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} R_{\mu\mu}) &= \frac{1}{\rho_\mu} \left[ \text{tr}(F_{\mu\mu}^{-1} B_{\mu\mu} F_{\mu\mu}^{-1} \Theta_{\mu\mu}) - n_\mu \right] \\ &\quad + \frac{1 - \rho_\mu^2}{\rho_\mu} \left[ \text{tr}(F_{\mu\mu} B_{\mu\mu}^{-1}) - \text{tr}(F_{\mu\mu}^{-1} \Theta_{\mu\mu}) \right] \\ &\quad + O(T^{-1}). \end{aligned} \quad (\text{A.53})$$

The results in (A.48) follow then by using the result given in page 389 of Magnus and Neudecker (1979).  $\square$

The stochastic expansion of the  $LS$  estimator of  $\rho_\mu$  is given in the next lemma:

**Lemma A.11.** *The  $LS$  estimator of  $\rho_\mu$ , denoted as  $\tilde{\rho}_\mu$ , admits a stochastic expansion of the form*

$$\tilde{\rho}_\mu = \rho_\mu + \tau \left( \rho_\mu^{(1)} + \tau \rho_\mu^{(2)} \right) + \omega(\tau^3), \quad (\text{A.54})$$

where

$$\rho_\mu^{(1)} = -\frac{u'_\mu R_2^{\mu\mu} u_\mu}{2\sqrt{T}\sigma_{u_\mu}^2}, \quad \rho_\mu^{(2)} = -\frac{u'_\mu \bar{P}_{X_\mu} R_2^{\mu\mu} \bar{P}_{X_\mu} u_\mu}{2\sigma_{u_\mu}^2} + \frac{u'_\mu u_\mu u'_\mu R_2^{\mu\mu} u_\mu}{2T\sigma_{u_\mu}^4}. \quad (\text{A.55})$$

**Proof of Lemma A.11.** To prove the lemma, we rely on the following results (see (A.56) – (A.59)): Let  $\varepsilon_{ti}$  be the  $(t, i)$ -th element of matrix  $E$ . Then, the  $(i, j)$ -th element of matrix  $E'E/T$  is

$$e_{ij} = \sum_{t=1}^T \varepsilon_{ti} \varepsilon_{tj} / T = \varepsilon'_i \varepsilon_j / T, \quad (\text{A.56})$$

where  $\varepsilon_i$  is the  $i$ -th column of matrix  $E$ . Since  $\sigma_{ij}$  and  $\sigma^{ij}$  are the  $(i, j)$ -th elements of matrices  $\Sigma$  and  $\Sigma^{-1}$ , respectively,  $\Sigma^{-1} = \Sigma^{-1} \Sigma \Sigma^{-1}$  implies that  $\sigma^{ij} = \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma_{kr} \sigma^{rj}$ . Hence, the  $(i, j)$ -th element of matrix  $\Sigma_1$  in Lemma A.7 is given as

$$\sigma_{ij}^{(1)} = \sqrt{T} (e_{ij} - \sigma_{ij}) \quad (\text{A.57})$$

and the  $(ij)$ -th element of the  $(M^2 \times 1)$  vector  $\text{vec}(S_1)$ , where  $S_1 = \Sigma^{-1} \Sigma_1 \Sigma^{-1}$ , is given as

$$s_{(ij)}^{(1)} = \sqrt{T} \left\{ \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} (\varepsilon'_k \varepsilon_r / T) \sigma^{rj} - \sigma^{ij} \right\}. \quad (\text{A.58})$$

Since  $u_\mu = P_\mu \varepsilon_\mu \Rightarrow u'_\mu R_2^{\mu\mu} u_\mu = \varepsilon'_\mu P'_\mu R_2^{\mu\mu} P_\mu \varepsilon_\mu$  and  $R_{\mu\mu} = P_\mu P'_\mu$ , we can show that

$$\begin{aligned} \mathcal{E}(u'_\mu R_2^{\mu\mu} u_\mu) &= \sigma_{\mu\mu} \text{tr}(R_2^{\mu\mu} R_{\mu\mu}) \Rightarrow \\ &\Rightarrow \mathcal{E}[(\varepsilon'_k \varepsilon_r / T) u'_\mu R_2^{\mu\mu} u_\mu] = \sigma_{kr} \sigma_{\mu\mu} \frac{2\rho_\mu}{1 - \rho_\mu^2} + O(T^{-1}) \\ &\Rightarrow \mathcal{E} \left( s_{(ij)}^{(1)} u'_\mu R_2^{\mu\mu} u_\mu \right) = \sqrt{T} \sigma_{\mu\mu} \frac{2\rho_\mu}{1 - \rho_\mu^2} \left\{ \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma_{kr} \sigma^{rj} - \sigma^{ij} \right\} + O(T^{-1/2}) \\ &\Rightarrow \lim_{T \rightarrow \infty} \mathcal{E} \left( s_{(ij)}^{(1)} u'_\mu R_2^{\mu\mu} u_\mu \right) = 0. \end{aligned} \quad (\text{A.59})$$

The rest of the proof follows using Lemma A.10.  $\square$

The stochastic expansions of the rest of the estimators of  $\rho_\mu$ , listed in footnote 2, are given in the next lemma:

**Lemma A.12.** *The GL, PW, ML and DW estimators of  $\rho_\mu$  admit the following stochastic expansions, respectively:*

$$\begin{aligned}\hat{\rho}_\mu^{GL} = \hat{\rho}_\mu^{PW} &= \tilde{\rho}_\mu - \tau^2 \frac{1 - \rho_\mu^2}{\sigma_{\mu\mu}} \left[ u'_\mu \bar{P}_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu \right. \\ &\quad \left. + u'_\mu R^{\mu\mu} V_\mu P_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu / 2 \right] + \omega(\tau^3), \\ \hat{\rho}_\mu^{ML} &= \hat{\rho}_\mu^{GL} + \tau^2 \left[ \rho_\mu \frac{1 - \rho_\mu^2}{\sigma_{\mu\mu}} (u_{1\mu}^2 + u_{T\mu}^2) - \rho_\mu \right] + \omega(\tau^3), \\ \hat{\rho}_\mu^{DW} &= \tilde{\rho}_\mu + \tau^2 \frac{1 - \rho_\mu^2}{2\sigma_{\mu\mu}} (u_{1\mu}^2 + u_{T\mu}^2) + \omega(\tau^3).\end{aligned}\quad (\text{A.60})$$

**Proof of Lemma A.12.** The results of the lemma can be easily proved based on Magee (1985, pages 279–281) for the GL and iterative PW estimators of  $\rho_\mu$ , Beach and MacKinnon (1978, pages 52–54) and Magee (1985, pages 281–284) for the ML estimator, and using Lemma A.11 and the definition of the DW estimator of  $\rho_\mu$ .  $\square$

The stochastic expansion of the elements of vector  $\delta_\varrho$ , are given in the next lemma:

**Lemma A.13.** *The  $(M \times 1)$  vector  $\delta_\varrho = \sqrt{T}(\hat{\varrho} - \varrho)/\tau$ , with elements  $\delta_{\rho_\mu}$  defined in (27), admits a stochastic expansion of the form*

$$\delta_\varrho = d_{1\varrho} + \tau d_{2\varrho} + \omega(\tau^2). \quad (\text{A.61})$$

For estimators  $\hat{\rho}_\mu^I$  ( $I = LS, GL, PW, ML, DW$ ), the elements of  $d_{1\varrho}$  and  $d_{2\varrho}$  in (A.61) are analytically given as follows:  $d_{(1)\rho_\mu}^{GL} = d_{(1)\rho_\mu}^{PW} = d_{(1)\rho_\mu}^{ML} = d_{(1)\rho_\mu}^{DW} = d_{(1)\rho_\mu}^{LS}$  and

$$\begin{aligned}d_{(1)\rho_\mu}^{LS} &= \rho_\mu^{(1)}, \\ d_{(2)\rho_\mu}^{LS} &= \rho_\mu^{(2)}, \\ d_{(2)\rho_\mu}^{GL} = d_{(2)\rho_\mu}^{PW} &= d_{(2)\rho_\mu}^{LS} - \frac{1 - \rho_\mu^2}{\sigma_{\mu\mu}} \left[ u'_\mu \bar{P}_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu \right. \\ &\quad \left. + u'_\mu R^{\mu\mu} V_\mu P_{X_\mu} R_2^{\mu\mu} P_{X_\mu} V_\mu R^{\mu\mu} u_\mu / 2 \right], \\ d_{(2)\rho_\mu}^{ML} &= d_{(2)\rho_\mu}^{GL} + \rho_\mu \frac{1 - \rho_\mu^2}{\sigma_{\mu\mu}} (u_{1\mu}^2 + u_{T\mu}^2) - \rho_\mu, \\ d_{(2)\rho_\mu}^{DW} &= d_{(2)\rho_\mu}^{LS} + \frac{1 - \rho_\mu^2}{2\sigma_{\mu\mu}} (u_{1\mu}^2 + u_{T\mu}^2).\end{aligned}\quad (\text{A.62})$$

**Proof of Lemma A.13.** The proof is straightforward using Lemmas A.11 and A.12.  $\square$

Next, we provide analytic forms of the elements of vectors  $l$  and  $c$ , and matrices  $L$ ,  $C$  and  $D_*$ , employed in the stochastic expansions of the tests statistics given in the

paper. To this end, we first derive the partial derivatives of matrix  $A$ , given in (28), with respect to the elements of  $\varrho$  and  $\varsigma$ . Using matrices  $B_{ij} = X'_i R^{ij} X_j / T$ , matrix  $A$  can be partitioned as follows:

$$A = [(\sigma^{ij} B_{ij})_{i,j=1, \dots, M}]. \quad (\text{A.63})$$

**Lemma A.14.** *The partial derivatives of matrix  $A$ , with respect to the elements of  $\varrho$  and  $\varsigma$ , can be analytically written as follows:*

$$\begin{aligned} A_{\rho\mu} &= [(\frac{\sigma^{ij}}{T} X'_i R_{\rho\mu}^{ij} X_j)_{i,j=1, \dots, M}], \quad A_{\rho\mu\rho_{\mu'}} = [(\frac{\sigma^{ij}}{T} X'_i R_{\rho\mu\rho_{\mu'}}^{ij} X_j)_{i,j=1, \dots, M}], \\ A_{\rho\mu\rho_{\mu'}}^* &= [(X'_i W_{ij} X_j / T)_{i,j=1, \dots, M}], \quad A_{\varsigma(\mu\mu')} = [(\delta_{\mu i} \delta_{j\mu'} B_{\mu\mu'})_{i,j=1, \dots, M}], \\ A_{\varsigma(\mu\mu')\varsigma(\nu\nu')} &= 0, \quad A_{\varsigma(\mu\mu')\varsigma(\nu\nu')}^* = \sigma_{\mu'\nu} A_{\varsigma(\mu\nu')}, \\ A_{\rho\mu\varsigma(\nu\nu')} &= [(\delta_{\nu i} \delta_{j\nu'} X'_i R_{\rho\mu}^{\nu\nu'} X_{\nu'} / T)_{i,j=1, \dots, M}], \\ A_{\rho\mu\varsigma(\nu\nu')}^* &= \left[ \left( \sum_{k=1}^M \frac{\delta_{j\nu'} \sigma^{ik} \sigma_{k\nu}}{T} X'_i R_{\rho\mu}^{ik} R_{k\nu} R^{\nu\nu'} X_{\nu'} \right)_{i,j=1, \dots, M} \right], \\ A_{\varsigma(\nu\nu')\rho\mu}^* &= \left[ \left( \sum_{r=1}^M \frac{\delta_{\nu i} \sigma_{\nu'r} \sigma^{rj}}{T} X'_i R^{\nu\nu'} R_{\nu'r} R_{\rho\mu}^{rj} X_j \right)_{i,j=1, \dots, M} \right]. \end{aligned} \quad (\text{A.64})$$

**Proof of Lemma A.14.** The proof follows immediately from equation (31), and Lemmas A.4 and A.5.  $\square$

Analytic formulae of the elements of vector  $l$  and matrix  $L$  are given in the following lemma:

**Lemma A.15.** *The elements of vector  $l$  and matrix  $L$  can be calculated as follows:*

$$l_{\rho\mu} = \sum_{i=1}^M \sum_{j=1}^M \sum_{k=1}^M \sum_{r=1}^M \sigma^{kr} h'_i G_{ik} X'_k R_{\rho\mu}^{kr} X_r G_{rj} h_j / T, \quad (\text{A.65})$$

$$l_{\varsigma(\mu\mu')} = \sum_{i=1}^M \sum_{j=1}^M h'_i G_{i\mu} B_{\mu\mu'} G_{\mu'j} h_j, \quad (\text{A.66})$$

$$\begin{aligned} l_{\rho\mu\rho_{\mu'}} &= \sum_{q=1}^M \sum_{s=1}^M \sum_{i=1}^M \sum_{j=1}^M \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma^{rj} \\ &\times h'_q G_{qi} X'_i R_{\rho\mu}^{ik} (\sigma_{kr} R_{kr} - 2X_k G_{kr} X'_r / T) R_{\rho\mu'}^{rj} X_j G_{js} h_s / T \\ &+ \sum_{q=1}^M \sum_{s=1}^M \sum_{i=1}^M \sum_{j=1}^M \sigma^{ij} \\ &\times h'_q G_{qi} X'_i R_{\rho\mu\rho_{\mu'}}^{ij} X_j G_{js} h_s / 2T, \end{aligned} \quad (\text{A.67})$$

$$l_{\varsigma(\mu\mu')\varsigma(\nu\nu')} = \sigma_{\mu'\nu} l_{\varsigma(\mu\nu')} - 2 \sum_{i=1}^M \sum_{j=1}^M h'_i G_{i\mu} B_{\mu\mu'} G_{\mu'\nu} B_{\nu\nu'} G_{\nu'j} h_j, \quad (\text{A.68})$$

$$\begin{aligned} l_{\rho\mu\varsigma(\nu\nu')} &= \sum_{q=1}^M \sum_{s=1}^M \sum_{i=1}^M \sum_{k=1}^M \sigma^{ik} h'_q G_{qi} X'_i R_{\rho\mu}^{ik} \\ &\times (\sigma_{k\nu} R_{k\nu} - 2X_k G_{k\nu} X'_{\nu'} / T) R^{\nu\nu'} X_{\nu'} G_{\nu's} h_s / T \\ &+ \sum_{q=1}^M \sum_{s=1}^M h'_q G_{q\nu} X'_\nu R_{\rho\mu}^{\nu\nu'} X_{\nu'} G_{\nu's} h_s / 2T, \end{aligned} \quad (\text{A.69})$$



$$\begin{aligned}
l_{\varsigma(\nu\nu')\rho\mu} &= \sum_{q=1}^M \sum_{s=1}^M \sum_{j=1}^M \sum_{r=1}^M \sigma^{rj} h'_q G_{qv} X'_\nu R^{\nu\nu'} \\
&\times (\sigma_{\nu'r} R_{\nu'r} - 2X_{\nu'} G_{\nu'r} X'_r / T) R_{\rho\mu}^{rj} X_j G_{js} h_s / T \\
&+ \sum_{q=1}^M \sum_{s=1}^M h'_q G_{qv} X'_\nu R_{\rho\mu}^{\nu\nu'} X_{\nu'} G_{\nu's} h_s / 2T. \tag{A.70}
\end{aligned}$$

**Proof of Lemma A.15.** The results of the lemma follow by using the definitions in (41), the partition of matrix  $G$  in (52) and Lemmas A.1 – A.14.  $\square$

Analytic formulae of the elements of vector  $c$  and matrices  $C$  and  $D_*$  are given in the following lemma:

**Lemma A.16.** *The elements of vector  $c$  and matrices  $C$  and  $D_*$  can be calculated as follows:*

$$c_{\rho\mu} = \sum_{i=1}^M \sum_{j=1}^M \sigma^{ij} \text{tr}(X'_i R_{\rho\mu}^{ij} X_j \Xi_{ji}) / T, \tag{A.71}$$

$$c_{\varsigma(\mu\mu')} = \text{tr}(B_{\mu\mu'} \Xi_{\mu'\mu}), \tag{A.72}$$

$$\begin{aligned}
c_{\rho\mu\rho\mu'} &= \sum_{i=1}^M \sum_{j=1}^M \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma_{kr} \sigma^{rj} \\
&\times \text{tr}(X'_i R_{\rho\mu}^{ik} R_{kr} R_{\rho\mu'}^{rj} X_j \Xi_{ji}) / T \\
&- 2 \sum_{i=1}^M \sum_{j=1}^M \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma^{rj} \\
&\times \text{tr}(X'_i R_{\rho\mu}^{ik} X_k G_{kr} X'_r R_{\rho\mu'}^{rj} X_j \Xi_{ji}) / T^2 \\
&+ \sum_{i=1}^M \sum_{j=1}^M \sigma^{ij} \text{tr}(X'_i R_{\rho\mu\rho\mu'}^{ij} X_j \Xi_{ji}) / 2T, \tag{A.73}
\end{aligned}$$

$$c_{\varsigma(\mu\mu')\varsigma(\nu\nu')} = \sigma_{\mu'\nu} c_{\varsigma(\mu\nu')} - 2\text{tr}(B_{\mu\mu'} G_{\mu'\nu} B_{\nu\nu'} \Xi_{\nu'\mu}), \tag{A.74}$$

$$\begin{aligned}
c_{\rho\mu\varsigma(\nu\nu')} &= \sum_{i=1}^M \sum_{k=1}^M \sigma^{ik} \sigma_{k\nu} \text{tr}(X'_i R_{\rho\mu}^{ik} R_{k\nu} R^{\nu\nu'} X_{\nu'} \Xi_{\nu'i}) / T \\
&- 2 \sum_{i=1}^M \sum_{k=1}^M \sigma^{ik} \text{tr}(X'_i R_{\rho\mu}^{ik} X_k G_{k\nu} B_{\nu\nu'} \Xi_{\nu'i}) / T \\
&+ \text{tr}(X'_\nu R_{\rho\mu}^{\nu\nu'} X_{\nu'} \Xi_{\nu'\nu}) / 2T, \tag{A.75}
\end{aligned}$$

$$\begin{aligned}
c_{\varsigma(\nu\nu')\rho\mu} &= \sum_{j=1}^M \sum_{r=1}^M \sigma_{\nu'r} \sigma^{rj} \text{tr}(X'_\nu R^{\nu\nu'} R_{\nu'r} R_{\rho\mu}^{rj} X_j \Xi_{j\nu}) / T \\
&- 2 \sum_{j=1}^M \sum_{r=1}^M \sigma^{rj} \text{tr}(B_{\nu\nu'} G_{\nu'r} X'_r R_{\rho\mu}^{rj} X_j \Xi_{j\nu}) / T \\
&+ \text{tr}(X'_\nu R_{\rho\mu}^{\nu\nu'} X_{\nu'} \Xi_{\nu'\nu}) / 2T, \tag{A.76}
\end{aligned}$$

$$\begin{aligned}
d_{\rho\mu\rho\mu'} &= \sum_{i=1}^M \sum_{j=1}^M \sum_{k=1}^M \sum_{r=1}^M \sigma^{ik} \sigma^{rj} \\
&\times \text{tr}(X'_i R_{\rho\mu}^{ik} X_k \Xi_{kr} X'_r R_{\rho\mu'}^{rj} X_j \Xi_{ji}) / 2T^2, \tag{A.77}
\end{aligned}$$

$$d_{\varsigma(\mu\mu')\varsigma(\nu\nu')} = \text{tr}(B_{\mu\mu'} \Xi_{\mu'\nu} B_{\nu\nu'} \Xi_{\nu'\mu}) / 2, \tag{A.78}$$

$$d_{\rho\mu\varsigma(\nu\nu')} = \sum_{i=1}^M \sum_{k=1}^M \sigma^{ik} \text{tr}(X'_i R_{\rho\mu}^{ik} X_k \Xi_{k\nu} B_{\nu\nu'} \Xi_{\nu'i}) / 2T, \tag{A.79}$$

$$d_{\varsigma(\nu\nu')\rho\mu} = \sum_{j=1}^M \sum_{r=1}^M \sigma^{rj} \text{tr}(B_{\nu\nu'} \Xi_{\nu'r} X'_r R_{\rho\mu}^{rj} X_j \Xi_{j\nu}) / 2T. \tag{A.80}$$

**Proof of Lemma A.16.** The results of the lemma can be easily calculated by using the definitions (56) and (57), partition of matrix  $\Xi$  in (52) and the following traces:

$$\begin{aligned} & \text{tr}(A_{\rho_\mu} \Xi), \text{tr}(A_{\rho_\mu \rho_{\mu'}} \Xi), \text{tr}(A_{\varsigma_{(\mu\mu')}} \Xi), \text{tr}(A_{\varsigma_{(\mu\mu')\varsigma_{(\nu\nu')}}} \Xi), \\ & \text{tr}(A_{\rho_\mu \varsigma_{(\nu\nu')}} \Xi), \text{tr}(A_{\rho_\mu \rho_{\mu'}}^* \Xi), \text{tr}(A_{\varsigma_{(\mu\mu')\varsigma_{(\nu\nu')}}}^* \Xi), \text{tr}(A_{\rho_\mu \varsigma_{(\nu\nu')}}^* \Xi), \quad (\text{A.81}) \\ & \text{tr}(A_{\rho_\mu} G A_{\rho_{\mu'}} \Xi), \text{tr}(A_{\rho_\mu} G A_{\varsigma_{(\nu\nu')}} \Xi), \text{tr}(A_{\varsigma_{(\mu\mu')}} G A_{\varsigma_{(\nu\nu')}} \Xi), \end{aligned}$$

with obvious modifications for

$$\begin{aligned} & \text{tr}(A_{\varsigma_{(\nu\nu')\rho_\mu}} \Xi), \text{tr}(A_{\varsigma_{(\nu\nu')\rho_\mu}}^* \Xi), \text{tr}(A_{\varsigma_{(\nu\nu')}} G A_{\rho_\mu} \Xi), \\ & \text{tr}(A_{\rho_\mu} \Xi A_{\rho_{\mu'}} \Xi), \text{tr}(A_{\rho_\mu} \Xi A_{\varsigma_{(\nu\nu')}} \Xi), \text{tr}(A_{\varsigma_{(\nu\nu')}} \Xi A_{\rho_\mu} \Xi), \text{tr}(A_{\varsigma_{(\mu\mu')}} \Xi A_{\varsigma_{(\nu\nu')}} \Xi). \end{aligned}$$

By using the above results and Lemmas A.1 – A.14, the proof completes.  $\square$

Analytic formulae of the scalars and vectors given in (33) are derived in the following lemma:

**Lemma A.17.** *Scalars  $\lambda_0$  and  $\kappa_0$ , vectors  $\lambda_\varrho$ ,  $\lambda_\varsigma$ ,  $\kappa_\varrho$  and  $\kappa_\varsigma$ , and matrices  $\Lambda_\varrho$ ,  $\Lambda_\varsigma$  and  $\Lambda_{\varrho\varsigma}$  can be calculated as follows:*

$$\lambda_0 = 0, \quad \lambda_\varrho = 0, \quad \lambda_\varsigma = 0, \quad (\text{A.82})$$

$$\Lambda_\varsigma = (\Sigma^{-1} \otimes \Sigma^{-1}) N (\Sigma^{-1} \otimes \Sigma^{-1}), \quad (\text{A.83})$$

where  $N$  is a  $(M^2 \times M^2)$  matrix whose  $((ij), (kr))$ -th element is

$$\nu_{(ij)(kr)} = \sigma_{ik} \sigma_{jr} + \sigma_{ir} \sigma_{jk} \quad (i, j, k, r = 1, \dots, M). \quad (\text{A.84})$$

The  $\mu$ -th diagonal element of the matrix  $\Lambda_\varrho$  is

$$\lim_{T \rightarrow \infty} \mathcal{E}(d_{(1)\rho_\mu}^2) = 1 - \rho_\mu^2, \quad (\text{A.85})$$

and its  $(\mu, \mu')$ -th off-diagonal element is

$$\lim_{T \rightarrow \infty} \mathcal{E}(d_{(1)\rho_\mu} d_{(1)\rho_{\mu'}}) = \frac{(1 - \rho_\mu^2)(1 - \rho_{\mu'}^2)}{(1 - \rho_\mu \rho_{\mu'})}, \quad (\text{A.86})$$

for  $\mu \neq \mu'$ . Further, we have

$$\Lambda_{\varrho\varsigma} = 0 \quad \text{and} \quad \Lambda_{\varsigma\varrho} = 0. \quad (\text{A.87})$$

For all estimators  $\hat{\sigma}_I$  and  $\hat{\varsigma}_I$  ( $I = UL, RL, GL, IG, ML$ ), we can compute the following  $(M \times M)$  matrices:

$$\begin{aligned} \Delta_{UL} &= 0, \quad \Delta_{GL} = \Delta_{IG} = \Delta_{ML} = K\Sigma - \left[ (\text{tr}(G_{ij} B_{ji}))_{i,j=1, \dots, M} \right], \\ \Delta_{RL} &= \left[ \left( (\text{tr}(B_{ii}^{-1} B_{ij} B_{jj}^{-1} B_{ji})) - n_i - n_j + K \right) \sigma_{ij} \right]_{i,j=1, \dots, M}. \end{aligned} \quad (\text{A.88})$$

Given them, we can calculate  $\kappa_0$  and  $\kappa_\varsigma$  as follows:

$$\kappa_0 = \text{tr} [\Sigma^{-1}(\Delta_{GL} - \Delta_I)] / M + n/M, \quad (\text{A.89})$$

and

$$\kappa_\varsigma = \text{vec} \{ (M + K + 1)\Sigma^{-1} - \Sigma^{-1}\Delta_I\Sigma^{-1} \}. \quad (\text{A.90})$$

Also, define scalars

$$c_1 = (1 - \rho_\mu^2)[(1 - \rho_\mu^2)\text{tr}(F_{\mu\mu}^{-1}\Theta_{\mu\mu}) + \text{tr}(F_{\mu\mu}^{-1}B_{\mu\mu}F_{\mu\mu}^{-1}\Theta_{\mu\mu})], \quad (\text{A.91})$$

and

$$c_2 = (1 - \rho_\mu^2)\text{tr}(F_{\mu\mu}B_{\mu\mu}^{-1}), \quad (\text{A.92})$$

where the  $(n_\mu \times n_\mu)$  matrices  $F_{\mu\mu}$ ,  $\Theta_{\mu\mu}$  and  $B_{\mu\mu}$  are defined in (A.46) and (A.47).

For all estimators  $\hat{\rho}_\mu^I$  ( $I = LS, GL, PW, ML, DW$ ), we calculate the elements  $\kappa_{\rho_\mu}$  of  $(M \times 1)$  vector  $\kappa_\varrho$  as follows:

$$\kappa_{\rho_\mu}^{LS} = -[(n_\mu + 3)\rho_\mu + (c_1 - 2n_\mu)/2\rho_\mu], \quad (\text{A.93})$$

and

$$\begin{aligned} \kappa_{\rho_\mu}^{GL} = \kappa_{\rho_\mu}^{PW} &= \kappa_{\rho_\mu}^{LS} + \frac{c_1 - (1 - \rho_\mu^2)(c_2 + n_\mu)}{2\rho_\mu}, \\ \kappa_{\rho_\mu}^{ML} &= \kappa_{\rho_\mu}^{GL} + \rho_\mu, \\ \kappa_{\rho_\mu}^{DW} &= \kappa_{\rho_\mu}^{LS} + 1. \end{aligned} \quad (\text{A.94})$$

**Proof of Lemma A.17.** From (33), (A.30), (A.35), and (A.61) we can easily show that

$$\lambda_0 = \lim_{T \rightarrow \infty} \mathcal{E}(\sigma_0^2), \quad \lambda_\rho = \lim_{T \rightarrow \infty} \mathcal{E}(\sigma_0 d_{1\rho}) \quad \text{and} \quad \lambda_\varsigma = \lim_{T \rightarrow \infty} \mathcal{E}(\sigma_0 d_{1\varsigma}). \quad (\text{A.95})$$

The results in (A.82) follows immediately since  $\sigma_0 = 0$  (see (A.36)). Equations (33) and (A.30) imply

$$\Lambda_\varsigma = \lim_{T \rightarrow \infty} \mathcal{E}(d_{1\varsigma} d'_{1\varsigma}). \quad (\text{A.96})$$

This result together with (A.31), (A.42) and (A.43) yield (A.83).

Since (33) and (A.61) imply that

$$\Lambda_\varrho = \lim_{T \rightarrow \infty} \mathcal{E}(d_{1\varrho} d'_{1\varrho}) \quad (\text{A.97})$$

and  $\sigma_{u_\mu}^2 = \sigma_{\mu\mu}^2 / (1 - \rho_\mu^2)$ , we can prove that the  $\mu$ -th diagonal element of the matrix  $\Lambda_\varrho$  is

$$\begin{aligned} \lim_{T \rightarrow \infty} \mathcal{E}(d_{1\varrho}^2)_{\rho_\mu} &= \lim_{T \rightarrow \infty} \mathcal{E}(u'_\mu R_2^{\mu\mu} u_\mu u'_\mu R_2^{\mu\mu} u_\mu) / 4T \sigma_{u_\mu}^2 \\ &= \lim_{T \rightarrow \infty} \left[ \frac{4T \sigma_{\mu\mu}^2}{1 - \rho_\mu^2} + O(1) \right] / 4T \sigma_{u_\mu}^2, \end{aligned} \quad (\text{A.98})$$

by combining the third result in Lemma A.10 with (A.55) and (A.62). The last result proves (A.85). Working along the same lines for  $\mu \neq \mu'$ , we can prove (A.86), for the  $(\mu, \mu')$ -th off-diagonal element of  $\Lambda_\varrho$ .

To prove (A.87), first note that (33), (A.30) and (A.61) imply

$$\Lambda_{\varrho\varsigma} = \lim_{T \rightarrow \infty} \mathcal{E}(d_{1\varrho} d'_{1\varsigma}). \quad (\text{A.99})$$

Substituting (A.31), (A.58) and (A.55) into (A.99), we can calculate the  $(\mu, (ij))$ -th element of  $(M \times M^2)$  matrix  $\Lambda_{\varrho\varsigma}$  as  $-d_{(1)\rho\mu} s_{(ij)}^{(1)}$ . Following the same steps to that of the proof of (A.59) we can show that

$$\lim_{T \rightarrow \infty} \mathcal{E} \left( -d_{(1)\rho\mu} s_{(ij)}^{(1)} \right) = 0. \quad (\text{A.100})$$

(A.87) can be proved immediately using  $\Lambda_{\varsigma\varrho} = \Lambda'_{\varrho\varsigma}$ .

For all estimators  $\hat{\sigma}_I$  ( $I = UL, RL, GL, IG, ML$ ), we can find that

$$\kappa_0 = \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} \sigma_0 + \sigma_1 \right) = \lim_{T \rightarrow \infty} \mathcal{E}(\sigma_1), \quad (\text{A.101})$$

by combining (33) with (A.40), (A.35) and (A.36). The last result proves (A.89). For all estimators  $\hat{\varsigma}_I$  ( $I = UL, RL, GL, IG, ML$ ), we can show that

$$\kappa_\varsigma = \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} d_{1\varsigma} + d_{2\varsigma} \right) = \text{vec} \left\{ \lim_{T \rightarrow \infty} \mathcal{E} \left( S_2^I \right) \right\}, \quad (\text{A.102})$$

since  $\mathcal{E}(S_1) = 0$  and  $\lim_{T \rightarrow \infty} \mathcal{E}(S_2^I) = M_I$  [see (A.40)], by using (33), (A.30), (A.31) and (A.40). This result implies (A.90).

Finally, we can calculate

$$\kappa_{\rho\mu}^{LS} = \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} d_{(1)\rho\mu}^{LS} + d_{(2)\rho\mu}^{LS} \right), \quad (\text{A.103})$$

by using (33) and (A.55), Lemmas A.10 and A.13. This yields (A.93). Along the same lines, we can calculate the following quantities:

$$\begin{aligned} \kappa_{\rho\mu}^{GL} = \kappa_{\rho\mu}^{PW} &= \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} d_{(1)\rho\mu}^{LS} + d_{(2)\rho\mu}^{GL} \right), \\ \kappa_{\rho\mu}^{ML} &= \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} d_{(1)\rho\mu}^{LS} + d_{(2)\rho\mu}^{ML} \right) \quad \text{and} \\ \kappa_{\rho\mu}^{DW} &= \lim_{T \rightarrow \infty} \mathcal{E} \left( \sqrt{T} d_{(1)\rho\mu}^{LS} + d_{(2)\rho\mu}^{DW} \right), \end{aligned} \quad (\text{A.104})$$

which proves (A.94). □

## Asymptotic expansions of size corrected tests: Proofs of theorems

Given the lemmas of the previous subsections, next we give the proofs of the theorems presented in the main text. These are based on known expansions of standard normal and chi-square distributed tests. We derive new expansions of the degrees-of-freedom-adjusted versions of these tests, by inverting their characteristic functions. These degrees-of-freedom-adjusted approximations of distribution functions are proved to be locally exact.

**Proof of Theorems 1 and 2.** Approximation (42) of Theorem 1 can be proved following the steps of the proof in Rothenberg (1988). The quantities in (40) can be obtained by expanding the corresponding quantities given by Rothenberg and retaining the first term in each of these expansions. The approximation (44) of Theorem 2 follows from the approximation (42) and the following asymptotic approximations of the Student- $t$  distribution and density functions, which are given in terms of the standard normal distribution and density functions, respectively (see Fisher (1925)):

$$I_{T-n}(x) = I(x) - (\tau^2/4)(1+x^2)xi(x) + O(\tau^4), \quad (\text{A.105})$$

$$i_{T-n}(x) = i(x) + O(\tau^2).$$

Note that approximation (44) of Theorem 2 is locally exact. This can be easily seen as follows: If parameter vector  $\gamma = (\varrho', \varsigma')'$  is known to belong to a ball of radius  $\vartheta$ , then, as  $\vartheta \rightarrow 0$ ,  $\gamma$  becomes a fixed known vector. By using (27), (29), (33) and (35) we can prove that

$$\Lambda = 0, \quad \lambda = \kappa = 0, \quad \lambda_0 = 2, \quad \kappa_0 = 0. \quad (\text{A.106})$$

Then, the analytic formulae of  $p_1$  and  $p_2$ , given in (43), become

$$p_1 = p_2 = 0. \quad (\text{A.107})$$

This result implies that, with an error of order  $O(\tau^3)$ , approximation (44) becomes the Student- $t$  distribution function with  $MT - n$  degrees of freedom.  $\square$

**Proof of Theorem 3.** We begin the proof by noticing that, under null hypothesis (36), the  $t$  statistic, given by (37), admits a stochastic expansion of the form

$$t = t_0 + \tau t_1 + \tau^2 t_2 + \omega(\tau^3), \quad (\text{A.108})$$

where the first term in the expansion is given as

$$t_0 = e'b/(e'Ge)^{1/2} = h'b, \quad \text{where } b = GX'\Omega u/\sqrt{T}.$$

The result given by equation (A.108) implies that the Cornish-Fisher corrected statistic  $t_*$ , given by (47), admits a stochastic expansion of the form

$$t_* = t_0 + \tau t_1 + \tau^2(t_2 - t_3) + \omega(\tau^3), \quad (\text{A.109})$$

where

$$t_3 = (p_1 + p_2 t_0^2)t_0/2.$$

Let  $s$  be an imaginary number, and  $\psi(s)$  and  $\phi(s)$  denote the characteristic functions of the  $t$  statistic, given by (37), and a standard normal random variable, respectively. Using (A.109) and the relationships:

$$E[\exp(st_0)t_0] = s\phi(s) \quad \text{and} \quad E[\exp(st_0)t_0^3] = (3s + s^3)\phi(s),$$

we can show that the characteristic function of the Cornish-Fisher corrected statistic  $t_*$ , denoted as  $\psi_*(s)$ , can be approximated as follows:

$$\begin{aligned} \psi_*(s) &= \psi(s) - \tau^2 s E[\exp(st_0)t_3] + O(\tau^3) \\ &= \psi(s) - \frac{\tau^2}{2} s [p_1 s + p_2(3s + s^3)]\phi(s) + O(\tau^3). \end{aligned}$$

Dividing  $\psi_*(s)$  by  $-s$ , applying the inverse Fourier transform and using Theorem 2, we can show that

$$\begin{aligned} \Pr\{t_* \leq x\} &= \Pr\{t \leq x\} + \frac{\tau^2}{2}(p_1 + p_2 x^2)xi_{T-n}(x) + O(\tau^3) \\ &= I_{T-n}(x) - \frac{\tau^2}{2}(p_1 + p_2 x^2)xi_{T-n}(x) \\ &\quad + \frac{\tau^2}{2}(p_1 + p_2 x^2)xi_{T-n}(x) + O(\tau^3) \\ &= I_{T-n}(x) + O(\tau^3). \end{aligned} \quad (\text{A.110})$$

The last result means that the Cornish-Fisher corrected statistic  $t_*$  is distributed as a Student- $t$  random variable with  $MT - n$  degrees of freedom.  $\square$

**Proof of Theorems 4 and 5.** Approximation (58) of Theorem 4 can be proved following the steps of the proof in Rothenberg (1984b). The quantities in (56) can be obtained by expanding the corresponding quantities given by Rothenberg and retaining the first term in each of these expansions. Approximation (60) of Theorem 5 follows

from approximation (58) and the following asymptotic approximations of the  $F$  distribution and density functions, which are given in terms of the chi-square distribution and density functions, respectively:

$$F_{T-n}^m(x) = F_m(mx) + (\tau^2/2)(m-2-mx)mf_m(mx) + O(\tau^4), \quad (\text{A.111})$$

$$f_{T-n}^m(x) = mf_m(mx) + O(\tau^2).$$

Note that approximation (60) of Theorem 5 can be easily seen to be locally exact. By using (A.106), (59), and (61), we can show that

$$\xi_1 = -m(m-2)/2 \quad \text{and} \quad \xi_2 = m(m+2)/2 \quad (\text{A.112})$$

$$\Rightarrow q_1 = q_2 = 0. \quad (\text{A.113})$$

This result means that, with an error of order  $O(\tau^3)$ , approximation (60) becomes the  $F$  distribution function with  $m$  and  $MT - n$  degrees of freedom.  $\square$

**Proof of Theorem 6.** To begin the proof, we first notice that, under null hypothesis (48), the  $F$  statistic, given by (50), admits a stochastic expansion of the form

$$F = F_0 + \tau F_1 + \tau^2 F_2 + \omega(\tau^3), \quad (\text{A.114})$$

where the first term in the expansion is

$$F_0 = b'Qb/m, \quad b = GX'\Omega u/\sqrt{T}.$$

Equation (A.114) implies that the Cornish-Fisher corrected statistic  $F_*$ , given by (64), admits a stochastic expansion of the form

$$F_* = F_0 + \tau F_1 + \tau^2(F_2 - F_3) + \omega(\tau^3), \quad (\text{A.115})$$

where

$$F_3 = (q_1 + q_2 F_0)F_0.$$

Let  $s$  be an imaginary number, and  $\psi(s)$  and  $\phi(s)$  now denote the characteristic functions of the  $F$  statistic, given by (50), and a chi-square random variable with  $m$  degrees of freedom, respectively. Using (A.115) and the following relationships:

$$E[\exp(sF_0)F_0] = \phi_{m+2}(s/m) \quad \text{and} \quad E[\exp(sF_0)F_0^2] = \frac{m+2}{m}\phi_{m+4}(s/m),$$

we can show that the characteristic function of the Cornish-Fisher corrected statistic  $F_*$ , denoted as  $\psi_*(s)$ , can be approximated as follows:

$$\begin{aligned}\psi_*(s) &= \psi(s) - \tau^2 s E[\exp(sF_0)F_3] + O(\tau^3) \\ &= \psi(s) - \tau^2 s [q_1 \phi_{m+2}(s/m) + q_2 \frac{m+2}{m} \phi_{m+4}(s/m)] + O(\tau^3). \quad (\text{A.116})\end{aligned}$$

For the chi-square density  $f_m(x)$ , the following results can be shown:

$$(mx)f_m(mx) = mf_{m+2}(mx) \quad \text{and} \quad (mx)^2 f_m(mx) = m(m+2)f_{m+4}(mx). \quad (\text{A.117})$$

Dividing (A.116) by  $-s$ , applying the inverse Fourier transform, and using Theorem 5 and the results of equations (A.111) and (A.117), we can show that

$$\begin{aligned}\Pr\{F_* \leq x\} &= \Pr\{F \leq x\} + \tau^2 [(q_1 m f_{m+2}(mx) + q_2 \frac{m+2}{m} m f_{m+4}(mx))] + O(\tau^3) \\ &= \Pr\{F \leq x\} + \tau^2 [(q_1 m x f_m(mx) + q_2 m x^2 f_m(mx))] + O(\tau^3) \\ &= \Pr\{F \leq x\} + \tau^2 (q_1 + q_2 x) m x f_m(mx) + O(\tau^3) \\ &= F_{T-n}^m(x) - \tau^2 (q_1 + q_2 x) x f_{T-n}^m(x) \\ &\quad + \tau^2 (q_1 + q_2 x) x f_{T-n}^m(x) + O(\tau^3) \\ &= F_{T-n}^m(x) + O(\tau^3). \quad (\text{A.118})\end{aligned}$$

The last result implies that the Cornish-Fisher corrected statistic  $F_*$  is distributed as an  $F$  random variable with  $m$  and  $MT - n$  degrees of freedom.  $\square$

## Bibliography

- Beach, C. M. and MacKinnon, J. G. (1978). A Maximum Likelihood Procedure for Regression with Autocorrelated Errors. *Econometrica*, 46(1):51–58.
- Breusch, T. S. (1980). Useful Invariance Results for Generalized Regression Models. *Journal of Econometrics*, 13:327–340.
- Cornish, E. A. and Fisher, R. A. (1937). Moments and Cumulants in The Specification of Distributions. *Revue del' Institute International de Statistic*, 4:1–14.
- Dhrymes, P. J. (1971). Equivalence of Iterative Aitken and Maximum Likelihood Estimators for A System of Regression Equations. *Australian Economic Papers*, 10:20–24.
- Dhrymes, P. J. (1978). *Introductory Econometrics*. Springer Verlag, New York.



- Fisher, R. A. (1925). Expansion of Student's Integral in Powers of  $n^{-1}$ . *Metron*, 5:109–120.
- Fisher, R. A. and Cornish, E. A. (1960). The Percentile of Distributions Having Known Cumulants. *Technometrics*, 2:209–225.
- Hill, G. W. and Davis, A. W. (1968). Generalized Asymptotic Expansions of Cornish-Fisher Type. *The Annals of Mathematical Statistics*, 39(4):1264–1273.
- Iglesias, E. M. and Phillips, G. D. (2010). The Bias to Order  $T^{-2}$  for the General k-class Estimator in A Simultaneous Equation Model. *Economics Letters*, 109:42–45.
- Iglesias, E. M. and Phillips, G. D. (2012). Almost Unbiased Estimation in Simultaneous Equations Models with Strong and / or Weak Instruments. *Journal of Business and Economic Statistics*, 30(4):505–520.
- Kiviet, J. and Phillips, G. D. (1996). The Bias of the Ordinary Least Squares Estimator in Simultaneous Equation Models. *Economics Letters*, (53):161–167.
- Kiviet, J., Phillips, G. D., and Schipp, B. (1995). The Bias of OLS, GLS, and ZEF Estimators in Dynamic Seemingly Unrelated Regression Models. *Journal of Econometrics*, 69:241–266.
- Magdalinos, M. A. (1985). Improving Some Instrumental Variables Test Procedures. *Econometric Theory*, 1:240–261.
- Magdalinos, M. A. (1992). Stochastic Expansions and Asymptotic Approximations. *Econometric Theory*, 8:343–367.
- Magdalinos, M. A. and Symeonides, S. D. (1995). Alternative Size Corrections for Some GLS Test Statistics: The Case of the AR(1) Model. *Journal of Econometrics*, 66:35–59.
- Magee, L. (1985). Efficiency of Iterative Estimators in the Regression Model with AR(1) Disturbances. *Journal of Econometrics*, 29:275–287.
- Magnus, J. R. and Neudecker, H. (1979). The Commutation Matrix: Some Properties and Applications. *The Annals of Statistics*, 7:381–394.
- McDonald, G. C. and Galarneau, D. I. (1975). A Monte Carlo Evaluation of Some Ridge-Type Estimators. *Journal of the American Statistical Association*, 70(350):407–416.

- Ogasawara, H. (2012). Cornish-Fisher Expansions Using Sample Cumulants and Monotonizing Transformations. *Journal of Multivariate Analysis*, 103:1–18.
- Parks, R. W. (1967). Efficient Estimation of a System of Regression Equations when Disturbances are Both Serially and Contemporaneously Correlated. *Journal of the American Statistical Association*, 62:500–509.
- Phillips, G. D. (2000). An Alternative Approach to Obtaining Nagar-type Moment Approximations in Simultaneous Equation Models. *Journal of Econometrics*, 97:345–364.
- Phillips, G. D. (2007). Nagar-type Moment Approximations in Simultaneous Equation Models: Some Further Results. In *The Refinement of Econometric Estimation and Test Procedures: Finite Sample and Asymptotic Analysis*, pages 1–1. Cambridge University Press, Cambridge.
- Prais, S. J. and Winsten, C. I. (1954). Trend Estimators and Serial Correlation. Cowles Commission Discussion Paper: Statistics No. 383.
- Rothenberg, T. J. (1984a). Approximate Normality of Generalized Least Squares Estimates. *Econometrica*, 52(4):811–825.
- Rothenberg, T. J. (1984b). Hypothesis Testing in Linear Models when the Error Covariance Matrix is Nonscalar. *Econometrica*, 52(4):827–842.
- Rothenberg, T. J. (1988). Approximate Power Functions for Some Robust Tests of Regression Coefficients. *Econometrica*, 56(5):997–1019.
- Symeonides, S. D., Tzavalis, E., and Kandiliorou, H. (2007). Cornish-Fisher Size Corrected t and F Statistics for the Linear Regression Model with Heteroskedastic Errors. In Philips, G. D. A. and Tzavalis, E., editors, *The Refinement of Econometric Estimation and Test Procedures: Finite Sample and Asymptotic Analysis*, pages 173–204. Cambridge University Press, Cambridge.
- Ullah, A. (2004). *Finite Sample Econometrics*. Oxford University Press.
- Zellner, A. (1962). An Efficient Method of Estimating Seemingly Unrelated Regressions and Tests for Aggregation Bias. *Journal of the American Statistical Association*, 57(298):348–368.
- Zellner, A. (1963). Estimators for Seemingly Unrelated Regression Equations: Some Exact Finite Sample Results. *Journal of the American Statistical Association*, 58(304):977–992.

Zellner, A. (1971). *An Introduction to Bayesian Inference in Econometrics*. John Wiley & Sons, New York.

Zellner, A. and Huang, D. S. (1962). Further Properties of Efficient Estimators for Seemingly Unrelated Regression Equations. *International Economic Review*, 3(3):300–313.

Zellner, A. and Theil, H. (1962). Three-Stage Least Squares: Simultaneous Estimation of Simultaneous Equations. *Econometrica*, 30(1):54–78.