Extensions of posets with an antitone involution to residuated structures

Ivan Chajda, Miroslav Kolařík and Helmut Länger

Abstract

We prove that every not necessarily bounded poset $\mathbf{P} = (P, \leq, ')$ with an antitone involution can be extended to a residuated poset $\mathbb{E}(\mathbf{P}) = (E(P), \leq, \odot, \rightarrow, 1)$ where $x' = x \to 0$ for all $x \in P$. If \mathbf{P} is a lattice with an antitone involution then $\mathbb{E}(\mathbf{P})$ is a lattice, too. We show that a poset can be extended to a residuated poset by means of a finite chain and that a Boolean algebra $(B, \lor, \land, ', p, q)$ can be extended to a residuated lattice $(Q, \lor, \land, \odot, \rightarrow, 1)$ by means of a finite chain in such a way that $x \odot y = x \land y$ and $x \to y = x' \lor y$ for all $x, y \in B$.

AMS Subject Classification: 03B52, 06A11, 06B05, 03B47

Keywords: Poset, bounded poset, residuated poset, residuated lattice, antitone involution, extension

Residuated posets in general and residuated lattices in particular form an algebraic axiomatization of certain substructural logics (see e.g. [4], [8], [9] and [10] and references therein), especially of fuzzy logic, see [1] for details. Residuated lattices were studied for a long time starting with the pioneering paper by Ward and Dilworth [12], see also [3] and [7]. Posets and lattices with an antitone involution can serve as a suitable model of such a logic because this involution can be considered as a negation and hence these logics satisfy the double negation law, see [3]. Let us mention that a kind of residuated posets were studied also in [5]. Moreover, residuated structures derived from semirings were treated in [6] and [9].

Recall that a poset with an antitone involution is an ordered triple $(P, \leq, ')$ such that (P, \leq) is a poset and ' is a unary operation on P satisfying

- if $a \leq b$ then $b' \leq a'$,
- $a'' \approx a$

for all $a, b \in P$. Recall further that a *residuated poset* is an ordered quintuple $(P, \leq, \odot, \rightarrow, 1)$ such that

• $(P, \leq, 1)$ is a poset with a greatest element,

¹Support of the research of the first and the third author by the Austrian Science Fund (FWF), project I 4579-N, and the Czech Science Foundation (GAČR), project 20-09869L, entitled "The many facets of orthomodularity", as well as by ÖAD, project CZ 02/2019, entitled "Function algebras and ordered structures related to logic and data fusion", and, concerning the first author, by IGA, project PřF 2020 014, is gratefully acknowledged.

- $(P, \odot, 1)$ is a commutative monoid,
- \rightarrow is a binary operation on P,
- $a \odot b \le c$ if and only if $a \le b \to c$

for all $a, b, c \in P$. The last property is called *adjointness*.

Unfortunately, not every lattice $(L, \lor, \land, ')$ with an antitone involution ' can be converted into a residuated lattice $(L, \lor, \land, \odot, \rightarrow, 1)$ satisfying $x' = x \rightarrow 0$ for all $x \in L$. For example, consider the non-modular lattice $\mathbf{N}_5 = (N_5, \lor, \land)$ whose elements are 0, a, b, c, 1where 0 is the least and 1 the greatest element, a < b and c is incomparable with a and b, see Figure 1:



It is easy to see that there exists exactly one antitone involution ' on (N_5, \leq) , namely 0' = 1, a' = b, b' = a, c' = c and 1' = 0. Suppose, \mathbf{N}_5 together with ' could be converted into a residuated poset $(N_5, \leq, \odot, \rightarrow, 1)$ satisfying $x' = x \rightarrow 0$ for all $x \in N_5$. Since $x \odot y \leq x$ and $x \odot y \leq y$ for all $x, y \in N_5$ (see Theorem 2.17 in [1]) we have $c \odot a \leq 0$ which implies $c \leq a \rightarrow 0 = a' = b$, a contradiction. Thus \mathbf{N}_5 together with ' cannot be converted into a residuated poset $(N_5, \leq, \odot, \rightarrow, 1)$ satisfying $x' = x \rightarrow 0$ for all $x \in N_5$.

Hence, it is a question whether such a lattice (or poset in general) can be extended to a residuated one by preserving the antitone involution. The aim of our paper is to show how such an extension can be constructed.

At first, we show that the unary operation $x' := x \to 0$ in a residuated lattice is antitone.

Lemma 1. Let $(P, \leq, \odot, \rightarrow, 1)$ be a residuated poset and put $x' := x \rightarrow 0$ for all $x \in P$ Then $x \leq x''$ for all $x \in P$ and ' is antitone.

Proof. Let $a, b \in P$. Then the every of the following assertions implies the next one:

$$a \to 0 \le a \to 0,$$

$$(a \to 0) \odot a \le 0,$$

$$a \odot (a \to 0) \le 0,$$

$$a \le (a \to 0) \to 0,$$

$$a \le a''.$$

Moreover, every of the following assertions implies the next one:

$$a \leq b,$$

$$a \leq (b \to 0) \to 0,$$

$$a \odot (b \to 0) \leq 0,$$

$$(b \to 0) \odot a \leq 0,$$

$$b \to 0 \leq a \to 0,$$

$$b' \leq a'.$$

Our first extension result is as follows.

Theorem 2. Let $\mathbf{P} = (P, \leq, ')$ be a poset with an antitone involution, assume $0 = c_1, \ldots, c_4 = 1 \notin P$, put $E(P) := P \cup \{c_1, \ldots, c_4\}$ and extend \leq and ' from P to E(P) by $0 < c_2 < x < c_3 < 1$ for all $x \in P$ and $c'_i := c_{5-i}$ for $i = 1, \ldots, 4$. Define binary operations \odot and \rightarrow on E(P) as follows:

$$0 \odot x = x \odot 0 := 0, 1 \odot x = x \odot 1 := x,$$

$$0 \rightarrow x = x \rightarrow 1 := 1, x \rightarrow 0 := x', 1 \rightarrow x := x$$

for $x \in E(P)$ and

$$x \odot y := \begin{cases} 0 & \text{if } x \le y', \\ c_2 & \text{otherwise} \end{cases} \qquad x \to y := \begin{cases} 1 & \text{if } x \le y, \\ c_3 & \text{otherwise} \end{cases}$$

for $x, y \in E(P) \setminus \{0, 1\}$. Then $\mathbb{E}(\mathbf{P}) := (E(P), \leq, \odot, \rightarrow, 1)$ is a residuated poset with the antitone involution ' satisfying $x' = x \rightarrow 0$ for all $x \in E(P)$. (If \mathbf{P} is already bounded then the least and greatest element of \mathbf{P} may be identified with c_2 and c_3 , respectively. If \mathbf{P} has elements a, b, c, d satisfying $a < b \leq x \leq c < d$ for all $x \in P \setminus \{a, d\}$ then a, b, c, d may be identified with c_1, \ldots, c_4 , respectively.)

Proof. Let $a, b, c \in E(P)$. Since x'' = x for all $x \in P$ and $c''_i = c'_{5-i} = c_{5-(5-i)} = c_i$ for all $i = 1, \ldots, 4, (E(P), \leq, ', 0, 1)$ is a bounded poset with an antitone involution. If $\{a, b, c\} \cap \{0, 1\} \neq \emptyset$ then, obviously, $(a \odot b) \odot c = a \odot (b \odot c)$. If $a, b, c \neq 0, 1, a \leq b'$ and b < c' then $(a \odot b) \odot c = 0 \odot c = 0 = a \odot 0 = a \odot (b \odot c)$. If $a, b, c \neq 0, 1, a \leq b'$ and $b \leq c'$ then $(a \odot b) \odot c = 0 \odot c = 0 = a \odot c_2 = a \odot (b \odot c)$. If $a, b, c \neq 0, 1, a \not\leq b'$ and $b \leq c'$ then $(a \odot b) \odot c = c_2 \odot c = 0 = a \odot 0 = a \odot (b \odot c)$. If $a, b, c \neq 0, 1, a \leq b'$ and $b \leq c'$ then $(a \odot b) \odot c = c_2 \odot c = 0 = a \odot c_2 = a \odot (b \odot c)$. Therefore, \odot is associative. Since $a \leq b'$ is equivalent to $b \leq a'$, \odot is commutative. If a = 0 then $a \odot b = 0$ and a < b'. If b = 0 then $a \odot b = 0$ and $a \le b'$. If a = 1 then $a \odot b = 0$ and a < b' are equivalent to b = 0. If b = 1 then $a \odot b = 0$ and a < b' are equivalent to a = 0. If $a, b \neq 0, 1$ then $a \odot b = 0$ is equivalent to $a \leq b'$. Hence $a \odot b = 0$ if and only if $a \leq b'$. If a = 0 then $a \to b = 1$ and $a \le b$. If b = 0 then $a \to b = 1$ and $a \le b$ are equivalent to a = 0. If a = 1 then $a \to b = 1$ and $a \le b$ are equivalent to b = 1.

If b = 1 then $a \to b = 1$ and $a \le b$. If $a, b \ne 0, 1$ then $a \to b = 1$ and $a \le b$ are equivalent. Hence $a \to b = 1$ if and only if $a \le b$. If a = 0 then $a \odot b \le c$ and $a \le b \to c$. If b = 0 then $a \odot b \le c$ and $a \le b \to c$. If c = 0 then $a \odot b \le c$ and $a \le b \to c$ are equivalent to $a \le b'$. If a = 1 then $a \odot b \le c$ and $a \le b \to c$ are equivalent to $b \le c$. If b = 1 then $a \odot b \le c$ and $a \le b \to c$ are equivalent to $a \le c$. If c = 1 then $a \odot b \le c$ and $a \le b \to c$. If $a, b, c \ne 0, 1$ then $a \odot b \le c_2$ and $c_3 \le b \to c$ and hence $a \odot b \le c$ and $a \le b \to c$. Thus the adjointness property holds.

As mentioned above, the non-modular lattice \mathbf{N}_5 with an antitone involution cannot be converted into a residuated lattice $(N_5, \lor, \land, \odot, \rightarrow, 1)$ with an antitone involution ' satisfying $x' = x \to 0$ for all $x \in N_5$. Using Theorem 2, we can extend \mathbf{N}_5 as follows.

Example 3. Theorem 2 applied to N_5 yields the residuated lattice depicted in Figure 2:



with operation tables

\odot	0	c_2	a	b	c	c_3	1	\rightarrow	0	c_2	a	b	c	c_3	1
0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1
c_2	0	0	0	0	0	0	c_2	c_2	c_3	1	1	1	1	1	1
a	0	0	0	0	c_2	c_2	a	a	b	C_3	1	1	C_3	1	1
b	0	0	0	c_2	c_2	c_2	b	b	a	c_3	c_3	1	c_3	1	1
c	0	0	c_2	c_2	0	c_2	c	c	С	c_3	c_3	c_3	1	1	1
c_3	0	0	c_2	c_2	c_2	c_2	c_3	c_3	c_2	c_3	c_3	c_3	c_3	1	1
1	0	c_2	a	b	c	c_3	1	1	0	c_2	a	b	С	c_3	1

Observe there is only one possibility for the antitone involution.

Let us note that if $\mathbf{P} = (P, \leq, ')$ is a finite chain containing at least three elements (with unique antitone involution) then, using the construction from Theorem 2, \mathbf{P} can be converted into a residuated chain $\mathbb{E}(\mathbf{P}) = (P, \leq, \odot, \rightarrow, 1)$ satisfying $x' = x \rightarrow 0$ for all $x \in P$.

Corollary 4. If $(C, \leq) = (\{c_1, \ldots, c_n\}, \leq)$ is a finite chain $0 = c_1 < c_2 < c_3 < \cdots < c_n = 1$ with $n \geq 3$ elements,

$$\begin{array}{l} 0 \odot c_i = c_i \odot 0 := 0, 1 \odot c_i = c_i \odot 1 := c_i, \\ 0 \to c_i = c_i \to 1 := 1, c_i \to 0 := c_{n+1-i}, 1 \to c_i := c_i \end{array}$$

for $i = 1, \ldots, n$ and

$$c_i \odot c_j := \begin{cases} 0 & \text{if } i+j \le n+1, \\ c_2 & \text{otherwise} \end{cases} \qquad c_i \to c_j := \begin{cases} 1 & \text{if } i \le j, \\ c_{n-1} & \text{otherwise} \end{cases}$$

for i, j = 2, ..., n-1 then $(C, \leq, \odot, \rightarrow, 1)$ is a residuated lattice where $x' := x \rightarrow 0$ is an antitone involution.

Example 5. Theorem 2 or Corollary 4 applied to the five-element chain yields the residuated lattice depicted in Figure 3:



with operation tables

\odot	0	c_2	c_3	c_4	1		\rightarrow	0	c_2	c_3	c_4	1
0	0	0	0	0	0	-	0	1	1	1	1	1
c_2	0	0	0	0	c_2		c_2	c_4	1	1	1	1
c_3	0	0	0	c_2	c_3		c_3	c_3	c_4	1	1	1
c_4	0	0	c_2	c_2	c_4		c_4	c_2	c_4	c_4	1	1
1	0	c_2	c_3	c_4	1		1	0	c_2	c_3	c_4	1

Here c_3 of Theorem 2 corresponds to c_4 of Example 5.

If the poset **P** in question is a lattice, we can also apply the construction of ', \odot and \rightarrow from Theorem 2 to obtain a residuated lattice $\mathbb{E}(\mathbf{P})$. Hence, we can state the following.

Corollary 6. Let $\mathbf{L} = (L, \lor, \land, ')$ be a lattice with an antitone involution '. Then \mathbf{L} can be extended to a residuated lattice $\mathbb{E}(\mathbf{L})$ with an antitone involution where the operations ', \odot and \rightarrow are constructed as in Theorem 2 and ' coincides in L with the original one.

Recall from [2] and [11] that a lattice $\mathbf{L} = (L, \vee, \wedge, ')$ with an antitone involution is called a *pseudo-Kleene algebra* if it satisfies the identities

- (1) $x \wedge x' \leq y \vee y'$,
- (2) $x \wedge (x' \vee y) \approx (x \wedge x') \vee (x \wedge y).$

L is called a *Kleene algebra* if it is a distributive pseudo-Kleene algebra. In this case (2) can be omitted since it follows by distributivity. Kleene algebras and pseudo-Kleene algebras are considered as an algebraic axiomatization of a propositional logic satisfying De Morgan's laws and the double negation law, but not necessarily the law of excluded middle because the antitone involution ' need not be a complementation, see also [9]. An example of a Kleene algebra is depicted in Figure 4:



A pseudo-Kleene algebra which is not a Kleene algebra is visualized in Figure 5:



One can easily check that if \mathbf{L} satisfies (1) and (2) then so does $\mathbb{E}(\mathbf{L})$ and if \mathbf{L} is distributive then also $\mathbb{E}(\mathbf{L})$ has this property, too. Hence, we can state

Corollary 7. Every pseudo-Kleene algebra or every Kleene algebra can be extended to a residuated pseudo-Kleene algebra or a residuated Kleene-algebra, respectively.

Hence, also the logic axiomatized by Kleene algebras or pseudo-Kleene algebras can be extended to a kind of fuzzy logics.

Motivated by the previous, we can extend every poset with an antitone involution to a residuated poset by means of a finite chain with an even number of at least four elements. The precise formulation is as follows:

Theorem 8. Let $(P, \leq, ')$ be a poset with an antitone involution, n an integer > 1, $0 = c_1, \ldots, c_{2n} = 1 \notin P$ and $Q := P \cup \{c_1, \ldots, c_{2n}\}$. Extend \leq and ' from P to Q by

$$c_1 < \dots < c_n < x < c_{n+1} < \dots < c_{2n}$$

for $x \in P$ and put $c'_i := c_{2n+1-i}$ for i = 1, ..., 2n. Define binary operations \odot and \rightarrow on Q as follows:

$$\begin{array}{l} 0 \odot x = x \odot 0 := 0, 1 \odot x = x \odot 1 := x, \\ 0 \rightarrow x = x \rightarrow 1 := 1, x \rightarrow 0 := x', 1 \rightarrow x := x \end{array}$$

for $x \in Q$ and

$$x \odot y := \begin{cases} 0 & \text{if } x \le y', \\ c_2 & \text{otherwise} \end{cases} \qquad x \to y := \begin{cases} 1 & \text{if } x \le y, \\ c_{2n-1} & \text{otherwise} \end{cases}$$
$$c_i \odot c_j := \begin{cases} 0 & \text{if } i+j \le 2n+1, \\ c_2 & \text{otherwise} \end{cases} \qquad c_i \to c_j := \begin{cases} 1 & \text{if } i \le j, \\ c_{2n-1} & \text{otherwise} \end{cases}$$
$$c_i \odot x = x \odot c_i := c_i \odot c_{n+1}, c_i \to x := c_i \to c_n, x \to c_i := c_{n+1} \to c_i \end{cases}$$

for $x, y \in P$ and i, j = 2, ..., 2n - 1. Then $(Q, \leq, \odot, \rightarrow, 1)$ is a residuated poset with the antitone involution ' satisfying $x' = x \rightarrow 0$ for all $x \in Q$.

Proof. We apply Theorem 2 to the poset $(P \cup \{c_3, \ldots, c_{2n-2}\}, \leq, ')$ with the antitone involution '. The element c_3 of Theorem 2 corresponds to the element c_{2n-1} of Theorem 8. According to Theorem 2 we have for all $x \in Q$

$$0 \odot x = x \odot 0 = 0, 1 \odot x = x \odot 1 = x,$$

$$0 \rightarrow x = x \rightarrow 1 = 1, x \rightarrow 0 = x', 1 \rightarrow x = x$$

Moreover, we have for all $x, y \in P$ and $i, j \in \{2, \dots, 2n-1\}$

$$x \odot y = \begin{cases} 0 & \text{if } x \le y', \\ c_2 & \text{otherwise} \end{cases} \qquad x \to y = \begin{cases} 1 & \text{if } x \le y, \\ c_{2n-1} & \text{otherwise} \end{cases}$$
$$c_i \odot c_j = \begin{cases} 0 & \text{if } i \le 2n+1-j, \\ c_2 & \text{otherwise} \end{cases} \qquad c_i \to c_j = \begin{cases} 1 & \text{if } i \le j, \\ c_{2n-1} & \text{otherwise} \end{cases}$$

$$c_i \odot x = x \odot c_i = \begin{cases} 0 & \text{if } i \le n, \\ c_2 & \text{otherwise} \end{cases}$$

$$c_i \to x = \begin{cases} 1 & \text{if } i \le n, \\ c_{2n-1} & \text{otherwise} \end{cases} \qquad x \to c_i = \begin{cases} 1 & \text{if } i > n, \\ c_{2n-1} & \text{otherwise} \end{cases}$$

The poset (Q, \leq) of Theorem 8 is visualized in Figure 6:

$$\begin{array}{c}
\bullet \ c_{2n} = 1 \\
\vdots \\
\bullet \ c_{n+1} \\
P \\
\bullet \ c_n \\
\vdots \\
\bullet \ c_1 = 0 \\
\end{array}$$
Fig. 6

Let P be a set and $\mathbf{P}_1 = (P \times \{1\}, \leq)$ and $\mathbf{P}_2 = (P \times \{2\}, \leq)$ posets. We call \mathbf{P}_2 the *dual* of \mathbf{P}_1 if for all $x, y \in P$ we have $(x, 2) \leq (y, 2)$ if and only if $(y, 1) \leq (x, 1)$. Similarly as before, a poset together with its dual can be extended to a residuated poset by means of a finite chain with at least four elements.

Theorem 9. Let $(P \times \{1\}, \leq)$ be a poset, $(P \times \{2\}, \leq)$ denote its dual, n an integer > 1, k a non-negative integer, $0 = c_1, \ldots, c_{2n+k} = 1 \notin P \times \{1, 2\}$ and $Q := (P \times \{1, 2\}) \cup \{c_1, \ldots, c_{2n+k}\}$. Extend \leq to Q by

$$c_1 < \dots < c_n < (x, 1) < c_{n+1} < \dots < c_{n+k} < (y, 2) < c_{n+k+1} < \dots < c_{2n+k}$$

for $x, y \in P$ and put $c'_i := c_{2n+k+1-i}$ for i = 1, ..., 2n+k and (x, i)' := (x, 3-i) for $x \in P$ and $i \in \{1, 2\}$. Define binary operations \odot and \rightarrow on Q as follows:

$$0 \odot x = x \odot 0 := 0, 1 \odot x = x \odot 1 := x,$$

$$0 \rightarrow x = x \rightarrow 1 := 1, x \rightarrow 0 := x', 1 \rightarrow x := x$$

for $x \in Q$,

$$(x,i) \odot (y,j) := \begin{cases} 0 & if \ i = j = 1 \ or \ ((i,j) = (1,2) \ and \ x \le y) \ or \\ (i,j) = (2,1) \ and \ y \le x, \\ c_2 & otherwise \end{cases}$$

$$(x,i) \rightarrow (y,j) := \begin{cases} 1 & if \ (i = j = 1 \ and \ x \le y) \ or \ (i,j) = (1,2) \ or \\ (i = j = 2 \ and \ y \le x), \\ c_{2n+k-1} & otherwise \end{cases}$$

for $x, y \in P$ and $i, j \in \{1, 2\}$ and

$$c_i \odot c_j := \begin{cases} 0 & \text{if } i+j \le 2n+k+1, \\ c_2 & \text{otherwise} \end{cases} \qquad c_i \to c_j := \begin{cases} 1 & \text{if } i \le j, \\ c_{2n+k-1} & \text{otherwise} \end{cases}$$

$$c_{i} \odot (x,1) = (x,1) \odot c_{i} := c_{i} \odot c_{n+1}, c_{i} \to (x,1) := c_{i} \to c_{n}, (x,1) \to c_{i} := c_{n+1} \to c_{i};$$

$$c_{i} \odot (x,2) = (x,2) \odot c_{i} := c_{i} \odot c_{n+k+1}, c_{i} \to (x,2) := c_{i} \to c_{n+k},$$

$$(x,2) \to c_{i} := c_{n+k+1} \to c_{i}$$

for $i, j \in \{2, ..., 2n + k - 1\}$ and $x \in P$. Then $(Q, \leq, \odot, \rightarrow, 1)$ is a residuated poset with an antitone involution ' satisfying $x' = x \rightarrow 0$ for all $x \in Q$.

Proof. We apply Theorem 2 to the poset $(P \cup \{c_3, \ldots, c_{2n+k-2}\}, \leq, \prime)$ with the antitone involution \prime . The element c_3 of Theorem 2 corresponds to the element c_{2n+k-1} of Theorem 9. According to Theorem 2 we have for all $x \in Q$

$$0 \odot x = x \odot 0 = 0, 1 \odot x = x \odot 1 = x,$$

$$0 \rightarrow x = x \rightarrow 1 = 1, x \rightarrow 0 = x', 1 \rightarrow x = x$$

Moreover, we have for all $x, y \in P$ and $i, j \in \{1, 2\}$

$$(x,i) \odot (y,j) = \begin{cases} 0 & \text{if } (x,i) \le (y,3-j), \\ c_2 & \text{otherwise} \end{cases}$$
$$(x,i) \to (y,j) = \begin{cases} 1 & \text{if } (x,i) \le (y,j), \\ c_{2n+k-1} & \text{otherwise} \end{cases}$$

Further, we have for all $i, j \in \{2, \ldots, 2n + k - 1\}$

$$c_i \odot c_j = \begin{cases} 0 & \text{if } i \le 2n+k+1-j, \\ c_2 & \text{otherwise} \end{cases} \qquad c_i \to c_j = \begin{cases} 1 & \text{if } i \le j, \\ c_{2n+k-1} & \text{otherwise} \end{cases}$$

Finally, we have for all $i \in \{2, \ldots, 2n + k - 1\}$ and $j \in \{1, 2\}$

$$c_i \odot (x, j) = (x, j) \odot c_i = \begin{cases} 0 & \text{if } c_i \le (x, 3 - j), \\ c_2 & \text{otherwise} \end{cases}$$

$$c_i \to (x, j) = \begin{cases} 1 & \text{if } c_i \le (x, j), \\ c_{2n+k-1} & \text{otherwise} \end{cases} \qquad (x, j) \to c_i = \begin{cases} 1 & \text{if } (x, j) \le c_i, \\ c_{2n+k-1} & \text{otherwise} \end{cases}$$

The poset (Q, \leq) of Theorem 8 is visualized in Figure 7:

•
$$c_{2n+k} =$$

: c_{n+k+1}
 P^d
• c_{n+k}
: c_{n+1}
 P
• c_n
: $c_1 = 0$
Fig. 7

1

It is obvious that the constructions described in Theorems 8 and 9 can be generalized to finitely many posets and finite chains in between them. For instance, if we have posets P_1, P_2, P_3 , their duals P_1^d, P_2^d, P_3^d and a poset P with an antitone involution then we can construct, similarly as before, a poset with an antitone involution of the form

$$\cdots < x_1 < \cdots < x_2 < \cdots < x_3 < \cdots < x < \cdots < y_3 < \cdots < y_2 < \cdots < y_1 < \cdots$$

for all $x_1 \in P_1, x_2 \in P_2, x_3 \in P_3, x \in P, y_3 \in P_3^d, y_2 \in P_2^d, y_1 \in P_1^d$ where the dots (from left to right) indicate finite chains with r, s, t, u, u, t, s, r elements, respectively, and then we can apply Theorem 2 to this poset with antitone involution.

The following result is well known, see e.g. [1] or [12].

Lemma 10. Let $(B, \lor, \land, ', p, q)$ be a Boolean algebra and put

$$x \odot y := x \land y,$$

$$x \to y := x' \lor y$$

for every $x, y \in B$. Then $(B, \leq, \odot, \rightarrow, q)$ is a residuated poset.

In the following we extend a Boolean algebra $(B, \vee, \wedge, ', p, q)$ to a residuated lattice $(Q, \vee, \wedge, \odot, \rightarrow, 1)$ with an antitone involution $x' = x \rightarrow 0$ by means of a (finite) chain such that the operations \odot and \rightarrow coincide on B with those mentioned in Lemma 10.

Theorem 11. Let $(B, \lor, \land, ', p, q)$ be a Boolean algebra, n a positive integer, $0 = c_1, \ldots$ $\ldots, c_{2n} = 1 \notin B$ and $Q := B \cup \{c_1, \ldots, c_{2n}\}$. Extend \leq and ' from B to Q by

$$c_1 < \dots < c_n < x < c_{n+1} < \dots < c_{2n}$$

for $x \in B$ and put $c'_i := c_{2n+1-i}$ for i = 1, ..., 2n. Define binary operations \odot and \rightarrow on Q as follows:

$$x \odot y := \begin{cases} 0 & \text{if } x \le y', \\ x \land y & \text{otherwise} \end{cases} \qquad x \to y := \begin{cases} 1 & \text{if } x \le y, \\ x' \lor y & \text{otherwise} \end{cases}$$

for $x, y \in Q$. Then $(Q, \leq, \odot, \rightarrow, 1)$ is a residuated lattice with an antitone involution $x' = x \rightarrow 0$ where for all $x, y \in B$ we have

$$x \odot y = x \land y,$$

$$x \to y = x' \lor y$$

Proof. Let $a, b, c \in Q$, $d, e \in B$ and $i, j, k \in \{1, \ldots, 2n\}$. Within this proof

$$a < B$$
 means $a < x$ for all $x \in B$,
 $a > B$ means $a > x$ for all $x \in B$,
 $i \lor j := \max(i, j)$,
 $i \land j := \min(i, j)$.

Obviously, $(Q, \leq, ', 0, 1)$ is a bounded lattice with an antitone involution. We have

$$a \odot b = b \odot a \text{ since } a \leq b' \text{ is equivalent to } b \leq a',$$

$$0 \odot a = 0,$$

$$1 \odot a = \begin{cases} 0 = a & \text{if } a = 0, \\ 1 \wedge a = a & \text{otherwise} \end{cases}$$

$$0 \rightarrow a = a \rightarrow 1 = 1,$$

$$a \rightarrow 0 = \begin{cases} 1 = 0' = a' & \text{if } a = 0, \\ a' \vee 0 = a' & \text{otherwise} \end{cases}$$

$$d \odot e = \begin{cases} c_1 = d \wedge e & \text{if } d \leq e', \\ d \wedge e & \text{otherwise} \end{cases}$$

$$d \rightarrow e = \begin{cases} 1 = d' \vee e & \text{if } d \leq e, \\ d' \vee e & \text{otherwise.} \end{cases}$$

This shows

$$0 \odot x = x \odot 0 = 0, 1 \odot x = x \odot 1 = x,$$

$$0 \rightarrow x = x \rightarrow 1 = 1, x \rightarrow 0 = x', 1 \rightarrow x = x$$

 $(x \in Q)$ and

$$\begin{aligned} x \odot y &= x \land y, \\ x \to y &= x' \lor y \end{aligned}$$

for all $x, y \in B$.

If $a \leq b'$ and $b \leq c'$ then $(a \odot b) \odot c = 0 \odot c = 0 = a \odot 0 = a \odot (b \odot c)$. If $a \leq b'$ and $b \nleq c'$ then $(a \odot b) \odot c = 0 \odot c = 0 = a \odot (b \land c) = a \odot (b \odot c)$. If $a \nleq b'$ and $b \leq c'$ then $(a \odot b) \odot c = (a \land b) \odot c = 0 = a \odot 0 = a \odot (b \odot c)$. Now consider the case $a \nleq b'$ and $b \nleq c'$. Then

$$(a \odot b) \odot c = (a \land b) \odot c,$$

$$a \odot (b \odot c) = a \odot (b \land c).$$

If $a, b, c \in B$ then $(a \wedge b) \odot c = (a \wedge b) \wedge c = a \wedge (b \wedge c) = a \odot (b \wedge c)$. If $a, b \in B$ and $c \notin B$ then c > B and hence $(a \wedge b) \odot c = (a \wedge b) \wedge c = a \wedge b = a \odot b = a \odot (b \wedge c)$. If $a \in B$, $b \notin B$ and $c \in B$ then b > B and hence $(a \wedge b) \odot c = a \odot c = a \odot (b \wedge c)$. If $a \in B$ and $b, c \notin B$ then b > B and $(a \leq c')$ if and only if $a \leq b' \vee c'$. Hence

If $a \in B$ and $b, c \notin B$ then b > B and $(a \le c)$ if and only if $a \le b \lor c$. Hence $(a \land b) \odot c = a \odot c = 0 = a \odot (b \land c)$ if $a \le c'$ and $(a \land b) \odot c = a \odot c = a \land c = (a \land b) \land c = a \land (b \land c) = a \odot (b \land c)$ otherwise.

If $a \notin B$ and $b, c \in B$ then a > B and hence $(a \land b) \odot c = b \odot c = b \land c = (a \land b) \land c = a \land (b \land c) = a \odot (b \land c).$

If $a \notin B$, $b \in B$ and $c \notin B$ then a, c > B and hence $(a \land b) \odot c = b \odot c = b \land c = b = a \land b = a \odot (b \land c)$.

If $a, b \notin B$ and $c \in B$ then b > B and $(a \le c' \text{ if and only if } a \land b \le c')$. Hence $(a \land b) \odot c = 0 = a \odot c = a \odot (b \land c)$ if $a \le c'$ and $(a \land b) \odot c = (a \land b) \land c = a \land (b \land c) = a \land c = a \odot (b \land c)$ otherwise.

Finally, assume $a, b, c \notin B$. Without loss of generality, $(a, b, c) = (c_i, c_j, c_k)$. Now we have

$$c_i \odot c_j = \begin{cases} 0 & \text{if } i+j \le 2n+1, \\ c_{i \land j} & \text{otherwise} \end{cases}$$

If $i + j, j + k \leq 2n + 1$ then $(c_i \odot c_j) \odot c_k = 0 \odot c_k = 0 = c_i \odot 0 = c_i \odot (c_j \odot c_k)$. If $i + j \leq 2n + 1 < j + k$ then $i + (j \land k) = (i + j) \land (i + k) \leq i + j \leq 2n + 1$ and hence $(c_i \odot c_j) \odot c_k = 0 \odot c_k = 0 = c_i \odot c_{j \land k} = c_i \odot (c_j \odot c_k)$. If $j + k \leq 2n + 1 < i + j$ then $(i \land j) + k = (i + k) \land (j + k) \leq j + k \leq 2n + 1$ and hence $(c_i \odot c_j) \odot c_k = c_{i \land j} \odot c_k = 0 = c_i \odot 0 = c_i \odot (c_j \odot c_k)$. If i + j, j + k > 2n + 1 then

$$(i \wedge j) + k = (i+k) \wedge (j+k) \le 2n+1$$
 if and only if $i+k \le 2n+1$,
 $i+(j \wedge k) = (i+j) \wedge (i+k) \le 2n+1$ if and only if $i+k \le 2n+1$

and hence $(c_i \odot c_j) \odot c_k = c_{i \wedge j} \odot c_k = c_i \odot c_{j \wedge k} = c_i \odot (c_j \odot c_k)$. This shows that \odot is associative. Since $a \leq b'$ is equivalent to $b \leq a'$, \odot is commutative. Therefore, $(Q, \odot, 1)$ is a commutative monoid. If $a \leq b'$ and $b \leq c$ then $a \odot b = 0 \leq c$ and $a \leq 1 = b \rightarrow c$. If $a \leq b'$ and $b \not\leq c$ then $a \odot b = 0 \leq c$, and $a \leq b \rightarrow c$ since $a \leq b' \lor c$. If $a \not\leq b'$ and $b \leq c$ then $a \odot b \leq c$ since $a \wedge b \leq c$, and $a \leq 1 = b \rightarrow c$. Now assume $a \not\leq b'$ and $b \not\leq c$. If $a, b, c \in B$ then $a \wedge b \leq c$ if and only if $a \leq b' \lor c$ because of Lemma 10. If $a, b \in B$ and $c \notin B$ then c < B and hence $a \wedge b \leq c$ and $a \leq b' \lor c$. If $b \notin B$ and $c \in B$ then b > B and hence $a \wedge b \leq c$ and $a \leq b' \lor c$ are both equivalent to $a \leq c$. If $a \in B$ and $b, c \notin B$ then b > B and hence $a \wedge b \leq c$ and $a \leq b' \lor c$ are both equivalent to $a \leq c$.

If $a \in B$ and $b, c \notin B$ then b > B and hence $a \land b \le c$ and $a \le b' \lor c$ are both equivalent to $a \le c$.

If $a \notin B$ and $b, c \in B$ then a > B and hence $a \wedge b = b \not\leq c$ and $a \not\leq b' \vee c$. If $a \notin B$, $b \in B$ and $c \notin B$ then c < B < a and hence $a \wedge b = b \not\leq c$ and $a \not\leq b' = b' \vee c$.

Finally, assume $a, b, c \notin B$. Without loss of generality, $(a, b, c) = (c_i, c_j, c_k)$. Now we have

$$c_i \to c_j = \begin{cases} 1 & \text{if } i \leq j, \\ c_{(2n+1-i)\lor j} & \text{otherwise} \end{cases}$$

If $i + j \leq 2n + 1$ and $j \leq k$ then $c_i \odot c_j = 0 \leq c_k$ and $c_i \leq 1 = c_j \rightarrow c_k$. If $i + j \leq 2n + 1$ and j > k then $c_i \odot c_j = 0 \leq c_k$ and $c_i \leq c_{(2n+1-j)\vee k}$ since $i \leq 2n + 1 - j$. If i + j > 2n + 1 and $j \leq k$ then $c_i \odot c_j = c_{i\wedge j} \leq c_k$ and $c_i \leq c_{2n} = c_j \rightarrow c_k$. Finally, assume i + j > 2n + 1 and j > k. Then the following are equivalent:

$$c_i \odot c_j \le c_k, c_{i \land j} \le c_k, i \land j \le k, i \le k.$$

Moreover, the following are equivalent:

$$c_i \leq c_j \rightarrow c_k,$$

$$c_i \leq c_{(2n+1-j)\vee k},$$

$$i \leq (2n+1-j) \vee k,$$

$$i+j \leq (2n+1) \vee (k+j),$$

$$i+j \leq k+j,$$

$$i \leq k.$$

This proves adjointness.

The poset (Q,\leq) of Theorem 8 is visualized in Figure 8:



Example 12. We consider the special case where $(B, \lor, \land, ', p, q)$ is the eight-element Boolean algebra and n = 2. The poset (Q, \leq) is visualized in Figure 9:



The corresponding residuated lattice has the operations

\odot	0	c_2	p	a	b	c a	b'	c'	q	c	3.	1
0	0	0	0	0	0	0 0	0	0	0	0	()
c_2	0	0	0	0	0	0 0	0	0	0	0	(C_2
p	0	0	0	0	0	0 0	0	0	0	p	1	p
a	0	0	0	a	0	0 0	a	a	a	a	. (a
b	0	0	0	0	b	0 b	0	b	b	b	l	5
c	0	0	0	0	0	c c	c	0	c	c	(c
a'	0	0	0	0	b	c a	c	b	a'	a	<i>'</i> (a'
b'	0	0	0	a	0	c c	b'	a	b'	b	' i	6'
c'	0	0	0	a	b	0 b	a	c'	c'	c	΄ (c'
q	0	0	0	a	b	c a	b'	c'	q	q	(7
c_3	0	0	p	a	b	c a	b'	c'	q	c	3 (C_3
1	0	c_2	p	a	b	c a	b'	c'	q	c	3 .	1
\rightarrow	0	c_2	p	a	b	С	a'	b'	c'	q	C_3	1
\rightarrow 0	0 1	$\frac{c_2}{1}$	<i>p</i> 1	$\frac{a}{1}$	$\frac{b}{1}$	<i>c</i> 1	$\frac{a'}{1}$	<i>b</i> ′	$\frac{c'}{1}$	<i>q</i> 1	$\frac{c_3}{1}$	1
\rightarrow 0 c_2	0 1 c ₃	$ \begin{array}{c} c_2 \\ 1 \\ 1 \end{array} $	p 1 1	$\frac{a}{1}$	<i>b</i> 1 1	$\frac{c}{1}$	a' 1 1	b' 1 1	<i>c</i> ′ 1 1	$\frac{q}{1}$	c_3 1 1	1 1 1
$\begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_3 \\ q \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \end{array}$	<i>p</i> 1 1 1	a 1 1 1	b 1 1 1	<i>c</i> 1 1 1	a' 1 1 1	b' 1 1 1	c' 1 1 1	q 1 1 1	<i>c</i> ₃ 1 1	1 1 1 1
$\begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \\ a \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_3 \\ q \\ a' \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \end{array}$	$\begin{array}{c} p \\ 1 \\ 1 \\ 1 \\ a' \end{array}$	a 1 1 1 1	b 1 1 1 a'	c 1 1 1 1 a'	a' 1 1 1 a'	b' 1 1 1 1	<i>c'</i> 1 1 1 1	q 1 1 1 1	c_3 1 1 1 1	1 1 1 1 1
$\begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \\ a \\ b \end{array}$	$egin{array}{c} 0 \ 1 \ c_3 \ q \ a' \ b' \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \end{array}$	$p \\ 1 \\ 1 \\ 1 \\ a' \\ b'$	$\begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ 1 \\ b' \end{array}$	$\begin{array}{c} b\\ 1\\ 1\\ 1\\ a'\\ 1\end{array}$	$\begin{array}{c} c\\ 1\\ 1\\ 1\\ a'\\ b' \end{array}$	a' 1 1 1 a' 1	b' 1 1 1 1 b'	<i>c'</i> 1 1 1 1	$egin{array}{c} q \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array}$	c_3 1 1 1 1 1 1	1 1 1 1 1 1
$\begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \\ a \\ b \\ c \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_{3} \\ q \\ a' \\ b' \\ c' \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \end{array}$	$p \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ c'$	$\begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ 1 \\ b' \\ c' \end{array}$	$\begin{array}{c} b\\ 1\\ 1\\ 1\\ a'\\ 1\\ c'\end{array}$	$\begin{array}{c} c \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ 1 \end{array}$	a' 1 1 1 a' 1 1	b' 1 1 1 1 b' 1	$\begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \end{array}$	$egin{array}{c} q \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array}$	c_3 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1
$ \begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \\ a \\ b \\ c \\ a' \end{array} $	$egin{array}{c} 0 \ 1 \ c_3 \ q \ a' \ b' \ c' \ a \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \\ a \end{array}$	$p \\ 1 \\ 1 \\ a' \\ b' \\ c' \\ a$	$\begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ b' \\ c' \\ a \end{array}$	$\begin{array}{c} b \\ 1 \\ 1 \\ 1 \\ a' \\ 1 \\ c' \\ c' \end{array}$	$c \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ 1 \\ b' \\ b'$	a' 1 1 1 a' 1 1 1	b' 1 1 1 b' 1 b'	$\begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \\ c' \end{array}$	$egin{array}{c} q \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array}$	c_3 1 1 1 1 1 1 1 1	$ \begin{array}{c} 1 \\ $
$\begin{array}{c} \rightarrow \\ 0 \\ c_2 \\ p \\ a \\ b \\ c \\ a' \\ b' \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_{3} \\ q \\ a' \\ b' \\ c' \\ a \\ b \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \\ a \\ b \end{array}$	$\begin{array}{c} p \\ 1 \\ 1 \\ a' \\ b' \\ c' \\ a \\ b \end{array}$	$egin{array}{c} a \\ 1 \\ 1 \\ 1 \\ b' \\ c' \\ a \\ c' \end{array}$	$\begin{array}{c} b \\ 1 \\ 1 \\ 1 \\ a' \\ 1 \\ c' \\ c' \\ b \end{array}$	$\begin{array}{c} c \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ 1 \\ b' \\ a' \end{array}$	$\begin{array}{c} a' \\ 1 \\ 1 \\ 1 \\ a' \\ 1 \\ 1 \\ 1 \\ a' \end{array}$	b' 1 1 1 1 b' 1 b' 1 b' 1	$\begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \\ c' \\ c' \\ c' $	$egin{array}{c} q \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\$	c_3 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1
$\begin{array}{c} \rightarrow \\ \hline 0 \\ c_2 \\ p \\ a \\ b \\ c \\ a' \\ b' \\ c' \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_{3} \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \end{array}$	$\begin{array}{c} p \\ 1 \\ 1 \\ a' \\ b' \\ c' \\ a \\ b \\ c \end{array}$	$\begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ 1 \\ b' \\ c' \\ a \\ c' \\ b' \end{array}$	$\begin{array}{c} b \\ 1 \\ 1 \\ a' \\ 1 \\ c' \\ c' \\ b \\ a' \end{array}$	$\begin{array}{c} c \\ 1 \\ 1 \\ a' \\ b' \\ 1 \\ b' \\ a' \\ c \end{array}$	$\begin{array}{c} a' \\ 1 \\ 1 \\ 1 \\ a' \\ 1 \\ 1 \\ 1 \\ a' \\ a'$	b' 1 1 1 1 b' 1 b' 1 b' 1 b'	$\begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \\ c' \\ c' \\ 1 \end{array}$	q 1 1 1 1 1 1 1 1 1 1 1 1 1 1	c_3 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1
$ \begin{array}{c} \rightarrow \\ \hline 0 \\ c_2 \\ p \\ a \\ b \\ c \\ a' \\ b' \\ c' \\ q \end{array} $	$\begin{array}{c} 0 \\ 1 \\ c_{3} \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \end{array}$	$\begin{array}{c} c_2 \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \end{array}$	$\begin{array}{c} p \\ 1 \\ 1 \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \end{array}$	$ \begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ 1 \\ b' \\ c' \\ a \\ c' \\ b' \\ a \\ a' \\ b' \\ a' \end{array} $	$\begin{array}{c} b \\ 1 \\ 1 \\ a' \\ 1 \\ c' \\ c' \\ b \\ a' \\ b \end{array}$	$\begin{array}{c} c \\ 1 \\ 1 \\ a' \\ b' \\ 1 \\ b' \\ a' \\ c \\ c \\ c \end{array}$	$egin{array}{c} a' \\ 1 \\ 1 \\ a' \\ 1 \\ 1 \\ 1 \\ a' \\ a' \\$	b' 1 1 1 1 b' 1 b' 1 b' b' b' b'	$ \begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \\ c' \\ c' \\ 1 \\ c' \\ c'$	$rac{q}{1}$ 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	c_3 1 1 1 1 1 1 1 1 1 1 1 1 1	$ \begin{array}{c} 1 \\ $
$\begin{array}{c} \rightarrow \\ \hline 0 \\ c_2 \\ p \\ a \\ b \\ c \\ a' \\ b' \\ c' \\ q \\ c_3 \end{array}$	$\begin{array}{c} 0 \\ 1 \\ c_{3} \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \\ c_{2} \end{array}$	$\begin{array}{c} c_{2} \\ 1 \\ 1 \\ q \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \\ c_{2} \end{array}$	$\begin{array}{c} p \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ c' \\ a \\ b \\ c \\ p \\ p \end{array}$	$\begin{array}{c} a \\ 1 \\ 1 \\ 1 \\ b' \\ c' \\ a \\ c' \\ b' \\ a \\ a \end{array}$	$\begin{array}{c} b \\ 1 \\ 1 \\ 1 \\ a' \\ 1 \\ c' \\ c' \\ b \\ a' \\ b \\ b \end{array}$	$\begin{array}{c} c \\ 1 \\ 1 \\ 1 \\ a' \\ b' \\ 1 \\ b' \\ a' \\ c \\ c \\ c \\ c \\ c \end{array}$	$ \begin{array}{c} a' \\ 1 \\ 1 \\ a' \\ 1 \\ 1 \\ a' \\ a'$	b' 1 1 1 1 b' 1 b' 1 b' b' b' b' b'	$ \begin{array}{c} c' \\ 1 \\ 1 \\ 1 \\ 1 \\ c' \\ c' \\ c' \\ 1 \\ c' \\ c'$	$rac{q}{1} \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \$	$egin{array}{c_3} 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 $	$ \begin{array}{c} 1 \\ $

References

- R. Bělohlávek, Fuzzy Relational Systems. Foundations and Principles. Springer, New York 2002. ISBN 978-1-4613-5168-9.
- [2] I. Chajda, A note on pseudo-Kleene algebras. Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math. 55 (2016), 39–45.
- [3] I. Chajda, A representation of residuated lattices satisfying the double negation law. Soft Computing 22 (2018), 1773–1776.
- [4] I. Chajda and D. Fazio, On residuation in paraorthomodular lattices. Soft Computing. DOI 10.1007/s00500-020-04699-w.
- [5] I. Chajda and H. Länger, Residuated operators in complemented posets. Asian-Eur. J. Math. 11 (2018), 1850097, 15 pp.

- [6] I. Chajda and H. Länger, Residuated structures derived from commutative idempotent semirings. Discuss. Math. Gen. Algebra Appl. 39 (2019), 23–33.
- [7] I. Chajda and H. Länger, Residuation in modular lattices and posets. Asian-Eur. J. Math. 12 (2019), 1950092, 10 pp.
- [8] L. C. Ciungu, Non-commutative Multiple-Valued Logic Algebras. Springer, Cham 2014. ISBN 978-3-319-01588-0.
- [9] P. Jipsen, From semirings to residuated Kleene lattices. Studia Logica 76 (2004), 291–303.
- [10] P. Jipsen and C. Tsinakis, A survey of residuated lattices. In J. Martínez (eds), Ordered Algebraic Structures, 19–56. Kluwer, Dordrecht 2002.
- [11] J. A. Kalman, Lattices with involution. Trans. Amer. Math. Soc. 87 (1958), 485–491.
- [12] M. Ward and R. P. Dilworth, Residuated lattices. Trans. Amer. Math. Soc. 45 (1939), 335–354.

Authors' addresses:

Ivan Chajda Palacký University Olomouc Faculty of Science Department of Algebra and Geometry 17. listopadu 12 771 46 Olomouc Czech Republic ivan.chajda@upol.cz Miroslav Kolařík Palacký University Olomouc Faculty of Science Department of Computer Science 17. listopadu 12 771 46 Olomouc Czech Republic miroslav.kolarik@upol.cz Helmut Länger TU Wien Faculty of Mathematics and Geoinformation Institute of Discrete Mathematics and Geometry Wiedner Hauptstraße 8-10 1040 Vienna Austria, and Palacký University Olomouc Faculty of Science Department of Algebra and Geometry 17. listopadu 12 771 46 Olomouc

Czech Republic helmut.laenger@tuwien.ac.at