The maximum regularity property of the steady Stokes problem associated with a flow through a profile cascade

Tomáš Neustupa

Received: date / Accepted: date

Abstract We deal with a steady Stokes-type problem, associated with a flow of a Newtonian incompressible fluid through a spatially periodic profile cascade. The used mathematical model is based on the reduction to one spatial period, represented by a bounded 2D domain Ω . The corresponding Stokes-type problem is formulated by means of the Stokes equation, equation of continuity and three types of boundary conditions: the conditions of periodicity on the curves Γ_0 and Γ_1 , the Dirichlet boundary conditions on $\Gamma_{\rm in}$ and Γ_p and an artificial "do nothing"-type boundary condition on $\Gamma_{\rm out}$. (See Fig. 1.) We explain on the level of weak solutions the sense in which the last condition is satisfied. We show that, although domain Ω is not smooth and different types of boundary conditions meet in the corners of Ω , the considered problem has a strong solution with the so called maximum regularity property.

Keywords The Stokes problem · Artificial boundary condition · Maximum regularity property

Mathematics Subject Classification (2010) 35Q30 · 76D03 · 76D07

1 Introduction

The profile cascade and reduction to one spatial period. The flow through a 3D turbine wheel is often being modelled by a flow through a 2D profile cascade, which consists of an infinite number of profiles that periodically repeat with the period τe_2 in the x_2 -direction, see Fig. 1. Here, we use the planar Cartesian coordinate system x_1, x_2 . Unit vectors in the directions of the x_1 and x_2 axes are denoted by e_1 and e_2 , respectively. The profile cascade consists of an infinite family of profiles $\{P_k\}_{k\in\mathbb{Z}}$ such that P_k are closed bounded sets in the stripe $\mathbb{R}^2_{(0,d)}:=\{\mathbf{x}\equiv(x_1,x_2)\in\mathbb{R}^2;\ 0< x_1< d\}$, with Lipschitzian boundaries, such that $P_k=P_0+k\tau e_2$ and $P_k\cap P_{k+1}=\emptyset$ for $k\in\mathbb{Z}$. As the set $\mathcal{O}:=\{(\mathbf{x}=(x_1,x_2)\in\mathbb{R}^2;\ x_1\in(0,d)\}\setminus\bigcup_{k=-\infty}^\infty P_k$, through which the fluid flows, is spatially periodic, it is natural to assume that, provided that the acting body force and the given boundary data are also spatially periodic with the same period τe_2 , the fluid flow is spatially periodic, too. This enables us to reduce the mathematical model of the flow through

Czech Technical University, Faculty of Mechanical Engineering, Department of Technical Mathematics, Karlovo nám. 13, 121 35 Praha 2, Czech Republic

E-mail: tomas.neustupa@fs.cvut.cz

the whole profile cascade to the flow through just one spatial period, which is denoted by Ω , see Fig. 1. This approach is used e.g. in papers [10]–[12] and [34]–[36], where the qualitative analysis of corresponding mathematical models is studied, and in papers [8], [22], [39], devoted to the numerical analysis of the models or corresponding numerical calculations.

Classical formulation of the problem in one spatial period. We assume that Ω is a Lipschitzian sub-domain of $\mathbb{R}^2_{(0,d)}$, such that its boundary consists of the line segment $\Gamma_{\rm in} \equiv A_0 A_1$ of length τ , the line segment $\Gamma_{\rm out} \equiv B_0 B_1$ of the same length τ , the closed




Fig. 1: The profile cascade with marked one spatial period

curve Γ_p (the boundary of profile P_0) and the curves Γ_0 , Γ_1 such that $\Gamma_1 = \Gamma_0 + \tau e_2$. (See Fig. 1.) As the curves Γ_0 and Γ_1 in fact represent artificial boundaries of Ω , chosen in \mathcal{O} , we may assume without loss of generality that both Γ_0 and Γ_1 are of the class C^2 .

The reduced mathematical problem consists of the equations

$$\partial_t \mathbf{u} - \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \mathbf{f} \tag{1}$$

$$\operatorname{div} \mathbf{u} = 0 \tag{2}$$

in the space-time cylinder $\Omega \times (0,T)$ (where T>0), completed by appropriate initial and boundary conditions. Here, $\mathbf{u}=(u_1,u_2)$ denotes the unknown velocity of the moving fluid, p denotes the unknown pressure, positive constant ν is the kinematic coefficient of viscosity and \mathbf{f} is the external body force. The density of the fluid (which is also supposed to be a positive constant) can be without loss of generality supposed to be equal to one. Equation (1) (the Navier–Stokes equation) expresses the conservation of momentum and equation (2) (the equation of continuity) expresses the conservation of mass.

We assume that the fluid flows into the cascade through the straight line $\gamma_{\rm in}$ (the x_2 -axis) and essentially leaves the cascade through the straight line $\gamma_{\rm out}$, whose equation is $x_1=d$. (By "essentially" we mean that possible reverse flows on $\gamma_{\rm out}$ are not excluded.) This is why we complete equations (1), (2) by the inhomogeneous Dirichlet boundary condition

$$\mathbf{u} = \mathbf{g} \quad \text{on } \Gamma_{\text{in}},$$
 (3)

the homogeneous Dirichlet boundary condition

$$\mathbf{u} = \mathbf{0} \quad \text{on } \Gamma_{\mathcal{D}} \tag{4}$$

and appropriate conditions on Γ_0 , Γ_1 and Γ_{out} . Due to the assumed spatial periodicity of the flow, it is reasonable to prescribe the boundary conditions of periodicity on Γ_0 and Γ_1 :

$$\mathbf{u}(x_1, x_2 + \tau) = \mathbf{u}(x_1, x_2)$$
 for $\mathbf{x} \equiv (x_1, x_2) \in \Gamma_0$, (5)

$$\frac{\partial \mathbf{u}}{\partial \mathbf{n}}(x_1, x_2 + \tau) = -\frac{\partial \mathbf{u}}{\partial \mathbf{n}}(x_1, x_2) \qquad \text{for } \mathbf{x} \equiv (x_1, x_2) \in \Gamma_0, \tag{6}$$

$$p(x_1, x_2 + \tau) = p(x_1, x_2)$$
 for $\mathbf{x} \equiv (x_1, x_2) \in \Gamma_0$. (7)

On $\Gamma_{\rm out}$, various authors use various artificial boundary conditions. One of the most popular ones is the condition

$$-\nu \frac{\partial \mathbf{u}}{\partial \mathbf{n}} + p\mathbf{n} = \mathbf{h}, \tag{8}$$

where **h** is a given vector–function on $\Gamma_{\rm out}$ and **n** denotes the unit outward normal vector, which is equal to \mathbf{e}_1 on $\Gamma_{\rm out}$. The boundary condition (8) (with $\mathbf{h} = \mathbf{0}$) is often called the "do nothing" condition, because it naturally follows from a weak formulation of the boundary–value problem, see e.g. [15] and [18].

On some previous related results. Since this condition does not enable one to control the amount of kinetic energy in Ω in the case of an possible backward flow on $\Gamma_{\rm out}$, many authors also use various modifications of condition (8). (See e.g. [4], [10], [11], [12], [34], [35].) The modified conditions enable one to derive a priori estimates of solutions and existence of weak solutions. In paper [10], the existence of a steady weak solution of the problem (1)–(8) was proven for "sufficiently small" velocity profile g on $\Gamma_{\rm in}$, while in [12] and [36], the function g can be arbitrarily large. The existence of a non-steady weak solution on an arbitrarily long time interval has been proven in [11]. In papers [26] and [27], the authors use the boundary condition (8) on an "outflow" part of the boundary for a flow in a channel, and they prove the existence of a weak solution for "small data". Possible backward flows on the "outflow" of the channel are controlled by means of additional conditions in [23], [24], [25], which consequently cause that the Navier–Stokes equations must be replaced by the Navier–Stokes variational inequalities.

There are no results in literature about the regularity up to the boundary of existing weak solutions. The question of higher regularity of a solution is closely connected with the so called $maximum\ regularity\ property$ of the associated steady Stokes problem, which we obtain from the Navier–Stokes problem if we neglect the derivative with respect to t and the nonlinear term. It consists of the equations

$$-\nu\Delta\mathbf{u} + \nabla p = \mathbf{f} \tag{9}$$

and (2) (in Ω), and the boundary conditions (3)–(8). The maximum regularity property roughly speaking means that the solution \mathbf{u} , respectively p, has by two, respectively one, spatial derivatives more than function f, and the derivatives are integrable with the same power as f. (See Theorem 2.) An analogous property of the steady Stokes problem is mostly known only in the case of a smooth domain Ω , see e.g. [37, Theorem I.2.2], [29, Theorem III.3], [13, Theorem IV.6.1] and [38, Theorem III.2.1.1] for the Stokes problem with the inhomogeneous Dirichlet boundary condition, [2], [5] for problems with the Navier-type boundary condition, [3], [6] for problems with Navier's boundary condition, [31] for the 2D Stokes problem with the Neumann boundary condition (i.e. prescribing the normal part of the stress tensor on the boundary) and [32] for the 2D Stokes problem, prescribing the normal component of velocity and the pressure on the boundary. Concerning the maximum regularity property of the Stokes problem in non-smooth domains, we can cite [17], [19] and [7], where the authors considered the Stokes problem in a 2D polygonal domain with the Dirichlet boundary condition. In paper [28], the authors studied the Stokes problem in a 2D channel D of a special geometry, considering the homogeneous Dirichlet boundary condition on the walls and the homogeneous condition (8) on the outflow, and proved that the velocity is in $\mathbf{W}^{2-\beta,2}(D)$ for certain $\beta \in (0,1)$, provided that $\mathbf{f} \in \mathbf{L}^2(D)$. (See [28, Theorem 2.1].)

On results of this paper. In this paper, we at first verify the existence of a weak solution u to the Stokes problem (9), (2)–(5) and we show that an appropriate pressure p can be chosen so that the pair (\mathbf{u},p) satisfies equations (9), (2) in the sense of distributions in Ω and the boundary condition (8) as an equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$. The boundary conditions (3)-(5) are satisfied in the usual sense of traces. (Theorem 1.) Then, for "smooth" input data, we prove the existence of a strong solution of the Stokes problem (9), (2)-(5) and its maximum regularity property. (Theorem 2.) This result cannot be simply deduced from the previous aforementioned papers, because our domain Ω is not smooth and we consider altogether three types of boundary conditions, two of whose "meet" at the corner points A_0, A_1, B_0 and B_1 of domain Ω . In order to prove the regularity "up to the boundary" in the neighborhood of Γ_{out} and Γ_p , we use the fact that the solution satisfies Dirichlet-type boundary conditions on $\Gamma_{\rm in}$ and Γ_p and we apply known results on the Stokes problem with Dirichlet's boundary condition. In the neighborhood of Γ_{out} , we use the fact that Γ_{out} is a part of a straight line and we apply the technique of the so called difference quotients, whose originality is usually attributed to L. Nirenberg and which is described e.g. in [1] and [16]. As to the curves Γ_0 and Γ_1 , we use the possibility of an appropriate extension of a solution in the x_2 -direction, which enables us to avoid problems in neighborhoods of the corner points A_0, A_1, B_0, B_1 and to study the regularity in the neighborhood of Γ_0 and Γ_1 as an interior

2 The weak steady Stokes problem

Notation. Recall that Ω is a domain in \mathbb{R}^2 , sketched on Fig. 1. Its boundary consists of the curves $\Gamma_{\rm in}$, $\Gamma_{\rm out}$, Γ_0 , Γ_1 and Γ_p , described in Section 1. We denote by $\mathbf{n}=(n_1,n_2)$ the outer normal vector field on $\partial\Omega$. Note that $\mathbf{n}=-\mathbf{e}_1$ on $\Gamma_{\rm in}$ and $\mathbf{n}=\mathbf{e}_1$ on $\Gamma_{\rm out}$.

- $\circ \Gamma_{\text{in}}^0$, respectively Γ_{out}^0 , denotes the open line segment without the end points A_0 , A_1 , respectively B_0 , B_1 . Similarly, Γ_0^0 , respectively Γ_1^0 denotes the curve Γ_0 , respectively Γ_1 , without the end points A_0 , B_0 , respectively A_1 , B_1 .
- \circ We denote vector functions and spaces of vector functions by boldface letters. Tensor functions are denoted e.g. by $\mathbb F$ or $\mathbb G$ and spaces of tensor functions are marked by a superscript 2×2 .
- We denote by $\|.\|_r$ the norm in $L^r(\Omega)$ or in $\mathbf{L}^r(\Omega)$ or in $L^r(\Omega)^{2\times 2}$. Similarly, $\|.\|_{r,s}$ is the norm in $W^{r,s}(\Omega)$ or in $\mathbf{W}^{r,s}(\Omega)$ or in $W^{r,s}(\Omega)^{2\times 2}$. The scalar product in $L^2(\Omega)$ or in $L^2(\Omega)$ or in $L^2(\Omega)^{2\times 2}$ is denoted by $(.,.)_2$.
- $\circ \mathcal{C}^{\infty}_{\sigma}(\overline{\Omega})$ denotes the linear space of infinitely differentiable divergence–free vector functions in $\overline{\Omega}$, whose support is disjoint with $\Gamma_{\text{in}} \cup \Gamma_p$ and that satisfy, together with all their derivatives (of all orders), the condition of periodicity (5). Note that each $\mathbf{w} \in \mathcal{C}^{\infty}_{\sigma}(\overline{\Omega})$ satisfies $\int_{\Gamma_{\text{out}}} \mathbf{w} \cdot \mathbf{n} \, dl = 0$.
- $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ is the closure of $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$ in $\mathbf{W}^{1,2}(\Omega)$. The space $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ can be characterized as a space of divergence–free vector functions $\mathbf{v} \in \mathbf{W}^{1,2}(\Omega)$, whose traces on $\Gamma_{\mathrm{in}} \cup \Gamma_{p}$ are equal to zero, the traces on Γ_{0} and Γ_{1} satisfy the condition of periodicity (5) and the

traces on Γ_{out} satisfy $\int_{\Gamma_{\mathrm{out}}} \mathbf{v} \cdot \mathbf{n} \; \mathrm{d}l = 0$. Note that as functions from $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ are equal to zero on $\Gamma_{\mathrm{in}} \cup \Gamma_p$ (in the sense of traces) and domain Ω is bounded, the norm in $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ is equivalent to $\|\nabla \cdot\|_2$.

- $\bullet \ \ \text{We denote by } \mathbf{W}_0^{-1,2}(\varOmega) \ \text{the dual space to} \ \mathbf{W}^{1,2}(\varOmega), \text{by } \mathbf{W}^{-1,2}(\varOmega) \ \text{the dual space to} \\ \mathbf{W}_0^{1,2}(\varOmega), \text{ and by } \| \, . \, \|_{\mathbf{W}_0^{-1,2}} \ \text{and} \ \| \, . \, \|_{\mathbf{W}^{-1,2}} \ \text{the corresponding norms}.$
- $\circ \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ is the dual space to $\mathbf{V}_{\sigma}^{1,2}(\Omega)$. The duality pairing between $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ and $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ is denoted by $\langle \, . \, , \, . \, \rangle_{\sigma}$. The norm in $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ is denoted by $\| \, . \, \|_{\mathbf{V}_{\sigma}^{-1,2}}$.
- \circ Denote by \mathcal{A} the linear mapping of $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ to $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$, defined by the equation

$$\langle \mathcal{A}\mathbf{v}, \mathbf{w} \rangle_{\sigma} := \int_{\Omega} \nabla \mathbf{v} : \nabla \mathbf{w} \, d\mathbf{x} \quad \text{for } \mathbf{v}, \mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega).$$

 \circ c denotes a generic constant, i.e. a constant whose values may change throughout the text.

Lemma 1 Operator \mathcal{A} is a one-to-one closed bounded operator from $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ to the dual space $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ with the domain $D(\mathcal{A}) = \mathbf{V}_{\sigma}^{1,2}(\Omega)$ and range $R(\mathcal{A}) = \mathbf{V}_{\sigma}^{-1,2}(\Omega)$. The inverse operator \mathcal{A}^{-1} is bounded, as an operator from $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ to $\mathbf{V}_{\sigma}^{1,2}(\Omega)$.

Proof Denote by N(A) the null space of A. Let $\mathbf{v} \in N(A)$. Then

$$\langle \mathcal{A}\mathbf{v}, \mathbf{w} \rangle_{\sigma} = (\nabla \mathbf{v}, \nabla \mathbf{w})_2 = 0$$

for all $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$. The choice $\mathbf{w} = \mathbf{v}$ yields $(\nabla \mathbf{v}, \nabla \mathbf{v})_2 = \|\nabla \mathbf{v}\|_2^2 = 0$. This (together with the boundary conditions on $\Gamma_{\mathrm{in}} \cup \Gamma_p$) implies that $\mathbf{v} = \mathbf{0}$. Thus, operator \mathcal{A} is injective. The boundedness of \mathcal{A} can be proven in this way::

$$\|\mathcal{A}\mathbf{v}\|_{\mathbf{V}_{\sigma}^{-1,2}} = \sup_{\substack{\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega), \\ \mathbf{w} \neq \mathbf{0}}} \frac{|\langle \mathcal{A}\mathbf{v}, \mathbf{w} \rangle_{\sigma}|}{\|\mathbf{w}\|_{1,2}} = \sup_{\substack{\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega), \\ \mathbf{w} \neq \mathbf{0}}} \frac{|(\nabla \mathbf{v}, \nabla \mathbf{w})_{2}|}{\|\mathbf{w}\|_{1,2}} \leq c \|\nabla \mathbf{v}\|_{2}.$$

The equality $D(\mathcal{A})=\mathbf{V}_{\sigma}^{1,2}(\Omega)$ follows from the definition of \mathcal{A} . The equality $R(\mathcal{A})=\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ follows from Riesz' theorem and the equivalence of the scalar products $(\cdot,\cdot)_{1,2}$ and $(\nabla,\nabla)_2$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega)$: if $\mathbf{f}\in\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ then there exists $\mathbf{v}\in\mathbf{V}_{\sigma}^{1,2}(\Omega)$ such that $\langle\mathbf{f},\mathbf{w}\rangle_{\sigma}=(\nabla\mathbf{v},\nabla\mathbf{w})_2$ for all $\mathbf{w}\in\mathbf{V}_{\sigma}^{1,2}(\Omega)$. Hence $\mathbf{f}=\mathcal{A}\mathbf{v}$.

Operator \mathcal{A} is closed, as a bounded linear operator, defined on the whole space $\mathbf{V}_{\sigma}^{1,2}(\Omega)$. Hence the inverse operator \mathcal{A}^{-1} is also closed. As a closed linear operator, defined on the whole space $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$, \mathcal{A}^{-1} is bounded from $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$ to $\mathbf{V}_{\sigma}^{1,2}(\Omega)$.

Assume that $\mathbb{F} \in L^2(\Omega)^{2 \times 2}$. Define a bounded linear functional $\mathbf{F} \in \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ by the formula

$$\langle \mathbf{F}, \mathbf{w} \rangle_{\sigma} := -\int_{\Omega} \mathbb{F} : \nabla \mathbf{w} \, d\mathbf{x} \quad \text{for all } \mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega).$$
 (10)

The next lemma comes from [12, Sec. 3].

Lemma 2 Assume that $\mathbf{g} \in \mathbf{W}^{1/2,2}(\Gamma_{\mathrm{in}})$ is a given function on Γ_{in} , such that it can be extended from Γ_{in} to γ_{in} as a function τ -periodic function from $\mathbf{W}^{1/2,2}_{loc}(\gamma_{\mathrm{in}})$. Then there exists a divergence-free extension \mathbf{g}_* of \mathbf{g} from Γ_{in} to Ω , such that $\mathbf{g}_* \in \mathbf{W}^{1,2}(\Omega)$,

- a) $\|\mathbf{g}_*\|_{1,2} \le c \|\mathbf{g}\|_{1/2,2; \Gamma_{in}}$ (where c is independent of \mathbf{g} and \mathbf{g}_*),
- b) \mathbf{g}_* satisfies the condition of periodicity (5) on $\Gamma_0 \cup \Gamma_1$,
- c) $\mathbf{g}_* = (\Phi/\tau) \mathbf{e}_1$ in a neighborhood of Γ_{out} , where $\Phi = \int_{\Gamma_{\text{in}}} \mathbf{g} \cdot \mathbf{e}_1 dl$,
- d) $\mathbf{g}_* = \mathbf{0}$ on Γ_p in the sense of traces.

Note that the function \mathbf{g} is assumed to be in $\mathbf{W}^{s,2}(\varGamma_{\mathrm{in}})$ for some $s>\frac{1}{2}$ and to satisfy the condition $\mathbf{g}(A_0)=\mathbf{g}(A_1)$ in [12]. However, this condition can be simply replaced by our assumption, i.e. that $\mathbf{g}\in\mathbf{W}^{1/2,2}(\varGamma_{\mathrm{in}})$ has a τ -periodic extension in $\mathbf{W}^{1/2,2}_{loc}(\gamma_{\mathrm{in}})$, with no affect on the proof in [12].

Let g_* be the function, provided by Lemma 2. Define a bounded linear functional G on $V^{1,2}_\sigma(\Omega)$ by the formula

$$\langle \mathbf{G}, \mathbf{w} \rangle_{\sigma} := -\int_{\Omega} \nabla \mathbf{g}_* : \nabla \mathbf{w} \, d\mathbf{x} \quad \text{for all } \mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega).$$
 (11)

Theorem 1 (on a weak solution of the Stokes problem (2)–(5), (8), (9))) Let \mathbf{F} , $\mathbf{G} \in \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ be defined by formulas (10) and (11), respectively. Then the equation $\nu A \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ has a unique solution $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$. Moreover, there exists $p \in L^2(\Omega)$ such that the functions $\mathbf{u} := \mathbf{g}_* + \mathbf{v}$, which is divergence–free, and p satisfy the equation

$$-\nu\Delta\mathbf{u} + \nabla p = \operatorname{div}\mathbb{F} \tag{12}$$

in the sense of distributions in Ω , the boundary condition

$$(-\nu\nabla\mathbf{u} + p\mathbb{I} - \mathbb{F}) \cdot \mathbf{n} = \mathbf{0} \tag{13}$$

as an equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$ and the estimate

$$||p||_2 \le c_1 (||\nabla \mathbf{u}||_2 + ||\mathbb{F}||_2),$$
 (14)

where $c_1 = c_1(\Omega, \nu)$.

Proof The existence and uniqueness of the solution \mathbf{v} of the equation $\nu A \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ follows from Lemma 1.

Denote by $\mathcal{C}_0^\infty(\Omega)$ the space of all infinitely differentiable vector functions in Ω with a compact support in Ω and put $\mathcal{C}_{0,\sigma}^\infty(\Omega) := \mathcal{C}_\sigma^\infty(\overline{\Omega}) \cap \mathcal{C}_0^\infty(\Omega)$. Suppose at first that $\mathbf{w} \in \mathcal{C}_{0,\sigma}^\infty(\Omega)$. Then $\langle A\mathbf{v}, \mathbf{w} \rangle_\sigma = -\langle\!\langle \Delta\mathbf{v}, \mathbf{w} \rangle\!\rangle$ for all $\mathbf{w} \in \mathcal{C}_{0,\sigma}^\infty(\Omega)$, where $\langle\!\langle \cdot, \cdot \rangle\!\rangle$ denotes the pairing between a distribution in Ω and a function from $\mathcal{C}_0^\infty(\Omega)$. Similarly, we can write $\langle \mathbf{F}, \mathbf{w} \rangle_\sigma = \langle\!\langle \operatorname{div} \mathbb{F}, \mathbf{w} \rangle\!\rangle$ and $\langle \mathbf{G}, \mathbf{w} \rangle_\sigma = \langle\!\langle \Delta\mathbf{g}_*, \mathbf{w} \rangle\!\rangle$. Then the equation $\langle \nu A\mathbf{v} - \mathbf{F} - \nu \mathbf{G}, \mathbf{w} \rangle_\sigma = 0$ and the identity $\mathbf{u} = \mathbf{g}_* + \mathbf{v}$ imply that

$$\langle\!\langle \nu \Delta \mathbf{u} + \operatorname{div} \mathbb{F}, \mathbf{w} \rangle\!\rangle = 0$$
 for all $\mathbf{w} \in \mathcal{C}_{0,\sigma}^{\infty}(\Omega)$.

Thus, $\nu \Delta \mathbf{u} + \operatorname{div} \mathbb{F}$ is a distribution that vanishes on all divergence–free functions $\mathbf{w} \in \mathcal{C}_0^{\infty}(\Omega)$. Due De Rham's lemma (see [37, p. 14]), there exists a distribution p_0 in Ω , such that

$$\nu \Delta \mathbf{u} + \operatorname{div} \mathbb{F} = \nabla p_0 \tag{15}$$

holds in Ω in the sense of distributions. As both $\nu\Delta \mathbf{u}$ and $\operatorname{div}\mathbb{F}$ can also be naturally identified with bounded linear functionals on $\mathbf{W}_0^{1,2}(\Omega)$, i.e. elements of the dual space $\mathbf{W}^{-1,2}(\Omega)$, ∇p_0 belongs to $\mathbf{W}^{-1,2}(\Omega)$, too. Applying [37, Proposition I.1.2], we deduce that $p_0 \in L^2(\Omega)$, it can be chosen so that $\int_{\Omega} p_0 \, \mathrm{d}\mathbf{x} = 0$, and

$$||p_0||_2 \le c ||\nu \Delta \mathbf{u} + \operatorname{div} \mathbb{F}||_{\mathbf{W}^{-1,2}}, \tag{16}$$

where c depends only on ν and Ω .

Since $-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F} \in L^2(\Omega)^{2\times 2}$ and $\operatorname{div}(-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F}) = \mathbf{0}$, we can apply [13, Theorem III.2.2] and deduce that $(-\nu \nabla \mathbf{u} + p_0 \mathbb{I} - \mathbb{F}) \cdot \mathbf{n} \in \mathbf{W}^{-1/2,2}(\Gamma_{\text{out}})$.

Let $\mathbf{w} \in \mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$. The equations (15) and $\nu \mathcal{A} \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$, the formula $\mathbf{u} = \mathbf{v} + \mathbf{g}_*$ and the generalized Gauss identity (see [13, p. 160]) imply that

$$\begin{split} 0 &= \left(\operatorname{div} \left[\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I} \right], \mathbf{w} \right)_2 \\ &= \left\langle \left(\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I} \right) \cdot \mathbf{n}, \mathbf{w} \right\rangle_{\Gamma_{\text{out}}} - \int_{\varOmega} \left[\nu \nabla \mathbf{u} + \mathbb{F} \right] : \nabla \mathbf{w} \, d\mathbf{x} - \int_{\varOmega} p_0 \, \operatorname{div} \mathbf{w} \, d\mathbf{x} \\ &= \left\langle \left(\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I} \right) \cdot \mathbf{n}, \mathbf{w} \right\rangle_{\Gamma_{\text{out}}} - \left\langle \nu \mathcal{A} \mathbf{v}, \mathbf{w} \right\rangle_{\sigma} + \left\langle \nu \mathbf{G}, \mathbf{w} \right\rangle_{\sigma} + \left\langle \mathbf{F}, \mathbf{w} \right\rangle_{\sigma} \\ &= \left\langle \left(\nu \nabla \mathbf{u} + \mathbb{F} - p_0 \mathbb{I} \right) \cdot \mathbf{n}, \mathbf{w} \right\rangle_{\Gamma_{\text{out}}}. \end{split}$$

It can be deduced e.g. from [12, Sec. 3] that the set of traces of all functions from $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$ on Γ_{out} is dense in the set of all functions $\mathbf{w} \in \mathbf{W}^{1/2,2}(\Gamma_{\mathrm{out}})$, such that $\int_{\Gamma_{\mathrm{out}}} \mathbf{w} \cdot \mathbf{n} \ \mathrm{d}l = 0$. Hence there exists $c_2 \in \mathbb{R}$ such that \mathbf{u} and p satisfy

$$(\nu \nabla \mathbf{u} - p_0 \mathbb{I} + \mathbb{F}) \cdot \mathbf{n} = c_2 \mathbf{n},$$

as an equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$. Put $p:=p_0+c_2$. Then \mathbf{u} and p satisfy the boundary condition (13) as an equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$. It follows from (15) that \mathbf{u} , p satisfy equation (12) in the sense of distributions in Ω . Finally, (16) implies that estimate (14) holds, too.

Function \mathbf{u} represents a weak solution of the Stokes problem (9), (2)–(5), where $\mathbf{f} = \operatorname{div} \mathbb{F}$, with the boundary condition (13) on Γ_{out} .

The next lemma follows from [14, Theorem 2.5]. It shows that it is not a loss of generality if we write the right hand side of equation (12) in the form $\operatorname{div} \mathbb{F}$ instead of just \mathbf{f} . On the other hand, considering the right hand side of (12) in the form $\operatorname{div} \mathbb{F}$ enables us to deduce that \mathbf{v} and p satisfy (13), as an equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$. An analogue, having just $\mathbf{f} \in \mathbf{W}^{-1,2}(\Omega)$ instead of $\operatorname{div} \mathbb{F}$, would not be possible.

Lemma 3 Let $\mathbf{f} \in \mathbf{W}_0^{-1,2}(\Omega)$. Then there exists $\mathbb{F} \in L^2(\Omega)^{2 \times 2}$, satisfying $\operatorname{div} \mathbb{F} = \mathbf{f}$ in the sense of distributions in Ω and

$$\|\mathbb{F}\|_{2} \le c \|\mathbf{f}\|_{\mathbf{W}_{0}^{-1,2}},$$
 (17)

where c is independent of \mathbf{f} and \mathbb{F} .

3 The strong steady Stokes problem

Notation. In section 3, we also use this notation:

 \circ $\mathbf{L}_{\sigma}^{2}(\Omega)$ is the closure of $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$ in $\mathbf{L}^{2}(\Omega)$. Functions from $\mathbf{L}_{\sigma}^{2}(\Omega)$ are divergence–free in the sense of distributions in Ω and their normal components (in the sense of traces) belong to the space $W^{-1/2,2}(\partial\Omega)$ (the dual to $W^{1/2,2}(\partial\Omega)$, see [13, Theorem III.2.2]). Moreover, $\mathbf{v} \cdot \mathbf{n} = \mathbf{0}$ holds as an equality in $W^{-1/2,2}(\Gamma_{\mathrm{in}} \cup \Gamma_{p})$,

$$\mathbf{v}(x_1, x_2 + \tau) \cdot \mathbf{n}(x_1, x_2 + \tau) = -\mathbf{v}(x_1, x_2) \cdot \mathbf{n}(x_1, x_2)$$

holds as an equality in $W^{-1/2,2}(\varGamma_0)$ and $\langle \mathbf{v}\cdot\mathbf{n},1\rangle_{\varGamma_{\mathrm{out}}}=0$, where $\langle\,.\,,\,.\,\rangle_{\varGamma_{\mathrm{out}}}$ denotes the duality pairing between $W^{-1/2,2}(\varGamma_{\mathrm{out}})$ and $W^{1/2,2}(\varGamma_{\mathrm{out}})$.

 $\circ \mathbb{R}^2_{d-}$ denotes the half-plane $x_1 < d$. Recall that $\mathbb{R}^2_{(0,d)} := \{ \mathbf{x} \equiv (x_1, x_2) \in \mathbb{R}^2; \ 0 < x_1 < d \}$ and $\mathcal{O} = \mathbb{R}^2_{(0,d)} \setminus \bigcup_{k=-\infty}^{\infty} P_k$.

 $\circ W^{k,2}_{\mathrm{per}}(\mathcal{O})$ (for $k \in \mathbb{N}$) denotes the space of functions from $W^{k,2}_{loc}(\mathcal{O})$, τ -periodic in variable x_2 .

- $\circ W^{k,2}_{\mathrm{per}}(\Omega)$ is the space of functions, that can be extended from Ω to \mathcal{O} as functions in $W^{k,2}_{\mathrm{per}}(\mathcal{O})$. (Obviously, the traces of these functions on Γ_0 and Γ_1 satisfy the condition of periodicity, analogous to (5).)
- $\circ \ W^{k-1/2,2}_{per}(\gamma_{\mathrm{out}}) \ (\text{for} \ k \in \mathbb{N}) \ \text{is the space of} \ \tau\text{-periodic functions in} \ W^{k-1/2,2}_{loc}(\gamma_{\mathrm{out}}).$
- o $W_{\rm per}^{k-1/2,2}(\varGamma_{\rm out})$ is the space of functions from $W^{k-1/2,2}(\varGamma_{\rm out})$, that can be extended from $\varGamma_{\rm out}$ to $\gamma_{\rm out}$ as functions in $W_{\rm per}^{k-1/2,2}(\gamma_{\rm out})$. The space $W_{\rm per}^{k-1/2,2}(\varGamma_{\rm in})$ is defined by analogy.
- \circ Spaces of corresponding vector functions are again denoted by boldface letters and spaces of corresponding tensor functions are marked by the superscript 2×2 .

Lemma 4 Let $n \in \mathbb{N}$ and the functions $g_0 \in W_{\mathrm{per}}^{n+1/2,2}(\Gamma_{\mathrm{out}})$, $g_1 \in W_{\mathrm{per}}^{n-1/2,2}(\Gamma_{\mathrm{out}})$, ..., $g_n \in W_{\mathrm{per}}^{1/2,2}(\Gamma_{\mathrm{out}})$ be given. Let $\delta > 0$ be so small that the profile P_0 (see Fig. 1) is on the left from the straight line $x_1 = d - \delta$. Then there exists $g_* \in W_{\mathrm{per}}^{n+1,2}(\Omega)$, such that

$$g_* = g_0, \quad \partial_1 g_* = g_1, \quad \dots, \quad \partial_1^n g_* = g_n \quad \text{on } \Gamma_{\text{out}}$$

in the sense of traces, supp $g_* = \{\mathbf{x} = (x_1, x_2) \in \Omega; d - \delta \le x_1 \le d\}$ and

$$||g_*||_{n+1,2} \le c \left(||g_0||_{n+1/2,2; \Gamma_{\text{out}}} + ||g_1||_{n-1/2,2; \Gamma_{\text{out}}} + \dots + ||g_n||_{1/2,2; \Gamma_{\text{out}}} \right),$$
 (18) where $c = c(\Omega, n, \delta)$.

Principles of the proof The functions g_0, g_1, \ldots, g_n can be extended from Γ_{out} to γ_{out} so that the extended functions (which we again denote by g_0, g_1, \ldots, g_n) are in the spaces $W^{n+1/2}_{\text{per}}(\gamma_{\text{out}}), W^{n-1/2,2}_{\text{per}}(\gamma_{\text{out}}), \ldots, W^{1/2,2}_{\text{per}}(\gamma_{\text{out}})$, respectively. Then we apply a variant of Theorem II.4.4 in [13], which enables us to deduce that there exists a function $\psi \in W^{n+1,2}_{loc}(\mathbb{R}^2_{d-}), \tau$ -periodic in variable x_2 , such that

$$\psi = g_0, \quad \partial_1 \psi = g_1, \quad \dots, \quad \partial_1^n \psi = g_n \quad \text{on } \Gamma_{\text{out}}$$

in the sense of traces and satisfying estimate (18). Note that Theorem II.4.4 from [13] in fact deals with functions $(g_0,g_1,\ldots,g_n)\in W^{n+1/2,2}(\gamma_{\rm out})\times W^{n-1/2,2}(\gamma_{\rm out})\times \ldots\times W^{1/2,2}(\gamma_{\rm out})$ and the extension ψ is in the space $W^{n+1,2}(\mathbb{R}^2_d)$. However, the proof (which is based on [33, Chap. 2, Theorems 5.5, 5.8]) can be modified so that the theorem can also be applied to $(g_0,g_1,\ldots,g_n)\in W^{n+1/2,2}_{\rm per}(\gamma_{\rm out})\times W^{n-1/2,2}_{\rm per}(\gamma_{\rm out})\times \cdots\times W^{1/2,2}_{\rm per}(\gamma_{\rm out})$ and the extension is in $W^{2,2}_{loc}(\mathbb{R}^2_d)$, τ -periodic in variable x_2 . Then multiplying ψ by an infinitely differentiable and τ -periodic in variable x_2 cut-off function η in \mathbb{R}^2_d , such that $\eta=1$ in some neighborhood of $\gamma_{\rm out}$ and $\eta=0$ in the neighborhood of $\gamma_{\rm in}$ and Γ_p , we obtain function g_* , whose restriction to Ω has the properties stated in the lemma.

Lemma 5 There exists a bounded bilinear operator $\mathcal{F}: \mathbf{L}^2(\Omega) \times \mathbf{W}^{1/2,2}_{per}(\Gamma_{out}) \to W^{1,2}_{per}(\Omega)^{2\times 2}$, such that if $\mathbf{f} \in \mathbf{L}^2(\Omega)$, $\mathbf{h} \in \mathbf{W}^{1/2,2}_{per}(\Gamma_{out})$ and $\mathbb{F} = \mathcal{F}(\mathbf{f}, \mathbf{h})$ then $\operatorname{div} \mathbb{F} = \mathbf{f}$ a.e. in Ω , $\mathbb{F} = \mathbb{O}$ (the zero tensor) on Γ_p and $\mathbb{F} \cdot \mathbf{n} = \mathbf{h}$ a.e. on Γ_{out} in the sense of traces.

Proof Denote by Ω_- the mirror image of Ω with respect to the line $x_1=0$. Thus, $\Omega_-:=\{(x_1,x_2)\in\mathbb{R}^2;\ (-x_1,x_2)\in\Omega\}$. Furthermore, put $\widetilde{\Omega}:=\Omega_-\cup \varGamma_{\mathrm{in}}^0\cup\Omega$. We will construct \mathbb{F} in the form $\mathbb{F}=\mathbb{F}_0+\mathbb{H}_0+\mathbb{H}_1+\mathbb{H}_2$, where the tensor functions $\mathbb{F}_0,\ldots,\mathbb{H}_2$ are described below.

1) Function \mathbb{F}_0 : Extend \mathbf{f} from Ω to $\widetilde{\Omega}$ so that the extended function (we denote it also by \mathbf{f}) is odd in variable x_1 . Then $\mathbf{f} \in \mathbf{L}^2(\widetilde{\Omega})$ and $\int_{\widetilde{\Omega}} \mathbf{f} \ d\mathbf{x} = \mathbf{0}$. Due to [13, Theorem III.3.3], there exists $\mathbb{F}_0 \in W_0^{1,2}(\widetilde{\Omega})^{2\times 2}$, such that $\operatorname{div} \mathbb{F}_0 = \mathbf{f}$ in $\widetilde{\Omega}$ and $\|\mathbb{F}_0\|_{1,2;\widetilde{\Omega}} \leq c \|\mathbf{f}\|_{2;\widetilde{\Omega}}$, where $c = c(\widetilde{\Omega})$. Hence we also have

$$\|\mathbb{F}_0\|_{1,2} \le c \|\mathbf{f}\|_2.$$
 (19)

- 2) Function \mathbb{H}_0 : Put $\overline{\mathbf{h}} \equiv (\overline{h}_1, \overline{h}_2) := \tau^{-1} \int_{\Gamma_{\mathrm{out}}} \mathbf{h} \, \mathrm{d}l$ and define $\mathbf{h}_0 \equiv (h_{01}, h_{02})^T := \mathbf{h} \overline{\mathbf{h}}$ on Γ_{out} . Naturally, the function \mathbf{h}_0 is in $\mathbf{W}^{1/2,2}_{\mathrm{per}}(\Gamma_{\mathrm{out}})$ and its advantage is that $\int_{\Gamma_{\mathrm{out}}} \mathbf{h}_0 \, \mathrm{d}l = \mathbf{0}$. We construct \mathbb{H}_0 so that its i-th row (for i = 1, 2) has the form $\nabla^{\perp} \psi_i$, where $\nabla^{\perp} = (\partial_2, -\partial_1)$ and ψ_i is an appropriate function from $W^{2,2}(\Omega)$, which is defined below, in two steps:
 - 2a) We define ψ_i at first on the line segment $\Gamma_{\rm out}$ by the formula

$$\psi_i(d, x_2) := \int_{b_{02}}^{x_2} h_{0i}(d, \vartheta) \, \mathrm{d}\vartheta,$$

where b_{02} is the x_2 -coordinate of point B_0 . (See Fig. 1.) Since $h_{0i} \in W^{1/2,2}_{per}(\Gamma_{\text{out}})$ and $\int_{b_{02}}^{b_{02}+\tau} h_{0i}(d,\vartheta) \, \mathrm{d}\vartheta = 0$, function ψ_i is in $W^{3/2,2}_{\text{per}}(\Gamma_{\text{out}})$. Obviously,

$$\partial_2 \psi_i = h_{0i}$$
 a.e. on Γ_{out} . (20)

2b) Applying Lemma 4, we deduce that there exists an extension of ψ_i from $\Gamma_{\rm out}$ to Ω (which we again denote by ψ_i), such that $\psi_i \in W^{2,2}_{\rm per}(\Omega)$, $\partial_1 \psi_i = 0$ on $\Gamma_{\rm out}$,

$$\|\psi_i\|_{2,2} \le c \|\psi_i\|_{3/2,2;\Gamma_{\text{out}}} \le c \|h_{0i}\|_{1/2,2;\Gamma_{\text{out}}}$$
 (21)

and ψ_i is supported in $\{\mathbf{x}=(x_1,x_2)\in\Omega;\ d-\delta\leq x_1\leq d\}$, where $\delta>0$ is so small that all points $\mathbf{x}=(x_1,x_2)$ on Γ_p satisfy $x_1< d-\delta$.

The scalar functions ψ_i (for i=1,2) satisfy (20). The vector functions $\nabla^{\perp}\psi_i$ are in $\mathbf{W}^{1,2}_{\mathrm{per}}(\Omega)$ and satisfy $\nabla^{\perp}\psi_i \cdot \mathbf{n} = h_{0i}$ a.e. on Γ_{out} . Since the i-the row in the tensor function \mathbb{H}_0 equals $\nabla^{\perp}\psi_i$, we have $\mathbb{H}_0 \in W^{1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$ and $\mathrm{div}\,\mathbb{H}_0 = \mathbf{0}$. Moreover, $\mathbb{H}_0 = \mathbb{O}$ on $\Gamma_{\mathrm{in}} \cup \Gamma_p$, $\mathbb{H}_0 \cdot \mathbf{n} = \mathbf{h}_0$ on Γ_{out} and

$$\|\mathbb{H}_0\|_{1,2} \le c \|\mathbf{h}_0\|_{1/2,2;\Gamma_{\text{out}}},$$
 (22)

where $c = c(\Omega)$.

- 3) Function \mathbb{H}_1 : Denote by $\overline{\mathbb{H}}$ the constant tensor with the entries $\overline{H}_{11} = \overline{h}_1$, $\overline{H}_{12} = 0$, $\overline{H}_{21} = \overline{h}_2$, $\overline{H}_{22} = 0$. Let ζ be an even infinitely differentiable function of one variable x_1 for $x_1 \in [-d,d]$, such that $\zeta(-d) = \zeta(d) = 1$ and $\mathrm{supp}\,\zeta \subset [-d,-d+\delta] \cup [d-\delta,d]$, where $\delta>0$ is so small that the profile P_0 lies on the left from the straight line $x_1=d-\delta$. Define $\mathbb{H}_1(x_1,x_2) := \zeta(x_1)\overline{\mathbb{H}}$. The tensor function \mathbb{H}_1 is in $W^{1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$ and satisfies $\mathbb{H}_1 \cdot \mathbf{n} = \overline{\mathbf{h}}$ on Γ_{out} , $\mathbb{H}_1 = \mathbb{O}$ on $\Gamma_{\mathrm{in}} \cup \Gamma_P$ and $\mathrm{div}\,\mathbb{H}_1 = \zeta'(x_1)\overline{\mathbf{h}}$ in $\widetilde{\Omega}$.
- 4) Function \mathbb{H}_2 : Since $\int_{\widetilde{\Omega}} \zeta'(x_1) \overline{\mathbf{h}} \, d\mathbf{x} = \mathbf{0}$, there exists (by [13, Theorem III.3.3]) a tensor function $\mathbb{H}_2 \in W_0^{1,2}(\widetilde{\Omega})^{2 \times 2}$, satisfying $\operatorname{div} \mathbb{H}_2 = -\zeta'(x_1) \overline{\mathbf{h}}$ in $\widetilde{\Omega}$. The restriction of \mathbb{H}_2 to Ω is in $W_{\operatorname{per}}^{1,2}(\Omega)^{2 \times 2}$ and \mathbb{H}_2 particularly satisfies $\mathbb{H}_2 \cdot \mathbf{n} = \mathbf{0}$ on $\Gamma_{\operatorname{out}}$.

Using the properties of \mathbb{F}_0 , \mathbb{H}_0 , \mathbb{H}_1 and \mathbb{H}_2 , we observe that $\mathbb{F} := \mathbb{F}_0 + \mathbb{H}_0 + \mathbb{H}_1 + \mathbb{H}_2$ has all the properties stated in the lemma. The whole procedure can be formalized so that the mapping $(\mathbf{f}, \mathbf{h}) \mapsto \mathbb{F}$ is a bilinear operator. One can simply derive from (19), (22) and the definition of \mathbb{H}_1 and \mathbb{H}_2 that this operator is bounded from $\mathbf{L}^2(\Omega) \times \mathbf{W}^{1/2,2}_{\mathrm{per}}(\Omega)$ to $W^{1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$.

The next lemma generalizes Lemma 2:

Lemma 6 Let $m \in \{0\} \cup \mathbb{N}$ and $\mathbf{g} \in \mathbf{W}^{m+1/2,2}_{per}(\Gamma_{in})$ be given. Then there exists a divergence–free extension $\mathbf{g}_* \in \mathbf{W}^{m+1,2}_{per}(\Omega)$ with the properties c) and d) from Lemma 2, such that

$$\|\mathbf{g}_*\|_{m+1,2} \le c \|\mathbf{g}\|_{m+1/2,2; \Gamma_m},$$
 (23)

where $c = c(\Omega, m)$.

Principles of the proof Let g_1 , g_2 be the components of ${\bf g}$. The function g_1 can be written in the form $g_1=g_1^0+\overline{g_1}$, where $\overline{g_1}:=\tau^{-1}\int_{\varGamma_{\rm in}}g_1\;{\rm d}l$. Then $\int_{\varGamma_{\rm in}}g_1^0\;{\rm d}l=0$.

Put $\psi(0,x_2):=\int_{a_{02}}^{x_2}g_1^0(0,\vartheta)\,\mathrm{d}\vartheta$ for $x_2\in(a_{02},a_{02}+\tau)$, where a_{02} is the x_2- component of point A_0 . (See Fig. 1.) Then $\psi\in W^{m+3/2,2}_{\mathrm{per}}(\varGamma_{\mathrm{in}})$. We may apply Lemma 4 (with \varGamma_{in} instead of \varGamma_{out} and n=m) to extend ψ from \varGamma_{in} to \varOmega so that the extended function ψ_* is in $W^{m+2,2}_{\mathrm{per}}(\varOmega)$ and

$$\partial_1 \psi = -g_2$$
 and $\partial_1^k \psi = 0$ $(k = 2, \dots, m+1)$

on $\gamma_{\rm in}$ in the sense of traces and

$$\|\psi\|_{m+2,2} \leq c \left(\|\psi\|_{m+3/2,2; \, \Gamma_{\text{in}}} + \|\partial_1\psi\|_{m+1/2,2; \, \Gamma_{\text{in}}} + \dots + \|\partial_1^{m+1}\psi\|_{1/2,2; \, \Gamma_{\text{in}}} \right)$$

$$\leq c \left(\|\partial_2\psi\|_{m+1/2,2; \, \Gamma_{\text{in}}} + \|\partial_1\psi\|_{m+1/2,2; \, \Gamma_{\text{in}}} \right)$$

$$= c \left(\|g_1^0\|_{m+1/2,2; \, \Gamma_{\text{in}}} + \|g_2\|_{m+1/2; \, \Gamma_{\text{in}}} \right).$$

Define $\mathbf{g}^0_* := \nabla^{\perp} \psi$. Then \mathbf{g}^0_* is a divergence–free function in Ω , belongs to $\mathbf{W}^{m+1,2}_{\mathrm{per}}(\Omega)$, satisfies the inequality $\|\mathbf{g}^0_*\|_{m+1,2} \leq c \|(g^0_1,g_2)\|_{m+1/2,2;\,\Gamma_{\mathrm{in}}}$ and its trace on Γ_{in} equals (g^0_1,g_2) . Put $\mathbf{g}_* := \mathbf{g}^0_* + (\overline{g_1},0)^T$.

Further steps, which modify function \mathbf{g}_* so that it also has the properties c) and d) from Lemma 2, can be made in the same way as in [12, Sec. 3].

Theorem 2 [on a strong solution of the Stokes problem (2)–(8), (9)] Let the curve Γ_p (which is the boundary of the profile) be of the class C^2 , $\mathbb{F} \in W^{1,2}_{per}(\Omega)^{2\times 2}$ and \mathbf{g} , \mathbf{g}_* be the functions from Lemma 6, corresponding to m=1. Let the functionals \mathbf{F} and \mathbf{G} be defined by formulas (10) and (11), respectively. Then

- (a) the unique solution \mathbf{v} of the equation $\nu A \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ belongs to $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{\mathrm{per}}^{2,2}(\Omega)$ and the associated pressure p is in $W_{\mathrm{per}}^{1,2}(\Omega)$,
- (b) the functions $\mathbf{u} := \mathbf{g}_* + \mathbf{v}$ and p satisfy equations (2) and (12) a.e. in Ω ,
- (c) \mathbf{u} , p satisfy the boundary conditions (3), (4) and

$$-\nu \frac{\partial \mathbf{u}}{\partial \mathbf{n}} + p\mathbf{n} = \mathbb{F} \cdot \mathbf{n}$$
 (24)

in the sense of traces on $\Gamma_{\rm in}$, Γ_p and $\Gamma_{\rm out}$, respectively,

(d) there exists a constant $c_3 = c_3(\nu, \Omega)$, such that

$$\|\mathbf{u}\|_{2,2} + \|\nabla p\|_2 \le c_3 (\|\mathbb{F}\|_{1,2} + \|\mathbf{g}_*\|_{2,2}).$$
 (25)

Note that the existence and uniqueness of the solution $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$ of the equation $\nu \mathcal{A} \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ follows from Lemma 2 or Theorem 1.

If $\mathbf{f} \in \mathbf{L}^2(\Omega)$ and $\mathbf{h} \in \mathbf{W}_{per}^{1/2,2}(\Omega)$ are given and $\mathbb F$ is the tensor function, provided by

If $\mathbf{f} \in \mathbf{L}^2(\Omega)$ and $\mathbf{h} \in \mathbf{W}_{per}^{1/2,2}(\Omega)$ are given and \mathbb{F} is the tensor function, provided by Lemma 7, then functions \mathbf{u} and p from Theorem 2 represent a strong solution to the Stokes problem (9), (2)–(8), where $\mathbf{f} = \operatorname{div} \mathbb{F}$ in equation (9).

The conclusions $\mathbf{u} \in \mathbf{W}^{2,2}(\Omega)$ and $p \in W^{1,2}(\Omega)$ of Theorem 2, together with inequality (25), represent the maximum regularity property of the studied problem.

Proof of Theorem 2 Put $\mathbf{h} \equiv (h_1, h_2)^T := \mathbb{F} \cdot \mathbf{n}$ on Γ_{out} . The tensor function \mathbb{F} can be written in the form $\mathbb{F} = \mathbb{F}_1 + \mathbb{F}_2$, where both \mathbb{F}_1 and \mathbb{F}_2 are in $W^{1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$, $\mathbb{F}_1 \cdot \mathbf{n} = h_1 \mathbf{e}_1$ on Γ_{out} and $\mathbb{F} \cdot \mathbf{n} = h_2 \mathbf{e}_2$ on Γ_{out} . Denote by \mathbf{F}_1 and \mathbf{F}_2 the functionals in $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$, related to \mathbb{F}_1 and \mathbb{F}_2 , respectively, through formula (10).

We split the proof of Theorem 2 to seven parts, where we successively prove

- 1) the implication (a) \Longrightarrow (b), (c),
- 2) the implication (a), (b), (c) \Longrightarrow (d),
- 3) the solvability of the equation $\nu A \mathbf{v} = \mathbf{f}$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$ for $\mathbf{f} \in \mathbf{L}_{\sigma}^{2}(\Omega)$,
- 4) the solvability of the equation $\nu \mathcal{A} \mathbf{v}_1 = \mathbf{F}_1$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$ and the inclusion $p_1 \in W^{1,2}(\Omega)$ for an associated pressure,
- 5) the solvability of the equation $\nu A \mathbf{v}_2 = \mathbf{F}_2$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$ and the inclusion $p_2 \in W^{1,2}(\Omega)$ for an associated pressure,
- 6) the solvability of the equation $\nu \mathcal{A} \mathbf{v}_3 = \nu \mathbf{G}$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$ and the inclusion $p_3 \in W^{1,2}(\Omega)$ for an associated pressure,
- 7) the validity of statement (a).

Note that this scheme does not create a logical circle, because we do not use the validity of statement (a) of Theorem 2 in parts 2) - 6). We use the validity of the implication (a) \Longrightarrow (b), (c), which is not the same as the validity of (a).

The most technical part of the proof is the derivation of inequality (25) in part 2). This inequality is further used in part 3) in order to show that a certain Stokes-type operator A is closed and its range is closed in $\mathbf{L}_{\sigma}^{2}(\Omega)$.

- 1) The implication (a) \Longrightarrow (b), (c). As equation (12) is satisfied in the sense of distributions in Ω (due to Theorem 1) and all terms in this equation are now in $\mathbf{L}^2(\Omega)$, the equation is satisfied a.e. in Ω . Clearly, \mathbf{u} also satisfies equation (2) a.e. in Ω and boundary conditions (3) and (4) in the sense of traces on Γ_{in} and Γ_p , respectively. Since \mathbf{u} and p satisfy the boundary condition (13) in the sense of equality in $\mathbf{W}^{-1/2,2}(\Gamma_{\mathrm{out}})$ (see Theorem 1) and all the functions $\nabla \mathbf{u}$, $p\mathbb{I}$ and \mathbb{F} have traces on Γ_{out} in $W^{1/2,2}(\Gamma_{\mathrm{out}})^{2\times 2}$, the boundary condition (13) holds on Γ_{out} in the sense of traces, too. It can now be written in the form (24).
- 2) The implication (a), (b), (c) \Longrightarrow (d). We split the proof of the implication to three lemmas, where we successively derive an inequality, analogous to (25), in the interior of Ω plus the neighborhood of Γ_p and $\Gamma_{\rm in}^0$ (Lemma 7), in the neighborhood of $\Gamma_{\rm out}^0$ (Lemma 8) and in the neighborhoods of Γ_0 and Γ_1 (Lemma 9).

Lemma 7 Let Ω' be sub-domain of Ω , such that $\overline{\Omega'} \subset \Omega \cup \Gamma_{in}^0 \cup \Gamma_{in}$. Then

$$\|\mathbf{v}\|_{2,2;\Omega'} + \|\nabla p\|_{2;\Omega'} \le c \left(\|\operatorname{div} \mathbb{F}\|_2 + \|\mathbf{g}_*\|_{2,2} + \|\mathbf{v}\|_{1,2}\right),$$
 (26)

where $c = c(\nu, \Omega, \Omega')$.

Proof Consider a C^2 sub-domain Ω'' of Ω , such that $\Omega' \subset \Omega''$, $\overline{\Omega''} \subset \Omega \cup \Gamma_{\mathrm{in}}^0 \cup \Gamma_p$ and the distance between $\partial \Omega'' \cap \Omega$ and $\partial \Omega' \cap \Omega$ is positive. Let η be an infinitely differentiable cut-off function in Ω such that supp $\eta \subset \overline{\Omega''}$ and $\eta = 1$ in Ω' . Put $\widetilde{\mathbf{v}} := \eta \mathbf{v}$ and $\widetilde{p} := \eta p$. Since \mathbf{v} , p satisfy (12) a.e. in Ω , the functions $\widetilde{\mathbf{v}}$, \widetilde{p} represent a strong solution of the problem

$$-\nu\Delta\widetilde{\mathbf{v}} + \nabla\widetilde{p} = \widetilde{\mathbf{f}} \qquad \text{in } \Omega'', \tag{27}$$

$$\operatorname{div} \widetilde{\mathbf{v}} = \widetilde{h} \qquad \qquad \operatorname{in} \Omega'', \qquad (28)$$

$$\widetilde{\mathbf{v}} = \mathbf{0}$$
 on $\partial \Omega''$, (29)

where

$$\widetilde{\mathbf{f}} := \eta \operatorname{div} \mathbb{F} - 2\nu \nabla \eta \cdot \nabla \mathbf{v} - \nu (\Delta \eta) \mathbf{v} - (\nabla \eta) p + \nu \eta \Delta \mathbf{g}_*$$
 and $\widetilde{h} := \nabla \eta \cdot \mathbf{v}$.

As $\operatorname{div} \mathbb{F} \in \mathbf{L}^2(\Omega)$, $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$ and $p \in L^2(\Omega)$ (satisfying (14)), we have $\widetilde{\mathbf{f}} \in \mathbf{L}^2(\Omega)$ and $\widetilde{h} \in W^{1,2}(\Omega)$. Moreover,

$$\|\widetilde{\mathbf{f}}\|_{2} \le c (\|\operatorname{div} \mathbb{F}\|_{2} + \|\mathbf{g}_{*}\|_{2,2} + \|\mathbf{v}\|_{1,2}),$$
 (30)

$$\|\widetilde{h}\|_{1,2} \le c \|\mathbf{v}\|_2 \le c \|\mathbf{v}\|_{1,2},$$
 (31)

where $c = c(\nu, \eta)$. Due to [37, Proposition I.2.3],

$$\|\widetilde{\mathbf{v}}\|_{2,2;\Omega''} + \|\nabla \widetilde{p}\|_{2;\Omega''} \le c \left(\|\widetilde{\mathbf{f}}\|_{2;\Omega''} + \|\widetilde{h}\|_{1,2;\Omega''} \right),$$

where $c = c(\nu, \Omega'')$. This inequality, together with (30) and (31), implies that \mathbf{v} and p satisfy (26).

Lemma 8 Let Γ'_{out} be a (closed) line segment on Γ^0_{out} . For $\rho > 0$, denote $\Omega' := \{ \mathbf{x} \in \mathbb{R}^2_{d-} ; \operatorname{dist}(\mathbf{x}; \Gamma'_{out}) < \rho \}$. Assume that ρ is so small that $\overline{\Omega'}$ is disjoint with Γ_0 , Γ_1 and Γ_p . Then \mathbf{v} and p satisfy estimate (26), where the constant c on the right hand side again depends only on ν , Ω and Ω' .

Proof The condition of smallness of ρ guarantees that $\overline{\Omega'} \subset \Omega \cup \Gamma^0_{\mathrm{out}}$.

Denote $U':=\{\mathbf{x}=(x_1,x_2)\in\mathbb{R}^2;\ \mathrm{dist}(\mathbf{x},\Gamma'_{\mathrm{out}})<\rho\}.$ (U') is the ρ -neighborhood of Γ'_{out} in \mathbb{R}^2 .) Let $\delta>0$. By analogy with U', denote by U'' the $(\rho+\delta)$ -neighborhood of Γ'_{out} in \mathbb{R}^2 . Denote further by U''_{d-} the intersection of U'' with the half-plane \mathbb{R}^2_{d-} and by U''_{d+} the intersection of U'' with the half-plane $\mathbb{R}^2_{d+}:=\{\mathbf{x}=(x_1,x_2)\in\mathbb{R}^2;\ x_1>d\}$. Assume that δ is so small that the distances between U'' and points B_0 , B_1 are positive and $U''_{d-}\subset\Omega$. (See Fig. 2.)

Let η be a C^{∞} -function in \mathbb{R}^2 , supported in $\overline{U''}$, such that $\eta=1$ in U' and η is symmetric with respect to the line $x_1=d$. (The last condition means that $\eta(d+\vartheta,x_2)=\eta(d-\vartheta,x_2)$ for all $\vartheta,\,x_2\in\mathbb{R}$.)




Fig. 2: The line segment $\Gamma'_{\rm out}$ and its neighborhoods U' and U''

Applying the results from [21], one can deduce that there exists a divergence-free extension \mathbf{v}'' of function \mathbf{v} from U''_{d-} to the whole set U'', such that $\mathbf{v}'' \in \mathbf{W}^{1,2}(U'')$ and $\|\mathbf{v}''\|_{1,2;U''} \leq c \|\mathbf{v}\|_{1,2;U''_{d-}}$, where c is independent of \mathbf{v} .

Since $\nabla \eta \cdot \mathbf{v}'' \in W_0^{1,2}(U'')$ and $\int_{U''} \nabla \eta \cdot \mathbf{v}'' \, d\mathbf{x} = 0$, there exists (by [13, Theorem III.3.3]) $\mathbf{v}_* \in \mathbf{W}_0^{2,2}(U'')$, such that $\operatorname{div} \mathbf{v}_* = \nabla \eta \cdot \mathbf{v}''$ in U'' and

$$\|\mathbf{v}_*\|_{2,2;\,U''} \, \leq \, c \, \|\nabla \eta \cdot \mathbf{v}''\|_{1,2;\,U''} \, \leq \, c \, \|\mathbf{v}''\|_{1,2;\,U''} \, \leq \, c \, \|\mathbf{v}\|_{1,2;\,U''_{d_-}},$$

where c is independent of \mathbf{v} . Extending \mathbf{v}_* by zero to $\Omega \setminus U''$, we have $\|\mathbf{v}_*\|_{2,2} \le c \|\mathbf{v}\|_{1,2}$. Put

$$\widetilde{\mathbf{v}} := \eta \mathbf{v}'' - \mathbf{v}_*, \qquad \widetilde{p} := \eta p.$$
 (32)

Function $\widetilde{\mathbf{v}}$ is divergence–free, belongs to $\mathbf{W}_0^{1,2}(U'')$ and satisfies the estimates

$$\|\widetilde{\mathbf{v}}\|_{1,2;U''} \le c (\|\mathbf{v}''\|_{1,2;U''} + \|\mathbf{v}_*\|_{1,2;U''}) \le c \|\mathbf{v}''\|_{1,2;U''} \le c \|\mathbf{v}\|_{1,2},$$

where c is independent of \mathbf{v} . The functions $\widetilde{\mathbf{v}}$, \widetilde{p} satisfy equation (27) a.e. in the half-plane \mathbb{R}^2_{d-} , where function $\widetilde{\mathbf{f}}$ now satisfies

$$\widetilde{\mathbf{f}} := \eta \operatorname{div} \mathbb{F} - 2\nu \nabla \eta \cdot \nabla \mathbf{v} - \nu (\Delta \eta) \mathbf{v} - (\nabla \eta) p + \nu \eta \Delta \mathbf{g}_* + \nu \Delta \mathbf{v}_* \quad \text{in } U_{d-1}''$$

and $\widetilde{\mathbf{f}} := \mathbf{0}$ in $\mathbb{R}^2_{d-} \setminus U''_{d-}$. Although this function differs from the function $\widetilde{\mathbf{f}}$ from the proof of Lemma 7, it satisfies the same estimate (30). Define

$$\widetilde{\mathbf{h}} := \eta \mathbb{F} \cdot \mathbf{n} + \nu \frac{\partial \mathbf{v}_*}{\partial \mathbf{n}}$$
 on Γ_{out} ,

where the right hand side is understood as a trace on Γ_{out} . The function $\tilde{\mathbf{h}}$ satisfies

$$\|\widetilde{\mathbf{h}}\|_{1/2,2;\,\Gamma_{\text{out}}} \leq c \|\mathbb{F}\|_{1/2,2;\,\Gamma_{\text{out}}} + \left\|\frac{\partial \mathbf{v}_*}{\partial \mathbf{n}}\right\|_{1/2,2;\,\Gamma_{\text{out}}} \leq c \|\mathbb{F}\|_{1,2} + c \|\mathbf{v}_*\|_{2,2;\,U''}$$

$$\leq c \|\mathbb{F}\|_{1,2} + c \|\mathbf{v}\|_{1,2;\,U''_{d}}. \tag{33}$$

Put $\widetilde{\mathbb{F}}:=\mathcal{F}(\widetilde{\mathbf{f}},\widetilde{\mathbf{h}})$, where \mathcal{F} is the operator from Lemma 5. Then $\widetilde{\mathbb{F}}\in W^{1,2}(\Omega)^{2\times 2}$, $\widetilde{\mathbf{f}}=\operatorname{div}\widetilde{\mathbb{F}}$ a.e. in Ω and $\widetilde{\mathbb{F}}\cdot\mathbf{n}=\widetilde{\mathbf{h}}$ a.e. on Γ_{out} . Moreover, due to (30) and (33),

$$\|\widetilde{\mathbb{F}}\|_{1,2} \le c \|\widetilde{\mathbf{f}}\|_{2} + c \|\widetilde{\mathbf{h}}\|_{1/2,2;T_{\text{out}}} \le c (\|\mathbb{F}\|_{1,2} + \|\mathbf{g}_{*}\|_{2,2} + \|\mathbf{v}\|_{1,2}). \tag{34}$$

Let the functional $\widetilde{\mathbf{F}} \in \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ be defined by the same formula as (10), where we only consider $\widetilde{\mathbb{F}}$ instead of \mathbb{F} . We claim that $\nu \mathcal{A}\widetilde{\mathbf{v}} = \widetilde{\mathbf{F}}$. Indeed, for any $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$, we have

$$\begin{split} \nu \, \langle \mathcal{A}\widetilde{\mathbf{v}}, \mathbf{w} \rangle_{\sigma} &= \nu \int_{\Omega} \nabla \widetilde{\mathbf{v}} : \nabla \mathbf{w} \, \mathrm{d}\mathbf{x} \, = \int_{\varGamma_{\mathrm{out}}} \nu \, \frac{\partial \widetilde{\mathbf{v}}}{\partial \mathbf{n}} \cdot \mathbf{w} \, \mathrm{d}l - \nu \int_{\Omega} \Delta \widetilde{\mathbf{v}} \cdot \mathbf{w} \, \mathrm{d}\mathbf{x} \\ &= \int_{\varGamma_{\mathrm{out}}} \nu \, \frac{\partial \widetilde{\mathbf{v}}}{\partial \mathbf{n}} \cdot \mathbf{w} \, \mathrm{d}l + \int_{\Omega} (-\nabla \widetilde{p} + \widetilde{\mathbf{f}}) \cdot \mathbf{w} \, \mathrm{d}\mathbf{x} \\ &= \int_{\varGamma_{\mathrm{out}}} \left[\nu \, \frac{\partial \widetilde{\mathbf{v}}}{\partial \mathbf{n}} - \widetilde{p} \, \mathbf{n} \right] \cdot \mathbf{w} \, \mathrm{d}l + \int_{\Omega} \mathrm{div} \, \widetilde{\mathbb{F}} \cdot \mathbf{w} \, \mathrm{d}\mathbf{x} \\ &= \int_{\varGamma_{\mathrm{out}}} \left[\eta \, \left(\nu \, \frac{\partial \mathbf{v}}{\partial \mathbf{n}} - p \, \mathbf{n} \right) - \nu \, \frac{\partial \mathbf{v}_*}{\partial \mathbf{n}} \right] \cdot \mathbf{w} \, \mathrm{d}l + \int_{\varGamma_{\mathrm{out}}} (\widetilde{\mathbb{F}} \cdot \mathbf{n}) \cdot \mathbf{w} \, \mathrm{d}l - \int_{\Omega} \widetilde{\mathbb{F}} : \nabla \mathbf{w} \, \mathrm{d}\mathbf{x} \\ &= \int_{\varGamma_{\mathrm{out}}} \left[-\eta \, \mathbb{F} \cdot \mathbf{n} - \nu \, \frac{\partial \mathbf{v}_*}{\partial \mathbf{n}} \right] \cdot \mathbf{w} \, \mathrm{d}l + \int_{\varGamma_{\mathrm{out}}} (\widetilde{\mathbb{F}} \cdot \mathbf{n}) \cdot \mathbf{w} \, \mathrm{d}l - \int_{\Omega} \widetilde{\mathbb{F}} : \nabla \mathbf{w} \, \mathrm{d}\mathbf{x} \\ &= -\int_{\varGamma_{\mathrm{out}}} \widetilde{\mathbf{h}} \cdot \mathbf{w} \, \mathrm{d}l + \int_{\varGamma_{\mathrm{out}}} (\widetilde{\mathbb{F}} \cdot \mathbf{n}) \cdot \mathbf{w} \, \mathrm{d}l - \int_{\Omega} \widetilde{\mathbb{F}} : \nabla \mathbf{w} \, \mathrm{d}\mathbf{x} \end{split}$$

$$= -\int_{\Omega} \widetilde{\mathbb{F}} : \nabla \mathbf{w} \, d\mathbf{x} = \langle \widetilde{\mathbf{F}}, \mathbf{w} \rangle_{\sigma}.$$

(We have used the identity $\nu \partial \mathbf{v}/\partial \mathbf{n} - p\mathbf{n} = -\mathbb{F} \cdot \mathbf{n}$ on $\Gamma_{\rm out}$, following from (24) and the fact that $\partial \mathbf{g}_*/\partial \mathbf{n} = \mathbf{0}$ on Γ_{out} .)

Let us summarize that we have constructed functions $\widetilde{\mathbf{v}}$, \widetilde{p} and $\widetilde{\mathbb{F}}$, such that $\widetilde{\mathbf{v}}$ satisfies the equation $\nu \mathcal{A}\widetilde{\mathbf{v}} = \widetilde{\mathbf{F}}$ and \widetilde{p} is an associated pressure. The functions $\widetilde{\mathbf{v}}$ and \widetilde{p} are supported in $\overline{U_{d-}''}$ and $\widetilde{\mathbf{v}}$, \widetilde{p} are related to \mathbf{v} , p through formulas (32).

Recall that $\widetilde{\mathbf{f}}$ is supported in $\overline{U''_{d-}}$ and $\widetilde{\mathbf{h}}$ is supported in $\overline{U''_{d-}} \cap \varGamma_{\mathrm{out}}$. For $\delta \in \mathbb{R}$, whose modulus is so small that $(x_1, x_2 + \delta) \in \varOmega$ for all $\mathbf{x} = (x_1, x_2) \in U''_{d-}$, denote

$$D_2^{\delta}\widetilde{\mathbf{f}}(x_1,x_2) := \frac{\widetilde{\mathbf{f}}(x_1,x_2+\delta) - \widetilde{\mathbf{f}}(x_1,x_2)}{\delta}, \quad D_2^{\delta}\widetilde{\mathbf{h}}(d,x_2) := \frac{\widetilde{\mathbf{h}}(d,x_2+\delta) - \widetilde{\mathbf{h}}(d,x_2)}{\delta}.$$

 $D_2^{\delta} \widetilde{\mathbf{f}}$ and $D_2^{\delta} \widetilde{\mathbf{h}}$ are the so called *difference quotients*, see [1], [16] and [38] for more details

regarding their properties and usage in studies of regularity of solutions of PDE's.

As $\widetilde{\mathbb{F}} \in W^{1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$ and $\widetilde{\mathbb{F}} = \mathbb{O}$ on Γ_p , it can be extended from Ω to $\mathbb{R}^2_{(0,d)}$ as a τ -periodic function in variable x_2 , lying in $W^{1,2}_{loc}(\mathbb{R}^2_{(0,d)})$ and being equal to \mathbb{O} in P_k (for all $k \in \mathbb{Z}$). Let us denote the extension again by $\widehat{\mathbb{F}}$ and define

$$D_2^{\delta}\widetilde{\mathbb{F}}(x_1, x_2) := \frac{\widetilde{\mathbb{F}}(x_1, x_2 + \delta) - \widetilde{\mathbb{F}}(x_1, x_2)}{\delta}.$$

Denote $\widetilde{\mathbb{F}}_{\delta}(x_1,x_2):=\delta^{-1}\int_0^{\delta}\widetilde{\mathbb{F}}(x_1,x_2+\vartheta)\;\mathrm{d}\vartheta.$ Then

$$D_2^{\delta}\widetilde{\mathbb{F}}(x_1, x_2) = \frac{1}{\delta} \int_0^{\delta} \partial_2 \widetilde{\mathbb{F}}(x_1, x_2 + \vartheta) d\vartheta = \partial_2 \widetilde{\mathbb{F}}_{\delta}(x_1, x_2).$$

Furthermore, using the τ -periodicity of the function $\widetilde{\mathbb{F}}_{\delta}$ in variable x_2 in $\mathbb{R}^2_{(0,d)}$, we get

$$\|\widetilde{\mathbb{F}}_{\delta}\|_{2}^{2} = \int_{\Omega} \left| \frac{1}{\delta} \int_{0}^{\delta} \widetilde{\mathbb{F}}(x_{1}, x_{2} + \vartheta) \, d\vartheta \right|^{2} d\mathbf{x}$$

$$= \int_{0}^{d} \int_{0}^{\tau} \left| \frac{1}{\delta} \int_{0}^{\delta} \widetilde{\mathbb{F}}(x_{1}, x_{2} + \vartheta) \, d\vartheta \right|^{2} dx_{2} dx_{1}$$

$$\leq \int_{0}^{d} \int_{0}^{\tau} \frac{1}{\delta} \int_{0}^{\delta} \left| \widetilde{\mathbb{F}}(x_{1}, x_{2} + \vartheta) \right|^{2} d\vartheta dx_{2} dx_{1}$$

$$= \int_{0}^{d} \frac{1}{\delta} \int_{0}^{\delta} \int_{0}^{\tau} \left| \widetilde{\mathbb{F}}(x_{1}, y_{2}) \right|^{2} dy_{2} d\vartheta dx_{1}$$

$$= \int_{0}^{d} \int_{0}^{\tau} \left| \widetilde{\mathbb{F}}(x_{1}, y_{2}) \right|^{2} dy_{2} dx_{1} = \int_{\Omega} \left| \widetilde{\mathbb{F}}(\mathbf{x}) \right|^{2} d\mathbf{x} = \|\widetilde{\mathbb{F}}\|_{2}^{2}.$$

We can similarly show that $\|\nabla \widetilde{\mathbb{F}}_{\delta}\|_{2}^{2} \leq \|\nabla \widetilde{\mathbb{F}}\|_{2}^{2}$. Consequently, $\|\widetilde{\mathbb{F}}_{\delta}\|_{1,2} \leq \|\widetilde{\mathbb{F}}\|_{1,2}$. Thus,

$$\|D_2^{\delta}\widetilde{\mathbb{F}}\|_2 = \|\partial_2\widetilde{\mathbb{F}}_{\delta}\|_2 \le \|\widetilde{\mathbb{F}}_{\delta}\|_{1,2} \le \|\widetilde{\mathbb{F}}\|_{1,2} \le c(\|\widetilde{\mathbf{f}}\|_2 + \|\widetilde{\mathbf{h}}\|_{1/2,2;\Gamma_{\text{out}}}).$$
(35)

Let $D_2^{\delta}\widetilde{\mathbf{v}}$ and $D_2^{\delta}\widetilde{p}$ be defined by analogy with $D_2^{\delta}\widetilde{\mathbf{f}}$ and $D_2^{\delta}\widetilde{\mathbb{F}}$. The functions $D_2^{\delta}\widetilde{\mathbf{v}}$, $D_2^{\delta}\widetilde{p}$ satisfy the equations

$$-\nu \Delta D_2^{\delta} \widetilde{\mathbf{v}} + \nabla D_2^{\delta} \widetilde{p} \; = \; \mathrm{div} \, D_2^{\delta} \widetilde{\mathbb{F}},$$

$$\operatorname{div} D_2^{\delta} \widetilde{\mathbf{v}} = 0$$

a.e. in Ω . Since

$$\nu\,\frac{\partial\widetilde{\mathbf{v}}}{\partial\mathbf{n}} - \widetilde{p}\,\mathbf{n} \;=\; \nu\,\left(\eta\,\frac{\partial\mathbf{v}}{\partial\mathbf{n}} - \frac{\partial\mathbf{v}_*}{\partial\mathbf{n}}\right) - \eta\,p\,\mathbf{n} \;=\; -\eta\,\mathbb{F}\cdot\mathbf{n} - \nu\,\frac{\partial\mathbf{v}_*}{\partial\mathbf{n}} \;=\; -\widetilde{\mathbf{h}}$$

on $\gamma_{\rm out}$, $D_2^{\delta}\widetilde{\mathbf{v}}$ and $D_2^{\delta}\widetilde{p}$ also satisfy the boundary condition

$$-\nu \frac{\partial D_2^{\delta} \widetilde{\mathbf{v}}}{\partial \mathbf{n}} + D_2^{\delta} \widetilde{p} \mathbf{n} = D_2^{\delta} \widetilde{\mathbf{h}}$$

on $\Gamma_{\rm out}$. From this, one can deduce that $\nu \mathcal{A} D_2^{\delta} \widetilde{\mathbf{v}} = \widetilde{\mathbf{F}}_{\delta}$. Here, the functional $\widetilde{\mathbf{F}}_{\delta}$, which is an element of $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$, is defined by the same formula as (10), where we only consider $D_2^{\delta} \widetilde{\mathbb{F}}$ instead of \mathbb{F} . It follows from Lemma 1 that

$$\|\nabla D_2^{\delta}\widetilde{\mathbf{v}}\|_2 \leq \|\widetilde{\mathbf{F}}_{\delta}\|_{\mathbf{V}_{\sigma}^{-1,2}}.$$

Since $\|\widetilde{\mathbf{F}}_{\delta}\|_{\mathbf{V}_{\sigma}^{-1,2}} \leq \|D_2^{\delta}\widetilde{\mathbb{F}}\|_2 \leq c \left(\|\widetilde{\mathbf{f}}\|_2 + \|\widetilde{\mathbf{h}}\|_{1/2,2; \Gamma_{\mathrm{out}}}\right)$, we obtain

$$\|\nabla D_2^{\delta}\widetilde{\mathbf{v}}\|_2 \le c \left(\|\widetilde{\mathbf{f}}\|_2 + \|\widetilde{\mathbf{h}}\|_{1/2,2;\Gamma_{\text{out}}}\right). \tag{36}$$

Applying further Theorem 1 (with $\mathbf{g}_* = \mathbf{0}$), (35) and (36), we obtain the estimate of $D_2^{\delta} \tilde{p}$:

$$\|D_2^{\delta}\widetilde{p}\|_2 \le c_1 \left(\|\nabla D_2^{\delta}\widetilde{\mathbf{v}}\|_2 + \|D_2^{\delta}\widetilde{\mathbb{F}}\|_2\right) \le c \left(\|\widetilde{\mathbf{f}}\|_2 + \|\widetilde{\mathbf{h}}\|_{1/2,2;T_{\text{out}}}\right). \tag{37}$$

As the right hand sides of (36) and (37) are independent of δ , we may let δ tend to 0 and we obtain

$$\|\nabla \partial_2 \widetilde{\mathbf{v}}\|_2 + \|\partial_2 \widetilde{p}\|_2 \le c \left(\|\widetilde{\mathbf{f}}\|_2 + \|\widetilde{\mathbf{h}}\|_{1/2,2; \Gamma_{\text{out}}}\right). \tag{38}$$

This shows that $\partial_1\partial_2\widetilde{v}_1$, $\partial_2^2\widetilde{v}_1$, $\partial_1\partial_2\widetilde{v}_2$, $\partial_2^2\widetilde{v}_2$ and $\partial_2\widetilde{p}$ are all in $L^2(\Omega)$ and their norms are less than or equal to the right hand side of (38). Consequently, as $\widetilde{\mathbf{v}}$ is divergence-free, the same statement also holds on $\partial_1^2\widetilde{v}_1$. Now, from equation (27) (considering just the first scalar component of this vectorial equation), we deduce that $\partial_1\widetilde{p}\in L^2(\Omega)$. Finally, considering the second scalar component in equation (27), we obtain $\partial_1^2\widetilde{v}_2\in L^2(\Omega)$, too. Thus, applying also (30) and (31), we obtain

$$\|\widetilde{\mathbf{v}}\|_{2,2} + \|\nabla \widetilde{p}\|_{2} \le c (\|\mathbb{F}\|_{1,2} + \|\mathbf{g}_{*}\|_{2,2} + \|\mathbf{v}\|_{1,2}).$$

This inequality, formulas (32),the estimate of $\|\mathbf{v}_*\|_{2,2}$ and the fact that $\eta=1$ on $\Omega'\equiv U'\cap\mathbb{R}^2_{d-}$ yield (26).

The next corollary is an immediate consequence of Lemmas 7 and 8.

Corollary 1 Let Ω' be a sub-domain of Ω , such that $\overline{\Omega'} \subset \Omega \cup \Gamma_{\rm in}^0 \cup \Gamma_p \cup \Gamma_{\rm out}^0$. Then \mathbf{v} and p satisfy estimate (26), where $c = c(\nu, \Omega, \Omega')$.

Lemma 9 Let Ω' be a sub-domain of Ω , such that $\overline{\Omega'} \cap \Gamma_p = \emptyset$ and $\Gamma_1 \subset \partial \Omega'$. Then \mathbf{v} and p satisfy estimate (26), where $c = c(\nu, \Omega, \Omega')$.

Proof Consider $\delta \in (0, \tau)$ and denote

$$A_0^{\delta} := A_0 + \delta \mathbf{e}_2, \quad A_1^{\delta} = A_1 + \delta \mathbf{e}_2, \quad B_0^{\delta} := B_0 + \delta \mathbf{e}_2, \quad B_1^{\delta} = B_1 + \delta \mathbf{e}_2,$$
$$\Gamma_{\text{in}}^{\delta} := \Gamma_{\text{in}} + \delta \mathbf{e}_2, \quad \Gamma_0^{\delta} = \Gamma_0 + \delta \mathbf{e}_2, \quad \Gamma_1^{\delta} := \Gamma_1 + \delta \mathbf{e}_2, \quad \Gamma_{\text{out}}^{\delta} = \Gamma_{\text{out}} + \delta \mathbf{e}_2,$$

where \mathbf{e}_2 is the unit vector in the direction of the x_2 -axis. Suppose that $\delta>0$ is so small that $\Omega'\cap \varGamma_0^\delta=\emptyset$ and the profile P_0 lies above $\varGamma_0^\delta,$ which means that $P_0\subset\{(x_1,y_2)\in\mathbb{R}^2;\ y_2>x_2\ \text{for}\ (x_1,x_2)\in\varGamma_0^\delta\}$. (Recall that $P_0=\overline{\mathrm{Int}\ \varGamma_p},$ see Fig. 1.) Denote by Ω^δ the domain bounded by the curves $\varGamma_{\mathrm{in}}^\delta, \varGamma_0^\delta, \varGamma_{\mathrm{out}}^\delta, \varGamma_1^\delta$ and \varGamma_p . Precisely,

$$\Omega^{\delta} := \{(x_1, y_2) \in \mathbb{R}^2; \ 0 < x_1 < d, \ x_2 < y_2 < x_2 + \tau \text{ for } (x_1, x_2) \in \Gamma_0^{\delta} \} \setminus P_0.$$

Denote by \mathbf{v}^{δ} the function, defined by the formulas

$$\mathbf{v}^{\delta}(x_1, x_2) := \begin{cases} \mathbf{v}(x_1, x_2) & \text{for } (x_1, x_2) \in \Omega^{\delta} \cap \Omega, \\ \mathbf{v}(x_1, x_2 - \tau) & \text{for } (x_1, x_2) \in \Omega^{\delta} \setminus \Omega. \end{cases}$$
(39)

By analogy, denote

$$\mathbb{F}^{\delta}(x_1, x_2) := \begin{cases}
\mathbb{F}(x_1, x_2) & \text{for } (x_1, x_2) \in \Omega^{\delta} \cap \Omega, \\
\mathbb{F}(x_1, x_2 - \tau) & \text{for } (x_1, x_2) \in \Omega^{\delta} \setminus \Omega,
\end{cases}$$

$$\mathbf{g}_*^{\delta}(x_1, x_2) := \begin{cases}
\mathbf{g}_*(x_1, x_2) & \text{for } (x_1, x_2) \in \Omega^{\delta} \cap \Omega, \\
\mathbf{g}_*(x_1, x_2 - \tau) & \text{for } (x_1, x_2) \in \Omega^{\delta} \setminus \Omega.
\end{cases}$$

Let the spaces $\mathbf{V}_{\sigma}^{1,2}(\Omega^{\delta})$ and $\mathbf{V}_{\sigma}^{-1,2}(\Omega^{\delta})$ be defined in the same way as $\mathbf{V}_{\sigma}^{1,2}(\Omega)$ and $\mathbf{V}_{\sigma}^{-1,2}(\Omega)$, respectively, and let operator \mathcal{A}^{δ} be defined in the same way as \mathcal{A} , with the only difference that it acts on functions from $\mathbf{V}_{\sigma}^{1,2}(\Omega^{\delta})$ to $\mathbf{V}_{\sigma}^{-1,2}(\Omega^{\delta})$. Obviously, $\mathbb{F}^{\delta} \in W^{1,2}(\Omega^{\delta})^{2\times 2}$ and $\|\mathbb{F}^{\delta}\|_{1,2;\,\Omega^{\delta}} = \|\mathbb{F}\|_{1,2}$. Similarly, the function \mathbf{g}_{*}^{δ} has the same norm and properties in Ω^{δ} as the function \mathbf{g}_{*} in Ω . Let the functionals \mathbf{F}^{δ} and \mathbf{G}^{δ} in the dual space $\mathbf{V}_{\sigma}^{-1,2}(\Omega^{\delta})$ be defined by analogous formulas as \mathbf{F} and \mathbf{G} .

Our next claim is to show that $\mathbf{v}^{\delta} \in \mathbf{V}_{\sigma}^{2}(\Omega^{\delta})$ and $\nu \mathcal{A}^{\delta} \mathbf{v}^{\delta} = \mathbf{F}^{\delta} + \nu \mathbf{G}^{\delta}$. Since $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$, there exists a sequence $\{\mathbf{v}_n\}$ in $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$, such that $\mathbf{v}_n \to \mathbf{v}$ in the norm of $\mathbf{W}^{1,2}(\Omega)$. Define

$$\mathbf{v}_n^{\delta}(x_1, x_2) := \begin{cases} \mathbf{v}_n(x_1, x_2) & \text{for } (x_1, x_2) \in \overline{\Omega^{\delta}} \cap \Omega, \\ \mathbf{v}_n(x_1, x_2 - \tau) & \text{for } (x_1, x_2) \in \overline{\Omega^{\delta}} \setminus \Omega. \end{cases}$$

Then $\mathbf{v}_n^{\delta} \in \mathcal{C}_{\sigma}^{\infty}(\overline{\Omega^{\delta}})$ and $\mathbf{v}_n^{\delta} \to \mathbf{v}^{\delta}$ in $\mathbf{W}^{1,2}(\Omega^{\delta})$. This confirms that $\mathbf{v}^{\delta} \in \mathbf{V}_{\sigma}^2(\Omega^{\delta})$. Furthermore, let $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$ and $\mathbf{w}^{\delta} \in \mathbf{V}_{\sigma}^{1,2}(\Omega^{\delta})$ be related in the same way as \mathbf{v} and \mathbf{v}^{δ} in (39). Then, denoting by $\langle \, . \, , \, . \, \rangle_{\sigma; \Omega^{\delta}}$ the duality pairing between $\mathbf{V}_{\sigma}^{-1,2}(\Omega^{\delta})$ and $\mathbf{V}_{\sigma}^{1,2}(\Omega^{\delta})$, we have

$$\begin{split} \langle \nu \mathcal{A}^{\delta} \mathbf{v}^{\delta}, \mathbf{w}^{\delta} \rangle_{\sigma; \, \Omega^{\delta}} &= \nu \, (\nabla \mathbf{v}^{\delta}, \nabla \mathbf{w}^{\delta})_{2; \, \Omega^{\delta}} \\ &= \nu \int_{\Omega^{\delta} \cap \, \Omega} \nabla \mathbf{v}^{\delta} : \nabla \mathbf{w}^{\delta} \, \, \mathrm{d}\mathbf{x} + \nu \int_{\Omega^{\delta} \smallsetminus \Omega} \nabla \mathbf{v}^{\delta} : \nabla \mathbf{w}^{\delta} \, \, \mathrm{d}\mathbf{x} \\ &= \nu \int_{\Omega^{\delta} \cap \, \Omega} \nabla \mathbf{v} : \nabla \mathbf{w} \, \, \mathrm{d}\mathbf{x} + \nu \int_{0}^{\delta} \int_{\Gamma_{1} + \vartheta \, \mathrm{e}_{2}} \nabla \mathbf{v}^{\delta} : \nabla \mathbf{w}^{\delta} \, \, \mathrm{d}l \, \, \mathrm{d}\vartheta \end{split}$$

$$= \nu \int_{\Omega^{\delta} \cap \Omega} \nabla \mathbf{v} : \nabla \mathbf{w} \, d\mathbf{x} + \nu \int_{0}^{\delta} \int_{\Gamma_{0} + \vartheta \, e_{2}} \nabla \mathbf{v} : \nabla \mathbf{w} \, dl \, d\vartheta$$

$$= \nu (\nabla \mathbf{v}, \nabla \mathbf{w})_{2} = \langle \nu A \mathbf{v}, \mathbf{w} \rangle_{\sigma} = \langle \mathbf{F}, \mathbf{w} \rangle_{\sigma} + \langle \mathbf{G}, \mathbf{w} \rangle_{\sigma}$$

$$= -\int_{\Omega} \mathbb{F} : \nabla \mathbf{w} \, d\mathbf{x} + \int_{\Omega} \nabla \mathbf{g}_{*} : \nabla \mathbf{w} \, d\mathbf{x}$$

$$= -\int_{\Omega^{\delta}} \mathbb{F}^{\delta} : \nabla \mathbf{w}^{\delta} \, d\mathbf{x} + \int_{\Omega^{\delta}} \nabla \mathbf{g}_{*}^{\delta} : \nabla \mathbf{w}^{\delta} \, d\mathbf{x}$$

$$= \langle \mathbf{F}^{\delta}, \mathbf{w}^{\delta} \rangle_{\sigma: \Omega^{\delta}} + \langle \mathbf{G}^{\delta}, \mathbf{w}^{\delta} \rangle_{\sigma: \Omega^{\delta}}.$$

This verifies that $\nu \mathcal{A}^{\delta} \mathbf{v}^{\delta} = \mathbf{F}^{\delta} + \nu \, \mathbf{G}^{\delta}$. Denote $(\Omega')^{\delta/2} := \Omega' \cup \{(x_1, y_2) \in \mathbb{R}^2; \ x_2 \leq y_2 < x_2 + \frac{1}{2}\delta \text{ for } (x_1, x_2) \in \varGamma_1\}$. Then $(\Omega')^{\delta/2}$ is a sub-domain of Ω^{δ} , such that $\varGamma_1^0 \subset (\Omega')^{\delta/2}$. The statements of Lemma 9 now follow from Corollary 1, applied to the equation $\nu \mathcal{A}^{\delta} \mathbf{v}^{\delta} = \mathbf{F}^{\delta} + \nu \, \mathbf{G}^{\delta}$ in domain Ω^{δ} , where we consider $(\Omega')^{\delta/2}$ instead of Ω' .

An analogue of Lemma 9 also holds if one considers Ω' , satisfying the condition $\Gamma_0 \subset$ $\partial\Omega'$ instead of $\Gamma_1\subset\partial\Omega'$. This, Corollary 1, Lemma 9 and Lemma 1 (which enables us to estimate $\|\mathbf{v}\|_{1,2}$ on the right hand side of (26)) now imply that (25) holds.

3) Solution of the equation $\nu A \mathbf{v} = \mathbf{f}$ for $\mathbf{f} \in \mathbf{L}^2_{\sigma}(\Omega)$. Denote by D(A) the set of functions $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{per}^{2,2}(\Omega)$, such that there exists $q \in W_{per}^{1/2,2}(\Gamma_{\mathrm{out}})$, satisfying $\partial \mathbf{v}/\partial \mathbf{n} =$ $q\mathbf{e}_1$ on Γ_{out} in the sense of an equality in $\mathbf{W}^{1/2,2}(\Gamma_{\mathrm{out}})$. The linear space $\widetilde{\mathcal{C}}_{\sigma}^{\infty}(\overline{\Omega}):=$ $\{\mathbf{w} \in \mathcal{C}_{\sigma}^{\infty}(\overline{\Omega}); \ \partial \mathbf{w}/\partial \mathbf{n} \perp \mathbf{e}_2 \text{ on } \Gamma_{\text{out}}\}$ contains D(A) as a dense subset. Since $\widetilde{\mathcal{C}}_{\sigma}^{\infty}(\overline{\Omega})$ is dense in $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$ in the L^2 -norm and $\mathbf{L}_{\sigma}^2(\Omega)$ is the closure of $\mathcal{C}_{\sigma}^{\infty}(\overline{\Omega})$ in $\mathbf{L}^2(\Omega)$, D(A) is dense in $\mathbf{L}_{\sigma}^2(\Omega)$. Put $A := \mathcal{A}|_{D(A)}$.

Let us at first show that R(A) (the range of A) is a subset of $\mathbf{L}^2_{\sigma}(\Omega)$. Thus, let $\mathbf{v} \in$ D(A) and q be a corresponding function in $W_{\rm per}^{1/2,2}(\Gamma_{\rm out})$. It follows from Lemma 4 that there exists an extension $q_* \in W^{1,2}_{\mathrm{per}}(\Omega)$ of q from Γ_{out} to Ω , which equals zero in the neighborhood of $\Gamma_{\rm in}$ and Γ_p and satisfies

$$||q_*||_{1,2} \le c ||q||_{1/2,2;\Gamma_{out}},$$
 (40)

where $c = c(\Omega)$. For any $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$, \mathbf{v} satisfies

$$\langle A\mathbf{v}, \mathbf{w} \rangle_{\sigma} = (\nabla \mathbf{v}, \nabla \mathbf{w})_{2} = \int_{\Gamma_{\text{out}}} \frac{\partial \mathbf{v}}{\partial \mathbf{n}} \cdot \mathbf{w} \, dl - (\Delta \mathbf{v}, \mathbf{w})_{2}$$
$$= \int_{\Gamma_{\text{out}}} q\mathbf{n} \cdot \mathbf{w} \, dl - (\Delta \mathbf{v}, \mathbf{w})_{2} = (\nabla q_{*} - \Delta \mathbf{v}, \mathbf{w})_{2}.$$

From this and the density of $V_{\sigma}^{1,2}(\Omega)$ in $L_{\sigma}^{2}(\Omega)$, we deduce that $A\mathbf{v}$ can be identified with a bounded linear functional on $L^2_{\sigma}(\Omega)$. Due to Riesz' theorem, it can be represented by an element of $L^2_{\sigma}(\Omega)$ (which is, in our case, the function $\nabla q_* - \Delta \mathbf{v}$). We have proven the inclusion $R(A) \subset \mathbf{L}^2_{\sigma}(\Omega)$.

Treating A as an operator in $\mathbf{L}_{\sigma}^{2}(\Omega)$, we easily verify that A is symmetric: let $\mathbf{v}^{(1)}$, $\mathbf{v}^{(2)} \in D(A)$. Then

$$\left(A{\bf v}^{(1)},{\bf v}^{(2)}\right)_2 \; = \; \left< \mathcal{A}{\bf v}^{(1)},{\bf v}^{(2)}\right>_\sigma \; = \; \left(\nabla {\bf v}^{(1)},\nabla {\bf v}^{(2)}\right)_2 \; = \; \left(\nabla {\bf v}^{(2)},\nabla {\bf v}^{(1)}\right)_2$$

$$= \langle A\mathbf{v}^{(2)}, \mathbf{v}^{(1)} \rangle_{\sigma} = (A\mathbf{v}^{(2)}, \mathbf{v}^{(1)})_{2} = (\mathbf{v}^{(1)}, A\mathbf{v}^{(2)})_{2}.$$

Further, we show that operator A is closed: let $\{\mathbf{v}_n\}$ be a sequence in D(A), such that $\mathbf{v}_n \to \mathbf{v}$ (for $n \to \infty$) in $\mathbf{L}^2_{\sigma}(\Omega)$. Put $\mathbf{f}_n := A\mathbf{v}_n$. Suppose that $\mathbf{f}_n \to \mathbf{f}$ in $\mathbf{L}^2_{\sigma}(\Omega)$. Put $\mathbb{F}_n := \mathcal{F}(\mathbf{f}_n, \mathbf{0})$, where \mathcal{F} is the operator from Lemma 5. As all functions \mathbf{v}_n $(n = 1, 2, \ldots)$ lie in $\mathbf{V}^{1,2}_{\sigma}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$, we may apply estimate (25) (where we consider $\mathbf{g}_* = \mathbf{0}$) to the difference $\mathbf{v}_m - \mathbf{v}_n$ (for any $m, n \in \mathbb{N}$) and afterwards use the boundedness of operator \mathcal{F} :

$$\|\mathbf{v}_m - \mathbf{v}_n\|_{2,2} \le c \|\mathbb{F}_m - \mathbb{F}_n\|_{1,2} = c \|\mathcal{F}(\mathbf{f}_m, \mathbf{0}) - \mathcal{F}(\mathbf{f}_n, \mathbf{0})\|_{1,2} \le c \|\mathbf{f}_m - \mathbf{f}_n\|_{2},$$

where c is independent of m, n. From this, we deduce that $\mathbf{v} \in \mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$. As $\partial \mathbf{v}_n/\partial \mathbf{n}$ is in $W_{per}^{1/2,2}(\Gamma_{\text{out}})$ and normal to Γ_{out} (for each $n \in \mathbb{N}$), there exists $q \in W_{per}^{1/2,2}(\Gamma_{\text{out}})$, such that $\partial \mathbf{v}/\partial \mathbf{n} = q\mathbf{n}$ on Γ_{out} . Hence $\mathbf{v} \in D(A)$ and $A\mathbf{v} = \mathbf{f}$. We have proven that A is a closed operator in $\mathbf{L}_{\sigma}^2(\Omega)$.

Operator A is positive, because $(A\mathbf{v},\mathbf{v}) = \|\nabla\mathbf{v}\|_2^2$ for $\mathbf{v} \in D(A)$. Consequently, A is a self-adjoint operator in $\mathbf{L}^2_{\sigma}(\Omega)$. Then $R(A)^{\perp}$ (the orthogonal complement to R(A) in $\mathbf{L}^2_{\sigma}(\Omega)$) is equal to N(A) (the null space of A), see [20, p. 168]. However, as $A \subset \mathcal{A}$ and $N(\mathcal{A}) = \{\mathbf{0}\}$, we also have $R(A)^{\perp} = N(A) = \{\mathbf{0}\}$. This shows that R(A) is dense in $\mathbf{L}^2_{\sigma}(\Omega)$. As all functions from D(A) are in $\mathbf{V}^{1,2}_{\sigma}(\Omega) \cap \mathbf{W}^{2,2}(\Omega)$, we may again apply estimate (25) (with $\mathbf{g}_* = \mathbf{0}$) and afterwards the open graph theorem and deduce that R(A) is closed in $\mathbf{L}^2_{\sigma}(\Omega)$. Thus, $R(A) = \mathbf{L}^2_{\sigma}(\Omega)$. As A coincides with A on D(A), we observe that if $\mathbf{f} \in \mathbf{L}^2_{\sigma}(\Omega)$ (which can be identified with a subspace of $\mathbf{V}^{-1,2}_{\sigma}(\Omega)$), the equation $\nu A \mathbf{v} = \mathbf{f}$ has a solution in D(A).

4) Solution of the equation $\nu A \mathbf{v}_1 = \mathbf{F}_1$. Recall that the functional $\mathbf{F}_1 \in \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ is defined by formula (10), where $\mathbb{F}_1 \cdot \mathbf{n} = h_1 \mathbf{n}$ on Γ_{out} and $h_1 \in W_{\text{per}}^{1/2,2}(\Gamma_{\text{out}})$. Extending function h_1 from Γ_{out} to Ω (by means of Lemma 4) so that the extended function h_{1*} is in $W_{\text{per}}^{1,2}(\Omega)$ and equals zero in the neighborhood of Γ_{in} and Γ_p , we obtain

$$\langle \mathbf{F}_{1}, \mathbf{w} \rangle_{\sigma} = -\int_{\Omega} \mathbb{F}_{1} : \nabla \mathbf{w} \, d\mathbf{x} = -\int_{\Gamma_{\text{out}}} (\mathbb{F}_{1} \cdot \mathbf{n}) \cdot \mathbf{w} \, dl + \int_{\Omega} \operatorname{div} \mathbb{F}_{1} \cdot \mathbf{w} \, d\mathbf{x}$$

$$= -\int_{\Gamma_{\text{out}}} h_{1} \mathbf{n} \cdot \mathbf{w} \, dl + \int_{\Omega} \operatorname{div} \mathbb{F}_{1} \cdot \mathbf{w} \, d\mathbf{x} = \int_{\Omega} [-\nabla h_{1*} + \operatorname{div} \mathbb{F}_{1}] \cdot \mathbf{w} \, d\mathbf{x}$$

$$= (-\nabla h_{1*} + \operatorname{div} \mathbb{F}_{1}, \mathbf{w})_{2}.$$

From this, we observe that \mathbf{F}_1 can be identified with a function from $\mathbf{L}^2_{\sigma}(\Omega)$. The inclusion $\mathbf{v}_1 \in \mathbf{V}^{1,2}_{\sigma}(\Omega) \cap \mathbf{W}^{2,2}_{\mathrm{per}}(\Omega)$ now follows from part 3) of this proof.

Due to Theorem 1, there exists $p_1 \in L^2(\Omega)$, such that \mathbf{v}_1 and p_1 satisfy equation (12), which now takes the form

$$-\nu\Delta\mathbf{v}_1 + \nabla p_1 + \operatorname{div}\mathbb{F}_1 = \mathbf{0}. \tag{41}$$

As $\Delta \mathbf{u}$ and div \mathbb{F}_1 belong to $\mathbf{L}^2(\Omega)$, ∇p_1 is in $\mathbf{L}^2(\Omega)$, too. Thus, $p_1 \in W^{1,2}(\Omega)$. Let us show that $p_1 \in W^{1,2}_{\mathrm{per}}(\Omega)$. Multiplying equation (41) by $\mathbf{w} \in \mathbf{V}^{1,2}_{\sigma}(\Omega)$, we obtain

$$0 = \int_{\Omega} \left[-\nu \Delta \mathbf{v}_{1} + \nabla p_{1} + \operatorname{div} \mathbb{F}_{1} \right] \cdot \mathbf{w} \, d\mathbf{x} = \int_{\partial \Omega} \left[-\nu \frac{\partial \mathbf{v}_{1}}{\partial \mathbf{n}} + p_{1} \mathbf{n} + \mathbb{F}_{1} \cdot \mathbf{n} \right] \cdot \mathbf{w} \, dl$$
$$= \int_{\Gamma_{0} \cup \Gamma_{1}} \left[-\nu \frac{\partial \mathbf{v}_{1}}{\partial \mathbf{n}} + p_{1} \mathbf{n} + \mathbb{F}_{1} \cdot \mathbf{n} \right] \cdot \mathbf{w} \, dl + \int_{\Gamma_{\text{out}}} \left[-\nu \frac{\partial \mathbf{v}_{1}}{\partial \mathbf{n}} + p_{1} \mathbf{n} + \mathbb{F}_{1} \cdot \mathbf{n} \right] \cdot \mathbf{w} \, dl$$

$$= \int_{\Gamma_0 \cup \Gamma_1} p_1 \, \mathbf{n} \cdot \mathbf{w} \, \mathrm{d}l = \int_{\Gamma_0} \left[p_1(x_1, x_2) - p(x_1, x_2 + \tau) \right] \mathbf{n} \cdot \mathbf{w} \, \mathrm{d}l.$$

Since this holds for all $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$, p_1 satisfies the condition of periodicity (7).

5) Solution of the equation $\nu A \mathbf{v}_2 = \mathbf{F}_2$. One can deduce by means of Lemma 4 that, there exists a function $\psi \in W^{3,2}_{\mathrm{per}}(\Omega)$, such that $\psi = \partial_1 \psi = 0$ on Γ_{out} and $\nu \partial_1^2 \psi = h_2$ on Γ_{out} . Function ψ equals zero in the neighborhood of Γ_{in} and Γ_p . Put $\mathbf{v}_2 := -\nabla^\perp \psi$. Then $\mathbf{v}_2 \in \mathbf{V}^{1,2}_{\sigma}(\Omega) \cap \mathbf{W}^{2,2}_{\mathrm{per}}(\Omega)$ and $\nu \partial \mathbf{v}_2 / \partial \mathbf{n} = -h_2 \, \mathbf{e}_2$ on Γ_{out} . Put $\mathbf{f}_2 := -\nu \Delta \mathbf{v}_2$ and $\mathbb{H} := \mathcal{F}(-\mathrm{div}\,\mathbb{F}_2 + \mathbf{f}_2, \mathbf{0})$, where \mathcal{F} is the operator from Lemma 5. Then $\mathrm{div}\,\mathbb{H} = -\mathrm{div}\,\mathbb{F}_2 + \mathbf{f}_2$ in Ω and $\mathbb{H} \cdot \mathbf{n} = \mathbf{0}$ on Γ_{out} . Now, for all $\mathbf{w} \in \mathbf{V}^{1,2}_{\sigma}(\Omega)$, we have

$$\nu \left\langle \nabla \mathbf{v}_{2}, \nabla \mathbf{w} \right\rangle_{2} = \int_{\Gamma_{\text{out}}} \nu \frac{\partial \mathbf{v}_{2}}{\partial \mathbf{n}} \cdot \mathbf{w} \, dl - \int_{\Omega} \nu \Delta \mathbf{v}_{2} \cdot \mathbf{w} \, d\mathbf{x}$$

$$= -\int_{\Gamma_{\text{out}}} h_{2} \, \mathbf{e}_{2} \cdot \mathbf{w} \, dl + \int_{\Omega} \mathbf{f}_{2} \cdot \mathbf{w} \, d\mathbf{x}$$

$$= -\int_{\Gamma_{\text{out}}} (\mathbb{F}_{2} \cdot \mathbf{n}) \cdot \mathbf{w} \, dl + \int_{\Omega} \operatorname{div} (\mathbb{F}_{2} + \mathbb{H}) \cdot \mathbf{w} \, d\mathbf{x}$$

$$= -\int_{\Gamma_{\text{out}}} (\mathbb{F}_{2} \cdot \mathbf{n}) \cdot \mathbf{w} \, dl + \int_{\Omega} \operatorname{div} \mathbb{F}_{2} \cdot \mathbf{w} \, d\mathbf{x} = -\int_{\Omega} \mathbb{F}_{2} : \nabla \mathbf{w} \, d\mathbf{x}$$

$$= \langle \mathbf{F}_{2}, \mathbf{w} \rangle_{\sigma}.$$

This shows that $\nu \mathcal{A} \mathbf{v}_2 = \mathbf{F}_2$. In other words, we have proven that this equation has a solution in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{\mathrm{per}}^{2,2}(\Omega)$. By analogy with p_1 , there exists an associated pressure $p_2 \in W_{\mathrm{per}}^{1,2}(\Omega)$.

6) Solution of the equation $\nu \mathcal{A} \mathbf{v}_3 = \nu \mathbf{G}$. As $\Delta \mathbf{g}_* \in \mathbf{L}^2(\Omega)$, we may put $\mathbb{F}_* := \mathcal{F}(\nu \Delta \mathbf{g}_*, \mathbf{0})$. Then $\operatorname{div} \mathbb{F}_* = \nu \Delta \mathbf{g}_*$ in Ω and $\mathbb{F}_* \cdot \mathbf{n} = \mathbf{0}$ on $\Gamma_{\operatorname{out}}$. Let the functional $\mathbf{F}_* \in \mathbf{V}_{\sigma}^{-1,2}(\Omega)$ be defined by formula (10), where we consider \mathbb{F}_* instead of \mathbb{F} . The solution \mathbf{v}_3 of the equation $\nu \mathcal{A} \mathbf{v}_3 = \mathbf{G}$ satisfies

$$\langle \nu \mathcal{A} \mathbf{v}_{3}, \mathbf{w} \rangle_{\sigma} = \langle \nu \mathbf{G}, \mathbf{w} \rangle_{\sigma} = -\int_{\Omega} \nu \nabla \mathbf{g}_{*} : \nabla \mathbf{w} \, d\mathbf{x} = \int_{\Omega} \nu \Delta \mathbf{g}_{*} \cdot \mathbf{w} \, d\mathbf{x}$$
$$= \int_{\Omega} \operatorname{div} \mathbb{F}_{*} \cdot \mathbf{w} \, d\mathbf{x} = -\int_{\Omega} \mathbb{F}_{*} : \nabla \mathbf{w} \, d\mathbf{x}$$

for all $\mathbf{w} \in \mathbf{V}_{\sigma}^{1,2}(\Omega)$. Since \mathbb{F}_* is in $W_{\mathrm{per}}^{1,2}(\Omega)^{2\times 2}$, we obtain the inclusion $\mathbf{v}_3 \in \mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{\mathrm{per}}^{2,2}(\Omega)$ from part 4) or 5) of this proof. The existence of an associated pressure $p_3 \in W_{\mathrm{per}}^{1,2}(\Omega)$ now follows by means of the same arguments as at the end of part 4).

7) The validity of statement (a). The solvability of the equation $\nu A \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ in $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{\mathrm{per}}^{2,2}(\Omega)$ and the existence of an associated pressure in $W_{\mathrm{per}}^{1,2}(\Omega)$ now follows from the decomposition of the right hand side to $\mathbf{F}_1 + \mathbf{F}_2 + \nu \mathbf{G}$ and from the parts 4), 5) and 6) of this proof.

The proof of Theorem 2 is completed.

Remark 1 Theorem 2 can be generalized so that instead of the functions $\mathbf{u} \in \mathbf{W}^{2,2}_{\mathrm{per}}(\Omega)$ and $p \in W^{1,2}_{\mathrm{per}}(\Omega)$, it yields $\mathbf{u} \in \mathbf{W}^{n+2,2}_{\mathrm{per}}(\Omega)$ and $p \in W^{n+1,2}_{\mathrm{per}}(\Omega)$ for $n \in \{0\} \cup \mathbb{N}$. The generalization says:

Let $n \in \mathbb{N} \cup \{0\}$. Let the closed curve Γ_p (the boundary of profile P_0) be of the class C^{n+2} , $\mathbb{F} \in W^{n+1,2}_{\mathrm{per}}(\Omega)^{2\times 2}$ and \mathbf{g} , \mathbf{g}_* be the functions from Lemma 6, where we consider m = n+1. Let the functionals \mathbf{F} and \mathbf{G} be defined by formulas (10) and (11), respectively. Then

- 1) the unique solution \mathbf{v} of the equation $\nu A \mathbf{v} = \mathbf{F} + \nu \mathbf{G}$ belongs to the space $\mathbf{V}_{\sigma}^{1,2}(\Omega) \cap \mathbf{W}_{\mathrm{per}}^{n+2,2}(\Omega)$ and the associated pressure p is in $W_{\mathrm{per}}^{n+1,2}(\Omega)$,
- 2) **u**, p satisfy statements (b) and (c) of Theorem 2,
- 3) there exists a constant $c_4 = c_4(\nu, \Omega, n)$, such that

$$\|\mathbf{u}\|_{n+2,2} + \|\nabla p\|_{n,2} \le c_4 \left(\|\mathbb{F}\|_{n+1,2} + \|\mathbf{g}_*\|_{n+2,2} \right). \tag{42}$$

As the complete proof of the generalization would be long and its steps would be just technical modifications of the steps from the proof of Theorem 2, we do not include it here. We only note that the corresponding analogue of Lemma 7 would use Proposition I.2.3 from [37] with n+2 instead of 2, the analogue of Lemma 8 would use Theorem III.3.3 from [13] in a subtler way and with m=n+1 instead of m=1 in order to obtain function \mathbf{v}_* (see the proof of Lemma 8), and it would be also necessary to use higher order difference quotients in the proof of the analogue of Lemma 8.

Acknowledgement. The author acknowledges the support of the European Regional Development Fund-Project "Center for Advanced Applied Science" No. CZ.02.1.01/0.0/0.0/16_019 /0000778.

References

- 1. S. Agmon: Lectures on Elliptic Boundary Value Problems. Van Nostrand Comp., New York 1965.
- H. Al Baba, Ch. Amrouche, M. Escobedo: Semi-group theory for the Stokes operator with Navier-type boundary condition in L^p-spaces. Arch. Rat. Mech. Anal. 223 (2017), No. 2, 881–940.
- 3. Ch. Amrouche, M. Escobedo, A. Ghosh: Semigroup theory for the Stokes operator with Navier boundary condition in L^p spaces. ArXiv: 1808.02001v1 [math.AP] 6 Aug 2018.
- C. H. Bruneau, P. Fabrie: New efficient boundary conditions for incompressible Navier–Stokes equations: A well–posedness result. *Math. Modelling and Num. Analysis* 30 (1996), No. 7, 815–840.
- G. Q. Chen, D. Osborne, Z. Qian: The Navier-Stokes equations with the kinematic and vorticity boundary conditions on non-flat boundaries. *Acta Math. Sci.* 29B (2009), No. 4, 919–948.
- G. Q. Chen, Z. Qian: A study of the Navier-Stokes equations with the kinematic and Navier boundary conditions. *Indiana Univ. Math. J.* 59 (2010), no. 2, 721–760.
- M. Dauge: Stationary Stokes and Navier–Stokes systems on two– or three-dimensional domains with corners. Part I: linearized equations. SIAM J. Math. Anal. 20 (1989), 74–97.
- 8. V. Dolejší, M. Feistauer, J. Felcman: Numerical simulation of compressible viscous flow through cascades of profiles. *Z. für Angew. Math. Mech.* **76** (1996), 301–304.
- M. Feistauer: Mathematical Methods in Fluid Dynamics. Pitman Monographs and Surveys in Pure and Applied Mathematics 67, Longman Scientific & Technical, Harlow 1993.
- M. Feistauer, T. Neustupa: On some aspects of analysis of incompressible flow through cascades of profiles. Operator Theory, Advances and Applications, Vol. 147, Birkhäuser, Basel, 2004, 257–276.
- M. Feistauer, T. Neustupa: On non-stationary viscous incompressible flow through a cascade of profiles. Math. Meth. Appl. Sci. 29 (2006), No. 16, 1907–1941.
- 12. M. Feistauer, T. Neustupa: On the existence of a weak solution of viscous incompressible flow past a cascade of profiles with an arbitrarily large inflow. *J. Math. Fluid Mech.* **15** (2013), 701–715.
- G. P. Galdi: An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Steady State Problems. Springer-Verlag, New York-Berlin-Heidelberg 2011.
- 14. M. Geissert, H. Heck, M. Hieber: On the equation div u=g and Bogovskii's operator in Sobolev spaces of negative order. In *Partial differential equations and functional analysis, Oper. Theory Adv. Appl.* **168**, Birkhüser, Basel 2006, pp. 113–121.

2.1

- R. Glowinski: Numerical Methods for Nonlinear Variational Problems. Springer-Verlag, New York-Berlin-Heidelberg-Tokyo, 1984.
- P. Grisvard: Elliptic Problems in non–Smooth Domains. Pitman Advanced Publishing Program, Boston– London–Melbourne 1985.
- P. Grisvard: Singularités des solutions du probléme de Stokes dans un polygone. Université de Nice, 1979.
- J. G. Heywood, R. Rannacher and S. Turek: Artificial boundaries and flux and pressure conditions for the incompressible Navier-Stokes equations. Int. J. for Numerical Methods in Fluids 22 (1996), 325–352.
- R. B. Kellog, J. E. Osborn: A regularity result for the Stokes problem in a convex polygon. *J. Functional Analysis* 21 (1976), 397–431.
- T. Kato: Perturbation Theory for Linear Operators. Springer-Verlag, Berlin-Heidelberg-New York, 1966.
- T. Kato, M. Mitrea, G. Ponce, M. Taylor: Extension and representation of divergence–free vector fields on bounded domains. *Mathematical Research Letters* 7 (2000), 643–650.
- 22. K. Kozel, P. Louda, J. Příhoda: Numerical solution of turbulent flow in a turbine cascade. *Proc. Appl. Math. Mech.* **6** (2006), 743–744.
- S. Kračmar, J. Neustupa: Modelling of flows of a viscous incompressible fluid through a channel by means of variational inequalities. ZAMM 74, No. 6, 637–639 (1994).
- S. Kračmar, J. Neustupa: A weak solvability of a steady variational inequality of the Navier–Stokes type with mixed boundary conditions. *Nonlinear Analysis* 47, No. 6, 4169–4180 (2001).
- S. Kračmar, J. Neustupa: Modeling of the unsteady flow through a channel with an artificial outflow condition by the Navier–Stokes variational inequality. *Math. Nachrichten* 291 (2018), Issue 11–12, 1– 14
- P. Kučera, Z. Skalák, Solutions to the Navier–Stokes equations with mixed boundary conditions. *Acta Appl. Math.* 54, No. 3, 275–288 (1998).
- 27. P. Kučera: Basic properties of the non-steady Navier–Stokes equations with mixed boundary conditions in abounded domain. *Ann. Univ. Ferrara* **55**, 289–308 (2009).
- P. Kučera, M. Beneš: Solution to the Navier–Stokes equatons with mixed boundary conditions in twodimensional bounded domains. *Math. Nachr.* 289 (2016), 2–3, 194–212.
- O. A. Ladyzhenskaya: The Mathematical Theory of Viscous Incompresible Flow. Gordon and Breach Science Publishers, Now York 1969.
- 30. J. L. Lions, E. Magenes: Problèmes aux limites non homogènes et applications. Dunod, Paris 1968.
- 31. D. Medková: The Neumann problem for the planar Stokes system. *Ann. Univ. Ferrara* **58** (2012), 307–329.
- D. Medková: One problem of the Navier type for the Stokes system inplanar domains. J. Diff. Equations 261 (2016), 5670–5689.
- J. Nečas: Les Méthodes Directes en Théorie des Equations Elliptiques. Masson et Cie, Editeurs, Paris 1967.
- T. Neustupa: Question of existence and uniqueness of solution for Navier–Stokes Equation with linear "do-nothing" type boundary condition on the outflow. *Lcture Notes in Computer Science* 5434 (2009), 431–438
- T. Neustupa: The analysis of stationary viscous incompressible flow through a rotating radial blade machine, existence of a weak solution. Appl. Math. and Computation 219 (2012), 3316–3322.
- 36. T. Neustupa: A steady flow through a plane cascade of profiles with an arbitrarily large inflow: the mathematical model, existence of a weak solution. Appl. Math. and Computation 272 (2016) 687-691.
- 37. R. Temam: Navier-Stokes Equations. North-Holland, Amsterdam-New York-Oxford 1977.
- 38. H. Sohr: The Navier–Stokes equations. The Eelementary Functional Analytic Approach. Birkhäuser Verlag, Basel–Boston–Berlin 2001.
- P. Straka, J. Příhoda, M. Kožíšek, J. Fürst: Simulation of transitional flows through a turbine blade cascade with heat transfer for various flow conditions. *EPJ Web of Conferences* 143 (2017), 02118, DOI: 10.1051/epjconf/201714302118.