ON THE GENUS AND AREA OF CONSTANT MEAN CURVATURE SURFACES WITH BOUNDED INDEX

ARTUR B. SATURNINO

ABSTRACT. Using the local picture of the degeneration of sequences of minimal surfaces developed by Chodosh, Ketover and Maximo [5] we show that in any closed Riemannian 3-manifold (M, g), the genus of an embedded CMC surface can be bounded only in terms of its index and area, independently of the value of its mean curvature. We also show that if M has finite fundamental group, the genus and area of any non-minimal embedded CMC surface can be bounded in term of its index and a lower bound for its mean curvature.

1. INTRODUCTION

Let Σ be a compact immersed two-sided constant mean curvature (CMC) surface in a Riemannian 3-manifold (M, g). The *strong Morse index* of Σ is defined as the index of the stability operator

$$Q(u,u) = \int_{\Sigma} |\nabla u|^2 - (|A|^2 + \operatorname{Ric}_M(N))u^2 dvol_{\Sigma}$$

where |A| is the norm of the second fundamental form of Σ and $u \in C^{\infty}(\Sigma)^{-1}$. We will refer to the strong Morse index of Σ as the index of Σ .

The index is a natural variational quantity associated to CMC surfaces, hence it is expected to be controlled in CMC surfaces produced through variational methods. For example, it has been shown that the index of minimal surfaces associated to the volume spectrum are controlled by the order of the associated spectrum element [13, 23], and analogous results are believed to hold for CMC surfaces produced trough the Zhou-Zhu min-max procedure [13, 24].

On the other hand, the geometry and topology of surfaces produced through variational methods tend to be hard to control directly. With the objective of bridging this gap, we are interested in the relation between the index of CMC surfaces and classical geometric and topological quantities. This relation has been well explored for minimal surfaces and hypersurfaces [2,5,6,17,20,22], although the following conjecture from Marques and Neves' 2014 ICM lectures is still open:

Conjecture 1 ([16]). If the ambient manifold has positive Ricci curvature, then an index I embedded orientable compact minimal hypersurface has first Betti number bounded by a fixed multiple of I.

The relation between the index and the geometry of CMC surfaces and hypersurfaces is well known in some special cases (e.g. [1,4,19]). Here we will study the

¹In the case where the surface is not minimal, it is natural to consider only variations of Σ that preserve volume to first degree. That means adding the restriction $\int_{\Sigma} u = 0$ to the domain of Q. The index of Q in this domain is called the *weak Morse index* of Σ . Note that the difference between the weak and strong Morse indices is at most one [3].

ARTUR B. SATURNINO

area and genus of CMC surfaces of bounded index by describing the degeneration of sequences of such surfaces in a similar way as Chodosh, Ketover and Maximo have done for minimal surfaces [5]. We will show that in any closed 3-manifold, the genus of a CMC surface is controlled by its index and area independently of the value of its mean curvature:

Theorem 2. Let $I, A_0 < \infty$ and suppose (M, g) is a closed 3-dimensional Riemannian manifold. There is a constant $C < \infty$ depending only on (M, g), I and A_0 such that any closed, connected, embedded CMC surface in (M, g) with area at most A_0 and index at most I has genus at most C.

If M is spherical, that is, M is closed with finite fundamental group, we will show that every sequence of closed connected CMC surfaces embedded in M with an uniform lower mean curvature bound and an uniform upper index bound has a subsequence that does not accumulate away from a finite set or points. Using this fact we are able to show:

Theorem 3. Let $I < \infty, \eta > 0$ and suppose (M, g) is a spherical 3-dimensional Riemannian manifold. There are constants $B, C < \infty$ depending only on (M, g), I and η such that any closed connected embedded CMC surface in (M, g) with index at most I and mean curvature at least η has area at most B and genus at most C.

The compactness assumption in Theorem 3 is necessary [11], as is the lower bound on the mean curvature [7]. However it is not clear if the assumption on the fundamental group of M is essential.

We will always assume the surfaces are connected and two-sided. Note that if Σ is a non-minimal CMC surface we can orient its normal bundle by the mean curvature vector. Then the scalar value of the mean curvature H = tr(A) of Σ is positive, we will often refer to H as the mean curvature of Σ . In the case Σ is minimal and one-sided, the proof of Theorem 2 is given in [5]. Now we will now present the main ideas in the proof of theorems 2 and 3.

1.1. Outline of the proofs of the main theorems. Theorems 2 and 3 are shown by studying the degeneration of sequences of closed embedded CMC surfaces, possibly with varying mean curvature, $\{\Sigma_n\}$ in (M, g) with index at most I. Assume by contradiction that the genus of the Σ_n form a divergent sequence, and assume for simplicity that, passing to a subsequence, the Σ_n have uniformly bounded mean curvature (if this is not the case an extra rescaling is needed).

By Theorem 4, we can pass $\{\Sigma_n\}$ to a subsequence with uniformly bounded second fundamental form away from a set of at most I points in M, called blow-up points. Near the blow-up points we follow the arguments of Chodosh, Ketover and Maximo [5] to show that a subsequence of rescaled Σ_n must converge to a minimal surface in \mathbb{R}^3 with index at most I. This fact allows us to bound the genus and area of the Σ_n near the blow-up points.

Away from the blow-up points the Σ_n must subconverge to a weak CMC lamination of M^2 . If the leaves of this lamination are proper we can use the limit lamination to bound the genus of the Σ_n away from the blow-up points. In order to show that the leaves of the limit lamination are proper we bound the number of

²Even though we will not use this fact, it is interesting to note that by Theorem 4 and the Local Removable Singularity Theorem for CMC laminations [15, Theorem 1.2] we can choose a subsequence so this limit lamination extends to the blow-up points.

sheets any Σ_n can have in any small region away from the blow-up points. At this point the proof of theorems 2 and 3 diverge.

In the case of Theorem 2 we have uniform area bound for the Σ_n , so the bound of the number of sheets follows from a standard argument. To prove Theorem 3 we will show that if M is spherical, the direction of the mean curvature vectors of these sheets must alternate in a certain sense. This, together with an uniform lower bound on the mean curvature of the Σ_n will give uniform bounds on the maximum number of sheets of any Σ_n that can pass through a small region away from the blow-up points.

1.2. **Outline of the paper.** In Section 2 we will use blow-up arguments to show curvature bounds for CMC surfaces with bounded index. Section 3 is dedicated to adapting results from Chodosh, Ketover and Maximo [5] to CMC surfaces. In Section 4 we show upper bounds on the number of sheets of a CMC surface that pass through a small region of a spherical 3-manifold. Finally, Section 5 contains the proof of the main theorems. Appendices A and B contain details about the local parametrization of CMC surfaces as graphs.

1.3. Acknowledgments. I would like to thank my advisor Davi Maximo for many insightful comments and conversations. I would also like to thank the reviewer for carefully reading the manuscript, pointing out typos and making observations that led to corrections to Proposition 10 and Lemma 11.

2. CURVATURE BOUNDS FOR CMC SURFACES WITH FINITE INDEX

The following scale-invariant inequality is a generalization of a classical inequality first discovered by Schoen [21] (c.f. [5, Lemma 2.2; 18, Main Theorem]). We will prove the theorem in full generality since we believe it may be useful in more contexts.

Theorem 4. For all $I < \infty$ there is a constant $C < \infty$ depending only on I such that the following holds: Let (M, g) be a complete Riemannian 3-manifold with absolute sectional curvature bounds $|K| \leq \Lambda < \infty$ for some $\Lambda > 0$ and suppose $\Sigma \to M$ is an immersed CMC surface with (strong) index at most I. Then there is a set $\mathcal{B} \subset \Sigma$ with at most I points such that for all $p \in \Sigma$

(1)
$$|A|(p)\min\left\{d_{\Sigma}(p,\partial\Sigma\cup\mathcal{B}),(\sqrt{\Lambda})^{-1}\right\} \leq C.$$

This Theorem will be shown in 2.2 after we introduce the language we will use to prove it. Note that since the constant C does not depend on the mean curvature H of Σ and $|A| \ge \sqrt{2}H$ we can conclude from (1) that if $\frac{\sqrt{2}H}{\sqrt{\Lambda}} \ge C$ we must have $d_{\Sigma}(p, \partial \Sigma \cup \mathcal{B}) \le \frac{C}{\sqrt{2}H}$ for all $p \in \Sigma$. So it follows:

Corollary 5. Take M, Λ, Σ, I an C as in Theorem 4 and let H be the mean curvature of Σ . If $\frac{\sqrt{2H}}{\sqrt{\Lambda}} \geq C$ we can conclude that for all $p \in \Sigma$:

(2)
$$d_{\Sigma}(p,\partial\Sigma) \le \left(I + \frac{1}{2}\right) \frac{\sqrt{2}C}{H}$$

if Σ has a boundary, and

(3)
$$\operatorname{diam}_{\Sigma}(\Sigma) \leq \frac{\sqrt{2IC}}{H}$$

if Σ is closed.

Proof. We will only show (2) as the proof of (3) is very similar. As observed above,

(4)
$$d_{\Sigma}(p, \partial \Sigma \cup \mathcal{B}) \le \frac{C}{\sqrt{2}H}$$

for all $p \in \Sigma$. So the distance between any point in \mathcal{B} and its complement in $\partial \Sigma \cup \mathcal{B}$ is at most $\frac{\sqrt{2}C}{H}$. Then an induction argument on I shows that any point in \mathcal{B} is at distance at most $I\frac{\sqrt{2}C}{H}$ from $\partial \Sigma$. Now (2) follows from this fact together with (4).

2.1. Harmonic coordinates and sequences of rescaled surfaces. Theorem 4 does not depend on the manifold M, only on its sectional curvature bounds. For this reason we will have to make use of harmonic coordinates. More specifically we will use the following result by Rosenberg, Souam and Toubiana:

Theorem 6 ([18, Theorem 2.1]). Let $\alpha \in (0,1)$ and $\delta > 0$. Let (M,g) be a Riemannian 3-manifold without boundary with absolute sectional curvature bounds $|K| \leq \Lambda < \infty$. Let Ω be a open subset of M and set

$$\Omega(\delta) = \{ x \in M : d_M(x, \Omega) < \delta \}$$

where d_M is the distance associated to g. Suppose there exists i > 0 such that for all $x \in \Omega(\delta)$ have $inj_{(M,g)}(x) \ge i$, where $inj_{(M,g)}(x)$ is the injectivety radius of M at x. Then there exists constants $Q_0, r_0 > 0$ depending only on i, δ, Λ and α , and not on M, such that for any $x \in \Omega$, there exists a harmonic coordinate chart $(U, \varphi, B_M(x, r_0))$, where $U \subset \mathbb{R}^3$ is an open set containing the origin, $B_M(x, r_0)$ is the geodesic ball in M centered at x with radius r_0 , and $\varphi : U \to B_M(x, r_0)$ is such that $\varphi(0) = x$, and such that the metric tensor $\varphi^* g$ is $C^{1,\alpha}$ - controlled. Namely the components $(\varphi^* g)_{ij}, i, j = 1, 2, 3$ of $\varphi^* g$ satisfy:

(5)
$$Q_0^{-1}\delta_{ij} \le (\varphi^*g)_{ij} \le Q_0\delta_{ij}$$

as quadratic forms, and

(6)
$$\|(\varphi^*g)_{ij}\|_{C^{1,\alpha}(U)} \le Q_0$$

The proofs of all theorems presented in this paper make use of sequences of rescaled surfaces. What we mean by rescaled surfaces will be slightly different for different proofs. Following [18], we will present here the definition used in the proof of Theorem 4, which is more involved because the surfaces might only be immersed and because the ambient space is not fixed.

Suppose $\{(M_n, g_n)\}_{n \in \mathbb{N}}$ is a sequence of complete 3-manifolds with uniformly bounded absolute sectional curvature $|K_n| \leq \Lambda < \infty$. For each $n \in \mathbb{N}$ let Σ_n be an immersed H_n -surface in M_n (that is, a constant mean curvature surface with mean curvature H_n), possibly with boundary, and take a sequence of points $p_n \in \Sigma_n \setminus \partial \Sigma_n$. For any sequence of positive numbers $\{\sigma_n\}_{n \in \mathbb{N}}$ with $\sigma_n \to \infty$ We will define a sequence of rescaled surfaces

$$\sigma_n(\Sigma_n - p_n)$$

immersed in \mathbb{R}^3 .

By abuse of notation, denote image of p_n under the immersion $\Sigma_n \to M_n$ by p_n . Since the exponential map of any of the M_n at any point is a local differomorphism when restricted to the euclidean ball of radius $\frac{\pi}{\sqrt{\lambda}}$ we may endow this ball with the

4

pull-back metric $\exp_{p_n}^* g_n$, which we will also denote by g_n . By [18, Lemma 2.2] the closed ball $\overline{B_{\mathbb{R}^3}(0, \frac{\pi}{2\sqrt{\Lambda}})} \subset (B_{\mathbb{R}^3}(0, \frac{\pi}{\sqrt{\Lambda}}), g_n)$ has injectivity radius at least $\frac{\pi}{4\sqrt{\Lambda}}$. Fix an $\alpha \in (0, 1)$ and consider r_0, Q_0 given by applying Theorem 6 with $\delta = \frac{\pi}{8\sqrt{\Lambda}}$ and $i = \frac{\pi}{4\sqrt{\Lambda}}$. For all $n \in \mathbb{N}$ take a harmonic chart $(U_n, \varphi_n, B_{\mathbb{R}^3}(0, r_0))$ with $0 \in U_n$ and $\varphi_n(0) = 0$. We define

$$\Sigma_n - p_n \subset (U_n, \varphi_n^* g_n)$$

to be the connected component of $(\exp_{p_n} \circ \varphi_n)^{-1}(\Sigma_n)$ passing trough the origin. Note that this is an immersed H_n -surface. For all $\sigma > 0$ let $\mu_{\sigma} : \mathbb{R}^3 \to \mathbb{R}^3$ be the dilation $x \mapsto \sigma x$. Define $\tilde{g}_n = (\mu_{\sigma_n} \circ \varphi_n)^* g_n$ It is clear that

$$\sigma_n(\Sigma_n - p_n) \subset (\mu_{\sigma_n}(U_n), \tilde{g}_n)$$

is an immersed $\frac{H_n}{\sigma_n}$ -surface. We will often use that $(\mu_{\sigma_n}(U_n), \tilde{g}_n) \to (\mathbb{R}^3, \delta_{ij})$ locally in $C^{1,\alpha}$ [18].

2.2. Proof of Theorem 4. Following the arguments of Chodosh, Ketover and Maximo [5, Lemma 2.2] we will use induction on I. The case I = 0 is done in [18]. Let I > 0 and assume by contradiction that the theorem does not hold for I. Then there is a sequence $\{(M_n, g_n)\}_{n \in \mathbb{N}}$ of complete 3-manifolds with absolute sectional curvature bounds $|K_n| \leq \Lambda_n < \infty$ and such that for each $n \in \mathbb{N}$ there is an $H_n \geq 0$ and an immersed H_n -surfaces $\Sigma_n \to M_n$ and points $q_n \in \Sigma_n$ such that

$$|A_n|(q_n)\min\{d_{\Sigma_n}(q_n,\partial\Sigma_n),(\sqrt{\Lambda_n})^{-1}\}>n$$

where $|A_n|$ is the norm of the second fundamental form of Σ_n . Since the left hand side of the inequality above is invariant under rescaling g_n we can assume that all Λ_n are equal to a constant Λ .

Let $R_n = \min\{d_{\Sigma_n}(q_n, \partial \Sigma_n), (\sqrt{\Lambda})^{-1}\}$ and denote $B_{\Sigma_n}(q_n, R_n)$ by D_n . Choose $p_n \in \Sigma_n$ so that

$$|A_n|(p_n)d_{D_n}(p_n,\partial D_n) = \max_{p \in D_n} |A_n|(p)d_{D_n}(p,\partial D_n).$$

Let $\tilde{R}_n = d_{D_n}(p_n, \partial D_n)$ and denote $B_{D_n}(p_n, \tilde{R}_n)$ by \tilde{D}_n . By construction

$$|A_n|(p_n)d_{\tilde{D}_n}(p_n,\partial\tilde{D}_n) = \max_{p\in\tilde{D}_n} |A_n|(p)d_{\tilde{D}_n}(p,\partial\tilde{D}_n).$$

It follows that for any $p \in B_{\tilde{D}_n}(p_n, \frac{\tilde{R}_n}{2})$ we have that $|A_n|(p) \leq 2|A_n|(p_n)$. Let $\lambda_n = |A_n|(p_n)$ and consider the sequence of rescaled surfaces

$$\lambda_n (\tilde{D}_n - p_n).$$

Note that $\lambda_n(\tilde{D}_n - p_n)$ is an $\frac{H_n}{\lambda_n}$ -surface and the norm of its second fundamental is bounded above by 2 in $B_{\mathbb{R}^3}(0, \frac{\lambda_n \tilde{R}_n}{2})$ with the dilated pull-back metric. The distance between the origin $0 \in \lambda_n(\tilde{D}_n - p_n)$ and the boundary of $\lambda_n(\tilde{D}_n - p_n)$ is at least $\lambda_n R_n$.

The arguments in [18] show that there is a complete immersed *H*-surface *S* in the euclidean 3-space (\mathbb{R}^3, g_{eucl}) such that:

- (1) $0 \in S;$
- (2) $|A_S|(0) = 1;$
- (3) $||A_S||_{C^0} \le 5;$

(4) passing to a subsequence, there are pieces $\Delta_n \subset \lambda_n(\tilde{D}_n - p_n)$ such that

 $\Delta_n \to S$

in the sense that there is a $\delta' > 0$ such that for any $x \in S$, and *n* large enough, the components of Δ_n close to $B_S(x, \delta)$ are graphical over $D(\delta) \subset T_x S$ and converge to $B_S(x, \delta)$ in C^2 -norm.

For R' large enough, the index of

 $B_{\Sigma_n}(p_n, R'/\lambda_n)$

is at least one for n sufficiently large, else S must be stable, contradicting the fact that S is complete and non-flat [18]. By induction there is a set

$$\hat{\mathcal{B}}_n \subset \hat{\Sigma}_n = \Sigma_n \setminus B_{\Sigma_n}(p_n, R'/\lambda_n)$$

with at most I - 1 points such that

$$|A_n|(x)\min\{d_{\hat{\Sigma}_n}(x,\partial\hat{\Sigma}_n\cup\hat{\mathcal{B}}_n),(\sqrt{\Lambda})^{-1}\}\leq C$$

We claim that we can take $\mathcal{B}_n = \hat{\mathcal{B}}_n \cup \{p_n\}$. Assume there is $z_n \in \Sigma_n$ such that

$$|A_n|(z_n)\min\{d_{\Sigma_n}(z_n,\partial\Sigma_n\cup\mathcal{B}_n),(\sqrt{\Lambda})^{-1}\}\to\infty.$$

We will consider two cases. First assume

$$\liminf_{n \to \infty} \lambda_n d_{\Sigma_n}(z_n, p_n) < \infty.$$

Passing to a subsequence we get that

$$\frac{d_{\Sigma_n}(z_n, p_n)}{\tilde{R}_n} = \frac{\lambda_n d_{\Sigma_n}(z_n, p_n)}{\lambda_n \tilde{R}_n} \to 0.$$

So, for n large enough $z_n \in B_{\Sigma_n}(p_n, \tilde{R}_n/2)$, and hence

$$|A_n|(z_n) \le 2\lambda_n.$$

We conclude that

$$\lim_{n \to \infty} |A_n|(z_n) d_{\Sigma_n}(z_n, p_n) \le \lim_{n \to \infty} 2\lambda_n d_{\Sigma_n}(z_n, p_n) < \infty.$$

Contradicting the choice of z_n . Now assume that

$$\liminf_{n \to \infty} \lambda_n d_{\Sigma_n}(z_n, p_n) = \infty.$$

It follows that for n large enough $d_{\Sigma_n}(z_n, p_n) > R'/\lambda_n$, hence $z_n \in \hat{\Sigma}_n$, contradicting the inductive hypothesis.

Remark 7. Note that this same proof works in the case where M has a boundary as long as Σ is away from this boundary.

3. PICTURE NEAR BLOW-UP POINTS

In this section we will consider sequences embedded CMC surfaces $\{\Sigma_n\}$ in a fixed closed 3-manifold (M, g). Since the Σ_n are embedded and the ambient manifold is fixed we can make sense of a sequences of rescaled surfaces

$$\sigma_n(\Sigma_n - p_n)$$

in a slightly different way to that presented in Subsection 2.1. Namely, we can take harmonic charts $(U_n, \varphi_n, B_M(x, p_n))$ and define

$$\Sigma_n - p_n \subset (U_n, \varphi_n^* g_n)$$

 $\mathbf{6}$

to be the pre-image of the component of $\Sigma_n \cap B_M(x, p_n)$ passing by p_n . Then $\sigma_n(\Sigma_n - p_n)$ has as the properties listed in Subsection 2.1 in addition to being embedded.

Following [5] we say that a sequence of finite sets of points $\mathcal{B}_n \subset \Sigma_n$ is a sequence of blow-up sets if:

(1) The curvature blows up at the \mathcal{B}_n , that is, taking $\lambda_n(p) = |A_n|(p)$ we have

$$\liminf_{n \to \infty} \min_{p \in \mathcal{B}_n} \lambda_n(p) \to \infty.$$

(2) Taking any sequence of points $p_n \in \mathcal{B}_n$ we can pass to a subsequence so

$$\lambda_n(p_n)(\Sigma_n - p_n) \to \mathcal{L}_{\infty}$$

where \mathcal{L}_{∞} is a weak CMC lamination of euclidean 3-space with $||A_{\mathcal{L}_{\infty}}||_{C^{0}} \leq 5$ and $|A_{\mathcal{L}_{\infty}}|(0) = 1$.

(3) The blow-up points do not appear in the blow-up limit of the other points, that is

$$\liminf_{n \to \infty} \min_{p, q \in \mathcal{B}_n, p \neq q} \lambda_n(p) d_M(p, q) = \infty$$

Proposition 8. Suppose that the index of the Σ_n is uniformly bounded above by a constant $I < \infty$. Then there is a sequence of blow-up sets $\mathcal{B}_n \subset \Sigma_n$ with cardinality at most I and a constant $C < \infty$ depending only on I such that

(7)
$$|A_n|(x)\min\{d_M(x,\partial\Sigma_n\cup\mathcal{B}_n),(\sqrt{\Lambda})^{-1}\}\leq C$$

for all n and all $x \in \Sigma_n$.

Proof. We will show that one can take the sets \mathcal{B}_n as in the proof of Theorem 4 by changing the distance function from the intrinsic distance to the extrinsic distance. It is clear that these sets have cardinality at most I and that the $\lambda_n(p_n)(\Sigma_n - p_n)$ are embedded with bounded second fundamental form and the distance from the origin to boundary of $\lambda_n(p_n)(\Sigma_n - p_n)$ diverges. So it follows that they must subconverge to a weak CMC lamination of euclidean 3-space. (see [9, Appendix B])

To show the third item in the definition holds, let $\hat{\mathcal{B}}_n$ be as constructed in the proof of Theorem 4 and assume by contradiction that

$$\liminf_{n \to \infty} \min_{q \in \hat{\mathcal{B}}_n} \lambda_n d_M(p_n, q) < \infty.$$

Take q_n to be a point in $\hat{\mathcal{B}}_n$ closest to p_n and let $\eta_n = |A_n|(q_n)$, repeating the arguments from the proof of Theorem 4 we have that, passing to a subsequence, for n large enough

$$\eta_n \le \frac{\lambda_n}{2}.$$

So $\liminf_{n\to\infty} \eta_n d_M(q_n, p_n) < \infty$. This clearly implies

$$\lim_{n \to \infty} |A_n|(q_n) d_M(q_n, \partial \hat{\Sigma}_n) < \infty$$

contradicting the choice of q_n .

In the case where the mean curvature of the Σ_n are uniformly bounded, repeating the arguments in [5, Section 4] we can conclude:

Lemma 9. Suppose $\{\Sigma_n\}$ is a sequence of CMC surfaces embedded in a complete Riemannian 3-manifold (M, g) with mean curvature uniformly bounded above by a constant H. Suppose the blow-up sets \mathcal{B}_n accumulate on a set \mathcal{B}_∞ . Then there is a $\delta > 0$ such that the set

$$B_M(\mathcal{B}_\infty,\delta) = \bigcup_{p_\infty \in \mathcal{B}_\infty} B_M(p_\infty,\delta)$$

is a union of disjoint balls where the following holds: Write Σ'_n for the connected components of $B_M(\mathcal{B}_{\infty}, \delta) \cap \Sigma_n$ containing points in \mathcal{B}_n and Σ''_n for the other connected components of $B_M(\mathcal{B}_{\infty}, \delta) \cap \Sigma_n$. Then there are functions m = m(I, H) and r = r(I, H) such that for n large:

- (1) The surfaces Σ''_n are CMC disks with second fundamental form uniformly bounded above.
- (2) The surfaces Σ'_n intersects $\partial B_M(\mathcal{B}_\infty, \delta)$ transversely in at most m simple closed curves.
- (3) The surfaces Σ'_n have genus at most r;
- (4) The surfaces Σ'_n have uniformly bounded area.

4. PICTURE AWAY FROM BLOW-UP POINTS

In this section we will show that a sequence of CMC surfaces embedded in a space with finite fundamental group and with mean curvature bounded away from zero cannot accumulate in a region where they have uniformly bounded second fundamental form (Lemma 11). The results presented here will be used to prove Theorem 3 and will also be needed to remove any upper mean curvature bound assumption on Theorem 2. We will state and prove some of our results for hypersurfaces in manifolds of arbitrary dimension, but for our applications we only need to consider surfaces in 3-manifolds.

4.1. Two-sided hypersurfaces in spaces with finite fundamental group. Let (M,g) be a Riemannian manifold (possibly with boundary) with finite fundamental group. Let $\Sigma \subset M$ be a complete properly embedded two-sided hypersurface and let N be a unit normal vector field of Σ . Suppose $\gamma : [0,1] \to M$ is a differentiable curve between points in $M \setminus \Sigma$ which is transversal to Σ . Write $\gamma^{-1}(\Sigma) = \{t_1, \dots, t_n\}$ where $t_1 < \dots < t_n$. We say that Σ is ℓ -alternating if for any such path γ one of the following holds:

- Either $\ell > n-1$;
- or there is a $j \leq \ell + 1$ such that

$$g(\gamma'(t_1), N(\gamma(t_1)))$$
 and $g(\gamma'(t_j), N(\gamma(t_j)))$

have opposite signs.

Proposition 10. Suppose $|\pi_1(M)| = \ell < \infty$. Then any complete properly embedded two-sided hypersurface $\Sigma \subset M$ is ℓ -alternating.

Proof. Let \tilde{M} be the universal cover of M and let $\tilde{\Sigma}$ be the lift of Σ in the sense that, for any evenly covered open set $U \subset M$ and any sheet $\tilde{U} \subset \tilde{M}$ over $U, \tilde{\Sigma} \cap \tilde{U}$ is mapped isometrically to $\Sigma \cap U$ by the covering map. Note that we can lift a unit normal vector field N of Σ to a unit normal vector field \tilde{N} of $\tilde{\Sigma}$.

Let $\gamma : [0,1] \to M$ be a differentiable path with ends away from Σ which is transversal to Σ and intersects Σ it in at least $\ell + 1$ points. Choose any lift $\tilde{\gamma}$ of γ . Note that $\gamma^{-1}(\Sigma) = \tilde{\gamma}^{-1}(\tilde{\Sigma})$ and for each t_j in this set

$$g(\gamma'(t_j), N(\gamma(t_j))) = \tilde{g}(\tilde{\gamma}'(t_j), \tilde{N}(\tilde{\gamma}(t_j))).$$

So we only need to show that there are $j, j' \leq \ell + 1$ such that

$$g(\tilde{\gamma}'(t_j), N(\gamma(t_j)))$$
 and $\tilde{g}(\tilde{\gamma}'(t_{j'}), N(\tilde{\gamma}(t_{j'})))$

have opposite sings.

Since the order of the cover $\tilde{M} \to M$ is ℓ , $\tilde{\Sigma}$ has at most ℓ connected components. So there must be a connected component $\tilde{\Sigma}_0$ of $\tilde{\Sigma}$ intersecting $\tilde{\gamma}$ at least twice. Let $\tilde{\gamma}(s_1) = \tilde{x}, \tilde{\gamma}(s_2) = \tilde{y} \in \tilde{\Sigma}_0$ be the first and second points of intersection of $\tilde{\gamma}$ with $\tilde{\Sigma}_0$ respectively. Assume by contradiction that the product of the velocity of $\tilde{\gamma}$ with the normal of $\tilde{\Sigma}_0$ has the same sign at both these points. By changing the direction of the normal of $\tilde{\Sigma}_0$ we can assume that these are both negative. Choose a path $\eta : [0, 1] \to \tilde{\Sigma}_0$ between \tilde{x} and \tilde{y} and for $r \geq 0$ set

$$\eta_r(t) = \exp_{\tilde{M}}(rN(\eta(t))).$$

Fix $\rho > 0$ small enough so that η_{ρ} does not intersect $\tilde{\Sigma}_0$. Taking $\epsilon > 0$ small enough, we can join $\gamma|_{[s_1-\epsilon,s_2-\epsilon]}$ to η_{ρ} by two smooth curves that do not intersect $\tilde{\Sigma}_0$. Let β be the concatenation of these 4 curves. Then β is a smooth by parts and intersects $\tilde{\Sigma}_0$ exactly at \tilde{x} . On the other hand, since \tilde{M} is simply-connected and $\tilde{\Sigma}_0$ is complete and proper, the intersection number of β and Σ_0 has to be even. \Box

4.2. CMC surfaces in manifolds with finite fundamental group. We finish this section with the proof of the lemma below.

Lemma 11. Let (M,g) be a 3-manifold (possibly with boundary) with absolute sectional curvature bounds $|K| \leq \Lambda < \infty$ and let $\Sigma \subset M$ be a properly embedded CMC surface. Fix any $\alpha \in (0,1)$ and let $\Omega \subset M$ be an open set such that there are $\delta, i > 0$ with the property that $\Omega(\delta)$ as defined in Theorem 6 does not intersect $\partial M \cup \partial \Sigma$ and $inj_{(M,g)}(p) \geq i$ for all $p \in \Omega(\delta)$. Suppose the second fundamental form of Σ in Ω is bounded above by a constant $C < \infty$. Then for all $p \in \Omega$ there is $a \epsilon > 0$ depending only on α, δ, i, C and Λ such that $\Sigma \cap B_M(p, \epsilon)$ is parametrized by graphs of functions with $C^{2,\alpha}$ bounds depending only on α, δ, i, C and Λ .

If we assume additionally that $\pi_1(M)$ is finite and that Σ is complete and has mean curvature mean curvature $H \ge \eta > 0$, then there is a $N < \infty$ depending only on $\alpha, \delta, i, C, \Lambda, \eta$ and $|\pi_1(M)|$ such that $\Sigma \cap B_M(p, \epsilon)$ is parametrized by no more than N graphs.

Proof. The first part of the statement is equivalent to the Uniform Graph Lemma presented in Appendix A. Now for the second part of the statement, note that by Proposition 10 the graphs that parametrize Σ must alternate. Since these graphs must have alternating mean curvature (in the sense of Appendix B) it is possible to bound the number N of graphs uniformly using the mean curvature equation, as shown in Proposition 14.

ARTUR B. SATURNINO

5. Proof of the main theorems

Now we are are ready to prove theorems 2 and 3. Since the proofs are very similar we present the proof of Theorem 2 in more detail and reference it in the proof of Theorem 3.

5.1. **Proof of Theorem 2.** Let (M, g) be a closed Riemannian 3-manifold and let $\{\Sigma_n\}$ be sequence of closed embedded CMC surfaces in M with index at most I and area at most A_0 . Suppose by contradiction that the genus of the Σ_n form a divergent sequence. Let H_n be the mean curvature of Σ_n . We will consider two cases depending on the behavior of the sequence $\{H_n\}$.

Case 1: $\liminf_{n\to\infty} H_n < \infty$. Pass to a subsequence so the H_n are uniformly bounded in n and divide the Σ_n into three parts: Σ'_n, Σ''_n as in Lemma 9 and

$$\Sigma_n^b = \Sigma_n \setminus B_M(\mathcal{B}_\infty, \delta/2).$$

By Theorem 4, the norm of the second fundamental form of the Σ_n is uniformly bounded away from $B_M(\mathcal{B}_{\infty}, \delta/3)$. Since the area of the Σ_n is uniformly bounded above the number of graphs in the local parametrization of Lemma 11 is uniformly bounded. So we can pass to a subsequence such that the Σ_n^b converge with finite multiplicity to a CMC surface Σ_{∞}^b (see [14, Theorem 4.37]). Hence for *n* large enough Σ_n^b is a cover of Σ_{∞}^b with uniformly bounded degree, so there are uniform bounds on the Betti numbers of the Σ_n^b .

Lemma 9 together with the area bound limits the Betti numbers of Σ'_n and Σ''_n . By [5, Lemma 3.1] the Σ^b_n intersects Σ'_n and Σ''_n on annuli, so we can use a Mayer-Vietoris sequence to bound on the Betti numbers of the Σ_n ³. This contradicts the assumption that the sequence of the genus of the Σ_n diverges.

Case 2: $H_n \to \infty$.

By Corollary 5 we can conclude that a subsequence of the Σ_n converges to a point $p_{\infty} \in M$ in the Hausdorff sense. So for n large enough

$$\Sigma_n \subset B_M(p_\infty, r)$$

where $r = inj_{(M,g)}(p_{\infty})/2$. Using the exponential map of M at p_{∞} we can pull Σ_n back to a properly embedded H_n -surface

$$\Sigma_n \subset (B_{\mathbb{R}^3}(p_\infty, r), \exp_{p_\infty}^* g)$$

Scaling the Σ_n by H_n we obtain a sequence of 1-surfaces

$$\overline{\Sigma}_n = H_n \tilde{\Sigma}_n$$

with uniformly bounded diameter in an increasing sequence of balls with metrics converging smoothly to the euclidean metric in compact sets.

Consider a decomposition of the $\overline{\Sigma}_n$ into $\overline{\Sigma}'_n$, $\overline{\Sigma}''_n$ and $\overline{\Sigma}^b_n$ analogous to the decomposition of the Σ_n in Case 1. By Lemma 11 there are uniform bounds on the number of sheets of the $\overline{\Sigma}^b_n$ on any small open set. Hence we can conclude that, passing to a subsequence, the $\overline{\Sigma}^b_n$ converge to a 1-surface Σ^b_∞ . These same arguments can be used to bound the number of connected components of the $\overline{\Sigma}''_n$. Finally we can conclude using the same arguments from Case 1.

³We can also use a surgery argument [5, Proposition 5.1]

5.2. **Proof of Theorem 3.** First note that by Theorem 2 we only have to show that there is an uniform area bound. Let (M,g) be a spherical Riemannian 3-manifold and let $\{\Sigma_n\}$ be a sequence of closed embedded CMC surfaces in M with index at most $I < \infty$ and mean curvature at least $\eta > 0$. Assume by contradiction that the area of the Σ_n form a divergent sequence.

Let H_n be the mean curvature of Σ_n . In the case where $\limsup_n H_n \to \infty$ we can pass to a subsequence so $H_n \to \infty$. Following the arguments used in Case 2 of the proof of Theorem 2 we can conclude that the area of the mean-curvature rescaled Σ_n is uniformly bounded, hence the area of the Σ_n must converge to 0, a contradiction.

So we may assume $\limsup_{n\to\infty} H_n < \infty$. In this case we can apply the same decomposition used in Case 1 on the proof of Theorem 2 and use Lemma 11 to bound the number of sheets of any Σ_n^b that can pass through small regions. A similar argument can be used to bound the number of connected components the Σ_n'' can have, and hence we can conclude that there are uniform area bounds for these two parts. Finally, Lemma 9 allows us to conclude that there are uniform area bounds for the Σ_n'' .

Appendix A. Uniform Graph Lemma

The following result is a version of the Uniform Graph Lemma (cf [14, Lemma 4.35]) which allows for $C^{2,\alpha}$ bounds for functions parametrizing CMC surfaces depending only on the second fundamental form of the surface and the ambient space sectional curvature. This lemma is slightly stronger than what we need in order to show our main theorems, however it is included here for the sake of completion.

Lemma 12. Suppose (M,g) is a 3-manifold with absolute sectional curvature bounds $|K| \leq \Lambda < \infty$ and let $\Sigma \subset M$ be a properly embedded CMC surface. Let $\Omega \subset M$ be an open set lying away from the boundary of M and suppose the second fundamental form of Σ in Ω is bounded above by a constant $C < \infty$. Fix any $\alpha \in (0,1)$ and suppose δ, i, r_0 and Q_0 are as in Theorem 6. Fix an $r \in (0, r_0)$ and let $x \in \Omega$ be such that $d_M(x, \partial \Sigma) > r$.

Choose a harmonic chart $(U, \varphi, B_M(x, r))$ as in Theorem 6. For any $\epsilon \in (0, r)$ let Σ' be the set of connected components of $\varphi^{-1}(\Sigma \cap B_M(x, r))$ intersecting the euclidean ball $B_{\mathbb{R}^3}(0, \epsilon)$. Then there are $\epsilon > 0$, $\rho \in (\epsilon, r)$ and $C' < \infty$ depending only on Λ, C, i and α , and a rotation $R \in O(\mathbb{R}^3)$ such that:

- (1) Every connected component of $R(\Sigma') \cap B(\rho) \times \mathbb{R}$ is the graph of a function u over $B(\rho)$.
- (2) For all such functions u we have

$$||u||_{C^{2,\alpha}(B(\rho))} \le C'.$$

Proof. From (5) and (6) we can conclude that there is a $K < \infty$ depending only on Q_0 and C such that the norm of the euclidean second fundamental form of $\varphi^{-1}(\Sigma \cap B_M(x,r))$ is at most K. For any $\epsilon > 0$ let Σ' be as above and let G': $\Sigma' \to \mathbb{RP}^2$ be the euclidean unoriented Gauss map of Σ' .

We first claim that there is an $\epsilon > 0$ depending only on K and r such that

$$\sup_{p,q\in\Sigma'\cap B_{\mathbb{R}^3}(0,\epsilon)} d_{\mathbb{P}^2}(G'(p),G'(q)) < \frac{\pi}{4}.$$

For p and q in the same connected component, the existence of ϵ follows immediately from the bounds on the second fundamental form. For p, q in distinct connected components we can use the bound on the second form to show that if ϵ is sufficiently small and G'(p) and G'(q) are too far apart then the connect components on Σ' containing p and q must cross.

Choose a rotation R so that the tangent planes to $R(\Sigma') \cap B_{\mathbb{R}^3}(0,\epsilon)$ are bounded away from any vertical plane by at least $\pi/6$. It follows that there is a constant $\tilde{\rho}$ depending only on K such that, for all $p = (p_1, p_2, p_3) \in R(\Sigma) \cap B_{\mathbb{R}^3}(0,\epsilon)$ the portion of $R(\Sigma')$ in the cylinder $B((p_1, p_2), \tilde{\rho}) \times \mathbb{R}$ is the graph of a function over $B(p, \tilde{\rho})$. The bounds on the euclidean second form and on the distance from the tangent planes to vertical planes give bounds on the C^2 norms for these functions.

By possibly making ϵ smaller, we can assume that $\epsilon < \tilde{\rho}/2$. So taking $\rho = \tilde{\rho}/2$, we have that every connected component of $R(\Sigma') \cap B(\rho) \times \mathbb{R}$ is a graph over $B(\rho)$ of a function with C^2 -control. Making ρ smaller and applying interior Schauder estimates [12, Theorem 6.2] using the mean curvature equation we can obtain $C^{2,\alpha}$ bounds for these functions.

APPENDIX B. ALTERNATING CMC GRAPHS

Here we will use a local version of the alternating property defined in 4.1 to bound the number of sheets any CMC hypersurface can have in a small region.

Let $B(r) \subset \mathbb{R}^d$ be an open ball of radius r and suppose g is a metric in $B(r) \times \mathbb{R}$ such that the g_{ij} have $C^{1,\alpha}$ -norm bounded by a constant Q_0 for some fixed $\alpha \in (0,1)$. Assume further that, as quadratic forms

$$Q_0^{-1}\delta_{ij} \le g_{ij} \le Q_0\delta_{ij}.$$

For any $u \in C^2(B(r))$ we will choose the normal N_u to the graph of u pointing upwards, that is $g(N_u, \partial_d) > 0$. For $x \in B(r)$ let $h_u(x)$ be the scalar mean curvature of the graph of u at (x, u(x)) with respect to the metric g and the normal N_u . We will refer to h_u as the mean curvature of u, and if h_u is constant we will say that u has constant mean curvature.

Proposition 13. Suppose $u, v \in C^{2,\alpha}(B(r))$ have constant mean curvature with opposite signs and

- (1) $u \leq v$ point-wise;
- (2) $|h_u| = |h_v| \ge \eta$ for some $\eta > 0$;
- (3) $||u||_{C^{2,\alpha}(B(r))}, ||v||_{C^{2,\alpha}(B(r))} \leq C$ for some C > 0.

Then there is a c > 0 depending only on Q_0, α, r, η and C such that

$$||u - v||_{C^0(B(r))} > c$$

Proof. Since u and v have $C^{2,\alpha}$ norm bounded above by C we can conclude that for all $\epsilon > 0$ there is a $\delta > 0$ depending only on C, α, ϵ and r such that

$$||u - v||_{C^2(B(r/2))} < \epsilon$$

whenever $||u - v||_{C^0(B(r))} < \delta$. So if u and v can be made arbitrarily close in B(r), their mean curvatures can also be made to be arbitrarily close in B(r/2).

On the other hand from the mean curvature equation (see [8, Chapter 5; 10, Chapter 7]) we can conclude that there is a $C' < \infty$ depending only on Q_0, α, r and C such that

$$2\eta \le \|h_u - h_v\|_{C^0(B(r/2))} \le C' \|u - v\|_{C^2(B(r/2))}$$

whenever u and v are C^2 -close enough. Contradicting the fact that u and v can be made arbitrarily close in the C^2 norm.

Assume $u_1, \dots, u_n \in C^{2,\alpha}(B(r))$ have non-zero constant mean curvature with the same magnitude and $u_1 \leq u_2 \leq \dots \leq u_n$ point-wise. We say that such a collection is ℓ -alternating for some $\ell \in \mathbb{N}$ if for all $j \leq n - \ell$ there is a $j' \leq j + \ell$ such that the mean curvature of u_j and $u_{j'}$ have opposite signs. Note that by Proposition 10 this property always holds for graphs parametrizing a complete properly embedded two-sided CMC hypersurface in a manifold with fundamental group of order at most ℓ .

Proposition 14. Let $u_1, \dots, u_n \in C^{2,\alpha}(B(r))$ be an ℓ -alternating collection and let $\eta > 0, C < \infty$ and $\epsilon \in (0, r)$ be such that for all $k = 1, \dots, n$:

- (1) $|h_{u_k}| \ge \eta;$
- (2) $||u_k||_{C^{2,\alpha}(B(r))} \le C;$
- (3) the graph of every u_k intersects an euclidean ball $B(\epsilon)$.

Then there is an $N < \infty$ depending only on $Q_0, \alpha, \eta, r, \ell, C$ and ϵ such that $n \leq N$.

Proof. By passing to a subset of $\{u_1, \dots, u_n\}$ with at least $\lfloor n/\ell \rfloor$ elements we can assume that the mean curvature of the u_k alternate (i.e. $\ell = 1$). The difference $w_{k,k+2} = u_{k+2} - u_k$ follows an elliptic equation with ellipticity and coefficients controlled by the C^1 norm of the metric and the C^1 norm of u_{k+2} and u_k (see [8, Chapter 5; 10, Chapter 7]). It follows that these functions obey a Harnack inequality with a constant C' that only depends on Q_0 and C [12, Theorem 8.20].

It follows from the C^1 bound on the u_k that $|u_k| < 3C\epsilon$ in $B(\epsilon)$. Hence for at least one of the $w_{k,k+2}$ must have

$$\inf_{x \in B(\epsilon)} w_{k,k+2}(x) \le \frac{6C\epsilon}{n}$$

So by the Harnack inequality $w_{k,k+2} \leq 6C' \frac{C\epsilon}{n}$ in $B(\epsilon/2)$, this implies that

$$||u_k - u_{k+1}||_{C^0(B(\epsilon/2))} \le 6C'\frac{C\epsilon}{n}$$

So we must have $n < 6\epsilon C'Cc^{-1}$ where c is as in Proposition 13 with $r = \epsilon/2$.

References

- Nicolau S. Aiex and Han Hong, Index estimates for surfaces with constant mean curvature in 3-dimensional manifolds, Calc. Var. Partial Differential Equations 60 (2021), no. 1, Paper No. 3, 20. MR4176856
- [2] Lucas Ambrozio, Alessandro Carlotto, and Ben Sharp, Comparing the Morse index and the first Betti number of minimal hypersurfaces, J. Differential Geom. 108 (2018), no. 3, 379– 410. MR3770846
- [3] Lucas Barbosa and Pierre Bérard, Eigenvalue and "twisted" eigenvalue problems, applications to CMC surfaces, J. Math. Pures Appl. (9) 79 (2000), no. 5, 427–450. MR1759435
- [4] Marcos Petrúcio Cavalcante and Darlan Ferreira de Oliveira, Lower bounds for the index of compact constant mean curvature surfaces in ℝ³ and S³, Rev. Mat. Iberoam. **36** (2020), no. 1, 195–206. MR4061986
- [5] Otis Chodosh, Daniel Ketover, and Davi Maximo, Minimal hypersurfaces with bounded index, Invent. Math. 209 (2017), no. 3, 617–664. MR3681392
- [6] Otis Chodosh and Davi Maximo, On the topology and index of minimal surfaces, J. Differential Geom. 104 (2016), no. 3, 399–418. MR3568626

ARTUR B. SATURNINO

- [7] Tobias H. Colding and William P. Minicozzi II, Examples of embedded minimal tori without area bounds, Internat. Math. Res. Notices 20 (1999), 1097–1100. MR1728019
- [8] ______, Minimal surfaces, Courant Lecture Notes in Mathematics, vol. 4, New York University, Courant Institute of Mathematical Sciences, New York, 1999. MR1683966
- [9] _____, The space of embedded minimal surfaces of fixed genus in a 3-manifold. IV. Locally simply connected, Ann. of Math. (2) 160 (2004), no. 2, 573–615. MR2123933
- [10] Tobias Holck Colding and William P. Minicozzi II, A course in minimal surfaces, Graduate Studies in Mathematics, vol. 121, American Mathematical Society, Providence, RI, 2011. MR2780140
- Baris Coskunuzer, H-surfaces with arbitrary topology in hyperbolic 3-space, J. Geom. Anal. 27 (2017), no. 2, 1211–1236. MR3625148
- [12] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition. MR1814364
- [13] Yangyang Li, An improved morse index bound of min-max minimal hypersurfaces (2020), available at arXiv:2007.14506v2.
- [14] W. H. Meeks III, A. Ros, and H. Rosenberg, The global theory of minimal surfaces in flat spaces, Lecture Notes in Mathematics, vol. 1775, Springer-Verlag, Berlin; Centro Internazionale Matematico Estivo (C.I.M.E.), Florence, 2002. Lectures given at the 2nd C.I.M.E. Session held in Martina Franca, July 7–14, 1999, Edited by Gian Pietro Pirola, Fondazione CIME/CIME Foundation Subseries. MR1901611
- [15] William H. Meeks III, Joaquín Pérez, and Antonio Ros, The classification of CMC foliations of ℝ³ and S³ with countably many singularities, Amer. J. Math. **138** (2016), no. 5, 1347– 1382. MR3553393
- [16] André Neves, New applications of min-max theory, Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II, 2014, pp. 939–957. MR3728646
- [17] Antonio Ros, One-sided complete stable minimal surfaces, J. Differential Geom. 74 (2006), no. 1, 69–92. MR2260928
- [18] Harold Rosenberg, Rabah Souam, and Eric Toubiana, General curvature estimates for stable H-surfaces in 3-manifolds and applications, J. Differential Geom. 84 (2010), no. 3, 623–648. MR2669367
- [19] Wayne Rossman and Nahid Sultana, Morse index of constant mean curvature tori of revolution in the 3-sphere, Illinois J. Math. 51 (2007), no. 4, 1329–1340. MR2417430
- [20] Alessandro Savo, Index bounds for minimal hypersurfaces of the sphere, Indiana Univ. Math. J. 59 (2010), no. 3, 823–837. MR2779062
- [21] Richard Schoen, Estimates for stable minimal surfaces in three-dimensional manifolds, Seminar on minimal submanifolds, 1983, pp. 111–126. MR795231
- [22] Antoine Song, Morse index, betti numbers and singular set of bounded area minimal hypersurfaces (2019), available at arXiv:1911.09166.
- [23] Xin Zhou, On the multiplicity one conjecture in min-max theory, Ann. of Math. (2) 192 (2020), no. 3, 767–820. MR4172621
- [24] Xin Zhou and Jonathan J. Zhu, Min-max theory for constant mean curvature hypersurfaces, Invent. Math. 218 (2019), no. 2, 441–490. MR4011704

Department of Mathematics, University of Pennsylvania, Philadelphia, PA 19104, USA

Email address: bsatur@sas.upenn.edu