GLOBAL SOLUTIONS AND STABILITY PROPERTIES OF THE 5TH ORDER GARDNER EQUATION

MIGUEL A. ALEJO AND CHULKWANG KWAK

ABSTRACT. In this work, we deal with the initial value problem of the 5th-order Gardner equation in \mathbb{R} , presenting the local well-posedness result in $H^2(\mathbb{R})$. As a consequence of the local result, in addition to H^2 -energy conservation law, we are able to prove the global well-posedness result in $H^2(\mathbb{R})$. Finally, we present a stability result for 5th order Gardner breather solution in the Sobolev space $H^2(\mathbb{R})$.

1. INTRODUCTION

In this work, we are concerned with the *focusing* 5th order Gardner equation

$$u_t + u_{5x} + 10\mu^2 u_{3x} + 20\mu u u_{3x} + 10u^2 u_{3x} + 120\mu^3 u u_x + 180\mu^2 u^2 u_x$$
(1.1)

 $+120\mu u^3 u_x + 10u_x^3 + 40\mu u_x u_{xx} + 40u u_x u_{xx} + 30u^4 u_x = 0, \ \mu \in \mathbb{R}^+.$

This higher order Gardner equation can be obtained from the corresponding 5th order *focusing* modified Korteweg-de Vries equation (shortly, 5th mKdV)

$$v_t + (v_{4x} + 10vv_x^2 + 10v^2v_{xx} + 6v^5)_x = 0, (1.2)$$

when one considers mKdV solutions of the form $v(t, x) = \mu + u(t, x)$, with $\mu \in \mathbb{R}^+$ and a suitable spatial translation¹.

The 5th order Gardner equation (1.1), as well as the 5th mKdV equation, is a well-known completely integrable model [18, 1, 45], with infinitely many conservation laws and well-known (long-time) asymptotic behavior of its solutions obtained with the help of the inverse scattering transform [24]. As a physical model, the 5th Gardner (1.1) and the 5th mKdV (1.2) equations describe large-amplitude internal solitary waves, showing a dynamics which can look rather different from the KdV form. On the other hand, solutions of (1.1) are invariant under space and time translations. Indeed, for any $t_0, x_0 \in \mathbb{R}$, $u(t - t_0, x - x_0)$ is also a solution of both equations. Beside that, the scaling invariance is not respected by (1.1).

As seen in (1.1), the 5th order Gardner equation (1.1) contains mixed nonlinearities of 5th KdV equation

$$v_t + (v_{4x} + 5v_x^2 + 10vv_{xx} + 10v^3)_x = 0, (1.3)$$

and 5th mKdV (1.2), and hence the well-posedness theory of (1.1) is highly relevant to the well-posedness of both equations. Ponce [53], first, showed the local well-posedness of 5th KdV in $H^s(\mathbb{R})$, $s \ge 4$ via the energy method in addition to the dispersive smoothing effect and a parabolic approximation method. Later, this local result has been improved by Kwon [43], precisely, the local well-posedness in $H^s(\mathbb{R})$, $s > \frac{5}{2}$. Thereafter, Guo, Kwon and the second author [22] and Kenig and Pilod [30], independently, proved the local well-posedness in $H^s(\mathbb{R})$, $s \ge 2$. Both works were based on the short time Fourier restriction norm method [28], while an additional weight and the (frequency localized) modified energy were used to prove the crucial energy estimates, respectively. Thanks to the H^2 -level energy conservation law, the local result extended to the global one.

On the other hand, the 5th mKdV (1.2) has been studied by Linares [46]. Linares proved the local well-posedness in $H^2(\mathbb{R})$ via the contraction mapping principle in addition to the dispersive smoothing effect [33, 34]. Later, Kwon [44] improved the local result in $H^s(\mathbb{R})$, $s > \frac{3}{4}$, by using the standard Fourier restriction norm method [7] in addition to Tao's [k, Z]-multiplier norm method [54].

We also refer to [32, 35, 52, 25, 29] for the local well-posedness of for higher order KdV and mKdV equations.

Date: January 16, 2019.

²⁰¹⁰ Mathematics Subject Classification. Primary 37K15, 35Q53; Secondary 35Q51, 37K10.

Key words and phrases. Higher order Gardner equation, Global well-posedness, Breather, stability, integrability.

M. A. was partially funded by Product. CNPq grant no. 305205/2016-1 and VI PPIT-US program ref. I3C.

C. K. is supported by FONDECYT Postdoctorado 2017 Proyect No. 3170067.

¹Such a spatial translation is performed in order to provide a simpler expression of the *N*-soliton solution in Section 3. On the other hand, it is known that not only the first order linear term but also the third order term of the linear part in (1.1) are negligible in the study of the well-posedness theory compared to the fifth order term.

It is known that the Initial value problem (IVP) of 5th KdV (1.3) is a quasilinear problem in the sense that the solution map is (not uniformly) continuous, while the Cauchy problem of 5th mKdV is a semilinear problem in the sense that the flow map is Lipschitz continuous (via the Picard iteration method, and hence, is analytic). Thus, one may expect that the IVP of the 5th Gardner equation (1.1) is also a quasilinear problem due to the strong (high-low) quadratic nonlinearity. Moreover, one expects to obtain the local well-posedness in $H^2(\mathbb{R})$ (and hence the global well-posedness in $H^2(\mathbb{R})$) from [22, 30]. However, to prove the local well-posedness of the 5th Gardner equation is definitely non-trivial, thus one of aims in this work is to indeed prove the local well-posedness.

As related problems, we also refer to [8, 42, 41, 31, 55] for the well-posedness of 5th KdV, 5th mKdV and higher order equations in KdV hierarchy under the periodic boundary condition.

Concerning explicit solutions of higher order mKdV and Gardner models, Matsuno [47] proved the existence and built explicitly the N-soliton solution of the focusing mKdV hierarchy of equations by using inverse scattering technics and the bilinear Hirota decomposition. Recently, Gomes et al [19] dealt with the defocusing mKdV with NVBC and the associated defocusing Gardner hierarchy, showing multisolitonic structures. Unfortunately, many of the solutions they obtained are singular solutions (up to the kink which is in L^{∞}).

The 5th order Gardner equation (1.1), as a completely integrable system, has an infinite set of conserved quantities. Indeed some of the (first) standard conservation laws of the (1.1) are the mass

$$M[u](t) := \frac{1}{2} \int_{\mathbb{R}} u^2(t, x) dx = M[u](0), \qquad (1.4)$$

the energy

$$E_{\mu}[u](t) := \int_{\mathbb{R}} \left(\frac{1}{2} u_x^2 - 2\mu u^3 - \frac{1}{2} u^4 \right) (t, x) dx = E[u](0), \tag{1.5}$$

and the higher order energy, defined respectively in $H^2(\mathbb{R})$

$$E_{5\mu}[u](t) := \int_{\mathbb{R}} \left(\frac{1}{2} u_{xx}^2 - 10\mu u u_x^2 + 10\mu^2 u^4 - 5u^2 u_x^2 + 6\mu u^5 + u^6 \right) (t, x) dx = E_5[u](0).$$
(1.6)

1.1. Main results. We are interested in the regularity properties of the 5th order Gardner equation (1.1) and long time behavior of H^2 global solutions to (1.1).

1.1.1. Well-posedness theory. In comparison with the 5th mKdV (1.2), the nonlinearity of (1.1) consists of more terms which break the balance with the 5th order linear dispersive part of (1.1). Precisely, additional quadratic terms with three derivatives, pose technical problems, for instance, the failure of bilinear $X^{s,b}$ estimates, see Remark 1.1 below (also see Remark 2.3 in [22]). However, an analogous argument used in [22, 30] enables us to attack the initial value problem of (1.1) in H^2 .

The notion of the well-posedness, which is taken into account in this paper, is as follows:

Definition 1.1 (Well-posedness). We say that the 5th Gardner equation (1.1) is local-in-time (or locally) well-posed in $H^{s}(\mathbb{R})$, if for any R > 0 and any $u_{0} \in \{f \in H^{s}(R) : \|f\|_{H^{s}} \leq R\}$, there exist a local time T = T(R) > 0 and a unique solution u to (1.1) in $C([0,T]; H^{s}(\mathbb{R})) \cap X_{T}$, for some auxiliary space X_{T} . Moreover, the solution map $u_{0} \mapsto u(t)$ is continuous from $\{f \in H^{s}(\mathbb{R}) : \|f\|_{H^{s}} \leq R\}$ to $C([0,T]; H^{s}(\mathbb{R}))$. The local result is extended to the global one, if T > 0 is independent of R.

We are, first, going to show that the 5th order Gardner equation (1.1) is locally well-posed in H^2 via the classical energy method in addition to the short time Fourier restriction norm method. We state the local well-posedness result as follows:

Theorem 1.2. The 5th order Gardner equation (1.1) is locally well-posed in $H^{s}(\mathbb{R}), s \geq 2$.

For the proof of Theorem 1.2, we use the short-time Fourier restriction norm method in a frequency dependent time interval. This is introduced by Ionescu, Kenig and Tataru [28] in the context of KP-I equation in the Besov-type space setting, see also [38, 16] for similar ideas in the different settings. The short-time Fourier restriction norm method has been further developed in, for instance, [20, 26, 21, 22, 30, 42, 41, 23].

The main difficulty arising in (1.1) is the strong high-low bilinear interaction component of the following type²

$$(P_{\leq 0}u) \cdot (P_{high}u_{xxx}) \tag{1.7}$$

²Here P is a appropriate truncation operator in the Fourier space, thus $P_{high}u$ means the high frequency ($||\xi| \gg 1$) localized portion of u, while the frequency support of $P_{<0}u$ is in [-1, 1].

newly generated from the map $v(t,x) = \mu + u(t,x)$. The standard bilinear $X^{s,b}$ -estimates³ ($||uu_{xxx}||_{X^{s,b-1}} \leq ||u||_{X^{s,b}}^2$) fails in usual $X^{s,b}$ spaces for any $s \in \mathbb{R}$ (see Remark 1.1 below), where the $X^{s,b}$ norm is defined in (2.7), since the dispersive smoothing effect in a coherent case occurring in (1.7) is not enough to control the three derivative in the high frequency mode. The following remark provides a counter-example to show the failure of the standard bilinear estimate:

Remark 1.1 (Remark 2.3 in [22]). Similarly as the 5th KdV case, also as mentioned before, the standard $X^{s,b}$ bilinear estimate fails to hold:

$$\left\| u \partial_x^3 v \right\|_{X^{s,b-1}} \le C \left\| u \right\|_{X^{s,b}} \left\| v \right\|_{X^{s,b}}, \tag{1.8}$$

due to the following *high-low* interactions causing the coherence, for instance,

$$u(t,x) = \mathcal{F}^{-1}[\mathbf{1}_{\Omega}(\tau,\xi)](t,x) \text{ and } v(t,x) = \mathcal{F}^{-1}[\mathbf{1}_{\Sigma}(\tau,\xi)],$$

where space-time frequency sets Ω and Σ are given by⁴

$$\Omega = \{(\tau,\xi) \in \mathbb{R}^2 : |\tau - \xi^5| \le 1, N \le |\xi| \le N + 1\} \text{ and } \Sigma = \{(\tau,\xi) \in \mathbb{R}^2 : |\tau - \xi^5| \le 1, |\xi| \le 1\},$$

for fixed large frequency $N \gg 1$. Indeed, a direct calculation gives LHS of $(1.8) = NN^s$, while RHS of $(1.8) = N^s$.

However, using $X^{s,b}$ structure in a short time interval (\approx (frequency)⁻²), one reduces the contribution of high frequency with low modulation, so that one handles *high-low* interaction component (1.7) (see Remark 1.2 below and Proposition 2.7).

Remark 1.2 (Remark 2.3 in [22]). The short time $X^{s,b}$ spaces (F^s and N^s to be introduced in Section 2.2) in the interval of the length (\approx (frequency)⁻²) resolves the low-high interaction counter-example presented in Remark 1.1. The corresponding sets in this setting are given by

$$\widetilde{\Omega} = \{(\tau,\xi) \in \mathbb{R}^2 : |\tau - \xi^5| \le N^2, N \le |\xi| \le N + N^{-2}\} \text{ and } \widetilde{\Sigma} = \{(\tau,\xi) \in \mathbb{R}^2 : |\tau - \xi^5| \le 1, |\xi| \le 1\},$$

and define u and v similarly as in Remark 1.1, but with respect to $\widetilde{\Omega}$ and $\widetilde{\Sigma}$, respectively. Then, one immediately obtains for any $s \in \mathbb{R}$ that

 $\|u\partial_x^3 v\|_{N^s} \sim N^s N^3 N^{-1} N^{-2} N \sim N^s N$ and $\|u\|_{F^s} \|v\|_{F^s} \sim N^s N.$

A price to pay for the profit of the short-time argument is an energy-type estimate. However, the strong highlow interactions, where the low frequency component has the largest modulation, cause a trouble in the energy estimates when following Ionescu-Kenig-Tataru's method. A way to treat this interaction is to use a weight, which was suggested in [27] to handle the same interaction for the Benjamin-Ono equation (see also [26]). Note that the modified energy, initially introduced in [43] and further developed in [30, 42, 41, 48, 49], plays a similar role as an additional weight. See [22] and [30] for a comparison.

Note moreover that a scaling equivalence enables us to focus on small solutions to (2.1) instead of (1.1) (see Section 2). To close the energy method argument for (2.1), we gather linear, nonlinear and energy estimates,

$$\begin{cases}
\|u\|_{F^{s}(T')} \lesssim \|u\|_{E^{s}(T')} + \|\mathcal{N}_{2}(u) + \mathcal{N}_{3}(u) + \mathcal{SN}(u)\|_{N^{s}(T')}, \\
\|\mathcal{N}_{2}(u) + \mathcal{N}_{3}(u) + \mathcal{SN}(u)\|_{N^{s}(T')} \lesssim \sum_{j=2}^{5} \|u\|_{F^{s}(T)}^{j}, \\
\|u\|_{E^{s}(T')}^{2} \lesssim \|u_{0}\|_{H^{s}}^{2} + \sum_{j=3}^{6} \|u\|_{F^{s}(T)}^{j}.
\end{cases}$$
(1.9)

The continuity argument ensures a priori bound of solutions to (2.1). Moreover, a similar estimate as in (1.9) for the difference of two solutions completes the limiting argument (compactness argument). We note that the energy estimate for the difference of two solutions does not hold true in F^s spaces due to the lack of the symmetry, but hold in the intersection of the weaker (F^0) and the stronger (F^{2s}) spaces, thus the Bona-Smith argument is essential to close the compactness method.

The global well-posedness follows immediately from the above local result and the conservation of the second order energy (1.6).

Theorem 1.3. The 5th order Gardner equation (1.1) is globally well-posed in the energy space $H^2(\mathbb{R})^5$.

$$\|f\|_{X^{s,b}} = \left\| \langle \xi \rangle^s \langle \tau - \xi^5 \rangle^b \widetilde{f} \right\|_{L^2_{\tau,\xi}},$$

³The $X^{s,b}$ spaces are equipped with the norm

where \tilde{f} is the space time Fourier coefficient (also denoted by $\mathcal{F}(f)$) and $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$. For more details, see Section 2.

⁴It suffices to regard only ∂_x^5 as a linear part of (1.1), since ∂_x^3 is negligible in a sense of the dispersion effect.

⁵The persistence of regularities ensures the global well-posedness in $H^{s}(\mathbb{R}), s \geq 2$.

Remark 1.3. It is well-known that local results can be extended to the global one in the energy space without the smallness assumption for defocusing equations (for simple models), while the smallness condition is necessary for the proof of the global well-posedness in the energy space for focusing equations (the large data global well-posedness for focusing equations has a different story). However, (1.1) admits the scaling equivalence, which is slightly different from the standard scaling symmetry (or invariance), but still plays an almost same role in the local (or perturbation) theory. Thus, one has Theorem 1.3 from Theorem 1.2 in addition to the (rescaled) conservation law (1.6). See Section 2, in particular Section 2.6, for more details.

On the other hand, an observation explained in Remark 1.1 above naturally poses an interesting question: Does the flow map from data to solutions fail to be (locally) uniformly continuous for all regularities? As an immediate answer to the question, we state the following (weak) ill-posedness result, which extends Cardoso and the first author's recent result [4] to all regularities:

Theorem 1.4. The 5th order Gardner equation (1.1) is weakly ill-posed in $H^s(\mathbb{R})$, for s > 0 in the following sense: there exist $c, C > 0, 0 < T \leq 1$, and two sequences u_n and v_n of solutions to (1.1) such that

$$\sup_{n} \|u_n(t)\|_{H^s} + \sup_{n} \|v_n(t)\|_{H^s} \le C, \quad t \in [0, T]$$

and initially

$$\lim_{n \to \infty} \|u_n(0) - v_n(0)\|_{H^s} = 0$$

but for every $t \in [0, T]$

$$\liminf_{n \to \infty} \|u_n(t) - v_n(t)\|_{H^s} \ge c |\sin t| \sim c |t|.$$

Theorem 1.4 can be expected from the observation in the linear local smoothing effect [33, 34]

$$\left\|\partial_x^2 e^{-t\partial_x^5} u_0\right\|_{L_x^\infty L_t^2} \lesssim \|u_0\|_{L^2}$$

compared to the three derivatives in the quadratic nonlinearity. In other words, the local smoothing effect, which recovers only two derivatives, is not enough to handle the nonlinear term $u\partial_x^3 u$, as already seen in Remark 1.1. Such a strong high-low interaction phenomenon can be seen in other dispersive equations, for instance, the Benjamin-Ono equation (BO) and the Kadomtsev-Petviashvili I equation (KP-I). Early, constructing examples reflecting (1.7), the flow map has been shown to be not C^2 continuous [50, 51], and uniformly continuous [39, 40].

To prove Theorem 1.4, we take an argument introduced in [39] (but essentially follows from [43]) in order to construct the approximate solutions, which indeed reveals the ill-posedness phenomenon. Using the local well-posedness theory, one shows the approximate solutions are indeed "good" approximate solutions in H^s sense, $s \ge 2$. Moreover, since the equation (1.1) is completely integrable (thus it admits infinitely many conservation laws), we are able to show the same conclusion in the regularity range not only $s \ge 2$, but also 0 < s < 2 by using L^2 and H^2 conservation laws.

The strategy employed in [4] was to use Gardner breather solutions as a way to measure the regularity of the associated Cauchy problem in H^s . This allowed to find the sharp Sobolev index under which the local well-posedness of the problem is lost, meaning that the dependence of 5th order Gardner solutions upon initial data fails to be continuous. We refer to, for instant, [37, 15, 44] for analogous arguments.

Finally, together with the result in [4], we get the following

Corollary 1.5. The 5th order Gardner equation (1.1) is (weakly) ill-posed in $H^s(\mathbb{R})$, for $s \in \mathbb{R}$, in the sense of the statement given in Theorem 1.4.

As already seen above, the 5th Gardner equation (1.1) contains the mixed nonlinearities of 5th order KdV and mKdV equations (1.3)-(1.2), so that one can see both ill-posedness nature of semilinear and quasilinear equations. In the proof of Theorem 1.4, approximate solutions are constructed in the following manner: the separation of the phase shift ($\mp t$) and the dispersion effect ($\Phi_N(t)$) in (A.1) inspired by the observation on the Burgers equation. However, in low regularity Sobolev space (L^2 or below), it is not clear to see such a phenomenon, see [39, 40, 43]. Nonetheless, the cubic nonlinearity (5th mKdV nonlinearity) reveals another ill-posedness phenomenon, breaking the uniform continuity of the flow map by the self-interaction of a single high frequency wave in low regularity spaces [4]. This nature can be seen in some semilinear equations, for instance [37, 9, 10, 15, 44, 2]. The mixed nonlinearities in (1.1), thus, ensure to claim the lack of the uniform continuity of the flow map of the 5th Gardner equation (1.1) in all regularity Sobolev spaces.

1.1.2. Global stability theory. Moreover, once we have characterized the IVP for (1.1) and with respect to stability properties of specific solutions of the (1.1), we present the following stability result for the 5th order breather solutions (3.2).

Theorem 1.6. Let $\alpha, \beta \in \mathbb{R} \setminus \{0\}$ be given. Breather solutions (3.2) of the 5th order Gardner equation (1.1) are orbitally stable for H^2 perturbations, whenever the parameter $\mu \in (0, \frac{\sqrt{\alpha^2 + \beta^2}}{2})$.

For more detailed statements and background about this stability property of breather solutions, see Section 3.

2. Well-posedness results

2.1. Setting. It is well-known that the integrability of equations (fixed coefficients of the nonlinearities) is no longer important for mathematical analysis in the local well-posedness theory.

Remark 2.1. As mentioned in Section 1, (1.1) does not allow the scaling invariance. However, defining $u_{\lambda} := \lambda u(\lambda^5 t, \lambda x), \lambda > 0$, ensures an equivalence between (1.1) and

$$w_t + w_{5x} + 10\mu^2 \lambda^2 w_{3x} + \mathcal{N}_2(w) + \mathcal{N}_3(w) + \mathcal{SN}(w) = 0, \qquad (2.1)$$

where $\mathcal{N}_2(w)$ is the nonlinearity from the fifth order KdV given by

$$\mathcal{N}_{2}(w) = 20\mu\lambda w_{x}w_{xx} + 40\mu\lambda ww_{xxx} + 180\mu^{2}\lambda^{2}w^{2}w_{x}, \qquad (2.2)$$

 $\mathcal{N}_3(w)$ is the nonlinearity from the fifth order mKdV given by

$$\mathcal{N}_3(w) = 10w^2 w_{3x} + 10w_x^3 + 40w w_x u_{xx} + 30w^4 w_x$$

and $\mathcal{SN}(w)$ is the rest terms generated from the transformation $u \mapsto \mu + u$, which is weaker compared to $\mathcal{N}_2(w)$ and $\mathcal{N}_3(w)$ in some sense, given by

$$\mathcal{SN}(w) = 120\mu^3 \lambda^3 w w_x + 120\mu \lambda w^3 w_x.$$
(2.3)

That is, u_{λ} , $\lambda > 0$ is a solution to (2.1), if and only if u is a solution to (1.1). See Section 2.6 for the details.

We use the notation \tilde{f} or $\mathcal{F}(f)$ for the space-time Fourier transform of f defined by

$$\widetilde{f}(\tau,\xi) = \int_{\mathbb{R}^2} e^{-ix\xi} e^{-it\tau} f(x,t) \, dxdt$$

for any $f \in \mathcal{S}'(\mathbb{R} \times \mathbb{R})$. Similarly, we use \mathcal{F}_x (or $\hat{}$) and \mathcal{F}_t to denote the Fourier transform with respect to space and time variable respectively.

Let \mathbb{Z}_+ denote the set of nonnegative integers. For $k \in \mathbb{Z}_+$, let define dyadic intervals $I_k, k \in \mathbb{Z}_+$ as

$$I_0 = \{\xi : |\xi| \le 2\}$$
 $I_k = \{\xi : |\xi| \in [2^{k-1}, 2^{k+1}]\}$ $k \ge 1.$

Let $\eta_0 : \mathbb{R} \to [0,1]$ denote a smooth bump function supported in [-2,2] and equal to 1 in [-1,1] with the following property of regularities:

$$\partial_n^j \eta_0(\xi) = O(\eta_0(\xi)/\langle \xi \rangle^j), \quad j = 0, 1, 2,$$
(2.4)

as ξ approaches end points of the support of η . For $k \in \mathbb{Z}_+$, let

$$\chi_0(\xi) = \eta_0(\xi)$$
 and $\chi_k(\xi) = \eta_0(\xi/2^k) - \eta_0(\xi/2^{k-1}), \quad k \ge 1,$ (2.5)

and

$$\chi_{[k_1,k_2]} = \sum_{k=k_1}^{k_2} \chi_k$$
 for any $k_1 \le k_2 \in \mathbb{Z}_+$.

For the time-frequency decomposition, we use the cut-off function η_j , but the same as $\eta_j = \chi_j$, $j \in \mathbb{Z}_+$. For $k \in \mathbb{Z}$ let P_k denote the (smooth) truncation operators on $L^2(\mathbb{R})$ defined by $\widehat{P_k u}(\xi) = \chi_k(\xi)\widehat{u}(\xi)$. We also define the operators P_k on $L^2(\mathbb{R} \times \mathbb{R})$ by formulas $\mathcal{F}(P_k u)(\xi, \tau) = \chi_k(\xi)\mathcal{F}(u)(\tau, \xi)$. For $l \in \mathbb{Z}$ let

$$P_{\leq l} = \sum_{k \leq l} P_k, \quad P_{\geq l} = \sum_{k \geq l} P_k$$

For $\xi \in \mathbb{R}$, $w(\xi) = -\xi^5$ is the dispersion relation associated to the equation $(2.1)^6$. For $k \in \mathbb{Z}$ and $j \in \mathbb{Z}_+$ let

$$D_{k,j} = \{ (\tau,\xi) \in \mathbb{R} \times \mathbb{R} : \xi \in [2^{k-1}, 2^{k+1}], \tau - w(\xi) \in I_j \}, \quad D_{k,\leq j} = \bigcup_{\ell \leq j} D_{k,\ell}.$$

For $f \in L^2(\mathbb{R})$, let $W(t)f \in C(\mathbb{R}:L^2)$ be the linear solution given by

$$\mathcal{F}_{x}[W(t)f](\xi,t) = e^{itw(\xi)}\hat{f}(\xi).$$
(2.6)

⁶Originally, we have $w(\xi) = -\xi^5 + 10\mu^2\xi^3$ corresponding to the linear part of (1.1). However, for fixed μ and for large frequency $|\xi| \gg 1$, the third order term are negligible compared to the fifth order term.

2.2. Function spaces. We introduce the $X^{s,b}$ spaces associated to (2.1), which is the completion of $\mathcal{S}'(\mathbb{R}^2)$ under the norm

$$\|f\|_{X^{s,b}} = \left\| \langle \tau - w(\xi) \rangle^b \langle \xi \rangle^s \widetilde{f} \right\|_{L^2(\mathbb{R}^2)},\tag{2.7}$$

where $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$. This Fourier restriction norm method was first implemented by in its current form by Bourgain [7] and further developed by Kenig, Ponce and Vega [36] and Tao [54]. The Fourier restriction norm method turns out to be very useful in the study of low regularity theory for the dispersive equations. We denote the localized space by X_T defined by standard localization to the interval [-T, T].

As already mentioned in Section 1, the 5th Gardner equation (1.1) is a quasilinear equation where the flow map is not uniformly continuous. This fact can be seen from [43], which proves that the 5th order KdV equation (1.3)is weakly ill-posed, since this phenomenon occurs precisely in a strong interaction between low and high frequencies localized data of the form

 $(u_{\leq 0}) \cdot (\partial_x^3 u_{\gg 1})$

which is also included in the nonlinearity of 5th order Gardner (1.1). For this reason, we must focus specifically on quadratic nonlinearity to prove the local well-posedness of the 5th Gardner equation (1.1). In what follows, we briefly introduce the functions spaces used in $[22]^7$.

One of the purposes in this paper, as mentioned in Section 1, is to obtain H^2 global solutions to (1.1). Moreover, this regularity threshold is determined by the estimates of quadratic terms with three derivatives, which is already known from [22, 30]. In what follows, we only focus on obtaining the estimates of cubic terms with three derivatives in H^2 , since the cubic terms are another nontrivial and strong nonlinearities in (1.1). On the other hand, we expect that all estimates of this cubic terms can be obtained below H^2 compared to the quadratic nonlinearities, since the degree 3 of nonlinearities allows more smoothing effects in *high-low* interactions. However, we do not here explore such estimates below H^2 for our purpose.

We fix $k \in \mathbb{Z}_+$, and define the weighted Besov-type $(X^{0,\frac{1}{2},1})$ space X_k for frequency localized functions in \widetilde{I}_k ,

$$X_k = \left\{ f \in L^2(\mathbb{R}^2) : \operatorname{supp} f \subset \mathbb{R} \times I_k, \quad \|f\|_{X_k} < \infty \right\},$$

equipped with the norm

$$\|f\|_{X_k} := \sum_{j=0}^{\infty} 2^{j/2} \beta_{k,j} \|\eta_j(\tau - w(\xi))f(\xi,\tau)\|_{L^2_{\xi,\tau}},$$

where

$$\beta_{k,j} = \begin{cases} 2^{j/2}, & k = 0, \\ 1 + 2^{(j-5k)/8}, & k \ge 1. \end{cases}$$
(2.8)

Remark 2.2. The use of the weight $\beta_{k,j}$ is essential to control the localized energy for the quadratic terms in $\mathcal{N}_2(u)$, in particular, the *high-low* interaction components, where the low frequency component has the largest modulation. See Lemma 2.10. Moreover, it enables us to avoid the logarithmic divergence in H^2 appearing in the energy estimates for the cubic nonlinearities in $\mathcal{N}_3(u)$, see Remark 2.10 and Propositions 2.12 and 2.13.

Remark 2.3. An opposite effect of the use of the weight is to worsen the high \times high \rightarrow low interactions in the nonlinear estimates for the quadratic terms in $\mathcal{N}_2(u)$

$$P_{\leq 0}(P_{high}u \cdot P_{high}v_{xxx}).$$

However, thanks to the representation of the quadratic nonlinearities as the compact, conservative form, i.e., $c_1 \partial_x u \partial_x^2 u + c_2 u \partial_x^3 u = c'_1 \partial_x (\partial_x u \partial_x u) + c'_2 \partial_x (u \partial_x^2 u)$, one derivative is removed, and hence we are able to balance both purposes.

Remark 2.4. Finally, the choice of a parameter $\frac{1}{8}$ in the weight for high frequency can be replaced by any parameter in [1/8, 3/16]. However, another choice of parameter is not able to improve the result, since the essential effect of the weight occurs in the *high-low* interactions, where the low frequency part has the largest modulation, as mentioned before.

At each frequency 2^k , we define functions spaces based on X_k , uniformly on the 2^{-2k} time scale.

$$F_k = \left\{ f \in L^2(\mathbb{R}^2) : \operatorname{supp} f \subset \mathbb{R} \times I_k, \quad \|f\|_{F_k} < \infty \right\},$$

equipped with the norm

$$\|f\|_{F_k} = \sup_{t_k \in \mathbb{R}} \left\| \mathcal{F}[f \cdot \eta_0(2^{2k}(t-t_k))] \right\|_{X_k}$$

⁷The basic method is similar to that used in [30], but it is chosen to avoid complicated calculations in the energy estimate.

and

$$N_k = \left\{ f \in L^2(\mathbb{R}^2) : \operatorname{supp} f \subset \mathbb{R} \times I_k, \quad \|f\|_{N_k} < \infty \right\},$$

equipped with the norm

$$\|f\|_{N_k} = \sup_{t_k \in \mathbb{R}} \left\| (\tau - \omega(\xi) + i2^{2k})^{-1} \mathcal{F}[f \cdot \eta_0(2^{2k}(t - t_k))] \right\|_{X_k}.$$

The standard way to construct localized spaces gives, for $T \in (0, 1]$, that

$$F_k(T) = \{ f \in C([-T,T]:L^2) : \|f\|_{F_k(T)} = \inf_{\widetilde{f}=f \text{ in } [-T,T]\times\mathbb{R}} \|\widetilde{f}\|_{F_k} \},$$

$$N_k(T) = \{ f \in C([-T,T]:L^2) : \|f\|_{N_k(T)} = \inf_{\widetilde{f}=f \text{ in } [-T,T]\times\mathbb{R}} \|\widetilde{f}\|_{N_k} \}.$$

We collect all pieces of spaces introduced above at dyadic frequency 2^k in the Littlewood-Paley way. For $s \ge 0$ and $T \in (0, 1]$, we define function spaces for solutions and nonlinear terms:

$$F^{s}(T) = \left\{ u: \|u\|_{F^{s}(T)}^{2} = \sum_{k=0}^{\infty} 2^{2sk} \|P_{k}(u)\|_{F_{k}(T)}^{2} < \infty \right\},$$
$$N^{s}(T) = \left\{ u: \|u\|_{N^{s}(T)}^{2} = \sum_{k=0}^{\infty} 2^{2sk} \|P_{k}(u)\|_{N_{k}(T)}^{2} < \infty \right\}.$$

In order to take the short time structure for IVP of (1.1), it is required to define the energy space as follows: for $s \ge 0$ and $u \in C([-T, T] : H^{\infty})$

$$\|u\|_{E^{s}(T)}^{2} = \|P_{\leq 0}(u(0))\|_{L^{2}}^{2} + \sum_{k \geq 1} \sup_{t_{k} \in [-T,T]} 2^{2sk} \|P_{k}(u(t_{k}))\|_{L^{2}}^{2}.$$

Remark 2.5. The short time Fourier restriction norm method used in this work was introduced by Ionescu, Kenig and Tataru [28], where the local well-posedness of KP-I equation in the energy space was proved, and further developed in [20, 26, 21, 22, 30, 42, 41, 23] and references therein. We also refer to [38, 16] for different formulas of short time analysis.

For the extension argument of functions in the spaces introduced above, we follow from [28] to define the set S_k of k-acceptable time multiplication factors for any $k \in \mathbb{Z}_+$:

$$S_{k} = \{m_{k} : \mathbb{R} \to \mathbb{R} : \|m_{k}\|_{S_{k}} = \sum_{j=0}^{10} 2^{-2jk} \|\partial^{j}m_{k}\|_{L^{\infty}} < \infty\}.$$

Direct estimates using the definitions and (2.10) show that for any $s \ge 0$ and $T \in (0, 1]$

$$\begin{cases} \left\| \sum_{k \in \mathbb{Z}_{+}} m_{k}(t) \cdot P_{k}(u) \right\|_{F^{s}(T)} \lesssim \left(\sup_{k \in \mathbb{Z}_{+}} \|m_{k}\|_{S_{k}} \right) \cdot \|u\|_{F^{s}(T)} ,\\ \left\| \sum_{k \in \mathbb{Z}_{+}} m_{k}(t) \cdot P_{k}(u) \right\|_{N^{s}(T)} \lesssim \left(\sup_{k \in \mathbb{Z}_{+}} \|m_{k}\|_{S_{k}} \right) \cdot \|u\|_{N^{s}(T)} ,\\ \left\| \sum_{k \in \mathbb{Z}_{+}} m_{k}(t) \cdot P_{k}(u) \right\|_{E^{s}(T)} \lesssim \left(\sup_{k \in \mathbb{Z}_{+}} \|m_{k}\|_{S_{k}} \right) \cdot \|u\|_{E^{s}(T)} .\end{cases}$$

We end this subsection with the following important lemma.

Lemma 2.1 (Properties of X_k). Let $k, l \in \mathbb{Z}_+$ with $l \leq 5k$ and $f_k \in X_k$. Then

$$\sum_{j=l+1}^{\infty} 2^{j/2} \beta_{k,j} \left\| \eta_j(\tau - \omega(\xi)) \int_{\mathbb{R}} |f_k(\tau',\xi)| 2^{-l} (1 + 2^{-l} |\tau - \tau'|)^{-4} d\tau' \right\|_{L^2}$$

$$+ 2^{l/2} \left\| \eta_{\leq l}(\tau - \omega(\xi)) \int_{\mathbb{R}} |f_k(\tau',\xi)| 2^{-l} (1 + 2^{-l} |\tau - \tau'|)^{-4} d\tau' \right\|_{L^2} \lesssim \|f_k\|_{X_k}.$$

$$(2.9)$$

In particular, if $t_0 \in \mathbb{R}$ and $\gamma \in \mathcal{S}(\mathbb{R})$, then

$$\left\| \mathcal{F}[\gamma(2^{l}(t-t_{0})) \cdot \mathcal{F}^{-1}(f_{k})] \right\|_{X_{k}} \lesssim \|f_{k}\|_{X_{k}}.$$
 (2.10)

Proof. See [22] for the proof.

2.3. L^2 -block estimates. For $x, y \in \mathbb{R}_+$, $x \leq y$ means that there exists C > 0 such that $x \leq Cy$, and $x \sim y$ means $x \leq y$ and $y \leq x$. We also use \leq_s and \sim_s as similarly, where the implicit constants depend on s. Let $a_1, a_2, a_3, a \in \mathbb{R}$. The quantities $a_{max} \geq a_{sub} \geq a_{thd} \geq a_{min}$ can be conveniently defined to be the maximum, sub-maximum, third-maximum and minimum values of a_1, a_2, a_3, a respectively.

For $\xi_1, \xi_2 \in \mathbb{R}$, let denote the (quadratic) resonance function by

$$H = H(\xi_1, \xi_2) = w(\xi_1) + w(\xi_2) - w(\xi_1 + \xi_2) = \frac{5}{2}\xi_1\xi_2(\xi_1 + \xi_2)(\xi_1^2 + \xi_2^2 + (\xi_1 + \xi_2)^2).$$

Similarly, for $\xi_1, \xi_2, \xi_3 \in \mathbb{R}$, let

$$G(\xi_1, \xi_2, \xi_3) = w(\xi_1) + w(\xi_2) + w(\xi_3) - w(\xi_1 + \xi_2 + \xi_3)$$

= $\frac{5}{2}(\xi_1 + \xi_2)(\xi_2 + \xi_3)(\xi_3 + \xi_1)(\xi_1^2 + \xi_2^2 + \xi_3^2 + (\xi_1 + \xi_2 + \xi_3)^2)$ (2.11)

be the (cubic) resonance function. Such resonance functions play an important role in the nonlinear $X^{s,b}$ -type estimates.

Let $f, g, h \in L^2(\mathbb{R}^2)$ be compactly supported functions. We define a quantity by

$$J_2(f,g,h) = \int_{\mathbb{R}^4} f(\zeta_1,\xi_1)g(\zeta_2,\xi_2)h(\zeta_1+\zeta_2+H(\xi_1,\xi_2),\xi_1+\xi_2) d\xi_1 d\xi_2 d\zeta_1 \zeta_2$$

The change of variables in the integration yields

$$J_2(f,g,h) = J_2(g^*,h,f) = J_2(h,f^*,g)$$

where $f^*(\zeta,\xi) = f(-\zeta,-\xi)$. From the identities

 $\xi_1 + \xi_2 = \xi_3$ and $(\tau_1 - w(\xi_1)) + (\tau_2 - w(\xi_2)) = (\tau_3 - w(\xi_3)) + H(\xi_1, \xi_2)$

on the support of $J_2(f^{\sharp}, g^{\sharp}, h^{\sharp})$, where $f^{\sharp}(\tau, \xi) = f(\tau - w(\xi), \xi)$ with the property $||f||_{L^2} = ||f^{\sharp}||_{L^2}$, we see that $J(f^{\sharp}, g^{\sharp}, h^{\sharp})$ vanishes unless

$$2^{k_{max}} \sim 2^{k_{med}} \gtrsim 1$$
 and $2^{j_{max}} \sim \max(2^{j_{med}}, |H|).$

For compactly supported functions $f_i \in L^2(\mathbb{R} \times \mathbb{R}), i = 1, 2, 3, 4$, we define

$$J_3(f_1, f_2, f_3, f_4) = \int_* f_1(\zeta_1, \xi_1) f_2(\zeta_2, \xi_2) f_3(\zeta_3, \xi_3) f_4(\zeta_1 + \zeta_2 + \zeta_3 + G(\xi_1, \xi_2, \xi_3), \xi_1 + \xi_2 + \xi_3)$$

where the $\int_* = \int_{\mathbb{R}^6} d\xi_1 d\xi_2 d\xi_3 d\zeta_1 d\zeta_2 d\zeta_3$. From the identities

$$\xi_1 + \xi_2 + \xi_3 = \xi_4 \quad \text{and} \quad (\tau_1 - w(\xi_1)) + (\tau_2 - w(\xi_2)) + (\tau_3 - w(\xi_3)) = (\tau_4 - w(\xi_4)) + G(\xi_1, \xi_2, \xi_3)$$

on the support of $J_3(f_1^{\sharp}, f_2^{\sharp}, f_3^{\sharp}, f_4^{\sharp})$, we see that $J_3(f_1^{\sharp}, f_2^{\sharp}, f_3^{\sharp}, f_4^{\sharp})$ vanishes unless

$$\frac{2^{k_{max}} \sim 2^{k_{sub}}}{2^{j_{max}} \sim \max(2^{j_{sub}}, |G|)},$$
(2.12)

where $|\xi_i| \sim 2^{k_i}$ and $|\zeta_i| \sim 2^{j_i}$, i = 1, 2, 3, 4. A direct calculation shows

$$|J_3(f_1, f_2, f_3, f_4)| = |J_3(f_2, f_1, f_3, f_4)| = |J_3(f_3, f_2, f_1, f_4)| = |J_3(f_1^*, f_2^*, f_4, f_3)|.$$

We give L^2 -block estimates for the quadratic and cubic nonlinearities. The bi- and tri-linear L^2 -block estimates for the 5th order equations have already been introduced and used in several works, we refer to [14, 13, 22, 30, 42, 41, 11, 12].

Lemma 2.2. Let $k_i \in \mathbb{Z}, j_i \in \mathbb{Z}_+, i = 1, 2, 3$. Let $f_{k_i, j_i} \in L^2(\mathbb{R} \times \mathbb{R})$ be nonnegative functions supported in $[2^{k_i-1}, 2^{k_i+1}] \times I_{j_i}$.

(a) For any $k_1, k_2, k_3 \in \mathbb{Z}$ with $|k_{max} - k_{min}| \leq 5$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$, then we have

$$J_2(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}) \lesssim 2^{j_{min}/2} 2^{j_{med}/4} 2^{-\frac{3}{4}k_{max}} \prod_{i=1}^3 \|f_{k_i,j_i}\|_{L^2}$$

(b) If $2^{k_{min}} \ll 2^{k_{med}} \sim 2^{k_{max}}$, then for all i = 1, 2, 3 we have

$$J_2(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}) \lesssim 2^{(j_1+j_2+j_3)/2} 2^{-3k_{max}/2} 2^{-(k_i+j_i)/2} \prod_{i=1}^3 \|f_{k_i,j_i}\|_{L^2}.$$

(c) For any $k_1, k_2, k_3 \in \mathbb{Z}$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$, then we have

$$J_2(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}) \lesssim 2^{j_{min}/2} 2^{k_{min}/2} \prod_{i=1}^3 \|f_{k_i,j_i}\|_{L^2}$$

Proof. We refer to [14, 13, 30] for the proof.

Corollary 2.3. Assume $k_i \in \mathbb{Z}$ and $j_i \in \mathbb{Z}_+$, i = 1, 2, 3 and $f_{k_i, j_i} \in L^2(\mathbb{R} \times \mathbb{R})$ be functions supported in D_{k_i, j_i} , i = 1, 2.

(a) For any $k_1, k_2, k_3 \in \mathbb{Z}$ with $|k_{max} - k_{min}| \leq 5$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$, then we have

$$\|\mathbf{1}_{D_{k_3,j_3}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2})\|_{L^2} \lesssim 2^{j_{min}/2} 2^{j_{med}/4} 2^{-\frac{3}{4}k_{max}} \prod_{i=1}^2 \|f_{k_i,j_i}\|_{L^2}$$

(b) If $2^{k_{min}} \ll 2^{k_{med}} \sim 2^{k_{max}}$, then for all i = 1, 2, 3 we have

$$\|\mathbf{1}_{D_{k_3,j_3}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2})\|_{L^2} \lesssim 2^{(j_1+j_2+j_3)/2} 2^{-3k_{max}/2} 2^{-(k_i+j_i)/2} \prod_{i=1}^2 \|f_{k_i,j_i}\|_{L^2}.$$

(c) For any $k_1, k_2, k_3 \in \mathbb{Z}$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$, then we have

$$\|\mathbf{1}_{D_{k_3,j_3}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2})\|_{L^2} \lesssim 2^{j_{min}/2} 2^{k_{min}/2} \prod_{i=1}^2 \|f_{k_i,j_i}\|_{L^2}$$

Lemma 2.4. Let $k_i \in \mathbb{Z}$ and $j_i \in \mathbb{Z}_+$, i = 1, 2, 3, 4. Let $f_{k_i, j_i} \in L^2(\mathbb{R} \times \mathbb{R})$ be nonnegative functions supported in $I_{j_i} \times [2^{k_i-1}, 2^{k_i+1}]$.

(a) For any $k_i \in \mathbb{Z}$ and $j_i \in \mathbb{Z}_+$, i = 1, 2, 3, 4, we have

$$J_3(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}, f_{k_4,j_4}) \lesssim 2^{(j_{min}+j_{thd})/2} 2^{(k_{min}+k_{thd})/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}.$$

(b) Let $k_{thd} \le k_{max} - 10$.

(b-1) If $(k_i, j_i) = (k_{thd}, j_{max})$ for i = 1, 2, 3, 4, we have

$$J_3(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}, f_{k_4,j_4}) \lesssim 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_{max}} 2^{k_{thd}/2} 2^{-j_{max}/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}$$

(b-2) If $(k_i, j_i) \neq (k_{thd}, j_{max})$ for i = 1, 2, 3, 4, we have

$$J_3(f_{k_1,j_1}, f_{k_2,j_2}, f_{k_3,j_3}, f_{k_4,j_4}) \lesssim 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_{max}} 2^{k_{min}/2} 2^{-j_{max}/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}.$$

Proof. We refer to [30, 41] for the proof. In [41], the second author established (cubic) L^2 -block estimates for functions $f_{k_i,j_i} \in L^2(\mathbb{R} \times \mathbb{Z})$, but the proof, here, is almost identical and easier, see [30].

Corollary 2.5. Let $k_i \in \mathbb{Z}$ and $j_i \in \mathbb{Z}_+$, i = 1, 2, 3, 4. Let $f_{k_i, j_i} \in L^2(\mathbb{R} \times \mathbb{R})$ be nonnegative functions supported in D_{k_i, j_i} .

(a) For any $k_i \in \mathbb{Z}$ and $j_i \in \mathbb{Z}_+$, i = 1, 2, 3, 4, we have

$$\|\mathbf{1}_{D_{k_4,j_4}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2}*f_{k_3,j_3})\|_{L^2} \lesssim 2^{(j_{min}+j_{thd})/2} 2^{(k_{min}+k_{thd})/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}$$

(b) Let $k_{thd} \leq k_{max} - 10$. (b-1) If $(k_i, j_i) = (k_{thd}, j_{max})$ for i = 1, 2, 3, 4, we have

$$\|\mathbf{1}_{D_{k_4,j_4}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2}*f_{k_3,j_3})\|_{L^2} \lesssim 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_{max}} 2^{k_{thd}/2} 2^{-j_{max}/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}.$$

(b-2) If $(k_i, j_i) \neq (k_{thd}, j_{max})$ for i = 1, 2, 3, 4, we have

$$\|\mathbf{1}_{D_{k_4,j_4}}(\tau,\xi)(f_{k_1,j_1}*f_{k_2,j_2}*f_{k_3,j_3})\|_{L^2} \lesssim 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_{max}} 2^{k_{min}/2} 2^{-j_{max}/2} \prod_{i=1}^4 \|f_{k_i,j_i}\|_{L^2}.$$

We end this subsection introducing the Strichartz estimates for the family of the fifth-order operators $\{e^{t\partial_x^5}\}_{t=-\infty}^{\infty}$.

Lemma 2.6 (Strichartz estimates for $e^{t\partial_x^5}$ operator [17]). Assume that $-1 < \sigma \leq \frac{3}{2}$ and $0 \leq \theta \leq 1$. Then there exists C > 0 depending on σ and θ such that

$$\left\| D^{\frac{\sigma\theta}{2}e^{t\partial_x^5}\varphi} \right\|_{L^q_tL^p_x} \le C \left\|\varphi\right\|_{L^2}$$

for $\varphi \in L^2$, where $p = \frac{2}{1-\theta}$ and $q = \frac{10}{\theta(\sigma+1)}$. In particular, we have $\left\| e^{t\partial_x^5} P_k \varphi \right\|_{L^6_{t,x}} \lesssim 2^{-k/2} \left\| P_k \varphi \right\|_{L^2}, \quad k \ge 1.$

2.4. Nonlinear estimates. We first recall from [22] the bilinear estimates as follows:

Proposition 2.7 (Nonlinear estimates for $\mathcal{N}_2(u)$, [22]). (a) If $s \ge 1$, $T \in (0, 1]$, and $u, v \in F^s(T)$ then $\|\mathcal{N}_2(u)\|_{N^s(T)} \lesssim \|u\|_{F^s(T)}^2 + \|u\|_{F^s(T)}^3$.

and

$$\begin{aligned} \|\mathcal{N}_{2}(u) - \mathcal{N}_{2}(v)\|_{N^{s}(T)} &\lesssim \left(\|u\|_{F^{s}(T)} + \|v\|_{F^{s}(T)}\right)\|u - v\|_{F^{s}(T)} + \left(\|u\|_{F^{s}(T)}^{2} + \|v\|_{F^{s}(T)}^{2}\right)\|u - v\|_{F^{s}(T)}. \end{aligned}$$
(b) If $T \in (0, 1]$, $u, v \in F^{0}(T) \cap F^{2}(T)$, then

$$\|\mathcal{N}_{2}(u) - \mathcal{N}_{2}(v)\|_{N^{0}(T)} &\lesssim \left(\|u\|_{F^{2}(T)} + \|v\|_{F^{2}(T)}\right)\|u - v\|_{F^{0}(T)} + \left(\|u\|_{F^{2}(T)}^{2} + \|v\|_{F^{2}(T)}^{2}\right)\|u - v\|_{F^{0}(T)}. \end{aligned}$$

Proof. The proof for the quadratic term, we refer to [22]. On the other hand, one can easily control the cubic term in $\mathcal{N}_2(u)$ rather than not only the quadratic term in $\mathcal{N}_2(u)$, but also the cubic term in $\mathcal{N}_3(u)$, since the cubic term in $\mathcal{N}_2(u)$ contains only one (total) derivative. Hence we omit the details, but one can capture the estimates in the proof of Proposition 2.8.

Proposition 2.8 (Nonlinear estimates for $\mathcal{N}_3(u)$). (a) If $s \ge 2$, $T \in (0, 1]$, and $u, v \in F^s(T)$ then

$$\|\mathcal{N}_{3}(u)\|_{N^{s}(T)} \lesssim \|u\|_{F^{s}(T)}^{3} + \|u\|_{F^{s}(T)}^{5}.$$
(2.13)

and

(b)

$$\|\mathcal{N}_{3}(u) - \mathcal{N}_{3}(v)\|_{N^{s}(T)} \lesssim \left(\|u\|_{F^{s}(T)}^{2} + \|v\|_{F^{s}(T)}^{2}\right)\|u - v\|_{F^{s}(T)} + \left(\|u\|_{F^{s}(T)}^{4} + \|v\|_{F^{s}(T)}^{4}\right)\|u - v\|_{F^{s}(T)}.$$

If $T \in (0, 1], u, v \in F^{0}(T) \cap F^{2}(T)$, then

$$\|\mathcal{N}_{3}(u) - \mathcal{N}_{3}(v)\|_{N^{0}(T)} \lesssim \left(\|u\|_{F^{2}(T)}^{2} + \|v\|_{F^{2}(T)}^{2}\right)\|u - v\|_{F^{0}(T)} + \left(\|u\|_{F^{2}(T)}^{4} + \|v\|_{F^{2}(T)}^{4}\right)\|u - v\|_{F^{0}(T)}.$$

Proof. We first consider the cubic term in $\mathcal{N}_3(u)$. From the support property (2.12) in addition to (2.11), we know that

$$\max(|\tau_j - w(\xi_j)|; j = 1, 2, 3, 4) \gtrsim |(\xi_1 + \xi_2)(\xi_1 + \xi_3)(\xi_2 + \xi_3)|(\xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2).$$
(2.14)

From the definition of N_k norm, the left-hand side of (cubic terms in) (2.13) is bounded by

$$\sup_{t_{k_4} \in \mathbb{R}} \left\| (\tau_4 - w(\xi_4) + i2^{2k_4})^{-1} 2^{3k_4} \mathbf{1}_{I_{k_4}}(\xi) \mathcal{F} \left[\eta_0 \left(2^{2k_4 - 2}(t - t_{k_4}) \right) P_{k_1} u \right] \right.$$

$$\left. \left. + \mathcal{F} \left[\eta_0 \left(2^{2k_4 - 2}(t - t_{k_4}) \right) P_{k_2} u \right] * \mathcal{F} \left[\eta_0 \left(2^{2k_4 - 2}(t - t_{k_4}) \right) P_{k_3} u \right] \right\|_{X_{k_4}}$$

$$(2.15)$$

We set $u_{k_i} = \mathcal{F}\left[\eta_0\left(2^{2k_4-2}(t-t_{k_4})\right)P_{k_i}u\right]$, i = 1, 2, 3. We decompose each u_{k_i} into $u_{k_i,j_i}(\tau,\xi) = u_{k_i}(\tau,\xi)\eta_{j_i}(\tau-w(\xi))$ with usual modification like $f_{\leq j}(\tau) = f(\tau)\eta_{\leq j}(\tau-\mu(n))$. Then, (2.15) is bounded by

$$\sum_{j_4 \ge 0} \frac{2^{3k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \sum_{j_1, j_2, j_3 \ge 2k_4} \left\| \mathbf{1}_{D_{k_4, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}$$

We, instead of Corollary 2.5, use the following observation to control

$$\left\| \mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3}) \right\|_{L^2}$$

for particular case: Lemma 2.6 yields

$$\begin{aligned} \left\| \mathcal{F}^{-1}[u_{k_{i},j_{i}}] \right\|_{L^{6}} &= \left\| \int e^{it\tau} e^{ix\xi} e^{itw(\xi)} u_{k_{i},j_{i}}^{\sharp}(\tau,\xi) \, d\xi d\tau \right\|_{L^{6}} \\ &\lesssim \int \left\| \int e^{ix\xi} e^{itw(\xi)} u_{k_{i},j_{i}}^{\sharp}(\tau,\xi) \, d\xi \right\|_{L^{6}} \, d\tau \\ &\lesssim 2^{-k_{i}/2} 2^{j_{i}/2} \| u_{k_{i},j_{i}}^{\sharp} \|_{L^{2}}, \end{aligned}$$

$$(2.16)$$

where $u_{k_i,j_i}^{\sharp}(\tau,\xi) = u_{k_i,j_i}(\tau+w(\xi),\xi)$ with $\|u_{k_i,j_i}^{\sharp}\|_{L^2} = \|u_{k_i,j_i}\|_{L^2}$. With this, Plancherel's theorem and the Hölder inequality give

$$\left\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\right\|_{L^2} \lesssim 2^{-(k_1+k_2+k_3)/2} 2^{(j_1+j_2+j_3)/2} \prod_{i=1}^3 \|u_{k_i,j_i}\|_{L^2}.$$
(2.17)

Case I. (high-high \Rightarrow high). Let $k_4 \geq 20$ and $|k_1 - k_4|, |k_2 - k_4|, |k_3 - k_4| \leq 5$. Applying (2.17) to $\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\|_{L^2}$, one has

$$(2.15) \lesssim \sum_{j_4 \ge 0} \frac{2^{3k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \sum_{j_1, j_2, j_3 \ge 2k_4} 2^{-k_4/2} 2^{(j_1+j_2+j_3)/2} \prod_{i=1}^3 \|u_{k_i, j_i}\|_{L^2}.$$

Note that $\beta_{k_4,j_4} \sim 1$ when $0 \leq j_4 \leq 5k$. Let denote the summand by \mathcal{M}_I , i.e.,

$$\mathcal{M}_{I} := \frac{2^{3k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}.$$
(2.18)

Then, we know

$$\mathcal{M}_I \lesssim 2^{j_4/2} 2^{-k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 0 \le j_4 \le 2k_4,$$
$$\mathcal{M}_I \lesssim 2^{-j_4/2} 2^{3k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 2k_4 \le j_4 \le 5k_4$$

and

$$\mathcal{M}_I \lesssim 2^{-j_4/2} 2^{(j_4 - 5k_4)/8} 2^{3k_4/2} 2^{(j_1 + j_2 + j_3)/2}$$
, when $5k_4 \le j_4$

Performing summations over j_i , i = 1, 2, 3, 4, we have

$$(2.15) \lesssim 2^{k_4/2} \prod_{i=1}^3 \|P_{k_i}u\|_{F_{k_i}}.$$

Indeed, we have from the definition of X_k -norm and (2.4) in [21] that

$$\sum_{j_1 \ge 2k_4} 2^{j_1/2} \|u_{k_1,j_1}\|_{L^2} \lesssim \sum_{j_1 > 2k_4} 2^{j_1/2} \beta_{k_1,j_1} \left\| \eta_{j_1}(\tau - \omega(\xi)) \int_{\mathbb{R}} |\bar{u}_{k_1}(\xi,\tau')| \cdot 2^{-2k_4} (1 + 2^{-2k_4} |\tau - \tau'|)^{-4} d\tau' \right\|_{L^2} \\ + 2^{(2k_4)/2} \left\| \eta_{\le 2k_4}(\tau - \omega(\xi)) \int_{\mathbb{R}} |\bar{u}_{k_1}(\xi,\tau')| 2^{-2k_4} (1 + 2^{-2k_4} |\tau - \tau'|)^{-4} d\tau' \right\|_{L^2} \\ \lesssim \|u_{k_1}\|_{X_{k_1}} \lesssim \|P_{k_1}u\|_{F_{k_1}},$$

where $\bar{u}_{k_1} = \mathcal{F}[P_{k_1}u \cdot \eta_0(2^{2k_1}(t-t_{k_1}))].$

Remark 2.6. As seen in the proof of **Case I** (also for other cases except for the case when the resulting frequency (ξ_4) is not the maximum frequency), the weight β_{k_4,j_4} does not play any role in the estimates, hence is negligible. See **Case I** and **Case III** (below) for comparison.

Case II. (high-high-low \Rightarrow high). Let $k_4 \ge 20$, $|k_2 - k_4|$, $|k_3 - k_4| \le 5$ and $k_1 \le k_4 - 10^8$. In this case, we have $j_{max} \ge 5k_4$ due to (2.14). The exactly same argument used in **Case I** (but use Corollary 2.5 (a) instead of (2.17) to control $\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\|_{L^2}$) gives better result as follows:

$$(2.15) \lesssim 2^{k_1/2} \prod_{i=1}^3 \|u_{k_i}\|_{X_{k_i}} \lesssim 2^{k_1/2} \prod_{i=1}^3 \|P_{k_i}u\|_{F_{k_i}}.$$

The last inequality holds true, thanks to (2.10), more precisely,

$$\|u_{k_1}\|_{X_{k_1}} = \left\| \mathcal{F}\left[\eta_0 \left((2^{2k_4-2}(t-t_{k_4})) \cdot P_{k_1} u \cdot \eta_0 \left(2^{2k_1}(t-t_{k_4}) \right) \right] \right\|_{X_{k_1}} \lesssim \|P_{k_1} u\|_{F_{k_1}} \,.$$

We omit the details.

Case III. (high-high \Rightarrow low). Let $k_3 \ge 20$, $|k_1 - k_3|, |k_2 - k_3| \le 5$ and $k_4 \le k_3 - 10$.

⁸Thanks to the symmetry of frequencies, our assumption that ξ_1 is the minimum frequency does not lose of the generality.

Remark 2.7. The trade-off of the use of the short time advantage (also, the use of the weight as in (2.8)) is to worsen some interactions for which the resulting frequency is lower than (at least) one of others, in particular, high-high-high \Rightarrow low and high-high-low \Rightarrow low interaction components. More precisely, in the case of the high-high-high \Rightarrow low, the time interval of length 2^{-2k_4} , on which the N_{k_4} -norm is taken, is longer than the interval of length 2^{-2k_4} , on which F_{k_i} -norm is taken, i = 1, 2, 3. In order to cover whole intervals of length 2^{-2k_4} in the estimates, one needs to divide the time interval of length 2^{-2k_4} into $2^{2k_3-2k_4}$ intervals of length $2^{-2^{2k_3}}$. Let choose $\gamma : \mathbb{R} \to [0, 1]$ (a kind of the partition of unity), which is a smooth function supported in [-1, 1] with $\sum_{m \in \mathbb{Z}} \gamma^3(x-m) \equiv 1$. Then, the left-hand side of (cubic terms in) (2.13) is bounded by (instead of (2.15))

$$\sup_{t_{k}\in\mathbb{R}} 2^{3k_{3}} \left\| (\tau_{4} - w(\xi_{4}) + i2^{2k_{4}})^{-1} \mathbf{1}_{I_{k_{4}}} \cdot \sum_{|m| \leq C2^{2k_{3}-2k_{4}}} \mathcal{F}[\eta_{0}(2^{2k_{4}}(t-t_{k}))\gamma(2^{2k_{3}}(t-t_{k})-m)P_{k_{1}}u] \\
\times \mathcal{F}[\eta_{0}(2^{2k_{4}}(t-t_{k}))\gamma(2^{2k_{3}}(t-t_{k})-m)P_{k_{2}}u] \times \mathcal{F}[\eta_{0}(2^{2k_{4}}(t-t_{k}))\gamma(2^{2k_{3}}(t-t_{k})-m)P_{k_{3}}u] \right\|_{X_{k_{4}}}.$$
(2.19)

The analogous procedure will be applied to the estimate of high-high-low \Rightarrow low interaction component below.

When $k_4 = 0$, (2.19) is bounded by

$$\sum_{j_4 \ge 0} 2^{5k_3} \sum_{j_1, j_2, j_3 \ge 2k_3} \left\| \mathbf{1}_{D_{0, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}$$

due to (2.8).

When $0 \le j_4 \le 5k_3 - 5$ or $\le 5k_3 + 5 \le j_4$, we apply Corollary 2.5 (a) to $\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\|_{L^2}$ to obtain

$$(2.19) \lesssim \left(\sum_{0 \le j_4 \le 5k_3 - 5} + \sum_{5k_3 + 5 \le j_4}\right) 2^{5k_3} \sum_{j_1, j_2, j_3 \ge 2k_3} 2^{(j_{min} + j_{thd})/2} 2^{k_3/2} \prod_{i=1}^3 \|u_{k_i, j_i}\|_{L^2}$$

We know $j_4 \neq j_{max}$ in the former case, while $j_4 = j_{max}$ and $2^{j_{max}} \sim 2^{j_{med}} \gg |H|$ in the latter case.

On the other hand, when $5k_3 - 5 \le j_4 \le 5k_3 + 5$ $(2^{j_4} = 2^{j_{max}} \sim |H| \gg 2^{j_{med}})$, we use (2.17). Then, similarly as the previous cases, we have (when $k_4 = 0$)⁹

$$(2.19) \lesssim 2^{\frac{7}{2}k_3} \prod_{i=1}^3 \|P_{k_i}u\|_{F_{k_i}} \,.$$

When $k_4 \neq 0$, similarly as above, (2.19) is bounded by

$$\sum_{j_4 \ge 0} \frac{2^{5k_3 - 2k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} 2^{(k_3 + k_4)/2} \sum_{j_1, j_2, j_3 \ge 2k_3} 2^{(j_{min} + j_{thd})/2} \prod_{i=1}^3 \|u_{k_i, j_i}\|_{L^2},$$

thanks to Corollary 2.5 (a), except for the case when $5k_3 - 5 \le j_4 \le 5k_3 + 5$. Let denote the summand by \mathcal{M}_{III} , similarly as in (2.18) i.e.,

$$\mathcal{M}_{III} := \frac{2^{5k_3 - 2k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} 2^{(k_3 + k_4)/2} 2^{(j_{min} + j_{thd})/2}.$$

If $2k_3 < 5k_4$, we know

$$\begin{aligned} \mathcal{M}_{III} &\lesssim 2^{j_4} 2^{2k_3} 2^{-7k_4/2} 2^{(j_1+j_2+j_3)/2}, & \text{when} \quad 0 \leq j_4 \leq 2k_4, \\ \mathcal{M}_{III} &\lesssim 2^{2k_3} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}, & \text{when} \quad 2k_4 \leq j_4 \leq 2k_3, \\ \mathcal{M}_{III} &\lesssim 2^{-j_4/2} 2^{3k_3} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}, & \text{when} \quad 2k_3 \leq j_4 \leq 5k_4, \\ \mathcal{M}_{III} &\lesssim 2^{-j_4/2} 2^{(j_4-5k_4)/8} 2^{3k_3} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}, & \text{when} \quad 5k_4 \leq j_4 \leq 5k_3 - 4 \end{aligned}$$

and

$$\mathcal{M}_{III} \lesssim 2^{-j_4/2} 2^{(j_4 - 5k_4)/8} 2^{3k_3} 2^{-3k_4/2} 2^{(j_1 + j_2 + j_3)/2}, \text{ when } 5k_3 + 4 \le j_4.$$

Otherwise (when $5k_4 < 2k_3$), the estimates of \mathcal{M}_{III} on $2k_4 \leq j_4 \leq 5k_3 - 4$ are replaced by

$$\mathcal{M}_{III} \lesssim 2^{2k_3} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 2k_4 \le j_4 \le 5k_4,$$
$$\mathcal{M}_{III} \lesssim 2^{(j_4-5k_4)/8} 2^{2k_3} 2^{-3k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 5k_4 \le j_4 \le 2k_3$$

$$\mathcal{M}_{III} \lesssim 2^{-j_4/2} 2^{(j_4 - 5k_4)/8} 2^{3k_3} 2^{-3k_4/2} 2^{(j_1 + j_2 + j_3)/2}, \quad \text{when} \quad 2k_3 \le j_4 \le 5k_3 - 4k_3 \le j_4 \le 2k_3 - 4k_4 \le 2k_3 \le 2k_4 \le 2k_3 + 2k_4 \le 2k_4 \le$$

⁹One can see that the worst bound comes from the low frequency with high modulation case $(j_4 = j_{max} > j_{med} + 5)$.

On the other hand, when $5k_3 - 5 \le j_4 \le 5k_3 + 5$, we use (2.17) to obtain

$$(2.20) \lesssim 2^{5k_3 - 2k_4} 2^{-5k_3/2} 2^{\frac{5}{8}(k_3 - k_4)} 2^{-3k_3/2} \sum_{j_1, j_2, j_3 \ge 2k_3} \prod_{i=1}^{3} 2^{j_i} \|u_{k_i, j_i}\|_{L^2}.$$

Summing over j_i , i = 1, 2, 3, 4, one has

$$(2.19) \lesssim C_1(k_3, k_4) \prod_{i=1}^3 \|P_{k_i} u\|_{F_{k_i}}, \qquad (2.20)$$

where

$$C_1(k_3, k_4) = \begin{cases} 2\frac{5}{5}k_3, & 2k_3 < 5k_4, \\ 2\frac{9}{4}k_3 2^{-\frac{17}{8}k_4}, & 5k_4 \le 2k_3. \end{cases}$$

Remark 2.8. A direct computation in the cubic term in $\mathcal{N}_3(u)$, one has

$$40uu_xu_{xx} + 10u^2u_{xxx} + 10u_x^3 = 10(u^2u_{xx})_x + 10(uu_x^2)_x.$$

Then, one can reduce 2^{3k_3} in (2.19) by $2^{2k_3+k_4}$, and hence obtain a better result. However, our regularity threshold is s = 2, and hence we, here, do not explore the trilinear estimates in lower regularity.

Case IV. (high-low-low \Rightarrow high). Let $k_4 \ge 20$, $|k_3 - k_4| \le 5$ and $k_1, k_2 \le k_4 - 10$. Without loss of generality, we may assume that $k_1 \le k_2$, thanks to the symmetry. Similarly as the **Case I**, it is enough to consider

$$\sum_{j_4 \ge 0} \frac{2^{3k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \sum_{j_1, j_2, j_3 \ge 2k_4} \left\| \mathbf{1}_{D_{k_4, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}.$$
(2.21)

Let denote the summand in (2.21) by \mathcal{M}_{IV} , i.e.,

$$\mathcal{M}_{IV} := \frac{2^{3k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \left\| \mathbf{1}_{D_{k_4, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}.$$

We further split this case into three cases: Case IV-a $k_2 = 0$, Case IV-b $k_1 = 0$ and $k_2 \neq 0$, and Case IV-c $k_1 \neq 0$.

Case IV-a. $k_2 = 0$. We do not distinguish Corollary (2.5) (b.1) and (b.2), since $2^{k_{min}} \leq 2^{k_{thd}} \leq 1$. Then, from Corollary 2.5 (b), we have¹⁰

$$\mathcal{M}_{IV} \lesssim 2^{j_4} 2^{-2k_4} \prod_{i=1}^3 2^{j_i/2} \|u_{k_i,j_i}\|_{L^2}, \text{ when } 0 \le j_4 \le 2k_4$$

and

$$\mathcal{M}_{IV} \lesssim 2^{-j_4/2} \beta_{k_4, j_4} \prod_{i=1}^3 2^{j_i/2} \|u_{k_i, j_i}\|_{L^2}, \text{ when } 2k_4 \le j_4.$$

Summing over j_i , i = 1, 2, 3, 4, one has

$$(2.15) \lesssim \prod_{i=1}^{3} \|u_{k_i}\|_{F_{k_i}}.$$

Case IV-b. $k_1 = 0$ and $k_2 \neq 0$. Note that we have $j_{max} \geq 4k_4 + k_2$ due to (2.14). We use Corollary 2.5 (b.1) (the worst case occurring in $j_2 = j_{max}$) when $0 \leq j_4 \leq 4k_4 + k_2 - 5$, and (b.2) ($j_2 = j_{max}$ never happens) when $4k_4 + k_2 - 5 \leq j_4$ to control $\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\|_{L^2}$ in \mathcal{M}_{IV} , then we have

$$\mathcal{M}_{IV} \lesssim 2^{j_4} 2^{-3k_4} \prod_{i=1}^3 2^{j_i/2} \|u_{k_i,j_i}\|_{L^2}, \quad \text{when} \quad 0 \le j_4 \le 2k_4,$$
$$\mathcal{M}_{IV} \lesssim 2^{-k_4} \prod_{i=1}^3 2^{j_i/2} \|u_{k_i,j_i}\|_{L^2}, \quad \text{when} \quad 2k_4 \le j_4 \le 4k_4 + k_2 - 5$$

$$\mathcal{M}_{IV} \lesssim 2^{-j_4/2} \beta_{k_4, j_4} 2^{k_4} \prod_{i=1}^3 2^{j_i/2} \|u_{k_i, j_i}\|_{L^2}, \text{ when } 4k_4 + k_2 - 5 \le j_4.$$

¹⁰We use, here, $2^{j_{max}} \ge 2^{2k_4}$ to deal with a maximum modulation, since our purpose is to obtain the local well-posedness only in $H^s(\mathbb{R})$, $s \ge 2$. However, one may obtain the better result by performing a delicate calculation in addition to $2^{j_{max}} \ge |H|$, instead of $2^{j_{max}} \ge 2^{2k_4}$. For the same reason, so the high-high-low \Rightarrow low case below as well.

Summing over j_i , i = 1, 2, 3, 4, one has

$$(2.15) \lesssim k_4 2^{-k_4} \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}$$

Case IV-c. $k_1 \neq 0$. Similarly as **Case IV-a** (if $|k_1 - k_2| < 5$ with Corollary 2.5 (b.2)) or **Case IV-b** (if $k_1 < k_2 - 5$), we have at most

$$(2.15) \lesssim 2^{k_1/2} \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}.$$

Case V. (high-high-low \Rightarrow low). Let $k_3 \ge 20$, $|k_2 - k_3| \le 5$ and $k_1, k_4 \le k_3 - 10$. We first divide this case into two cases: **Case V-a** $k_4 = 0$ and **Case V-b** $k_4 \ne 0$.

Case V-a. $k_4 = 0$. From Remark 2.7, it suffices to consider

$$\sum_{j_4 \ge 0} 2^{5k_3} \sum_{j_1, j_2, j_3 \ge 2k_3} \left\| \mathbf{1}_{D_{0,j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}$$
(2.22)

due to (2.8). If $k_1 = 0$, by using Corollary 2.5 (a), we have

$$(2.22) \lesssim 2^{4k_3} \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}.$$
(2.23)

More precisely, when $|\xi_2 + \xi_3| \ll \xi_3^{-2}$ (equivalently $|H| \ll 2^{2k_3}$) we know

$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_3}$$
, when $0 \le j_4 \le 2k_3$

and

$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3)/2} 2^{-j_4/2}$$
, when $2k_3 \le j_4$.

When $|\xi_2 + \xi_3| \gtrsim \xi_3^{-2}$ (equivalently $|H| \gtrsim 2^{2k_3}$), we know

$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3+j_4)/2} 2^{-2k_3}, \text{ when } 1 \le 2^{j_4} \le 2^{2k_3}$$
$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3)/2} 2^{-j_4/2}, \text{ when } 2^{2k_3} \le 2^{j_4} \le |H|/2$$
$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3)/2} 2^{-k_3}, \text{ when } |H|/2 \le 2^{j_4} \le 3|H|/2$$

and

$$2^{(j_{min}+j_{thd})/2} \le 2^{(j_1+j_2+j_3)/2} 2^{-j_4/2}$$
, when $3|H|/2 \le 2^{j_4}$.

Note that the number of j_4 is finite (≤ 10) when $|H|/2 \leq 2^{j_4} \leq 3|H|/2$. Thus, the summation over j_i , i = 1, 2, 3, 4, yields (2.23).

Otherwise $(k_1 \neq 0)$, similarly as **Case IV-b**, we have

$$(2.22) \lesssim 2^{3k_3} 2^{k_1/2} \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}.$$

Case V-b. $k_4 \neq 0$. Similarly, it is enough to consider

$$\sum_{j_4 \ge 0} \frac{2^{5k_3 - 2k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \sum_{j_1, j_2, j_3 \ge 2k_3} \left\| \mathbf{1}_{D_{k_4, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}.$$
(2.24)

Let denote the summand in (2.24) by \mathcal{M}_V , i.e.,

$$\mathcal{M}_V := \frac{2^{5k_3 - 2k_4} 2^{j_4/2} \beta_{k_4, j_4}}{\max(2^{j_4}, 2^{2k_4})} \left\| \mathbf{1}_{D_{k_4, j_4}} \cdot (u_{k_1, j_1} * u_{k_2, j_2} * u_{k_3, j_3}) \right\|_{L^2}$$

If $k_1 = 0$, we use Corollary 2.5 (b) to control $\|\mathbf{1}_{D_{k_4,j_4}} \cdot (u_{k_1,j_1} * u_{k_2,j_2} * u_{k_3,j_3})\|_{L^2}$. Then, we know

$$\mathcal{M}_V \lesssim 2^{j_4} 2^{k_3} 2^{-9k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 0 \le j_4 \le 2k_4,$$
$$\mathcal{M}_V \lesssim 2^{k_3} 2^{-5k_4/2} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 2k_4 \le j_4 \le 5k_4,$$
$$\mathcal{M}_V \lesssim 2^{3k_3/2} 2^{-3k_4} 2^{(j_1+j_2+j_3)/2}, \quad \text{when} \quad 5k_4 \le j_4 \le 4k_3 + k_4 - 5$$

$$\mathcal{M}_V \lesssim 2^{-j_4/2} 2^{(j_4 - 5k_4)/8} 2^{3k_3} 2^{-3k_4/2} 2^{(j_1 + j_2 + j_3)/2}$$
, when $4k_3 + k_4 - 5 \le j_4$

Summing over j_i , i = 1, 2, 3, 4, one has

$$(2.22) \lesssim \max\left(k_3 2^{3k_3/2} 2^{-3k_4}, 2^{3k_3/2} 2^{-2k_4}\right) \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}.$$

Otherwise $(k_1 \neq 0)$, analogous arguments as **Case V-b** (for $|k_1 - k_4| \ge 5$ case) and **Case V-a**, in particular $k_1 = 0$ case, (for $|k_1 - k_4| \le 5$ case) can be applied, and hence we have (but, omit the details)

$$(2.22) \lesssim C_2(k_1, k_3, k_4) \prod_{i=1}^3 \|u_{k_i}\|_{F_{k_i}}$$

where

$$C_{2}(k_{1},k_{3},k_{4}) = \begin{cases} 2^{3(k_{3}-k_{4})/2} \max\left(k_{3}2^{-k_{4}},1\right), & k_{1} \le k_{4}-5, \\ 2^{3(k_{3}-k_{4})/2} \max\left(k_{3}2^{-k_{4}/2}2^{k_{1}/8},2^{-k_{4}}2^{(k_{1}-k_{4})/8}\right), & k_{4} \le k_{1}-5, \\ 2^{4k_{3}}2^{k_{1}}, & |k_{1}-k_{4}| \le 5 \end{cases}$$

The estimate of *low-low* \Rightarrow *low* interaction component can be easily obtained, and hence we omit the details. On the other hand, the estimate of quintic term in $\mathcal{N}_3(u)$ will be taken into account in the estimate of $\mathcal{SN}(u)$ below. Thus, by collecting all, we complete the proof.

Proposition 2.9 (Nonlinear estimates for $\mathcal{SN}(u)$). (a) If $s \ge 2$, $T \in (0,1]$, and $u, v \in F^s(T)$ then

$$\|\mathcal{SN}(u)\|_{N^{s}(T)} \lesssim \|u\|_{F^{s}(T)}^{2} + \|u\|_{F^{s}(T)}^{4}.$$

and

(b)

$$\|\mathcal{SN}(u) - \mathcal{SN}(v)\|_{N^{s}(T)} \lesssim \left(\|u\|_{F^{s}(T)} + \|v\|_{F^{s}}(T) + \|u\|_{F^{s}(T)}^{3} + \|v\|_{F^{s}(T)}^{3}\right) \|u - v\|_{F^{s}(T)}$$

If $T \in (0, 1], u, v \in F^{0}(T) \cap F^{2}(T)$, then

$$\|\mathcal{SN}(u) - \mathcal{SN}(v)\|_{N^0(T)} \lesssim \left(\|u\|_{F^2(T)} + \|v\|_{F^2}(T) + \|u\|_{F^2(T)}^3 + \|v\|_{F^2(T)}^3 \right) \|u - v\|_{F^0(T)}.$$

Proof. The quadratic term in $\mathcal{SN}(u)$ can be easily treated compared to one in $\mathcal{N}_2(u)$, due to a less number of derivatives, similarly as the cubic term in $\mathcal{N}_2(u)$ compared to one in $\mathcal{N}_3(u)$. The rest of the proof (also for the quadratic and cubic terms in $\mathcal{SN}(u)$ and $\mathcal{N}_3(u)$, respectively) is based on the following direct computation

$$\left\|\mathbf{1}_{D_{k_{\ell},j_{\ell}}}\cdot(u_{k_{1},j_{1}}*\cdots*u_{k_{\ell-1},j_{\ell-1}})\right\|_{L^{2}} \lesssim 2^{-(k_{max}+k_{med})/2}2^{-(j_{max}+j_{med})/2}2^{k_{\ell}/2}\prod_{i=1}^{\ell-1}2^{j_{i}/2}2^{k_{i}/2}\left\|u_{k_{i},j_{i}}\right\|_{L^{2}},$$

which can be obtained by Cauchy-Schwarz inequality. Due to a less number of derivatives (indeed, one (total) derivative) in SN(u), the analogous (but much simpler) argument used in the proof of Proposition 2.8 immediately yields Proposition 2.9. In particular, the total derivative form enables us to drop one derivative taken in a high frequency mode, see Remark 2.8. We omit the details.

2.5. Energy estimates. Assume that $u, \mathcal{G} \in C([-T, T]; L^2)$ satisfy

$$\begin{cases} u_t + u_{5x} = \mathcal{G}, & (x,t) \in \mathbb{R} \times (-T,T) \\ u(0,x) = u_0(x) \end{cases}$$

A direct calculation gives

$$\sup_{|t_k| \le T} \|u(t_k)\|_{L^2}^2 \le \|u_0\|_{L^2}^2 + \sup_{|t_k| \le T} \left| \int_{\mathbb{R} \times [0, t_k]} u \cdot \mathcal{G} \, dx dt \right|.$$
(2.25)

To control the second term (for $\mathcal{N}_2(u)$, $\mathcal{N}_3(u)$ and $\mathcal{SN}(u)$) of the right-hand side of (2.25), we need following lemmas.

Lemma 2.10 ([22]). Let $T \in (0, 1]$ and $k_1, k_2, k_3 \in \mathbb{Z}_+$.

(a) Assume $k_1 \le k_2 \le k_3$ and $|k_3 - k_1| \le 5$, $u_i \in F_{k_i}(T)$, i = 1, 2, 3. Then

$$\left| \int_{\mathbb{R}\times[0,T]} u_1 u_2 u_3 \, dx dt \right| \lesssim 2^{-\frac{7}{4}k_3} \prod_{i=1}^3 \|u_i\|_{F_{k_i}(T)} \, .$$

(b) Assume $k_1 \leq k_2 \leq k_3$ and $k_3 \geq 10$, $2^{k_1} \ll 2^{k_2} \sim 2^{k_3}$ and $u_i \in F_{k_i}(T)$, i = 1, 2, 3. If $k_1 \geq 1$, then

$$\left| \int_{\mathbb{R} \times [0,T]} u_1 u_2 u_3 \, dx dt \right| \lesssim 2^{-2k_3 - \frac{1}{2}k_1} \prod_{i=1}^3 \|u_i\|_{F_{k_i}(T)}$$

If $k_1 = 0$, then

$$\left| \int_{\mathbb{R}\times[0,T]} u_1 u_2 u_3 \, dx dt \right| \lesssim 2^{-k_3} \prod_{i=1}^3 \|u_i\|_{F_{k_i}(T)},$$
$$\int_{\mathbb{R}\times[0,T]} (\partial_x u_1) u_2 u_3 \, dx dt \right| \lesssim 2^{-2k_3} \prod_{i=1}^3 \|u_i\|_{F_{k_i}(T)}.$$

(c) Assume $k_1 \leq k - 10$. Then

$$\left| \int_{\mathbb{R}\times[0,T]} P_k(u) P_k(\partial_x^3 u \cdot P_{k_1} v) \, dx dt \right| \lesssim 2^{\frac{1}{2}k_1} \, \|P_{k_1} v\|_{F_{k_1}(T)} \sum_{|k'-k| \le 10} \|P_{k'} u\|_{F_{k'}(T)}^2 \, .$$

(d) Under the same condition as in (c), we have

$$\left| \int_{\mathbb{R}\times[0,T]} P_k(u) P_k(\partial_x^2 u \cdot P_{k_1} \partial_x v) \, dx dt \right| \lesssim 2^{\frac{1}{2}k_1} \, \|P_{k_1}v\|_{F_{k_1}(T)} \sum_{|k'-k| \le 10} \|P_{k'}u\|_{F_{k'}(T)}^2$$

Lemma 2.11. Let $T \in (0,1]$, $k_1, k_2, k_3, k_4 \in \mathbb{Z}_+$, and $u_i \in F_{k_i}(T)$, i = 1, 2, 3, 4. We further assume $k_1 \le k_2 \le k_3 \le k_4$ with $k_4 \ge 10$. Then

(a) For $|k_1 - k_4| \le 5$, we have

$$\left| \int_{[0,T] \times \mathbb{R}} u_1 u_2 u_3 u_4 \, dx dt \right| \lesssim 2^{-k_4/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.26}$$

(b) For $|k_2 - k_4| \le 5$ and $k_1 \le k_4 - 10$, we have

$$\left| \int_{[0,T] \times \mathbb{R}} u_1 u_2 u_3 u_4 \, dx dt \right| \lesssim 2^{-k_4} 2^{k_1/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.27}$$

(c) Let $|k_3 - k_4| \leq 5$ and $k_2 \leq k_4 - 10$. In general, we have

$$\left| \int_{[0,T] \times \mathbb{R}} u_1 u_2 u_3 u_4 \, dx dt \right| \lesssim 2^{-k_4} 2^{k_1/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.28}$$

In particular, if $k_1 = 0$ and $k_2 \ge 1$, we have

$$\left| \int_{[0,T]\times\mathbb{R}} u_1 u_2 u_3 u_4 \, dx dt \right| \lesssim 2^{-2k_4} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.29}$$

If $0 < k' \le k_4 - 10$, we have

$$\left| \int_{[0,T]\times\mathbb{R}} P_{k'}(u_1u_2)u_3u_4 \, dxdt \right| \lesssim 2^{-2k_4} 2^{k_1/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.30}$$

If k' = 0, we have

$$\left| \int_{[0,T]\times\mathbb{R}} P_0(\partial_x(u_1u_2)) u_3 u_4 \, dx dt \right| \lesssim 2^{-2k_4} 2^{k_1/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)} \,. \tag{2.31}$$

Remark 2.9. In [30], a weaker estimate (2.28) is enough to control the cubic term with one derivative, while, in this paper, (2.29)-(2.31) are necessary to control the cubic terms with three derivatives. On the other hand, under the periodic boundary condition, (2.28) is optimal, due to the lack of *smoothing effect*. We refer to [30] and [41] for a part of proof and the periodic case (also for the comparison), respectively.

Proof of Lemma 2.11. We only prove (a) and (c). The proof of (b) is analogous to the proof of (c), thus we omit it. For part (b), see [30, 41].

(a) We apply a similar argument as in Remark 2.7 to the interval [0, T]. Let choose $\rho : \mathbb{R} \to [0, 1]$ to make a partition of unity, that is, $\sum_{m \in \mathbb{Z}} \rho^4(x - m) \equiv 1$, for all $x \in \mathbb{R}$. It follows that

$$\left| \int_{[0,T]\times\mathbb{R}} u_1 u_2 u_3 u_4 \, dx dt \right| \lesssim \sum_{|m| \lesssim 2^{2k_4}} \left| \int_{\mathbb{R}^2} \prod_{i=1}^4 \left(\rho(2^{2k_4}t - m) \mathbf{1}_{[0,T]}(t) u_i \right) \, dx dt \right|.$$

 Set

$$A := \left\{ m : \rho(2^{2k_4}t - m)\mathbf{1}_{[0,T]}(t) \text{ non-zero and } \neq \rho(2^{2k_4}t - m) \right\}$$

Note that $|A| \leq 4$. We split

$$\sum_{|m| \lesssim 2^{2k_4}} = \sum_{m \in A} + \sum_{m \in A^c}$$

It suffices to show (a) on the second summation, since otherwise, the same argument in addition to

$$\sup_{j \in \mathbb{Z}_+} 2^{j/2} \left\| \eta_j(\tau - w(\xi)) \cdot \mathcal{F}[\mathbf{1}_{[0,1]}(t)\rho(2^{2k}t - m)u] \right\|_{L^2} \lesssim \left\| \rho(2^{2k}t - m)\widetilde{u} \right\|_{X_k}$$

gives better result, thanks to the absence of 2^{2k_4} (see [21, 30] for the details).

On the second summation $(\sum_{m \in A^c})$, we can ignore $\mathbf{1}_{[0,T]}(t)$. Similarly as in Section 2.4, let $u_{k_i} = \mathcal{F}[\rho(2^{2k_4}t - m)\hat{u}_i(\xi_i)]$ and $u_{k_i,j_i} = \eta_{j_i}(\tau_i - w(\xi_i))u_{k_i}$, i = 1, 2, 3, 4. Parseval's identity and (2.9) yield

$$\sum_{m \in A^c} \left| \int_{\mathbb{R}^2} \prod_{i=1}^4 \left(\rho(2^{2k_4}t - m) \mathbf{1}_{[0,T]}(t) u_i \right) \, dx dt \right| \lesssim \sup_{m \in A^c} 2^{2k_4} \sum_{j_1, j_2, j_3, j_4 \ge 2k_4} \left| \int_{\mathbb{R}^2} \prod_{i=1}^4 \mathcal{F}^{-1}[u_{k_i, j_i}](t, x) \, dx dt \right|.$$
(2.32)

Hölder inequality and (2.16) ensure

$$\left| \int_{\mathbb{R}^2} \prod_{i=1}^4 \mathcal{F}^{-1}[u_{k_i,j_i}](t,x) \, dx dt \right| \lesssim \|u_{k_1,j_1}\|_{L^2} \prod_{i=2}^4 \|\mathcal{F}^{-1}[u_{k_i,j_i}]\|_{L^6} \\ \lesssim 2^{-3k_4/2} 2^{-j_1/2} \prod_{i=1}^4 2^{j_i/2} \|u_{k_i,j_i}\|_{L^2} \,,$$

together with (2.32), one concludes (2.26) and we complete the proof.

(c) The proof of (2.28) can be found in [30]. The proof of (2.29) follows the proof of (2.28) with a modification $j_{max} \ge 4k_4 + k_2 - 10$ instead of $j_{max} \ge 2k_4$. Thus we omit the detail. Note that

$$\begin{cases} 2^{k'} \sim 2^{k_2}, & \text{if } k_1 \le k_2 - 4, \\ 2^{k'} \ll 2^{k_2}, & \text{if } |k_1 - k_2| \le 4, \end{cases}$$
(2.33)

since $2^{k'} \sim |\xi_1 + \xi_2| \lesssim 2^{k_2 11}$.

We first show (2.30). Similarly as above, it suffices to estimate on $\sum_{m \in A^c}$. Using $2^{j_{max}} \gtrsim 2^{4k_4} 2^{k'}$ in addition to (2.33), one immediately obtains from Lemma 2.4 (b) that

LHS of (2.30)
$$\lesssim 2^{-2k_4} 2^{k_1/2} \sum_{j_1, j_2, j_3, j_4 \ge 2k_4} \prod_{i=1}^4 2^{j_i/2} \|u_{k_i, j_i}\|_{L^2} \lesssim 2^{-2k_4} 2^{k_1/2} \prod_{i=1}^4 \|u_i\|_{F_{k_i}(T)}.$$

The proof of (2.31) is analogous to the proof of Lemma 4.1 (b) (in particular (4.6)) in [22]. The left-hand side of (2.31) can be replaced by

$$\sum_{\ell \leq 0} \left| \int_{[0,T] \times \mathbb{R}} P_{\ell}(u_1 u_2) u_3 u_4 \, dx dt \right|.$$

If $k_2 = 0$, similarly as the proof of (2.30), we have

LHS of (2.31)
$$\lesssim 2^{-2k_4} \prod_{i=1}^{4} \|u_i\|_{F_{k_i}(T)}$$

Otherwise $(k_1 \ge 1)^{12}$, the same argument in the proof of (2.30) yields

LHS of (2.31)
$$\lesssim 2^{-2k_4} 2^{k_1/2} \prod_{i=1}^4 ||u_i||_{F_{k_i}(T)}$$

Thus, we complete the proof.

¹¹The case $|\xi_1 + \xi_2| \sim 2^{k_2}$, when $|k_1 - k_2| \leq 4$, exists, if both ξ_1 and ξ_2 have same sign. However, under this condition, one has the same conclusion as (2.29).

¹²The case $|\xi_1 + \xi_2| \leq 1$ cannot happen when $k_1 = 0$ and $k_2 \geq 1$.

Remark 2.10. Using the weight, one can have at least $2^{k_4/4}$ more derivative gain, while $2^{\frac{5}{8}k_1}$ derivative loss occurs. Indeed, a direct computation gives

$$\sum_{j_1 \ge 2k_4} 2^{j_1/2} \|u_{k_1,j_1}\|_{L^2} \lesssim 2^{-\frac{2k_4-5k_1}{8}} \|u_1\|_{F_{k_1}(T)}.$$

Such derivative gain may be helpful to avoid the occurrence of the logarithmic divergence in H^2 -energy estimates (see [41]). Moreover, the derivative loss in low frequencies is not big in H^2 , so be handled in H^2 . This approach may be applied to LWP of the fifth-order mKdV $H^2(\mathbb{T})$ (improvement of [41], in authors' forthcoming project).

Proposition 2.12. Let $s \ge 2$ and $T \in (0,1]$. Then, for the solution $u \in C([-T,T]; H^{\infty}(\mathbb{T}))$ to (2.1), we have

$$\|u\|_{E^{s}(T)}^{2} \lesssim \|u_{0}\|_{H^{s}}^{2} + \sum_{j=3}^{6} \|u\|_{F^{s}(T)}^{j}.$$

$$(2.34)$$

Proof of Proposition 2.12. The definition of the $E^{s}(T)$ norm says

$$||u||_{E^{s}(T)}^{2} - ||P_{\leq 0}(u_{0})||_{L^{2}}^{2} = \sum_{k \geq 1} \sup_{t_{k} \in [-T,T]} 2^{2sk} ||P_{k}(u(t_{k}))||_{L^{2}}^{2}.$$

Then, we immediately have

$$2^{2sk} \|P_k(u(t_k))\|_{L^2}^2 - 2^{2sk} \|P_k(u_0)\|_{L^2}^2 \lesssim 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(u) P_k(\mathcal{N}_2(u)) \, dx dt \right| + 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(u) P_k(\mathcal{N}_3(u)) \, dx dt \right| + 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(u) P_k(\mathcal{SN}(u)) \, dx dt \right| =: I_1(k) + I_2(k) + I_3(k),$$

thanks to (2.25). Proposition 4.2 (in addition to Remark 4.3) in [22] yields

$$\sum_{k \ge 1} I_1(k) \lesssim \|u\|_{F^s(T)}^3 + \|u\|_{F^s(T)}^4$$

for $s \geq \frac{5}{4}$.

We now focus on $I_2(k)$. Note that a direct calculation gives

$$40uu_x u_{xx} + 10u^2 u_{xxx} + 10u_x^3 = 10(u^2)_x u_{xx} + 10(u^2 u_{xx})_x + 10u_x^3$$

We split $I_2(k)$ (in particular cubic part in $\mathcal{N}_3(u)$) into $I_{2,1} + I_{2,2} + I_{2,3}$, where

$$I_{2,1} := 2^{2sk} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(u) P_k((u^2)_x u_{xx}) \, dx dt \right|,$$
$$I_{2,2} := 2^{2sk} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(u) P_k((u^2 u_{xx})_x) \, dx dt \right|,$$

and

$$I_{2,3} := 2^{2sk} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(u) P_k(u_x^3) \, dx dt \right|.$$

We first estimate $I_{2,1}$. We further decompose $I_{2,1}$ as follows:

$$\begin{split} I_{2,1} \lesssim & 2^{2sk} \sum_{k' \le k-10} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(u) P_k(P_{k'}(u^2)_x u_{xx}) \, dx dt \right| \\ & + 2^{2sk} \sum_{k' > k-10, k_3 \ge 0} \left| \int_{\mathbb{R} \times [0,t_k]} P_k^2(u) P_{k'}((u^2)_x) P_{k_3}(u_{xx}) \, dx dt \right| \\ & =: I_{2,1,1} + I_{2,1,2}. \end{split}$$

Note that $k' \leq k_2 + 10$ in $I_{2,1,1}$. Lemma 2.11 yields

$$I_{2,1,1} \lesssim 2^{2sk} \sum_{k_1 \le k_2 \le k-10} 2^{k_1/2} 2^{k_2} \|P_{k_1}u\|_{F_{k_1}(T)} \|P_{k_2}u\|_{F_{k_2}(T)} \sum_{|k_0-k| \le 10} \|P_{k_0}u\|_{F_{k_0}(T)}^2$$

$$\lesssim 2^{2sk} \sum_{|k_0-k| \le 10} 2^{\frac{5}{2}k_0} \|P_{k_0}u\|_{F_{k_0}(T)}^4$$

$$\lesssim 2^{2sk} \sum_{k_1 \ge k+10} 2^{\frac{3}{2}k_1} \|P_{k_1}u\|_{F_{k_1}(T)}^2 \sum_{|k_0-k| \le 10} \|P_{k_0}u\|_{F_{k_0}(T)}^2$$

$$\lesssim 2^{2sk} \|P_ku\|_{F_{k}(T)}^2 \|u\|_{F_{k_1}(T)}^2$$

We divide the summation over k_3 in $I_{2,1,2}$ by $\sum_{k_3 \leq k-10} + \sum_{|k_3-k| \leq 10} + \sum_{k_3 \geq k+10}$. Then, by the support property, we know that the integral vanishes unless $|k'-k| \leq 10$ on the first and second summations and $k' \geq k+10$ on the last summation. Note on the last summation that $|k'-k_3| \leq 10$.

On the first summation, the following cases of k_1 and k_2 (assuming $k_1 \leq k_2$ by the symmetry) are possible:

- (1) $k_1 \le k 10$ and $|k_2 k| \le 10$ (2) $|k_1 - k_2| \le 10$ and $k_2 \ge k + 10$
- (3) $|k_1 k_2| \le 10$ and $|k_2 k| \le 10$

It suffices to assume in the first case that $k_1 \leq k_3$, since two derivatives are taken in the k_3 -frequency mode. We use (2.29) and (2.28) with the use of the weight (see Remark 2.10) when $k_1 = 0$ and $k_1 \geq 1$, respectively, to estimate $I_{2,1,2}$, precisely, $I_{2,1,2}$ is bounded by

$$2^{2sk} \|P_0 u\|_{F_0(T)} \sum_{k_3 \le k-10} 2^{k_3} \|P_{k_3} u\|_{F_{k_3}(T)} \sum_{|k_2 - k| \le 10} \|P_{k_2} u\|_{F_{k_2}(T)}^2 + 2^{2sk} \sum_{1 \le k_1 \le k_3 \le k-10} 2^{\frac{3}{2}k_1} 2^{2k_3} 2^{-\frac{1}{8}k} \|P_{k_1} u\|_{F_{k_1}(T)} \|P_{k_3} u\|_{F_{k_3}(T)} \sum_{|k_2 - k| \le 10} \|P_{k_2} u\|_{F_{k_2}(T)}^2 \lesssim 2^{2sk} \|P_k u\|_{F_k(T)}^2 \|u\|_{F^2(T)}^2$$

Under the second case, by (2.30), $I_{2,1,2}$ is bounded by

$$2^{2sk} \sum_{\substack{k_3 \le k - 10 \\ |k_1 - k_2| \le 10 \\ k_2 \ge k + 10}} 2^{\frac{5}{2}k_3} 2^{k_2 - 2k_2} \prod_{j=1}^{3} \left\| P_{k_j} u \right\|_{F_{k_j}(T)} \left\| P_k u \right\|_{F_k(T)}$$

$$\lesssim \max(2^{\frac{3}{2}k}, 2^{(s-1)k}) \left\| P_k u \right\|_{F_k(T)} \left\| u \right\|_{F^s(T)}^3,$$

for $s \ge 0$.

Under the last case, by (2.27), $I_{2,1,2}$ is bounded by

$$2^{2sk} \sum_{\substack{k_3 \le k - 10 \\ |k_1 - k_2| \le 10 \\ |k_2 - k| \le 10}} 2^{\frac{5}{2}k_3} \prod_{j=1}^{3} \left\| P_{k_j} u \right\|_{F_{k_j}(T)} \left\| P_k u \right\|_{F_k(T)} \\ \lesssim 2^{2sk} \left\| P_k u \right\|_{F_k(T)}^2 \left\| u \right\|_{F^{\frac{5}{4}}(T)}^2.$$

On the second summation, the possible cases of k_1 and k_2 are same as before. Using (2.27), (2.30) and (2.26), one concludes that $I_{2,1,2}$ is bounded by

$$2^{2sk} \sum_{\substack{k_1 \leq k-10 \\ |k_2-k_3| \leq 10 \\ |k_2-k| \leq 10}} 2^{2k} 2^{k_1/2} \prod_{j=1}^3 \left\| P_{k_j} u \right\|_{F_{k_j}(T)} \left\| P_k u \right\|_{F_k(T)} + 2^{2sk} \sum_{\substack{|k_3-k| \leq 10 \\ |k_1-k_2| \leq 10 \\ k_2 \geq k+10}} 2^{\frac{3}{2}k_2} \prod_{j=1}^3 \left\| P_{k_j} u \right\|_{F_{k_j}(T)} \left\| P_k u \right\|_{F_k(T)} + 2^{2sk} \sum_{\substack{|k_1-k_2| \leq 10 \\ |k_3-k| \leq 10 \\ |k_2-k| \leq 10}} 2^{\frac{5}{2}k} \prod_{j=1}^3 \left\| P_{k_j} u \right\|_{F_{k_j}(T)} \left\| P_k u \right\|_{F_k(T)} \lesssim 2^{2sk} \left\| P_k u \right\|_{F_k(T)}^2 \left\| u \right\|_{F_2(T)}^2.$$

On the last summation, the following cases of k_1 and k_2 (assuming $k_1 \leq k_2$ by the symmetry) are possible:

- (1) $k_1 \le k_2 10$ and $|k_2 k_3| \le 10$
- (2) $|k_1 k_2| \le 10$ and $k_2 \ge k_3 10$
- (3) $|k_1 k_2| \le 10$ and $|k_2 k_3| \le 10$

Since k-frequency is the lowest frequency, hence one similarly or easily has

$$2^{2sk} \sum_{\substack{k_1 \le k_2 - 10 \\ |k_2 - k_3| \le 10 \\ k_3 \ge k + 10}} 2^{2k_3} 2^{\frac{3}{2}k_1} 2^{-k_3/8} \prod_{j=1}^{3} \|P_{k_j}u\|_{F_{k_j}(T)} \|P_ku\|_{F_k(T)} \\ + 2^{2sk} \sum_{\substack{k_2 \ge k_3 + 10 \\ |k_1 - k_2| \le 10 \\ k_3 \ge k + 10}} 2^{k_3} 2^{k/2} \prod_{j=1}^{3} \|P_{k_j}u\|_{F_{k_j}(T)} \|P_ku\|_{F_k(T)} \\ + 2^{2sk} \sum_{\substack{k_2 \ge k_3 + 10 \\ |k_2 - k_3| \le 10 \\ k_3 \ge k + 10}} 2^{2k_3} 2^{k/2} \prod_{j=1}^{3} \|P_{k_j}u\|_{F_{k_j}(T)} \|P_ku\|_{F_k(T)} \\ \lesssim \max\left(2^{\frac{15}{8}}, 2^{(\frac{27}{8} - s)k}\right) \|P_ku\|_{F_k(T)} \|u\|_{F_s(T)}^3,$$

for $s \ge \frac{9}{8}$, thanks to Lemma 2.11 (c) and (b).

The estimate of $I_{2,2}$ is very similar as before. In view of the estimate of $I_{2,1}$, one knows that the worst case appears when the frequency support of u_{xx} is I_k . However, a direct calculation (integration by parts) gives

$$\begin{aligned} \left| \int_{\mathbb{R}\times[0,t_k]} P_k(u) (P_{k'}(u^2) P_k(u_{xx}))_x \, dx dt \right| &= \left| \int_{\mathbb{R}\times[0,t_k]} (P_k(u))_x (P_{k'}(u^2) P_k(u_{xx})) \, dx dt \right| \\ &= \frac{1}{2} \left| \int_{\mathbb{R}\times[0,t_k]} ((P_k(u))_x)^2 (P_{k'}(u^2))_x \, dx dt \right|, \end{aligned}$$

which is exactly same as $I_{2,1}$ (in particular, $I_{2,1,1}$ and $I_{2,1,2}$ under $|k_3 - k| \leq 10$). The rigorous justification of this observation can be seen in the commutator estimates, see the proof of Lemma 4.1 (c) in [22] for the details or see the proof of Proposition 2.13 below. Moreover, one can see that the derivatives are fairly distributed in $I_{2,3}$, and hence it can be easily or similarly controlled as the estimate of $I_{2,1}$. We omit the details.

On the other hand, the rest part,

$$2^{2sk} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(u) P_k(F(u)) \, dx dt \right|,$$

where $F(u) = (u^p)_x$, p = 2, 4, 5, can be immediately handled by using

$$\left| \int_{\mathbb{R}\times[0,t_k]} \prod_{j=1}^{p+1} u_j \, dx dt \right| \lesssim 2^{(k_1+\dots+k_{p-1})/2} \prod_{j=1}^{p+1} \|u_j\|_{F_{k_j}(T)} \,,$$

where $u_j = P_{k_j} u \in F_{k_j}(T), \ j = 1, \dots, p+1$ and assuming that $k_1 \leq \dots \leq k_{p+1}$, for p = 2, 4, 5.

Collecting all, we have

$$\sum_{k\geq 1} (I_2(k) + I_3(k)) \lesssim \|u\|_{F^s(T)}^2 + \|u\|_{F^s(T)}^4 + \|u\|_{F^s(T)}^5 + \|u\|_{F^s(T)}^6,$$

for $s \ge 2$, thus we complete the proof of (2.34).

Let u_1 and u_2 be solutions to (2.1). Define $v = u_1 - u_2$, then v solves

$$v_t + v_{5x} + \mathcal{N}_2(u_1, u_2) + \mathcal{N}_3(u_1, u_2) + \mathcal{SN}(u_1, u_2) = 0, \qquad v(0, x) = u_1(0, x) - u_2(0, x), \tag{2.35}$$

where

$$\mathcal{N}_2(u_1, u_2) = \mathcal{N}_2(u_1) - \mathcal{N}_2(u_2), \quad \mathcal{N}_3(u_1, u_2) = \mathcal{N}_3(u_1) - \mathcal{N}_3(u_2) \text{ and } \mathcal{SN}(u_1, u_2) = \mathcal{SN}(u_1) - \mathcal{SN}(u_2).$$

Proposition 2.13. Let $s \ge 2$ and $T \in (0,1]$. Then, for solutions $v \in C([-T,T]; H^{\infty}(\mathbb{T}))$ to (2.35) and $u_1, u_2 \in C([-T,T]; H^{\infty}(\mathbb{T}))$ to (2.1), we have

$$\|v\|_{E^{0}(T)}^{2} \lesssim \|v_{0}\|_{L^{2}}^{2} + \left(\sum_{j=1}^{2} \left(\|u_{j}\|_{F^{s}(T)} + \|u_{j}\|_{F^{s}(T)}^{2} + \|u_{j}\|_{F^{s}(T)}^{3} + \|u_{j}\|_{F^{s}(T)}^{4} + \|u_{j}\|_{F^{s}(T)}^{4}\right)\right) \|v\|_{F^{0}(T)}^{2},$$

and

$$\begin{aligned} \|v\|_{E^{s}(T)}^{2} &\lesssim \|v_{0}\|_{H^{s}}^{2} + \left(\sum_{j=1}^{2} \left(\|u_{j}\|_{F^{s}(T)} + \|u_{j}\|_{F^{s}(T)}^{2} + \|u_{j}\|_{F^{s}(T)}^{3} + \|u_{j}\|_{F^{s}(T)}^{4}\right)\right) \|v\|_{F^{s}(T)}^{2} \\ &+ \left(\|u_{1}\|_{F^{2s}(T)} + \|u_{2}\|_{F^{2s}(T)}\right) \|v\|_{F^{0}(T)} \|v\|_{F^{s}(T)} \,. \end{aligned}$$

Remark 2.11. One can see that the cubic terms with three derivatives are not harmful even in F^s , while the same terms are the main enemy under the periodic boundary condition. The principal reason is due to the lack of the smoothing effect under the periodic condition. Compare Lemma 2.11 (c) and Lemma 6.4 (c) and (d) in [41].

Proof. We first concentrate on the estimate on $||v||_{E^0(T)}^2$. From the definition of $||v||_{E^s(T)}^2$ and (2.25), it suffices to control

$$2^{2sk} \|P_k(v(t_k))\|_{L^2}^2 - 2^{2sk} \|P_k(v_0)\|_{L^2}^2 \lesssim 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(v) P_k(\mathcal{N}_2(u_1, u_2)) \, dx dt \right| + 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(v) P_k(\mathcal{N}_3(u_1, u_2)) \, dx dt \right| + 2^{2sk} \left| \int_{\mathbb{R} \times [0, t_k]} P_k(v) P_k(\mathcal{SN}(u_1, u_2)) \, dx dt \right| =: 2^{2sk} \widetilde{I}_1(k) + 2^{2sk} \widetilde{I}_2(k) + 2^{2sk} \widetilde{I}_3(k).$$

Proposition 4.4 in [22] yields

$$\sum_{k\geq 1} \widetilde{I}(k) \lesssim \left(\sum_{j=1}^{2} (\|u_j\|_{F^s(T)} + \|u_j\|_{F^s(T)}^2) \right) \|v\|_{F^0(T)}^2$$

and

$$\sum_{k\geq 1} 2^{2sk} \widetilde{I}(k) \lesssim \left(\sum_{j=1}^{2} (\|u_j\|_{F^s(T)} + \|u_j\|_{F^s(T)}^2) \right) \|v\|_{F^s(T)}^2 + \left(\|u_1\|_{F^{2s}(T)} + \|u_2\|_{F^{2s}(T)} \right) \|v\|_{F^0} \|v\|_{F^s(T)},$$

for $s \geq 2$. Moreover, since the quintic term in $\mathcal{N}_3(u_1, u_2)$ and $\mathcal{SN}(u_1, u_2)$ contains only one derivative, one can easily handle them compared to the cubic term in $\mathcal{N}_3(u_1, u_2)$. Thus, in what follows, we only focus on $\widetilde{I}_2(k)$ (in particular, the cubic terms), similarly as the proof of Proposition 2.12.

We write $\widetilde{I}_2(k) = \widetilde{I}_{2,1} - \widetilde{I}_{2,2} + \widetilde{I}_{2,3}$, where

and

$$\widetilde{I}_{2,3} := \left| \int_{\mathbb{R} \times [0,t_k]} P_k(v) P_k(v_x(u_{1,x}^2 + u_{1,x}u_{2,x} + u_{2,x}^2)) \, dx dt \right|.$$

We, here, only consider $\tilde{I}_{2,2}$ in order to provide a rigorous proof of the estimate of $I_{2,2}$ in the proof of Proposition 2.12. Moreover, it is easier to handle $\tilde{I}_{2,1,2}$ than $\tilde{I}_{2,1,1}$ (or similar), since less derivatives are taken in v, hence it is enough to estimate only $\widetilde{I}_{2,1,1}$. We reduce $\widetilde{I}_{2,2,1}$ as

$$\int_{\mathbb{R}\times[0,t_k]} P_k(v_x) P_k(u^2 v_{xx}) \, dx dt$$

A direct calculation gives

$$\begin{aligned} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(v_x) P_k(u^2 v_{xx}) \, dx dt \right| &\lesssim \sum_{k' \leq k-10} \left| \int_{\mathbb{R} \times [0,t_k]} P_k(v_x) P_k(P_{k'}(u^2) v_{xx}) \, dx dt \right| \\ &+ \sum_{k' \geq k-9, k_3 \geq 0} \left| \int_{\mathbb{R} \times [0,t_k]} P_k^2(v_x) P_{k'}(u^2) P_{k_3}(v_{xx}) \, dx dt \right| \end{aligned}$$

Since

$$P_{k}(v_{x})P_{k}(P_{k'}(u^{2})v_{xx}) = P_{k}(v_{x})P_{k}(v_{xx})P_{k'}(u^{2}) + P_{k}(v_{x})[P_{k}, P_{k'}(u^{2})]v_{xx}$$
$$= \frac{1}{2}((P_{k}(v_{x}))^{2})_{x}P_{k'}(u^{2}) + P_{k}(v_{x})[P_{k}, P_{k'}(u^{2})]v_{xx}$$

where [A, B] = AB - BA, the integration by parts yields

$$\sum_{k' \le k-10} \left| \int_{\mathbb{R} \times [0,t_k]} ((P_k(v_x))^2)_x P_{k'}(u^2) \, dx dt \right| = \sum_{k' \le k-10} \left| \int_{\mathbb{R} \times [0,t_k]} ((P_k(v_x))^2) P_{k'}((u^2)_x) \, dx dt \right|,$$

which is already dealt with in the proof of Proposition 2.12 (in particular, $I_{2,1,1}$). Thus, we have

$$\sum_{k\geq 1} \sum_{k'\leq k-10} \left| \int_{\mathbb{R}\times[0,t_k]} ((P_k(v_x))^2) P_{k'}((u^2)_x) \, dx dt \right| \lesssim \|u\|_{F^s(T)}^2 \|v\|_{F^0(T)}^2$$

and

$$\sum_{k\geq 1} 2^{2sk} \sum_{k'\leq k-10} \left| \int_{\mathbb{R}\times[0,t_k]} ((P_k(v_x))^2) P_{k'}((u^2)_x) \, dx dt \right| \lesssim \|u\|_{F^s(T)}^2 \|v\|_{F^s(T)}^2,$$

for $s \geq 2$.

On the other hand, a direct computation, in addition to the mean value theorem, (2.5) and (2.4), gives

$$\mathcal{F}\left([\widetilde{P}_k,\widetilde{P}_{k'}(u^2)](v_{xx})\right)(\tau,\xi) = C \int_{\mathbb{R}^2} \mathcal{F}(\widetilde{P}_{k'}((u^2)_x))(\tau',\xi') \cdot \mathcal{F}(v_x)(\tau-\tau',\xi-\xi') \cdot m(\xi,\xi') \, d\xi' d\tau',$$

where,

$$m(\xi,\xi')| = \left|\frac{(\xi-\xi')(\chi_k(\xi)-\chi_k(\xi-\xi'))}{\xi'}\right| \lesssim |(\xi-\xi_1)\chi'_k(\xi-\theta\xi_1)| \lesssim \sum_{|k-k'|\leq 4} \chi_{k'}(\xi-\xi_1),$$

for $0 \le \theta \le 1$. Thus, an analogous argument yields

$$\sum_{k\geq 1} \sum_{k'\leq k-10} \left| \int_{\mathbb{R}\times[0,t_k]} P_k(v_x) [P_k, P_{k'}(u^2)] v_{xx} \, dx dt \right| \lesssim \|u\|_{F^s(T)}^2 \|v\|_{F^0(T)}^2$$

$$\sum_{k\geq 1} 2^{2sk} \sum_{k'\leq k-10} \left| \int_{\mathbb{R}\times[0,t_k]} P_k(v_x) [P_k, P_{k'}(u^2)] v_{xx} \, dx dt \right| \lesssim \|u\|_{F^s(T)}^2 \|v\|_{F^s(T)}^2 \,,$$

for $s \geq 2$.

The rest of the proof, which is the estimate of

$$\sum_{k' \ge k-9, k_3 \ge 0} \left| \int_{\mathbb{R} \times [0, t_k]} P_k^2(v_x) P_{k'}(u^2) P_{k_3}(v_{xx}) \, dx dt \right|, \tag{2.36}$$

is almost identical to the proof of the estimate of $I_{2,1,2}$ in the proof of Proposition 2.12. Thus, we have

$$\sum_{k\geq 1} (2.36) \lesssim \|u\|_{F^{s}(T)}^{2} \|v\|_{F^{0}(T)}^{2}$$
$$\sum_{k\geq 1} 2^{2sk}(2.36) \lesssim \|u\|_{F^{s}(T)}^{2} \|v\|_{F^{s}(T)}^{2},$$

and

for
$$s > 2$$
. Thus, we complete the proof.

2.6. Local and Global well-posedness. The local-well-posedness argument (the classical energy method) is now standard. We refer the readers to [28, 26, 22, 30, 41] and references therein, for more details, and we, here, give a sketch of proof.

We first state fundamental properties of $X^{s,b}$ -type norms.

Proposition 2.14. Let $s \ge 0$, $T \in (0, 1]$, and $u \in F^s(T)$, then

$$\sup_{[-T,T]} \|u(t)\|_{H^s} \lesssim \|u\|_{F^s(T)}$$

Proposition 2.15. Let $T \in (0, 1]$, $u, v \in C([-T, T] : H^{\infty})$ and

 $t \in$

$$\partial_t u + \partial_x^5 u = v \text{ on } \mathbb{R} \times (-T, T)$$

Then we have

$$\|u\|_{F^{s}(T)} \lesssim \|u\|_{E^{s}(T)} + \|v\|_{N^{s}(T)}$$

for any $s \geq 0$.

See Appendix in [22] for the proofs. We also refer to [28, 30, 26].

One can observe that (1.1) admits the scaling equivalence with (2.1): For $\lambda > 0$, if u is a solution to (1.1), then u_{λ} , defined by

$$u_{\lambda}(t,x) := \lambda u(\lambda^5 t, \lambda x),$$

is a solution to (2.1). Moreover, a direct calculation yields $||u_{0,\lambda}||_{\dot{H}^s} = \lambda^{s+\frac{1}{2}} ||u_0||_{\dot{H}^s}$, which says the scaling exponent $s_c = -\frac{1}{2}$. Thus, a small data local well-posedness of (2.1) ensures the local-in-time well-posedness of (1.1) for an arbitrary data.

From Duhamel's principle, we know that the solution to (2.1) is of the following integral form:

$$u(t) = W(t)u_0 + \int_0^t W(t-s) \left(\mathcal{N}_2(u)(s) + \mathcal{N}_3(u)(s) + \mathcal{SN}(u)(s)\right) ds,$$

where W(t) is defined as in (2.6). We assume

$$\|u_0\|_{H^s} \le \epsilon \ll 1. \tag{2.37}$$

Remark that for any fixed $\mu \in \mathbb{R}^+$, we can choose $0 < \lambda_0 \ll 1$ sufficiently small such that the initial data satisfy (2.37) and $\mu \lambda \leq 1$ for all $\lambda \leq \lambda_0$. The second condition ensures that our local well-posedness argument does not depend on μ .

We fix $s \ge 2$. Proposition 2.15, Propositions 2.7, 2.8 and 2.9, and Proposition 2.12 ensures

$$\left\{ \begin{array}{l} \|u\|_{F^{s}(T')} \lesssim \|u\|_{E^{s}(T')} + \|\mathcal{N}_{2}(u) + \mathcal{N}_{3}(u) + \mathcal{SN}(u)\|_{N^{s}(T')}; \\ \|\mathcal{N}_{2}(u) + \mathcal{N}_{3}(u) + \mathcal{SN}(u)\|_{N^{s}(T')} \lesssim \sum_{j=2}^{5} \|u\|_{F^{s}(T)}^{j}; \\ \|u\|_{E^{s}(T')}^{2} \lesssim \|u_{0}\|_{H^{s}}^{2} + \sum_{j=3}^{6} \|u\|_{F^{s}(T)}^{j}, \end{array} \right.$$

for any $T' \in [0, T]$, which, in addition to the smallness condition (2.37) and continuity argument (see Lemma 6.3 in [30] for the details), implies a priori bound:

$$\sup_{t \in [-T,T]} \|u(t)\|_{H^s} \lesssim \|u(t)\|_{F^s(T)} \lesssim \|u_0\|_{H^s} \,.$$
(2.38)

To complete the proof, we need

Proposition 2.16. Assume $s \ge 2$. Let $u_1, u_2 \in F^s(T)$ be solutions to (2.1) with small initial data $u_{1,0}, v_{2,0} \in H^{\infty}$. Let $v = u_1 - u_2$ and $v_0 = u_{1,0} - u_{2,0}$. Then we have

$$\|v\|_{F^0(T)} \lesssim \|v_0\|_{L^2}$$

and

$$\|v\|_{F^{s}(T)} \lesssim \|v_{0}\|_{H^{s}} + \|u_{1,0}\|_{H^{2s}} \|v_{0}\|_{L^{2}}$$

It immediately follows from Proposition 2.15, Propositions 2.7, 2.8 and 2.9, and Proposition 2.13 under (2.38).

For fixed $u_0 \in H^s$, a density argument enables us to choose a sequence $\{u_{0,n}\}_{n=1}^{\infty} \subset H^{\infty}$ such that $u_{0,n} \to u_0$ in H^s as $n \to \infty$. Let $u_n(t) \in H^{\infty}$ is a solution to (2.1) with initial data $u_{0,n}$. Using a similar argument as above and Proposition 2.16, one shows $\{u_n\}$ is a Cauchy sequence. Indeed, for $K \in \mathbb{Z}_+$, let $u_{0,n}^K = P_{\leq K}(u_{0,n})$. Then, $u_n^K = P_{\leq K}u_n$ satisfies the frequency localized equation $(P_{\leq K}(2.1))$ with the initial data $u_{0,n}^K$. We have from the triangle inequality that

$$\sup_{t \in [-T,T]} \|u_m - u_n\|_{H^s} \lesssim \sup_{t \in [-T,T]} \|u_m - u_m^K\|_{H^s} + \sup_{t \in [-T,T]} \|u_m^K - u_n^K\|_{H^s} + \sup_{t \in [-T,T]} \|u_n^K - u_n\|_{H^s}.$$

The first and last terms are bounded by ϵ , thanks to *a priori* bound, and the second term is bounded by ϵ , thanks to Proposition 2.13, precisely,

$$\sup_{t \in [-T,T]} \left\| v_m^K - v_n^K \right\|_{H^s(\mathbb{T})} \lesssim \left\| v_{0,m}^K - v_{0,n}^K \right\|_{H^s} + K^s \left\| v_{0,m}^K - v_{0,n}^K \right\|_{L^2} \lesssim \epsilon.$$

Hence the 3ϵ -argument completes the proof and we obtain a solution as the limit. The uniqueness and the continuity of dependence follow from an analogous argument.

Remark 2.12. In view of all analyses above, we do not use the integrability of the Gardner equation (2.1) to prove the local well-posedness, and thus we can apply our argument to prove the local result of (2.1) with arbitrary coefficients.

Small solutions u to (2.1) satisfies the (rescaled) conservation laws (1.4)-(1.5)-(1.6), namely

$$M[u](t) := \frac{1}{2} \int_{\mathbb{R}} u^2(t, x) dx = M[u](0),$$

$$E_{\mu}[u](t) := \int_{\mathbb{R}} \left(\frac{1}{2}u_x^2 - 2\mu\lambda u^3 - \frac{1}{2}u^4\right)(t, x) dx = E_{\mu}[u](0),$$

$$\int_{\mathbb{R}} \left(\frac{1}{2}u_x^2 - 2\mu\lambda u^3 - \frac{1}{2}u^4\right)(t, x) dx = E_{\mu}[u](0),$$

and

$$E_{5\mu}[u](t) := \int_{\mathbb{R}} \left(\frac{1}{2} u_{xx}^2 - 10\mu\lambda u u_x^2 + 10\mu^2\lambda^2 u^4 - 5u^2 u_x^2 + 6\mu\lambda u^5 + u^6 \right)(t,x) dx = E_{5\mu}[u](0).$$
(2.39)

Using above conserved quantities and the Sobolev embedding in addition to the smallness condition, one proves Theorem 1.3.

3. Stability of breathers of the 5th order Gardner equation

Once we have shown the existence of global solutions of the Cauchy problem for the 5th order Gardner equation (1.1), we study now the stability properties of a special solution of (1.1). Before dealing with this stability result, we present basic facts on solutions of (1.1). The simplest solution of the 5th order *focusing* Gardner equation is a traveling wave like solution, usually called as soliton solution, and explicitly defined as follows

Definition 3.1. The 1-soliton solution $Q_{\mu} \equiv Q_{\mu,c}$ of the 5th order focusing Gardner equation (1.1) is given by

$$Q_{\mu}(t,x) := Q_{\mu,c}(x - v_{\mu,c}t + x_1), \quad Q_{\mu,c}(z) := \frac{c}{2\mu + \sqrt{4\mu^2 + c}\cosh(\sqrt{c}z)}, \quad c > 0, \ \mu, \ x_1 \in \mathbb{R},$$
with $v_{\mu,c} := c^2 + 10\mu^2 c,$
(3.1)

which indeed has a completely similar profile to the well known Gardner soliton profile [3]. The 1-soliton solution $Q_{\mu,c}$ (3.1) of the 5th order Gardner equation (1.1) satisfies the nonlinear second order ODE:

$$Q''_{\mu} - c Q_{\mu} + 6\mu Q_{\mu}^2 + 2Q_{\mu}^3 = 0, \quad Q_{\mu} > 0, \quad Q_{\mu} \in H^1(\mathbb{R}).$$

Note that, as a solution of (1.1), Q_{μ} also satisfies naturally the fourth order ODE

$$\begin{aligned} Q_{\mu}^{''''} &- v_{\mu,c} \, Q_{\mu} + \tilde{f}_5(Q_{\mu}) = 0, \quad \text{with} \\ \tilde{f}_5(Q_{\mu}) &= 10(\mu + Q_{\mu})^2 Q_{\mu}^{''} + 10(\mu + Q_{\mu})(Q_{\mu}^{'})^2 + 60\mu^3 Q_{\mu}^2 + 60\mu^2 Q_{\mu}^3 + 30\mu Q_{\mu}^4 + 6Q_{\mu}^5 \end{aligned}$$

Moreover, it is possible to build a solution made of the composition of N of such solitons, which is usually called as the *N*-soliton solution for the 5th order Gardner equation

$$u_{\mu}(t,x) := i\partial_x \log(g_{\mu}/f_{\mu}), \quad \text{where}$$

$$f_{\mu}(t,x) := \sum_{\sigma=0,1} \exp\left[\sum_{k=1}^N \sigma_k \left(\theta_k + \rho_k\right) + \sum_{k< m}^N \sigma_k \sigma_m A_{km}\right], \quad \text{and} \quad g_{\mu}(t,x) := f_{\mu}^*(t,x),$$

$$i\sqrt{c_k} + 2\mu \qquad (\sqrt{c_k} - \sqrt{c_m})^2$$

with

$$\exp[\rho_k] := \frac{i\sqrt{c_k} + 2\mu}{\sqrt{c_k}}, \quad \exp[A_{km}] = \left(\frac{\sqrt{c_k} - \sqrt{c_m}}{\sqrt{c_k} + \sqrt{c_m}}\right)^2, \ k, m = 1, 2, \dots, N,$$
$$\theta_k = \sqrt{c_k}(x - v_{5,k}t - \varrho_k) \quad \text{and} \quad v_{5,k} := c_k^2 + 10\mu^2 c_k,$$

where ς^* means the complex conjugate of ς , here c_k is the scaling and ϱ_k an arbitrary constant phase, $\sum_{\mu=0,1}$ means here the summation over all possible combinations of $\sigma_k = 0, 1, \ k = 1, 2, ... N$ and $\sum_{k < m}^{N}$ the summation over all possible combinations of the N elements under the constraint k < m.

From that multi-soliton solution, the 2-soliton solution is apparent. Now, if we take this 2-soliton solution of (1.1), and transforming its corresponding scalings c_1, c_2 to complex ones $c_1 = c_2^* := (\beta + i\alpha)^2$, $\alpha, \beta \in \mathbb{R} \setminus \{0\}$, it allows us to build a new solution of (1.1) named as the *breather solution*. This is a localized in space and periodic in time (modulo symmetries of the equation) solution, and it is defined as follows:

Definition 3.2 (5th order Gardner breather). Let $\alpha, \beta \in \mathbb{R} \setminus \{0\}$, $\mu \in \mathbb{R}^+ \setminus \{0\}$ such that $\Delta = \alpha^2 + \beta^2 - 4\mu^2 > 0$, and $x_1, x_2 \in \mathbb{R}$. The 5th order breather solution $B_{\mu} \equiv B_{\mu,5}$ of the 5th order Gardner equation (1.1), is given explicitly by the formula

$$B_{\mu} \equiv B_{\alpha,\beta,\mu,5}(t,x;x_1,x_2) := 2\partial_x \left[\arctan\left(\frac{G_{\mu}(t,x)}{F_{\mu}(t,x)}\right) \right], \tag{3.2}$$

where

$$G_{\mu}(t,x) := \frac{\beta\sqrt{\alpha^2 + \beta^2}}{\alpha\sqrt{\Delta}}\sin(\alpha y_1) - \frac{2\mu\beta[\cosh(\beta y_2) + \sinh(\beta y_2)]}{\Delta},$$

$$F_{\mu}(t,x) := \cosh(\beta y_2) - \frac{2\mu\beta[\alpha\cos(\alpha y_1) - \beta\sin(\alpha y_1)]}{\alpha\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}},$$

with y_1 and y_2

 $y_1 = x + \delta_5 t + x_1, \quad y_2 = x + \gamma_5 t + x_2,$

and with velocities

$$\delta_5 := -\alpha^4 + 10\alpha^2\beta^2 - 5\beta^4 + 10(\alpha^2 - 3\beta^2)\mu^2, \gamma_5 := -\beta^4 + 10\alpha^2\beta^2 - 5\alpha^4 + 10(3\alpha^2 - \beta^2)\mu^2.$$
(3.3)

First of all, we remember the following identity for solutions of the of 5th order Gardner equation (1.1) (see [6, Appendix A] for a detailed proof of a similar identity for the classical Gardner equation)

Lemma 3.3. Let $u(t,x) = \partial_x \log(\frac{F_\mu - iG_\mu}{F_\mu + iG_\mu})$ be any solution of the 5th order Gardner equation (1.1). Then

 $u^2 = \partial_x^2 \log(G_\mu^2 + F_\mu^2) - 2\mu u.$

Now, we can compute explicitly the mass of such breather solution:

Lemma 3.4. Let $B_{\mu} \equiv B_{\mu,5}$ be the breather solution (3.2) of the 5th order Gardner equation (1.1). Then the mass of B_{μ} is

$$M[B_{\mu}] := 2\beta + 2\mu \arctan\left[\frac{4\mu\beta}{\Delta}\right]$$

Proof. The proof follows directly from Lemma 3.3, by using the breather solution (3.2).

Moreover, the breather solution (3.2) of the 5th order Gardner equation (1.1) satisfies the following nonlinear identities:

Lemma 3.5. Let $B_{\mu} \equiv B_{\mu,5}$ be the breather solution (3.2) of the 5th order Gardner equation (1.1). Then (1) $B_{\mu} = \tilde{B}_{\mu,x}$, with $\tilde{B}_{\mu} = \tilde{B}_{\alpha,\beta,\mu}$ given by the smooth L^{∞} -function

$$\tilde{B}_{\mu}(t,x) := 2 \arctan\left(\frac{G_{\mu}}{F_{\mu}}\right).$$

(2) For any fixed $t \in \mathbb{R}$, we have $(\tilde{B}_{\mu})_t$ well-defined in the Schwartz class, satisfying

$$B_{\mu,4x} + \tilde{B}_{\mu,t} + 10(\mu + B_{\mu})^2 B_{\mu,xx} + 10(\mu + B_{\mu}) B_{\mu,x}^2 + 6(10\mu^3 B_{\mu}^2 + 10\mu^2 B_{\mu}^3 + 5\mu B_{\mu}^4 + B_{\mu}^5) = 0.$$
(3.4)

Proof. The first item above is a direct consequence of the definition of B_{μ} in (3.2). On the other hand, (3.4) is a consequence of (1.1) and integration in space (from $-\infty$ to x) of (1.1).

Finally, we show that breather solutions (3.2) of the 5th order Gardner equation (1.1) satisfy the following identity: **Lemma 3.6.** Let $B_{\mu} \equiv B_{\mu,5}$ be the breather solution (3.2) of the 5th order Gardner equation (1.1). Then, for all $t \in \mathbb{R}$,

$$\tilde{B}_{\mu,t} = (\alpha^2 + \beta^2)^2 B_{\mu} + 2(\alpha^2 - \beta^2 - 5\mu^2) (B_{\mu,xx} + 2B_{\mu}^3 + 6\mu B_{\mu}^2).$$
(3.5)

Proof. See appendix B for a detailed proof of this nonlinear identity.

Now, we prove that breather solutions (3.2) of the 5th order Gardner equation (1.1) satisfy a fourth order ODE, which indeed is the same as the one satisfied by *classical* Gardner breather solutions (see [6, Theorem 3.5] for further details)

Theorem 3.7. Let $B_{\mu} \equiv B_{\mu,5}$ be the breather solution (3.2) of the 5th order Gardner equation (1.1). Then, for any fixed $t \in \mathbb{R}$, B_{μ} satisfies the nonlinear stationary equation

$$\mathcal{W}(B_{\mu}) := B_{\mu,4x} - 2(\beta^2 - \alpha^2)(B_{\mu,xx} + 6\mu B_{\mu}^2 + 2B_{\mu}^3) + (\alpha^2 + \beta^2)^2 B_{\mu} + 10B_{\mu}B_{\mu,x}^2 + 10B_{\mu}^2 B_{\mu,xx} + 6B_{\mu}^5 + 10\mu B_{\mu,xx}^2 + 20\mu B_{\mu}B_{\mu,xx} + 40\mu^2 B_{\mu}^3 + 30\mu B_{\mu}^4 = 0.$$
(3.6)

Proof. We use the identity (3.4) to substitute the $B_{\mu,4x}$ term in the left-hand side of (3.6), simplifying it as:

$$\mathcal{W}(B_{\mu}) = -\left(\tilde{B}_{\mu,t} + 10(\mu + B_{\mu})^{2}B_{\mu,xx} + 10(\mu + B_{\mu})B_{\mu,x}^{2} + 6(10\mu^{3}B_{\mu}^{2} + 10\mu^{2}B_{\mu}^{3} + 5\mu B_{\mu}^{4} + B_{\mu}^{5})\right) - 2(\beta^{2} - \alpha^{2})(B_{\mu,xx} + 6\mu B_{\mu}^{2} + 2B_{\mu}^{3}) + (\alpha^{2} + \beta^{2})^{2}B_{\mu} + 10B_{\mu}B_{\mu,x}^{2} + 10B_{\mu}^{2}B_{\mu,xx} + 6B_{\mu}^{5} + 10\mu B_{\mu,xx}^{2} + 20\mu B_{\mu}B_{\mu,xx} + 40\mu^{2}B_{\mu}^{3} + 30\mu B_{\mu}^{4} = -\tilde{B}_{t} + (\alpha^{2} + \beta^{2})^{2}B_{\mu} + 2\left(\alpha^{2} - \beta^{2} - 5\mu^{2}\right)(B_{\mu,xx} + 2B_{\mu}^{3} + 6\mu B_{\mu}^{2}) = 0,$$

where in the last line we have used the identity (3.5).

Note that being the shift parameters x_1, x_2 in (3.2) selected as independents of time, a simple argument guarantees that the previous Theorem 3.7 still holds under time dependent, translation parameters $x_1(t)$ and $x_2(t)$.

Corollary 3.8. Let $B^0_{\mu} \equiv B^0_{\alpha,\beta,\mu}(t,x;0,0)$ be any Gardner breather as in (3.2), and $x_1(t), x_2(t) \in \mathbb{R}$ two continuous functions, defined for all t in a given interval. Consider the modified breather

$$B_{\mu}(t,x) := B^{0}_{\alpha,\beta,\mu}(t,x;x_{1}(t),x_{2}(t)), \qquad (cf. (3.2))$$

Then B_{μ} satisfies (3.6), for all t in the given interval.

Proof. From the invariance of the equation (3.6) under spatial translations, we conclude.

Even more, we can characterize variationally these breather solutions of the 5th order Gardner equation. Explicitly, considering the $H^2(\mathbb{R})$ conserved quantity (1.6)

$$E_{5\mu}[u](t) := \int_{\mathbb{R}} \left(\frac{1}{2} u_{xx}^2 - 10\mu u u_x^2 + 10\mu^2 u^4 - 5u^2 u_x^2 + 6\mu u^5 + u^6 \right) \, dx, \tag{3.7}$$

we can introduce a H^2 functional, associated to the breather solution. Namely, we define this functional as a linear combination of the energy (1.5), the mass (1.4) and (3.7) in the following way

$$\mathcal{H}_{\mu}[u](t) := E_{5\mu}[u](t) + 2(\beta^2 - \alpha^2)E_{\mu}[u](t) + (\alpha^2 + \beta^2)^2M[u](t).$$
(3.8)

Therefore, $\mathcal{H}_{\mu}[u]$ is a conserved quantity, well-defined for H^2 -solutions of (1.1). Additionally, we have that

Lemma 3.9. Breather solutions B_{μ} (3.2) of the 5th order Gardner equation (1.1) are critical points of the Lyapunov functional \mathcal{H}_{μ} (3.8). In fact, for any $z \in H^2(\mathbb{R})$ with sufficiently small H^2 -norm, and $B_{\mu} = B_{\alpha,\beta,\mu}$ any 5th Gardner breather solution, one has, for all $t \in \mathbb{R}$, that

$$\mathcal{H}_{\mu}[B_{\mu}+z] - \mathcal{H}_{\mu}[B_{\mu}] = \frac{1}{2}\mathcal{Q}_{\mu}[z] + \mathcal{N}_{\mu}[z]$$

with \mathcal{Q}_{μ} being the quadratic form defined in (3.9) below, and $\mathcal{N}_{\mu}[z]$ satisfying $|\mathcal{N}_{\mu}[z]| \leq K ||z||^{3}_{H^{2}(\mathbb{R})}$.

Proof. A direct computation with the integration by parts yields

$$\begin{split} E_{5\mu}[B_{\mu}+z] &= E_{5\mu}[B_{\mu}+z] \\ &+ \int \left(B_{\mu,4x} + 10\mu B_{\mu,x}^{2} + 20\mu B_{\mu}B_{\mu,xx} + 10B_{\mu}B_{\mu,x}^{2} + 10B_{\mu}^{2}B_{\mu,x} + 40\mu^{2}B_{\mu}^{3} + 30\mu B_{\mu}^{4} + 6B_{\mu}^{5} \right) z \\ &+ \frac{1}{2} \int \left(\partial_{x}^{4} + (20\mu B_{\mu} + 10B_{\mu}^{2})\partial_{x}^{2} - 20(\mu B_{\mu,x} + B_{\mu}B_{\mu,x})\partial_{x} \\ &+ (-10B_{\mu,x}^{2} + 120\mu^{2}B_{\mu}^{2} + 120\mu B_{\mu}^{3} + 30B_{\mu}^{4}) \right) z \cdot z \\ &+ \int \left(-10\mu z z_{x}^{2} - 10B_{\mu} z z_{x} - 10B_{\mu,x} z_{x} z^{2} + 40\mu^{2}B_{\mu} z^{3} + 60\mu B_{\mu}^{2} z^{3} + 20B_{\mu}^{3} z^{3} \right) , \end{split}$$

$$E_{\mu}[B_{\mu}+z] = E_{\mu}[B_{\mu}] - \int \left(B_{\mu,xx} + 6\mu B_{\mu}^{2} + 2B_{\mu}^{3} \right) z - \frac{1}{2} \int \left(\partial_{x}^{2} + 12\mu B_{\mu} + 6B_{\mu}^{2} \right) z \cdot z - \int \left(2\mu z^{3} + 2B_{\mu} z^{3} + \frac{1}{2} z^{4} \right) dz \end{split}$$

and

$$M[B_{\mu} + z] = M[B_{\mu}] + \int B_{\mu}z + \frac{1}{2}\int z \cdot z.$$

Collecting all, one obtain

$$\mathcal{H}_{\mu}[B_{\mu}+z] = \mathcal{H}_{\mu}[B_{\mu}] + \int_{\mathbb{R}} \mathcal{W}(B_{\mu})z + \frac{1}{2}\mathcal{Q}_{\mu}[z] + \mathcal{N}_{\mu}[z],$$

where the quadratic form

$$\mathcal{Q}_{\mu}[z] := \int \mathcal{L}_{\mu} z \cdot z, \qquad (3.9)$$

associated to the linearized operator \mathcal{L}_{μ} given by

$$\mathcal{L}_{\mu} := \partial_{x}^{4} + (20\mu B_{\mu} + 10B_{\mu}^{2} - 2(\beta^{2} - \alpha^{2}))\partial_{x}^{2} - 20(\mu B_{\mu,x} + B_{\mu}B_{\mu,x})\partial_{x} + (-10B_{\mu,x}^{2} + 120\mu^{2}B_{\mu}^{2} + 120\mu B_{\mu}^{3} + 30B_{\mu}^{4} - 2(\beta^{2} - \alpha^{2})(12\mu B_{\mu} + 6B_{\mu}^{2}) + (\alpha^{2} + \beta^{2})^{2})$$
(3.10)

and the collection of higher order terms (with respect to z) $N_{\mu}[z]$ is given by

$$\mathcal{N}_{\mu}[z] := \int \left(-10\mu z z_x^2 - 10B_{\mu} z z_x - 10B_{\mu,x} z_x z^2 + 40\mu^2 B_{\mu} z^3 + 60\mu B_{\mu}^2 z^3 + 20B_{\mu}^3 z^3\right) \\ - 2(\beta^2 - \alpha^2) \int \left(2\mu z^3 + 2B_{\mu} z^3 + \frac{1}{2} z^4\right).$$

Theorem 3.7 ensures $\int_{\mathbb{R}} \mathcal{W}(B_{\mu})z = 0$, and hence one has $\mathcal{H}_{\mu}'[B_{\mu}] = 0$. Moreover, from direct estimates, one has $\mathcal{N}_{\mu}[z] = O(\|z\|^{3}_{H^{2}(\mathbb{R})})$, as desired.

3.1. Spectral analysis. As a direct consequence of the already studied spectral properties of the linearized operator \mathcal{L}_{μ} , associated to the *classical* Gardner breather solution B_{μ} , in [6], we obtain the same spectral results for breather solutions of the 5th order Gardner equation (1.1). In fact, all statements on spectral properties and the main Theorem in [6] are valid for the 5th order Gardner equation (1.1), even if explicit coefficients are different. Therefore in the following lines and for the sake of completeness, we only summarize and list the main features of \mathcal{L}_{μ} (3.10). Let B_{μ} as introduced in (3.2). Consider now the two directions associated to spatial translations. We define

$$B_1(t;x_1,x_2) := \partial_{x_1} B_\mu(t;x_1,x_2), \quad \text{and} \quad B_2(t;x_1,x_2) := \partial_{x_2} B_\mu(t;x_1,x_2).$$
(3.11)

Moreover, we compute and denote as *scaling directions*, the derivatives

$$\Lambda_{\alpha}B_{\mu} = \frac{\partial B_{\mu}}{\partial \alpha}, \quad \Lambda_{\beta}B_{\mu} = \frac{\partial B_{\mu}}{\partial \beta}.$$
(3.12)

We get the following (see [6] for more details)

Lemma 3.10. For any breather solution B_{μ} (3.2) of the 5th order Gardner equation (1.1), we get that

(1) (Continuous spectrum) \mathcal{L}_{μ} is a linear, unbounded operator in $L^{2}(\mathbb{R})$, with dense domain $H^{4}(\mathbb{R})$. Moreover, \mathcal{L}_{μ} is self-adjoint, and is a compact perturbation of the constant coefficients operator

$$\mathcal{L}_{\mu,0} := \partial_x^4 - 2(\beta^2 - \alpha^2)\partial_x^2 + (\alpha^2 + \beta^2)^2.$$

In particular, the continuous spectrum of \mathcal{L}_{μ} is the closed interval $[(\alpha^2 + \beta^2)^2, +\infty)$ in the case $\beta \geq \alpha$, and $[4\alpha^2\beta^2, +\infty)$ in the case $\beta < \alpha$, with no embedded eigenvalues are contained in this region.

(2) (Kernel) For each $t \in \mathbb{R}$, one has

$$\ker \mathcal{L}_{\mu} = \operatorname{span} \{ B_1(t; x_1, x_2), B_2(t; x_1, x_2) \}.$$

(3) Consider the scaling directions $\Lambda_{\alpha}B$ and $\Lambda_{\beta}B$ introduced in (3.12). Then, given $\alpha, \beta > 0$ and $\forall \mu \in (0, \frac{\sqrt{\alpha^2 + \beta^2}}{2})$, we have

$$\int_{\mathbb{R}} \Lambda_{\alpha} B_{\mu} \mathcal{L}_{\mu}[\Lambda_{\alpha} B_{\mu}] = 16\alpha^{2}\beta \left[1 + \frac{4\mu^{2}\Delta}{\Delta^{2} + 16\mu^{2}\beta^{2}}\right] > 0,$$

and

$$\int_{\mathbb{R}} \Lambda_{\beta} B_{\mu} \mathcal{L}_{\mu}[\Lambda_{\beta} B_{\mu}] = -16\beta \left[\alpha^2 + 2\mu^2 \left(1 + \frac{(\Delta - 2\beta^2)(\alpha^2 + \beta^2 + 4\mu^2)}{\Delta^2 + 16\mu^2\beta^2} \right) \right] < 0.$$

(4) Let

$$B_{0,\mu} := \frac{\alpha \Lambda_{\beta} B_{\mu} + \beta \Lambda_{\alpha} B_{\mu}}{8\alpha \beta (\alpha^2 + \beta^2)},$$

Then $B_{0,\mu}$ is in the Schwartz class, satisfying $\mathcal{L}_{\mu}[B_{0,\mu}] = -B_{\mu}$ and

$$\int_{\mathbb{R}} B_{0,\mu} B_{\mu} = \frac{1}{4\beta(\alpha^2 + \beta^2)} \left(\frac{\Delta^2 + 4\mu^2 \Delta}{\Delta^2 + 16\mu^2 \beta^2}\right) > 0. \quad \forall \mu \in \left(0, \frac{\sqrt{\alpha^2 + \beta^2}}{2}\right)$$

(5) Let B_1 and B_2 be the kernel elements defined in (3.11), $D_{\mu} = F_{\mu}^2 + G_{\mu}^2$ and W be the Wronskian matrix of the functions B_1 and B_2 , precisely given by

$$W[B_1, B_2](t, x) := \begin{bmatrix} B_1 & B_2 \\ (B_1)_x & (B_2)_x \end{bmatrix} (t, x).$$

Then

$$\det W[B_1, B_2](t, x) := \frac{2\beta^3(\alpha^2 + \beta^2)^2((\alpha^2 + \beta^2)^2 - 8\mu^2(\alpha^2 - 2\mu^2))}{\Delta^3 D_{\mu}^2} \times \left[\sinh(2\beta y_2) + \frac{8\beta^2\mu^2\cosh(2\beta y_2)}{(\alpha^2 + \beta^2)^2 - 8\mu^2(\alpha^2 - \mu^2)} - \frac{\beta\Delta((\alpha^2 + \beta^2)^2 - 4\mu^2(\alpha^2 - \beta^2))\sin(2\alpha y_1)}{\alpha(\alpha^2 + \beta^2)((\alpha^2 + \beta^2)^2 - 8\mu^2(\alpha^2 - \mu^2))} + \frac{8\beta^2\mu^2\Delta\cos(2\alpha y_1)}{(\alpha^2 + \beta^2)((\alpha^2 + \beta^2)^2 - 8\mu^2(\alpha^2 - 2\mu^2))} \right].$$

- (6) For every $\mu \in (0, \frac{\sqrt{\alpha^2 + \beta^2}}{2})$, the operator \mathcal{L}_{μ} defined in (3.10) has a unique negative eigenvalue $-\lambda_0^2 < 0$, of multiplicity one, where λ_0 depends on α , β , μ , x_1 , x_2 and t.
- (7) (Coercivity) Let $\alpha, \beta > 0$ and $\mu \in (0, \frac{\sqrt{\alpha^2 + \beta^2}}{2})$. For the quadratic from $Q_{\mu}[z]$ as in (3.9), associated to \mathcal{L}_{μ} (3.10), there exists a well-defined and positive continuous function $\nu_0 = \nu_0(\alpha, \beta, \mu)$ such that, for all $z_0 \in H^2(\mathbb{R})$ satisfying

$$\int_{\mathbb{R}} z_0 B_{-1} = \int_{\mathbb{R}} z_0 B_1 = \int_{\mathbb{R}} z_0 B_2 = 0,$$

the following Coercivity condition holds true:

$$Q_{\mu}[z_0] \ge \nu_0 \|z_0\|_{H^2(\mathbb{R})}^2.$$
(3.13)

For the proof of this Lemma, we refer the interested reader to [6, Lemma 5.10]. Finally, we present the stability result for breather solutions (3.2) of the 5th order Gardner equation (1.1):

Theorem 3.11 (H^2 -stability of 5th order Gardner breathers). Let $\alpha, \beta \in \mathbb{R} \setminus \{0\}$ and $\mu \in (0, \frac{\sqrt{\alpha^2 + \beta^2}}{2})$. Let $B_\mu \equiv B_{\mu,5}$ the breather solution (3.2) of the 5th order Gardner equation (1.1). Then, there exist positive parameters η_0, A_0 , depending on α, β and μ , such that the following holds: Consider $u_0 \in H^2(\mathbb{R})$, and assume that there exists $\eta \in (0, \eta_0)$ such that

$$||u_0 - B_\mu(t=0;0,0)||_{H^2(\mathbb{R})} \le \eta.$$

Then there exist $x_1(t), x_2(t) \in \mathbb{R}$ such that the solution u(t) of the Cauchy problem for the 5th order Gardner equation (1.1) with initial data u_0 , satisfies

$$\sup_{t \in \mathbb{R}} \|u(t) - B_{\mu}(t; x_1(t), x_2(t))\|_{H^2(\mathbb{R})} \le A_0 \eta$$

with

$$\sup_{t\in\mathbb{R}} |x_1'(t)| + |x_2'(t)| \le KA_0\eta,$$

for a constant K > 0.

Proof. We take $u = u(t) \in H^2(\mathbb{R})$ as the corresponding local in time solution of the Cauchy problem associated to (1.1), with initial condition $u(0) = u_0 \in H^2(\mathbb{R})$. Therefore, once we guaranteed for the case of the breather solution of the 5th order Gardner equation, that it satisfies the same 4th order ODE (3.6) as the *classical* Gardner breather, that a suitable coercivity property holds for the bilinear form \mathcal{Q}_{μ} associated to the breather solution of (1.1) (see (3.13)), and the existence of a unique negative eigenvalue (Lemma 3.10 (6)) of the linearized operator \mathcal{L}_{μ} given in (3.10), the stability proof follows the same steps as the H^2 -stability of *classical* Gardner breathers [6, Theorem 6.1] (see also [5, Theorem 6.1]). Namely, we proceed assuming that the maximal time of stability T is finite and we arrive to a contradiction.

Appendix A. Proof of Theorem 1.4

The aim of this section is to prove the ill-posedness of (1.1) for s > 0, which, in addition to the first author's recent work [4], completely justifies that the 5th Gardner equation (1.1) is the quasilinear equation in the sense that the flow map from data to solutions is not (locally) uniformly continuous for all regularities, see Corollary 1.5. Since the weak-illposedness phenomenon occurs due to the strong high-low interaction in the quadratic nonlinearity with three derivatives, Theorem 1.2 in [43] seems to guarantee the lack of uniform continuity of the flow map associated to (1.1) for s > 0. This section contributes to prove that the equation (1.1) is indeed weakly ill-posed for s > 0.

The proof basically follow the argument used in [43], initially introduced by Koch-Tzvetkov [39]. Since the (weak) ill-posedness phenomenon arises from the strong high-low quadratic nonlinearity (high frequency waves with low frequency perturbations), the main part of the proof is identical to the argument in [43]. Thus, we, here, provide an additional estimate to be needed for the other nonlinearities.

In view of the argument presented in Section 2.6, it suffices to show the ill-posedness of (2.1) with small initial data.

A.1. Setting. We first define the approximate solution, which is an ansatz to cause the (weak) ill-posedness phenomenon. Let $\phi, \phi \in C_0^{\infty}(\mathbb{R})$ be smooth bump functions satisfying

 $\phi \equiv 1$, |x| < 1, and $\phi \equiv 0$, |x| > 2

and

$$\widetilde{\phi} \equiv 1, \quad x \in \operatorname{supp}(\phi) \quad \text{and} \quad \widetilde{\phi}\phi \equiv \phi,$$

respectively. For $N \geq 1$ and $0 < \delta < 1$, set

$$\phi_N(x) := \phi\left(\frac{x}{N^{4+\delta}}\right), \quad \widetilde{\phi}_N(x) := \widetilde{\phi}\left(\frac{x}{N^{4+\delta}}\right)$$

Let $\epsilon > 0$ be a sufficiently small for the initial data to satisfy (2.37). Let

$$u_{0,l}^{\pm}(x) := \pm \epsilon N^{-3} \widetilde{\phi}_N(x)$$

and $u_l^{\pm}(t,x)$ be the solution to (2.1) with the initial data $u_{0,l}^{\pm}(x)$. Let $\Phi_N(t) := (N^5 - 10\mu^2\lambda^2 N^3)t$ and

$$u_h^{\pm}(t,x) := N^{-\frac{4+\delta}{2}-s} \phi_N(x) \cos(Nx - \Phi_N(t) \mp t)$$
(A.1)

be a high frequency part of the approximate solution, and thus define the approximate solution as

$$u_{ap}^{\pm}(t,x) := u_l^{\pm}(t,x) + u_h^{\pm}(t,x)$$

Then the main task is to prove the following proposition:

Proposition A.1 (Proposition 6.2 in [43]). Let $\max(0, 2-2s) < \delta < 1$. Let u_N^{\pm} be the unique solution to (2.1) with initial data $u_N^{\pm}(0, x) = \pm \epsilon N^{-3} \widetilde{\phi}_N(x) + N^{-\frac{4+\delta}{2}-s} \phi_N(x) \cos(Nx).$

$$\|u_{N}^{\pm} - u_{ap}^{\pm}\|_{H^{s}} = o(1),$$
(A.2)

for s > 0 and |t| < 1, as $N \to \infty$.

Then, we have

Once (A.2) holds true, one conclude that

$$\begin{aligned} \left\| u_N^+ - u_N^- \right\|_{H^s} &= N^{-\frac{4+\delta}{2}-s} \left\| \phi_N(x) \left(\cos\left(Nx - \Phi_N(t) + t\right) - \cos\left(Nx - \Phi_N(t) - t\right) \right) \right\|_{H^s} + o(1) \\ &= 2N^{-\frac{4+\delta}{2}-s} \left\| \phi_N(x) \sin\left(Nx - \Phi_N(t)\right) \right\|_{H^s} \left| \sin t \right| + o(1), \end{aligned}$$

which, in addition to Lemma A.2 below, implies

$$\lim_{N \to \infty} \left\| u_N^+ - u_N^- \right\|_{H^s} \ge c |\sin t| \sim c |t|$$

for |t| < 1. This completes the proof of Theorem 1.4.

We recall from [39, 43] the following useful lemmas to prove Proposition A.1.

Lemma A.2 (Lemma 2.3 in [39]). Let $s \ge 0$, $\delta > 0$ and $\gamma \in \mathbb{R}$. Then,

$$\lim_{N \to \infty} N^{-\frac{4+\delta}{2}-s} \|\phi_N(x)\sin(Nx+\gamma)\|_{H^s} = c_0 \|\phi\|_{L^2},$$

for some $c_0 > 0$.

Lemma A.3 (Lemma 6.3 in [43]). Let K be a positive integer and $K - 2 - s \ge k \ge 0$. Then, we have

$$\left\|\partial_x^k u_l^{\pm}(t,\cdot)\right\|_{L^2} \lesssim_K N^{-\frac{2-\delta}{2}-k(4+\delta)} \tag{A.3}$$

$$\left\|\partial_x^k u_l^{\pm}(t,\cdot)\right\|_{L^{\infty}} \lesssim_K N^{-3-k(4+\delta)} \tag{A.4}$$

$$\left\| u_l^{\pm}(t,\cdot) - u_{0,l}^{\pm}(\cdot) \right\|_{L^2} \lesssim_K N^{-15-3\delta}$$
(A.5)

Proof. The proof of (A.3) and (A.4) follows from a direct computation and Theorem 1.2, in particular, a priori bound (2.38). Moreover, the proof of (A.5) follows from a direct calculation in (2.1) and (A.3)–(A.4). The proof is almost identical to the proof of Lemma 6.3 in [43], thus we omit the details. \Box

Lemma A.4. Let

$$\mathcal{P}^{\pm}(t,x) := u_{ap,t}^{\pm} + u_{ap,5x}^{\pm} + 10\mu^2 \lambda^2 u_{ap,3x}^{\pm} + \mathcal{N}_2(u_{ap}^{\pm}) + \mathcal{N}_3(u_{ap}^{\pm}) + \mathcal{SN}(u_{ap}^{\pm}), \tag{A.6}$$

where $\mathcal{N}_2(\cdot)$, $\mathcal{N}_3(\cdot)$ and $\mathcal{SN}(\cdot)$ are defined as in (2.2)–(2.3), respectively. Let s > 0, $0 < \delta < 2$ and $|t| \leq 1$. Then, we have

$$\left\|\mathcal{P}^{\pm}(t,\cdot)\right\|_{L^{2}} \lesssim N^{-s-\delta} + N^{\frac{2-\delta}{2}-2s} + N^{-1-\delta-3s} + N^{1-\frac{3(4+\delta)}{2}-4s} + N^{1-2(4+\delta)-5s}.$$
(A.7)

Moreover, if $\sigma > 0$, we have

$$\left\|\mathcal{P}^{\pm}(t,\cdot)\right\|_{H^{\sigma}} \lesssim N^{-s-\delta+\sigma} + N^{\frac{2-\delta}{2}-2s+\sigma} + N^{-1-\delta-3s+\sigma} + N^{1-\frac{3(4+\delta)}{2}-4s+\sigma} + N^{1-2(4+\delta)-5s+\sigma}.$$
 (A.8)

Proof. It suffices to consider \mathcal{P}^+ , since an identical argument holds true for \mathcal{P}^- . We drop the super-index +. We decompose \mathcal{P} into $\mathcal{P}_1 + \mathcal{P}_2$, where $\mathcal{P}_2 = \mathcal{N}_3(u_{ap}) - \mathcal{N}_3(u_l) + \mathcal{SN}(u_{ap}) - \mathcal{SN}(u_l)$ and $\mathcal{P}_1 = \mathcal{P} - \mathcal{P}_2$. Lemma 6.4 in [43] exactly shows (A.7) and (A.8) for \mathcal{P}_1^{-13} . Our setting of ϕ , $\tilde{\phi}$ and u_l is essential to deal with

$$\Lambda := \epsilon N^{-\frac{4+\delta}{2}-s} \phi_N(x) (\partial_t + \partial_x^5 + 10\mu^2 \lambda^2 \partial_x^3 + \epsilon^{-1} u_l \partial_x^3) \cos\left(Nx - \Phi_N(t) - t\right)$$

contained in \mathcal{P}_1 (compared to F_4 in the proof of Lemma 6.4 in [43]). Indeed, a direct calculation in addition to $u_{0,l}(x) := \epsilon N^{-3} \widetilde{\phi}_N(x)$ and $\phi \widetilde{\phi} = \phi$ gives

$$\Lambda = N^{-\frac{4+\delta}{2}-s} \phi_N(x) \left(u_l N^3 - \epsilon \right) \sin \left(Nx - \Phi_N(t) - t \right) = N^{-\frac{4+\delta}{2}-s} N^3 \phi_N(x) \left(u_l - u_{0,l} \right) \sin \left(Nx - \Phi_N(t) - t \right),$$

which is handled by using (A.5). Thus, it suffices to show (A.7) and (A.8) for \mathcal{P}_2 . Putting first $u_{ap} = u_l + u_h$ into $10u_{ap}^2 u_{ap,3x} - 10u_l^2 u_{l,x}$ in \mathcal{P}_2 , one has

$$10u_l^2 u_{h,xxx} + 20u_l u_h u_{l,xxx} + 20u_l u_h u_{h,xxx} + 10u_h^2 u_{l,xxx} + 10u_h^2 u_{h,xxx}.$$
(A.9)

Note that

$$u_{h,x} = N^{-\frac{4+\delta}{2}-s} \left(\partial_x \phi_N(x) \cos\left(Nx - \Phi_N(t) - t\right) + \phi_N(x)\partial_x \cos\left(Nx - \Phi_N(t) - t\right)\right) \\ = N^{-\frac{4+\delta}{2}-s} \left(N^{-(4+\delta)}\phi_{N,x}(x) \cos\left(Nx - \Phi_N(t) - t\right) - N\phi_N(x) \sin\left(Nx - \Phi_N(t) - t\right)\right).$$

¹³A small difference between \mathcal{P}_1 and F in Lemma 6.4 in [43] does not make any trouble. Indeed, our setting of u_h corresponds to (2.1), so that one can immediately apply the argument in the proof of Lemma 6.4 in [43] to our case. Moreover, the cubic term with one derivative in $\mathcal{N}_2(u_{ap})$ can be dealt with similarly as $\mathcal{SN}(u_{ap})$.

Thus, one can see that the worst term arises from the case when the derivative acts on $\cos(Nx - \Phi_N(t) - t)$. Using Lemmas A.2 and A.3, one estimates

$$\|(\mathbf{A}.9)\|_{L^2} \lesssim N^{-3-s} + N^{-\frac{4+\delta}{2}-2s} + N^{-1-\delta-3s}.$$

An analogous argument yield

$$\begin{split} \left\| u_{ap,x}^{3} - u_{l,x}^{3} \right\|_{L^{2}} &\lesssim N^{-5-2(4+\delta)-s} + N^{-1-\frac{3(4+\delta)}{2}-2s} + N^{-1-\delta-3s}, \\ \\ \left\| u_{ap}u_{ap,x}u_{ap,xx} - u_{l}u_{l,x}u_{l,xx} \right\|_{L^{2}} &\lesssim N^{-8-\delta-s} + N^{-\frac{4+\delta}{2}-2s} + N^{-1-\delta-3s}, \\ \\ \left\| u_{ap}^{4}u_{ap,x} - u_{l}^{4}u_{l,x} \right\|_{L^{2}} &\lesssim N^{-11-s} + N^{-8-\frac{4+\delta}{2}-2s} + N^{-5-(4+\delta)-3s} + N^{-2-\frac{3(4+\delta)}{2}-4s} + N^{1-2(4+\delta)-5s}, \\ \\ \\ \left\| u_{ap}u_{ap,x} - u_{l}u_{l,x} \right\|_{L^{2}} &\lesssim N^{-2-s} + N^{1-\frac{4+\delta}{2}-2s} \end{split}$$

and

$$\left\|u_{ap}^{3}u_{ap,x}-u_{l}^{3}u_{l,x}\right\|_{L^{2}} \lesssim N^{-8-s} + N^{-5-\frac{4+\delta}{2}-2s} + N^{-2-(4+\delta)-3s} + N^{1-\frac{3(4+\delta)}{2}-4s}.$$

Collecting all, we completes the proof of (A.7). Moreover, the fractional Leibniz rule ensure at least $\|\mathcal{P}\|_{\dot{H}^{\sigma}} \lesssim_{\sigma} N^{\sigma} \|\mathcal{P}\|_{L^2}$, which in addition to (A.7) implies (A.8), since $u_{l,t} + u_{l,5x} + 10\mu^2\lambda^2 u_{l,3x} + 30\mu^4\lambda^4 u_{l,x} + \mathcal{N}_2(u_l) + \mathcal{N}_3(u_l) + \mathcal{SN}(u_l) = 0$ and the others contains at least one u_h . We complete the proof.

A.2. Proof of Proposition A.1. Let $w^{\pm} := u_N^{\pm} - u_{ap}^{\pm}$. We only show $||w^+||_{H^s} = o(1)$ as $N \to \infty$ and drop the super-index +. For $s \ge 2$, the local well-posedness theory is available. A direct calculation gives

$$\Gamma w + \mathcal{N}_2(u_N) - \mathcal{N}_2(u_{ap}) + \mathcal{N}_3(u_N) - \mathcal{N}_3(u_{ap}) + \mathcal{SN}(u_N) - \mathcal{SN}(u_{ap}) + \mathcal{P} = 0,$$

where $\Gamma := \partial_t + \partial_x^5 + 10\mu^2\lambda^2\partial_x^3$ and \mathcal{P} is as in (A.6). For $2 \leq \sigma$, the local well-posedness, in particular (2.38), ensures

$$\|u_N\|_{C_T H^{\sigma}} + \|u_N\|_{F^{\sigma}(T)} \lesssim \|u_N(0)\|_{H^{\sigma}} \lesssim N^{\sigma-s}.$$
(A.10)

Moreover, a direct calculation and the local theory (for u_l) gives

$$||u_{ap}||_{C_T H^{\sigma}} + ||u_{ap}||_{F^{\sigma}(T)} \lesssim N^{-\frac{2-\delta}{2}} + N^{\sigma-s}.$$
 (A.11)

Using Propositions 2.15, Propositions 2.7, 2.8, 2.9 and 2.13, and (A.7) under (A.10) and (A.11), one concludes

$$\|w\|_{F^0(T)} \lesssim \|\mathcal{P}\|_{L^1_T L^2_x} = O(N^{-s-\beta})$$

for $\beta = \min(\delta, -\frac{2-\delta}{2} + s) > 0$, which, in addition to Proposition 2.14, implies

$$\|w\|_{L^{\infty}_{T}L^{2}_{x}} = O(N^{-s-\beta}). \tag{A.12}$$

Furthermore, an analogous argument (but using (A.8) instead of (A.7)) in addition to

$$||u_{ap}||_{F^{2}s(T)} ||w||_{F^{0}(T)} = O(N^{s}N^{-s-\beta}) = O(N^{-\beta}),$$

ensures $||w||_{F^s(T)} = O(N^{-\beta})$, which concludes (A.2) as $N \to \infty$ for $s \ge 2$.

To fill the regularity range 0 < s < 2, we use the conservation law and the interpolation theorem. H^2 conservation law (2.39) and a direct calculation yield

$$||u_N||_{H^2} \lesssim N^{2-s}$$
 and $||u_{ap}||_{H^2} \lesssim N^{2-s}$,

respectively, which concludes

$$\|w\|_{H^2} \lesssim N^{2-s}.$$
 (A.13)

The interpolation between (A.12) and (A.13) ensures

$$\|w\|_{H^s} \lesssim \|w\|_{L^2}^{1-\frac{s}{2}} \|w\|_{H^3}^{\frac{s}{2}} \lesssim N^{-\frac{\beta(2-s)}{2}},$$

which proves (A.2) as $N \to \infty$ for 0 < s < 2.

Appendix B. Proof of Lemma 3.6.

We are going to prove the identity (3.5)

$$\tilde{B}_{\mu,t} = (\alpha^2 + \beta^2)^2 B_{\mu} + 2(\alpha^2 - \beta^2 - 5\mu^2) (B_{\mu,xx} + 2B_{\mu}^3 + 6\mu B_{\mu}^2).$$

Firstly and for the sake of simplicity, we will use the following notation:

$$\begin{split} A_1 &:= (\alpha^2 + \beta^2)^2, \quad A_2 := 2(\alpha^2 - \beta^2 - 5\mu^2), \\ \Delta &= \alpha^2 + \beta^2 - 4\mu^2, \quad e^z = \cosh(z) + \sinh(z), \\ D &:= f^2 + g^2, \quad \text{where} \quad f, g \text{ and its derivatives are given by:} \end{split}$$

$$f = \cosh(\beta y_2) - \frac{2\beta\mu}{\alpha\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}} (\alpha\cos(\alpha y_1) - \beta\sin(\alpha y_1)),$$

$$f_1 := f_x = \beta\sinh(\beta y_2) + \frac{2\beta\mu}{\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}} (\beta\cos(\alpha y_1) + \alpha\sin(\alpha y_1)),$$

$$f_2 := f_t = \beta\gamma_5\sinh(\beta y_2) + \frac{2\beta\delta_5\mu}{\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}} (\beta\cos(\alpha y_1) + \alpha\sin(\alpha y_1)),$$

(B.1)

$$f_3 := f_{xx} = \beta^2 \cosh(\beta y_2) + \frac{2\alpha\beta\mu}{\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}} (-\alpha\cos(\alpha y_1) + \beta\sin(\alpha y_1)),$$

$$f_4 := f_{xxx} = \beta^3 \sinh(\beta y_2) - \frac{2\alpha^2\beta\mu}{\sqrt{\alpha^2 + \beta^2}\sqrt{\Delta}} (\beta\cos(\alpha y_1) + \alpha\sin(\alpha y_1))$$

and

$$g = \frac{\beta\sqrt{\alpha^2 + \beta^2}}{\alpha\sqrt{\Delta}}\sin(\alpha y_1) - \frac{2\beta\mu e^{\beta y_2}}{\Delta},$$

$$g_1 := g_x = \frac{\beta\sqrt{\alpha^2 + \beta^2}}{\sqrt{\Delta}}\cos(\alpha y_1) - \frac{2\beta^2\mu e^{\beta y_2}}{\Delta},$$

$$g_2 := g_t = \frac{\beta\delta_5\sqrt{\alpha^2 + \beta^2}}{\sqrt{\Delta}}\cos(\alpha y_1) - \frac{2\beta^2\gamma_5\mu e^{\beta y_2}}{\Delta},$$

$$g_3 := g_{xx} = -\frac{\alpha\beta\sqrt{\alpha^2 + \beta^2}}{\sqrt{\Delta}}\sin(\alpha y_1) - \frac{2\beta^3\mu e^{\beta y_2}}{\Delta},$$

$$g_4 := g_{xxx} = -\frac{\alpha^2\beta\sqrt{\alpha^2 + \beta^2}}{\sqrt{\Delta}}\cos(\alpha y_1) - \frac{2\beta^4\mu e^{\beta y_2}}{\Delta},$$
(B.2)

where velocities (γ_5, δ_5) are given in (3.3). From the explicit expression of the breather solution (3.2) but now written in terms of the above derivatives (B.1)–(B.2), we obtain that:

$$B_{\mu} = 2 \frac{g_1 f - f_1 g}{D}$$
 and $\tilde{B}_{\mu,t} = 2 \frac{g_2 f - f_2 g}{D}$. (B.3)

Moreover we get

$$B_{\mu}^{2} = 4\left(\frac{g_{1}f - f_{1}g}{D}\right)^{2}$$
 and $B_{\mu}^{3} = 8\left(\frac{g_{1}f - f_{1}g}{D}\right)^{3}$. (B.4)

Now, we compute $B_{\mu,xx}$. First we get

$$B_{\mu,x} = -\frac{2}{D^2} \left(f^3 g_3 - f^2 (2f_1 g_1 + f_3 g) + fg \left(2f_1^2 + gg_3 - 2g_1^2 \right) + g^2 (2f_1 g_1 - f_3 g) \right),$$

and then

$$B_{\mu,xx} = 2\frac{M_1}{D^3},$$
 (B.5)

where

$$M_{1} := \left(f^{5}g_{4} - f^{4}(3f_{1}g_{3} + 3f_{3}g_{1} + f_{4}g) + 2f^{3} \left(3f_{1}^{2}g_{1} + 3f_{1}f_{3}g + g^{2}g_{4} - 3gg_{1}g_{3} - g_{1}^{3} \right) - 2f^{2}g \left(3f_{1}^{3} - 9f_{1}g_{1}^{2} + f_{4}g^{2} \right) + fg^{2} \left(-18f_{1}^{2}g_{1} + 6f_{1}f_{3}g + g^{2}g_{4} - 6gg_{1}g_{3} + 6g_{1}^{3} \right) + g^{3} \left(2f_{1}^{3} + f_{1} \left(3gg_{3} - 6g_{1}^{2} \right) + g(3f_{3}g_{1} - f_{4}g) \right) \right),$$
(B.6)

and therefore from (B.3), (B.4), (B.5) and (B.6), we get

$$A_1 B_\mu + A_2 (B_{\mu,xx} + 2B_\mu^3 + 6\mu B_\mu^2) = \frac{M_2}{D^3},$$
(B.7)

where

$$M_2 := 2\Big(A_1 D^2 (fg_1 - f_1g) + A_2 (8(fg_1 - f_1g)^3 + 12\mu D(f_1g - fg_1)^2 + M_1)\Big),\tag{B.8}$$

Now, we verify by using the symbolic software *Mathematica* that, after expanding f's and g's terms (B.1)–(B.2) and lengthy rearrangements, the above term (B.8) simplifies as follows:

$$M_2 = 2D^2(g_2f - gf_2)$$

Finally, remembering (B.7), we have that

$$A_1B_{\mu} + A_2(B_{\mu,xx} + 2B_{\mu}^3 + 6\mu B_{\mu}^2) = \frac{M_2}{D^3} = \frac{2D^2(g_2f - gf_2)}{D^3} = \tilde{B}_{\mu,t}$$

and we conclude.

References

- M. Ablowitz and P. Clarkson, Solitons, nonlinear evolution equations and inverse scattering, London Mathematical Society Lecture Note Series, 149. Cambridge University Press, Cambridge, 1991.
- [2] M.A. Alejo, On the ill-posedness of the Gardner equation, J. Math. Anal. Appl., 396 no. 1, 256-260 (2012).
- [3] M.A. Alejo, Well-posedness and Stability results for solitons of the Gardner equation, NoDEA/Nonlinear Differential Equations and Applications, Volume 19, Number 4 (2012), 503–520.
- [4] M.A. Alejo and E. Cardoso, On the ill-posedness of the 5th-order Gardner equation, preprint, arXiv:1810.10434 [math.AP]
- [5] M.A. Alejo and C. Muñoz, Nonlinear stability of mKdV breathers, Comm. Math. Phys., 37 (2013), 2050–2080.
- [6] M.A. Alejo, Nonlinear stability of Gardner breathers, Jour. Diff. Equat. 264, n.2, 1192-1230 (2018).
- J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. Parts I, II, Geom. Funct. Anal. 3 (1993) 107–156, 209–262.
- [8] J. Bourgain, On the Cauchy problem for periodic KdV-type equations, Proceedings of the Conference in Honor of Jean-Pierre Kahane (Orsay, 1993). J. Fourier Anal. Appl. 1995, Special Issue, 17–86.
- [9] N. Burq, P. Gérard and N. Tzvetkov, An instability property of the nonlinear Schrödinger equation on S^d, Math. Res. Lett. 9 (2002), no. 2-3, 323–335.
- [10] N. Burq, P. Gérard and N. Tzvetkov, Two singular dynamics of the nonlinear Schrödinger equation on a plane domain, Geom. Funct. Anal. 13 (2003), no. 1, 1–19.
- M. Cavalcante and C. Kwak, The initial-boundary value problem for the Kawahara equation on the half-line, preprint, arXiv:1805.05229 [math.AP].
- [12] M. Cavalcante and C. Kwak, Local well-posedness of the fifth-order KdV-type equations on the half-line, accepted for the publication in CPAA, http://arxiv.org/abs/1808.06494.
- [13] W. Chen and Z. Guo, Global well-posedness and I-method for the fifth-order Korteweg-de Vries equation, J. Anal. Math. 114 (2011) 121–156.
- [14] W. Chen, J. Li, C. Miao and J. Wu, Low regularity solutions of two fifth-order KdV type equations, J. Anal. Math. 107 (2009) 221–238.
- [15] M. Christ, J. Colliander, T. Tao, Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations, Amer. J. Math. 125 (2003), no. 6, 1235–1293.
- [16] M. Christ, J. Colliander, T. Tao, A priori bounds and weak solutions for the nonlinear Schrödinger equation in Sobolev space of negative order, Journal of Functional Analysis 254 (2008), 368–395.
- [17] S. Cui and S. Tao, Strichartz estimates for dispersive equations and solvability of the Kawahara equation, J. Math. Anal. Appl. 304 (2005) 683–702.
- [18] C.S. Gardner, M.D. Kruskal and R. Miura, Korteweg-de Vries equation and generalizations. II. Existence of conservation laws and constants of motion, J. Math. Phys. 9, no. 8, 1204–1209 (1968).
- [19] J.F. Gomes, G.S. França and A.H. Zimerman, Nonvanishing boundary condition for the mKdV hierarchy and the Gardner equation 2012 J. Phys. A: Math. Theor. 45 015207.
- [20] Z. Guo, Local well-posedness and a priori bounds for the modified Benjamin-Ono equation, Advances in Differential Equations, 16/11-12 (2011), 1087–1137.
- [21] Z. Guo, Local well-posedness for dispersion generalized Benjamin-Ono equations in Sobolev spaces, J. Differential Equations 252 (2012) 2053–2084.
- [22] Z.Guo, C. Kwak, S. Kwon, Rough solutions of the fifth-order KdV equations, J. Funct. Anal. 265 (2013) 2791–2829.
- [23] Z. Guo, T. Oh, Non-existence of solutions for the periodic cubic NLS below L², Int. Math. Res. Not. IMRN (2018), no. 6, 16561729.
- [24] R. Grimshaw, A. Slunyaev and E. Pelinovsky, Generation of solitons and breathers in the extended Korteweg-de Vries equation with positive cubic nonlinearity. Chaos 20 (2010), n.1, 01310201–01310210.
- [25] A. Grünrock, On the hierarchies of higher order mKdV and KdV equations, Cent. Eur. J. Math. Vol. 8(3), 500-536, (2010).

- [26] Z. Guo, L. Peng, B. Wang and Y. Wang, Uniform well-posedness and inviscid limit for the Benjamin-Ono-Burgers equation, Advances in Mathematics 228 (2011) 647–677.
- [27] A. Ionescu, C. Kenig, Global well-posedness of the Benjamin-Ono equation in low-regularity spaces, J. Amer. Math. Soc. 20 (3) (2007) 753–798.
- [28] A. Ionescu, C. Kenig, D. Tataru, Global well-posedness of the KP-I initial-value problem in the energy space, Invent. Math. 173 (2) (2008) 265–304.
- [29] T. Kato, Well-posedness for the fifth order KdV equation, Funkcialaj Ekvacioj 55 (1) (2012) 17–53.
- [30] C. Kenig and D. Pilod, Well-posedness for the fifth-order KdV equation in the energy space, Trans. Amer. Math. Soc. 367 (2015) 2551-2612.
- [31] C. Kenig and D. Pilod, Local well-posedness for the KdV hierarchy at high regularity, Adv. Diff. Eq., 21 (2016), 801–836.
- [32] C. E. Kenig, G. Ponce and L. Vega, On the hierarchy of the generalized KdV equations, Singular limits of dispersive waves (Lyon, 1991), NATO Adv. Sci. Inst. Ser. B Phys., vol. 320, Plenum, New York, 1994, pp. 347356.
- [33] C. E. Kenig, G. Ponce and L. Vega, Well-posedness of the initial value problem for the Korteweg-de Vries equation, J. Amer. Math. Soc. 4 (1991), no. 2, 323–347.
- [34] C. E. Kenig, G. Ponce and L. Vega, Oscillatory integrals and regularity of dispersive equations, Indiana U. Math. J 40 (1991) 33–69.
- [35] C. E. Kenig, G. Ponce and L. Vega, Higher-order nonlinear dispersive equations, Proc. Amer. Math. Soc. 122 (1994), no. 1, 157166, DOI 10.2307/2160855.
- [36] C. Kenig, G. Ponce, L. Vega, A bilinear estimate with applications to the KdV equation, J. Amer. Math. Soc. 9 (1996) 573–603.
- [37] C.E. Kenig, G. Ponce and L. Vega, On the ill-posedness of some canonical dispersive equations, Duke Math. J. 106, no. 3, 617–633 (2001).
- [38] H. Koch, D. Tataru, A priori bounds for the 1D cubic NLS in negative Sobolev spaces, Int. Math. Res. Not. IMRN 16 (2007), Art. ID rnm053, 36, DOI 10.1093/imrn/rnm053. MR2353092 (2010d:35307)
- [39] H. Koch, N. Tzvetkov, Nonlinear wave interactions for the Benjamin-Ono equation, Int. Math. Res. Not. 30 (2005) 1833–1847.
- [40] H. Koch, N. Tzvetkov, On finite energy solutions of the KP-I equation, Math. Z. 258 (2008), no. 1, 55-68.
- [41] C. Kwak Low regularity Cauchy problem for the fifth-order modified KdV equations on T, Journal of Hyperbolic Differential Equations Vol. 15, No. 3 (2018) 463–557. http://dx.doi.org/10.1142/S0219891618500170.
- [42] C. Kwak Local well-posedness for the fifth-order KdV equations on T, J. Differential Equations 260 (2016) 7683–7737. http://dx.doi.org/10.1016/j.jde.2016.02.001.
- [43] S. Kwon, On the fifth order KdV equation: Local well-posedness and lack of uniform continuity of the solution map J. Differential Equations, 245 (2008) 2627–2659.
- [44] S. Kwon, Well posedness and Ill-posedness of the Fifth-order modified KdV equation, Electr. Journal Diff. Equations. vol. 2008, n.1, 1–15 (2008).
- [45] G.L. Lamb, *Elements of Soliton Theory*, Pure Appl. Math., Wiley, New York, 1980.
- [46] F. Linares, A higher order modified Korteweg-de Vries equation, Comp. Appl. Math. 14, n.3, 253-267, (1995).
- [47] Y. Matsuno, Bilinearization of Nonlinear Evolution Equations: Higher Order mKdV, Jour. Phys. Soc.Japan, 49, n.2 (1980).
- [48] L. Molinet, D. Pilod and S. Vento, Unconditional uniqueness for the modified Korteweg- de Vries equation on the line, to appear in Rev. Mat. Iber. (2018).
- [49] L. Molinet, D. Pilod and S. Vento, On unconditional well-posedness for the periodic modified Korteweg-De Vries equation, to appear in J. Math. Soc. Japan (2018)
- [50] L. Molinet, J.C. Saut and N. Tzvetkov, Ill-posedness issues for the BenjaminOno and related equations, SIAM J.Math. Anal. 33 (2001) 982–988.
- [51] L. Molinet, J.C. Saut and N. Tzvetkov, Well-posed and ill-posedness results for the KadomtsevPetviashvili-I equation, Duke Math. J. 115 (2) (2002) 353–384.
- [52] D. Pilod, On the Cauchy problem for higher-order nonlinear dispersive equations, J. Differential Equations 245 (2008), no. 8, 20552077, DOI 10.1016/j.jde.2008.07.017.
- [53] G. Ponce, Lax pairs and higher order models for water waves, J. Differential Equations 102 (2) (1993) 360-381.
- [54] T. Tao, Multilinear weighted convolution of L^2 functions and applications to nonlinear dispersive equations, Amer. J. Math. 123 (5) (2001) 839–908.
- [55] K. Tsugawa, Parabolic smoothing effect and local well-posedness of fifth-order semilinear dispersive equations on torus, Harmonic analysis and nonlinear partial differential equations, 177193, RIMS Kkyroku Bessatsu, B60, Res. Inst. Math. Sci. (RIMS), Kyoto, 2016, arXiv:1707.09550 [math.AP].

Departamento de Matemática, Universidade Federal de Santa Catarina, Brasil

FACULTAD DE MATEMÁTICAS, PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, CAMPUS SAN JOAQUÍN. AVDA. VICUÑA MACKENNA 4860, SANTIAGO, CHILE AND INSTITUTE OF PURE AND APPLIED MATHEMATICS, CHONBUK NATIONAL UNIVERSITY

E-mail address: miguel.alejo@ufsc.br

E-mail address: chkwak@mat.uc.cl