Automatic complexity of Fibonacci and Tribonacci words

Bjørn Kjos-Hanssen* October 15, 2020

Abstract

For a complexity function C, the lower and upper C-complexity rates of an infinite word ${\bf x}$ are

$$\underline{C}(\mathbf{x}) = \liminf_{n \to \infty} \frac{C(\mathbf{x} \upharpoonright n)}{n}, \quad \overline{C}(\mathbf{x}) = \limsup_{n \to \infty} \frac{C(\mathbf{x} \upharpoonright n)}{n}$$

respectively. Here $\mathbf{x} \upharpoonright n$ is the prefix of x of length n. We consider the case $C = \mathbf{A_N}$, the nondeterministic automatic complexity. If these rates are strictly between 0 and 1/2, we call them intermediate. Our main result is that words having intermediate $\mathbf{A_N}$ -rates exist, viz. the infinite Fibonacci and Tribonacci words.

1 Introduction

The automatic complexity of Shallit and Wang [10] is the minimal number of states of an automaton accepting only a given word among its equal-length peers. This paper continues a line of investigation into the automatic complexity of particular words of interest such as

- maximal length sequences for linear feedback shift registers [7],
- overlap-free and almost square-free words [3], and
- random words [8].

All these examples have high complexity: to be precise, they have maximal automatic complexity rate (Definition 8). On the other hand, a periodic word has low complexity and a rate of 0. In the present paper we give the first examples of infinite words with intermediate automatic complexity rate: the infinite Fibonacci and Tribonacci words.

^{*}This work was partially supported by grants from the Simons Foundation (#315188 and #704836 to Bjørn Kjos-Hanssen) and Decision Research Corporation (Automatic Complexity of Fibonacci arrays, University of Hawai'i Foundation Account #129-4770-4).

Automatic complexity is an automata-based and length-conditional analogue of Sipser's CD complexity [11] which is in turn a computable analogue of the noncomputable Kolmogorov complexity. The nondeterministic case was taken up by Hyde and Kjos-Hanssen [3]. We recall our basic notions. Let |x| denote the length of a word x.

Definition 1 ([10]). The nondeterministic automatic complexity $A_N(x)$ of a word x is the minimal number of states of a nondeterministic finite automaton M (without ϵ -transitions) such that M accepts x and moreover there is only one accepting path in M of length |x|. We let A^- denote the deterministic but non-total automatic complexity, defined as follows: automata are required to be deterministic, but the transition functions need not be total: there does not need to be a transition for every symbol at every state.

Shallit and Wang's original automatic complexity $\mathbf{A}(x)$ does have the totality requirement.

1.1 Fibonacci and Tribonacci words

Definition 2 (k-bonacci numbers). For $k \geq 2$ and $n \geq 0$, the nth k-bonacci number $w_n = w_n^{(k)}$ is defined by $w_n = 0$ if $n \leq k - 1$, $w_k = 1$, and $w_n = \sum_{i=n-k}^{n-1} w_i$ for $n \geq k + 1$.

In particular, the Fibonacci numbers are the 2-bonacci numbers.

Let $\Sigma_k = \{a_0, \dots, a_{k-1}\}$ be an alphabet of cardinality $k \geq 0$. We shall denote specific symbols in the alphabet as $0 = a_0$, $1 = a_1$ and so on.

Definition 3 (k-bonacci words). We define the k-morphism $\varphi_k: \Sigma_k \to \Sigma_k^*$ by

$$\varphi_k(\mathbf{a}_i) = \mathbf{a}_0 \mathbf{a}_{i+1}, \quad 0 \le i \le k-2,
\varphi_k(\mathbf{a}_{k-1}) = \mathbf{a}_0.$$

We also let φ act on words of length greater than 1, by the morphism property

$$\varphi(uv) = \varphi(u)\varphi(v).$$

Let ε be the empty word. The k-bonacci word $W_n = W_n^{(k)}$ is defined by

$$\begin{array}{rcl} W_n & = & \varepsilon, \quad 0 \leq n \leq k-2, \\ W_{k-1} & = & \mathtt{a}_{k-1}, \\ W_n & = & \varphi_k(W_{n-1}), \quad n \geq k. \end{array}$$

Lemma 4. The length of the nth k-bonacci word is equal to the nth k-bonacci number: $|W_n^{(k)}| = w_n^{(k)}$.

Lemma 5. For all $k \geq 2$ and $n \geq 2k-1$, if W_n is the nth k-bonacci word then

$$W_n = W_{n-1} \dots W_{n-k}.$$

We then say Fibonacci for 2-bonacci and Tribonacci for 3-bonacci. Thus the finite Tribonacci words T_n are defined by

$$T_0 = \varepsilon$$
 $T_1 = \varepsilon$
 $T_2 = 2$
 $T_3 = 0$
 $T_4 = 01$
 $T_n = T_{n-1}T_{n-2}T_{n-3}, n \ge 5$

and the finite Fibonacci words F_n by

$$F_0 = \varepsilon$$
 $F_1 = 1$ $F_2 = 0$ $F_n = F_{n-1}F_{n-2}, n \ge 3.$

Definition 6. The infinite k-bonacci word $\mathbf{f}^{(k)}$ is the fixed point $\varphi_k^{(\infty)}(0)$ of the morphism φ_k .

Thus, the infinite Tribonacci word

$$T = \mathbf{f}^{(3)} = 0102010010201...$$

is a fixed point of the morphism $0 \to 01, 1 \to 02, 2 \to 0$. It is a variant of the Fibonacci word obtained from the morphism $0 \to 01, 1 \to 0$.

2 Lower bounds from critical exponents

A word is square-free if it does not contain any subword of the form xx (denoted x^2), |x| > 0. Shallit and Wang showed that a square-free word has high automatic complexity, and we shall show that integrality of powers is not crucial: that is, we shall use critical exponents.

Definition 7. Let \mathbf{w} be an infinite word over the alphabet Σ , and let x be a finite word over Σ . Let $\alpha > 0$ be a rational number. The word x is said to occur in \mathbf{w} with exponent α if there is a subword y of \mathbf{w} with $y = x^a x_0$ where x_0 is a prefix of x, a is the integer part of α , and $|y| = \alpha |x|$. We say that y is an α -power. The word \mathbf{w} is α -power-free if it contains no subwords which are α -powers.

The critical exponent for \mathbf{w} is the supremum of the α for which \mathbf{w} has α -powers, or equivalently the infimum of the α for which \mathbf{w} is α -power-free.

Definition 8. Fix a finite alphabet Σ . For an infinite word $\mathbf{w} \in \Sigma^{\infty}$, let $\mathbf{w} \upharpoonright n$ denote the prefix of \mathbf{w} of length n. Let $C: \Sigma^* \to \mathbb{N}$. The lower C-complexity rate of \mathbf{w} is

$$\underline{C}(\mathbf{w}) = \liminf_{n \to \infty} \frac{C(\mathbf{w} \upharpoonright n)}{n}.$$

The upper C-complexity rate of \mathbf{w} is

$$\underline{C}(\mathbf{w}) = \limsup_{n \to \infty} \frac{C(\mathbf{w} \upharpoonright n)}{n}.$$

If these are equal we may speak simply of the C-complexity rate. In the case where $C = A_N$ we may speak of automatic complexity rate.

Definition 9 (Fibonacci constant). Let $\phi = \frac{1+\sqrt{5}}{2}$, the positive root of $\phi^2 = \phi + 1$.

Definition 10 (Tribonacci constant). Let

$$\xi = \frac{1}{3} \left(1 + \sqrt[3]{19 - 3\sqrt{33}} + \sqrt[3]{19 + 3\sqrt{33}} \right).$$

Theorem 11 ([9]). The critical exponent of the infinite Fibonacci word $\mathbf{f}^{(2)}$ is $2 + \phi \approx 3.6$.

Tan and Wen [12] studied critical exponents, calling them free indices.

Theorem 12 (Tan and Wen [12, Theorem 4.5]). The critical exponent of the infinite Tribonacci word $\mathbf{f}^{(3)}$ is $3 + \frac{1}{2}(\theta^2 + \theta^4)$, where

$$\theta = \frac{1}{3} \left(-1 - \frac{2}{(17 + 3\sqrt{33})^{1/3}} + (17 + 3\sqrt{33})^{1/3} \right)$$

$$\approx 0.543689012692076...$$

is the unique real root of the equation $\theta^3 + \theta^2 + \theta = 1$.

Lemma 13. The critical exponent of $\mathbf{f}^{(3)}$ is the real zero

$$2 + \frac{1}{6}\sqrt[3]{54 - 6\sqrt{33}} + \frac{\sqrt[3]{9 + \sqrt{33}}}{6^{2/3}}$$

= 3.19148788...

of the polynomial $2x^3 - 12x^2 + 22x - 13$.

Lemma 13 follows from Theorem 12 by computer software (Wolfram Alpha).

Definition 14. Let x be a word of length n, $x = x_1, \ldots, x_n$. Two occurrences of words a (starting at position i) and b (starting at position j) in a word x are disjoint if x = uavbw where u, v, w are words and |u| = i - 1, |uav| = j. If in addition |v| > 0 then we say that these occurrences of a and b are strongly disjoint.

Theorem 15 ([8]). If the critical exponent of a word x is at most $\gamma \geq 2$ then there is an $m \geq 0$ and a set of m many strongly disjoint at-least-square powers in x with $A_N(x) \geq \frac{n+1-m}{\gamma}$.

Theorem 16. If the critical exponent of a word x is at most $\gamma \geq 2$ then $A_N(x) \geq \frac{n+1-\sqrt{2n}}{\gamma}$.

Proof. By uniqueness of path the m many powers in Theorem 15 must have distinct base lengths. Thus the base lengths add up to at least $\sum_{k=1}^{m} k = m(m+1)/2$, which implies $m(m+1)/2 \le n$. Consequently $m \le \sqrt{2n}$ and

$$A_{N}(x) \ge \frac{n+1-m}{\gamma} \ge \frac{n+1-\sqrt{2n}}{\gamma}.$$

Theorem 17. The A_N -complexity rate of the infinite Fibonacci word $\mathbf{f}^{(2)}$ is at least

$$\frac{2}{5+\sqrt{5}} = 0.27639\dots$$

The A_N -rate of the infinite Tribonacci number $\mathbf{f}^{(3)}$ is at least

the real root of $-2 + 12x - 22x^2 + 13x^3$.

Proof. These two facts now follow by applying Theorem 16 with Theorems 11 and 12, respectively. \Box

Karhumäki [4] showed that the Fibonacci words contain no 4th power and this implies (the deterministic version is in Shallit and Wang 2001 [10, Theorem 9]) $A_N(x) \ge (n+1)/4$.

Theorem 18. The A_N -complexity rate of the infinite k-bonacci word $\mathbf{f}^{(k)}$ is at least 1/4 for any $k \geq 2$.

Proof. Glen [2] showed that the k-bonacci word has no fourth power for any $k \geq 2$. Thus, the critical exponent is at most 4 and by Theorem 16 we are done.

3 Upper bounds from factorizations

There are many factorization result possible for k-bonacci words. Even their definitions like $F_n = F_{n-1}F_{n-2}$ are factorizations. We shall prove some such results that help us obtain upper bounds on automatic complexity: Theorem 20 and 21. In the following, for convenience we renumber by defining $\tilde{T}_n = T_{n+3}$.

Definition 19. For $n \geq 0$ and $0 \leq k \leq n$, we let $\langle k \rangle_n = \tilde{T}_{n-k}$. We also write $||k||_n = |\langle k \rangle_n|$, the length of $\langle k \rangle_n$. When n is understood from context we write $\langle k \rangle = \langle k \rangle_n$ and $||k|| = ||k||_n$.

Theorem 20. For large enough n, $\tilde{T}_{n-2}^2 \prod_{k=6}^{3\lfloor n/3 \rfloor + 1} \tilde{T}_{n-k}$ is a prefix of \tilde{T}_n .

Proof. Note that the equation $\tilde{T}_n = \tilde{T}_{n-1}\tilde{T}_{n-2}\tilde{T}_{n-3}$ holds for $n \geq 2$. Thus we can write

$$\langle n - 2 \rangle = \langle n - 1 \rangle \langle n \rangle \langle n + 1 \rangle \tag{1}$$

but we cannot expand $\langle n-1 \rangle$. The idea now is to use a loop of length 13 followed by one of length 6:

$$\begin{split} \tilde{T}_n &= \langle 1 \rangle \langle 2 \rangle \langle 3 \rangle = (\langle 2 \rangle \langle 3 \rangle \langle 4 \rangle) \langle 2 \rangle \langle 3 \rangle = \langle 2 \rangle \langle 3 \rangle \langle 4 \rangle (\langle 3 \rangle \langle 4 \rangle \langle 5 \rangle) \langle 3 \rangle \\ &= \langle 2 \rangle \langle 3 \rangle \langle 4 \rangle (\langle 4 \rangle \langle 5 \rangle \langle 6 \rangle) \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle = \langle 2 \rangle \langle 3 \rangle \langle 4 \rangle (\langle 5 \rangle \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle \\ &= \langle 2 \rangle^2 \langle 6 \rangle \langle 7 \rangle \ \langle 5 \rangle \ \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \ \langle 3 \rangle \\ &= \langle 2 \rangle^2 \langle 6 \rangle \langle 7 \rangle \ (\langle 8 \rangle \langle 9 \rangle \langle 10 \rangle \langle 8 \rangle \langle 9 \rangle \langle 7 \rangle \langle 8 \rangle) \ \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \ \langle 3 \rangle \\ &= \langle 2 \rangle^2 \left(\prod_{k=6}^{10} \langle k \rangle \right) \langle 8 \rangle \ \langle 9 \rangle \langle 7 \rangle \langle 8 \rangle \ \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \ \langle 3 \rangle \\ &= \langle 2 \rangle^2 \left(\prod_{k=6}^{13} \langle k \rangle \right) \ \langle 11 \rangle \langle 12 \rangle \langle 10 \rangle \langle 11 \rangle \langle 9 \rangle \langle 7 \rangle \langle 8 \rangle \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle. \end{split}$$

Thus for m = 4 we have

$$\langle 0 \rangle = \langle 2 \rangle^2 \left(\prod_{k=6}^{3m+1} \langle k \rangle \right) \langle 3m-1 \rangle \left(\prod_{M=m}^0 \langle 3M \rangle \langle 3M-2 \rangle \langle 3M-1 \rangle \right) \langle 3 \rangle. \tag{2}$$

Here we use the notation $\prod_{M=m}^{0} a_M = a_M a_{M-1} \dots a_0$. To prove (2) for $m \ge 4$ by induction we expand:

$$\langle 3m-1\rangle = \langle 3m+2\rangle\langle 3m+3\rangle\langle 3m+4\rangle\langle 3m+2\rangle\langle 3m+3\rangle\langle 3m+1\rangle\langle 3m+2\rangle$$

This is valid as long as $3m+1 \le n-2$ by (1), i.e., as long as $3(m+1) \le n$. \square

Theorem 21. For any $m \ge 4$, $\prod_{k=6}^{3m+1} \langle k \rangle$ is a prefix of $\langle 2 \rangle$.

Proof. We have

$$\langle 2 \rangle = \langle 3 \rangle \langle 4 \rangle \langle 5 \rangle = \langle 4 \rangle \langle 5 \rangle \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle = \langle 6 \rangle \langle 7 \rangle \langle 8 \rangle \langle 6 \rangle \langle 7 \rangle \langle 5 \rangle \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle$$

By substitution into the proof it will suffice to show that $\langle 4 \rangle \langle 5 \rangle \dots$ is a prefix of $\langle 0 \rangle$.

To show that, we first show that $\langle 3 \rangle \langle 4 \rangle \dots$ is a prefix of $\langle 0 \rangle$:

$$\begin{array}{rcl} \langle 0 \rangle & = & \langle 1 \rangle \langle 2 \rangle \langle 3 \rangle = \langle 2 \rangle \langle 3 \rangle \langle 4 \rangle \langle 2 \rangle \langle 3 \rangle \\ & = & \langle 3 \rangle \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle \langle 4 \rangle \langle 2 \rangle \langle 3 \rangle \end{array}$$

Now by substitution, $\langle 6 \rangle \langle 7 \rangle \langle 8 \rangle \dots$ is a prefix of $\langle 3 \rangle$ and we are done. (Incidentally this can now be used to show that $\langle 2 \rangle \langle 3 \rangle \dots$ is a prefix of $\langle 0 \rangle$.) Finally, let us show that $\langle 4 \rangle \langle 5 \rangle \dots$ is a prefix of $\langle 0 \rangle$:

$$\begin{array}{lll} \langle 0 \rangle & = & \langle 1 \rangle \langle 2 \rangle \langle 3 \rangle = \langle 2 \rangle \langle 3 \rangle \langle 4 \rangle \langle 2 \rangle \langle 3 \rangle = \langle 3 \rangle \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle \langle 4 \rangle \langle 2 \rangle \langle 3 \rangle \\ & = & \langle 4 \rangle \langle 5 \rangle \langle 6 \rangle \langle 4 \rangle \langle 5 \rangle \langle 3 \rangle \langle 4 \rangle \langle 2 \rangle \langle 3 \rangle \end{array}$$

 $^{^1\}mathrm{It}$ does not hold for $n=1\colon$ we have $\tilde{T}_2=(\mathtt{01})(\mathtt{0})(\mathtt{2})=\tilde{T}_1\tilde{T}_0\tilde{T}_{-1},$ but $\tilde{T}_1=\mathtt{01}\neq(\mathtt{0})(\mathtt{2})()=\tilde{T}_0\tilde{T}_{-1}\tilde{T}_{-2}.$

By substitution and Theorem 20, $\langle 7 \rangle \langle 8 \rangle \langle 9 \rangle \dots$ is a prefix of $\langle 4 \rangle$, and we are done.

We can also expand the tail of \tilde{T}_n :

$$\langle 0 \rangle = \langle 1 \rangle \langle 3 \rangle \langle 4 \rangle \langle 5 \rangle \langle 4 \rangle \langle 5 \rangle \langle 6 \rangle$$

which ends in $(\langle 4 \rangle \langle 5 \rangle)^*$ with * a non-integer exponent.

Theorem 22. For $n \geq 6$,

$$A^{-}(\tilde{T}_n) \le ||1|| - \sum_{k=6}^{3\lfloor n/3\rfloor + 1} ||k||.$$

For example, when n = 6 this is 24 - (1 + 1) = 22, which fits with an 8-day long computation we performed.

Proof. Using now a ||2||-cycle followed by a path and then a ||4|| + ||5||-cycle, we can subtract the extra prefix from Theorem 20 and use only

$$||2|| + ||4|| + ||5|| + \left(||0|| - 2||2|| - 2(||4|| + ||5||) - ||6|| - \sum_{k=6}^{3\lfloor n/3\rfloor + 1} ||k||\right)$$

$$= ||1|| - \sum_{k=6}^{3\lfloor n/3\rfloor + 1} ||k||$$

(since ||0|| = ||6|| + 2||5|| + 3||4|| + 2||3|| + ||2||) states. Uniqueness is guaranteed by Theorem 24.

Lemma 23 ([1]). The Tribonacci constant (Definition 10) satisfies

$$\xi = \lim_{n \to \infty} \frac{\|1\|_n}{\|0\|_n} = 1.83929\dots$$

and is the unique real root of $\xi^3 = \xi^2 + \xi + 1$.

In particular $\xi = 1/\theta$ with θ as in Theorem 12.

Theorem 24. For large enough n, the equation

$$x\|2\| + y(\|4\| + \|5\|) = 2(\|2\| + \|4\| + \|5\|) \tag{3}$$

for nonnegative integers x, y, has the unique solution x = y = 2.

Proof. Suppose (x, y) is a solution, not equal to (2, 2). Then we have x = 0, x = 1, y = 0, or y = 1. If x = 0 then

$$\mathbb{Z} \ni y = 2\left(\frac{\|2\| + \|4\| + \|5\|}{\|4\| + \|5\|}\right) = 2\left(\frac{\|6\|}{\|4\| + \|5\|} + 3\right)$$

which is impossible as soon as $\|6\| > 0$ since $2\|6\| < \|4\| + \|5\|$. Similarly, if x = 1 then

$$\mathbb{Z} \ni y = \frac{2(\|2\| + \|4\| + \|5\|) - \|2\|}{\|4\| + \|5\|}$$
$$= \frac{\|2\|}{\|4\| + \|5\|} + 2 = \frac{\|6\|}{\|4\| + \|5\|} + 4.$$

If y = 0 then

$$\mathbb{Z}\ni x=2\frac{\|2\|+\|4\|+\|5\|}{\|2\|}=2\left(1+\frac{\|2\|-\|3\|}{\|2\|}\right)=2\left(2-\frac{\|3\|}{\|2\|}\right)$$

which is impossible as soon as ||3|| > 0 since

$$2 > \frac{2\|3\|}{\|2\|} = \frac{\|3\| + \|4\| + \|5\| + \|6\|}{\|3\| + \|4\| + \|5\|} > 1.$$

Finally, if y = 1 then

$$\mathbb{Z}\ni x=\frac{2(\|2\|+\|4\|+\|5\|)-\|4\|-\|5\|}{\|2\|}=2+\frac{\|4\|+\|5\|}{\|2\|}$$

which is impossible as soon as 0 < ||4|| + ||5|| < ||2||, i.e., ||4|| > 0.

Theorem 25. The A⁻-complexity rate of the Tribonacci word satisfies

$$\limsup_{n \to \infty} \frac{A^{-}(T_n)}{|T_n|} \le 1/6(-8 + (586 - 102\sqrt{33})^{1/3} + (2(293 + 51\sqrt{33}))^{1/3})$$

$$\approx 0.4870856...$$

Proof. We calculate

$$\limsup_{n \to \infty} \frac{A^{-}(\tilde{T}_n)}{|\tilde{T}_n|} \le \lim_{n \to \infty} \frac{\|1\| - \sum_{k=6}^{3\lfloor n/3 \rfloor + 1} \|k\|}{\|0\|} = \frac{1}{\xi} - \sum_{k=6}^{\infty} \frac{1}{\xi^k}$$

$$= \frac{1}{\xi} - \frac{1}{3\xi^2 + 3\xi + 2}$$

$$= \frac{1}{6} (-8 + (586 - 102\sqrt{33})^{1/3} + (2(293 + 51\sqrt{33}))^{1/3}). \quad \Box$$

Definition 26. $A_N^{lower}(x)$ is the minimal q such that for all sequences of strongly disjoint powers

$$x_1^{\alpha_1}, \ldots, x_m^{\alpha_m},$$

in x, with the uniqueness condition that

$$\sum \alpha_i |x_i| = \sum \alpha_i y_i \implies y_i = |x_i|, \text{all } i,$$

we have

$$2q \ge n + 1 - m - \sum_{i=1}^{m} (\alpha_i - 2)|x_i|. \tag{4}$$




Figure 1: An automaton witnessing the automatic complexity of the Fibonacci word of length 55.

The definition of $A_N^{lower}(x)$ may seem very technical. The point is that

- ullet A_N^{lower} appears to be faster to compute than $A_N,$
- by [8, Theorem 19], we have $A_N^{lower}(x) \leq A_N(x)$ for all words x, and
- $A_N^{lower}(x)$ is a better lower bound than that obtained simply by the critical exponent considerations in Theorem 15.

We have implemented A_N and A_N^{lower} ([5]) with results in Table 1 and Table 2. (A lookup tool is also available for automatic complexity [6].) Note that

$$A^{-}(T_{n}) = A^{-}(\tilde{T}_{n-3}) \leq \|1\| - \sum_{k=6}^{3\lfloor (n-3)/3\rfloor + 1} \|k\|$$

$$= \|1\| - \sum_{k=6}^{3\lfloor n/3\rfloor - 2} \|k\|$$

$$= \|1\| - \sum_{k=6}^{7} \|k\| = t_{8} - t_{3} - t_{2} = 24 - 1 - 1 = 22, \quad (n = 9),$$

$$= \|1\| - \sum_{k=6}^{7} \|k\| = t_{9} - t_{4} - t_{3} = 44 - 2 - 1 = 41, \quad (n = 10).$$

Remark 27. Witnessing automata for A_N are conveniently generated by state sequences. A state sequence is the sequence of states visited by the unique accepting path of length n+1 (having potentially up to n edges and n+1 states). A week-long computer search for the length 55 Fibonacci word

revealed the witnessing state sequence, where states are given numerical labels, using letters A, B, C, \ldots for the numbers $10, 11, 12, \ldots$:

We illustrate the automaton induced by this state sequence in Figure 1. Generalizing this example gives

$$\overline{A_N}(\mathbf{f}^{(2)}) \le \frac{1}{\varphi^2} + \frac{1}{\varphi^7} = 0.41640786499$$

Lemma 28. For $n \geq 6$, the equation

$$xf_{n-2} + yf_n = 2(f_{n-2} + f_n)$$

for nonnegative integers x, y, has the unique solution x = y = 2.

Proof. If x = 0 then

$$y = 2\frac{f_{n-2} + f_n}{f_n} = 2 + \frac{2f_{n-2}}{f_n} = 2 + \frac{2f_{n-2}}{f_{n-1} + f_{n-2}} \in (2,3)$$

is not an integer as long as $n \ge 4$. If x = 1 then

$$y = \frac{2(f_{n-2} + f_n) - f_{n-2}}{f_n} = 2 + \frac{f_{n-2}}{f_n} \in (2,3)$$

is not an integer as long as $n \geq 3$. If y = 0 then

$$x = 2\frac{f_{n-2} + f_n}{f_{n-2}} = 2\frac{3f_{n-2} + f_{n-3}}{f_{n-2}} = 6 + \frac{f_{n-3} + f_{n-4} + f_{n-5}}{f_{n-3} + f_{n-4}} \in (7,8)$$

is not an integer as long as $f_{n-5} > 0$, i.e., n > 6. If y = 1 then

$$x = \frac{2(f_{n-2} + f_n) - f_n}{f_{n-2}} = 2 + \frac{f_n}{f_{n-2}} = 2 + \frac{f_{n-2} + f_{n-2} + f_{n-3}}{f_{n-2}} \in (4, 5)$$

as long as $f_{n-3} > 0$.

Theorem 29. The upper automatic complexity rate of the infinite Fibonacci word $\overline{A_N}(\mathbf{f}^{(2)})$ is at most $\frac{2}{\varphi^3}$.

Proof. We exploit the lengths of Fibonacci words.

$$f_{n} = f_{n-1} + f_{n-2} = f_{n-2} + f_{n-3} + f_{n-2}$$

$$= f_{n-3} + f_{n-4} + (f_{n-3} + f_{n-3} + f_{n-4})$$

$$= f_{n-4} + f_{n-5} + f_{n-4} + (f_{n-3} + f_{n-3} + f_{n-4})$$

$$= \underbrace{f_{n-4}}_{\text{hardcode this}} + \underbrace{(f_{n-5} + f_{n-5} + f_{n-6})}_{f_{n-5}\text{-state cycle}} + \underbrace{(f_{n-3} + f_{n-3} + f_{n-4})}_{f_{n-3}\text{-state cycle}}.$$

n	t_n	T_n	$.313t_n$	${\rm A_N^{lower}}$	A^-	(5)	$.487t_n$
0	0		0	1	1		0
1	0		0	1	1		0
2	1	2	.3	1	1		0.49
3	1	0	.3	1	1		0.49
4	2	01	.6	2	2		0.97
5	4	0102	1.3	3	3		1.95
6	7	0102010	2.2	4	4		3.4
7	13	0102010010201	4.1	7	7		6.3
8	24	01020100102	7.5	12	13		11.7
9	44	01020102010	13.8	21		22	21.4
10	81	01020010201	25.4	36		41	39.4

Table 1: Lower and limiting upper bounds on $A_N(T_n)$ and $A^-(T_n)$.

This way we obtain for a Fibonacci word x of length f_n that

$$A_N(x) \le f_{n-4} + f_{n-5} + f_{n-3} = 2f_{n-3}.$$

In the limit, $f_n/f_{n-1} \sim \varphi = \frac{1+\sqrt{5}}{2} \approx 1.6$, so $f_n/f_{n-3} \sim \varphi^3 = 4.236$, so

$$A_{\mathcal{N}}(x) \le \frac{2}{\varphi^3} f_n = 0.47 f_n.$$

The cycles give a unique path of length f_n for large enough n, since

$$f_{n-5}x + f_{n-3}y = 2(f_{n-5} + f_{n-3})$$

has a unique solution x = y = 2 by Lemma 28.

4 Conclusion

More can be done on automatic complexity rates of k-bonacci words. For instance, we conjecture that $\overline{A_N}(\mathbf{f}^{(2)}) \leq 1/\varphi^2 = .382$. More precisely, we conjecture that this can be shown by analyzing the decomposition

$$\begin{aligned} \langle 0 \rangle &= & \langle 1 \rangle \langle 2 \rangle = \langle 2 \rangle \langle 3 \rangle \langle 2 \rangle = \langle 3 \rangle \langle 4 \rangle \langle 3 \rangle \langle 3 \rangle \langle 4 \rangle = \langle 4 \rangle \langle 5 \rangle \langle 4 \rangle \langle 4 \rangle \langle 5 \rangle \langle 4 \rangle \langle 5 \rangle \langle 4 \rangle \langle 6 \rangle \langle$$

However, in the present article we are content to have proven that the Fibonacci word has intermediate automatic complexity rate in Theorem 29.

n	f_n	F_n	$.276f_n$	$A_{\rm N}^{\rm lower}$	$.382f_n$
0	0		0	1	0
1	1	1	.3	1	0.4
2	1	0	.3	1	0.4
3	2	01	.6	2	0.8
4	3	010	.8	2	1.1
5	5	01001	1.4	3	1.9
6	8	01001010	2.2	4	3.1
7	13	0100101001001	3.6	6	5.0
8	21	010010100100101001010	5.8	9	8.0
9	34	0100101001001010010101001001	9.4	14	13.0
10	55	01001010	15.2	21	21.0

Table 2: Lower and limiting upper bounds on $A_N(F_n)$.

References

- [1] Steven R. Finch. *Mathematical constants*, volume 94 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2003.
- [2] Amy Glen. Powers in a class of \mathscr{A} -strict standard episturmian words. Theoret. Comput. Sci., 380(3):330–354, 2007.
- [3] Kayleigh K. Hyde and Bjørn Kjos-Hanssen. Nondeterministic automatic complexity of overlap-free and almost square-free words. *Electron. J. Com*bin., 22(3), 2015. Paper 3.22, 18.
- [4] Juhani Karhumäki. On cube-free ω -words generated by binary morphisms. Discrete Appl. Math., 5(3):279–297, 1983.
- [5] Sun Young Kim, Bjørn Kjos-Hanssen, and Clyde James Felix. Code for "Automatic complexity of Fibonacci and Tribonacci words". https://github.com/bjoernkjoshanssen/tetranacci, 2019.
- [6] Bjørn Kjos-Hanssen. Complexity lookup. http://math.hawaii.edu/wordpress/bjoern/complexity-of-0110100110110/.
- [7] Bjørn Kjos-Hanssen. Automatic complexity of shift register sequences. *Discrete Math.*, 341(9):2409–2417, 2018.
- [8] Bjørn Kjos-Hanssen. An incompressibility theorem for automatic complexity, 2019. arXiv preprint 1908.10843v1.
- [9] F. Mignosi and G. Pirillo. Repetitions in the Fibonacci infinite word. RAIRO Inform. Théor. Appl., 26(3):199–204, 1992.

- [10] Jeffrey Shallit and Ming-Wei Wang. Automatic complexity of strings. *J. Autom. Lang. Comb.*, 6(4):537–554, 2001. 2nd Workshop on Descriptional Complexity of Automata, Grammars and Related Structures (London, ON, 2000).
- [11] Michael Sipser. A complexity theoretic approach to randomness. In *Proceedings of the Fifteenth Annual ACM Symposium on Theory of Computing*, STOC '83, pages 330–335, New York, NY, USA, 1983. ACM.
- [12] Bo Tan and Zhi-Ying Wen. Some properties of the Tribonacci sequence. *European J. Combin.*, 28(6):1703–1719, 2007.